Predicting clinical endocrine response in advanced breast cancers using a reproducible

low-dimensional biomarker

Aritro Nath, Ph.D.[1], Adam L. Cohen, M.D.[2], Jeffrey T. Chang, Ph.D.[3] and Andrea H. Bild,

Ph.D.[1*]

[1] Division of Molecular Pharmacology, Department of Medical Oncology & Clinical

Therapeutics, City of Hope, Monrovia, California

[2] Division of Oncology, Department of Medicine, Huntsman Cancer Institute, University of

Utah, Salt Lake City, Utah

[3] Department of Integrative Biology & Pharmacology, University of Texas Health Science

Center at Houston, Houston, Texas

*Correspondence: Andrea H. Bild, Ph.D.

Professor, Department of Medical Oncology and Therapeutics

City of Hope Comprehensive Cancer Institute

1218 S Fifth Ave, Monrovia, CA 91016

(626) 218-6052

abild@coh.org

Conflict of interest statement: The authors declare no potential conflicts of interest.

Abstract

In the absence of reliable and effective prognostic biomarkers, endocrine therapy remains the

standard of care for all advanced and metastatic estrogen receptor-positive (ER+) breast cancers.

Attempts to develop biomarkers using the baseline tumor transcriptome or genome of advanced

ER+ breast cancers have so far been unsuccessful due to predictive models with poor

reproducibility in independent studies. Here we present an approach to develop a low-

dimensional biomarker that estimates the risk of adverse events on endocrine therapy using the

baseline tumor transcriptome of patients. Using a framework for supervised dimensionality

reduction of the gene expression feature space, we constructed an endocrine response signature

(ENDORSE) modeled on the survival outcomes of ER+ breast cancers from METABRIC.

ENDORSE outperformed transcriptome-wide and knowledge-based signature models while

significantly improving upon routine histopathological and genomic classifiers in cross-

validation analyses. The ENDORSE risk estimate accurately predicted the outcomes for

endocrine therapy in three independent clinical trials for ER+ breast cancers. Further, analysis of

the phenotypes enriched in high-risk categories show endocrine resistance was not associated

with rates of proliferation, but instead with a potential loss of DNA damage repair and cell-

matrix interaction pathways.

Introduction

Estrogen receptor (ER)-positive tumors are the most common form of breast cancer [1]. Endocrine therapy remains the backbone of initial therapy for metastatic ER+ breast cancers, which are now defined as tumors with 1-100% of cell nuclei stained positive for the ER [2,3]. However, ER+ tumors are heterogeneous, both in terms of dependence on estrogen signaling for growth and survival and intrinsic or acquired resistance to endocrine therapy [4,5]. Therefore, optimal clinical management of each ER+ breast cancer depends on accurate prediction of response to endocrine therapy and selection of companions for endocrine therapy. Several genomic tests are available for classifying breast cancers into molecular subtypes [6] or assessing the likelihood of benefit from chemotherapy in early-stage, node-negative ER+ breast cancers [7,8]. However, these assays are not used in advanced or metastatic breast cancer. Therefore, the goal of our study is to develop a tumor expression-based prognostic biomarker for ER+ breast cancers on endocrine therapy, independent of the tumor stage or lymph node status.

Complex models, where the number of candidate features (genes) are much greater than the number of available samples ($p >> n$), can be difficult to interpret, are easy to overfit, and often overperform in training datasets with poor predictive ability in independent validations [9,10]. Consequently, few biomarkers developed using the baseline tumor transcriptome are translated into clinical practice [11]. To address these challenges, we developed a framework to model endocrine therapeutic response using a low-dimensional biomarker. We first systematically reduced the dimensionality of the feature space to derive a gene signature and model the survival outcomes of patients on endocrine therapy using a univariate gene set enrichment (GES) score calculated for each sample. The resulting endocrine response signature (ENDORSE) biomarker

classified ER+ breast cancers into three risk categories. We demonstrate the robustness and

reproducibility of ENDORSE through cross-validation analyses in the METABRIC dataset, in

comparison with existing molecular classifiers PAM50 [6] and IntClust [12], knowledge-based

molecular signatures, and transcriptome-wide predictors. Further, we validated the performance

of ENDORSE in three independent clinical datasets with advanced or metastatic ER+ breast

cancers. As negative controls, we also tested ENDORSE in ER- METABRIC breast cancers and

in an independent clinical trial with HER2+ breast cancers to demonstrate the specificity of the

biomarker. Comparison of biological phenotypes showed that the biomarkers was not merely a

proxy for proliferation while also outperforming proliferation-based signatures. Overall, our

analyses demonstrate that the low dimensional ENDORSE biomarker is a reliable prognostic

biomarker for advanced and metastatic ER+ breast cancers.

RESULTS

Developing a low-dimensional endocrine response signature

To develop the endocrine response biomarker, we used the baseline transcriptome from the

METABRIC[13,14] ER+ breast cancer cohort as the training dataset. The selection criteria for

inclusion in the training data were hormone receptor status (ER+/HER2-), no treatment with

chemotherapy, and cancer as the cause of death (Supplementary Table 1).

Supplementary Table 1: ER+ METABRIC patient characteristics			
Variable	Mean	95% C.I.	N available
Time to event (in months)	135	130 - 140	833
Events	0.409	0.376 - 0.443	833
Age at diagnosis	61.5	60.7 - 62.3	833
Tumor size	24.3	23.4 - 25.1	828
Tumor stage *Stage 0-1 (n=270)* *Stage 2 (n=324)* *Stage >3 (n=40)*	1.64	1.59 - 1.69	634
Mutation count	5.55	5.31 - 5.79	809
Number of positive lymph nodes detected *0 (n=491)* *1-3 (n=228)* *4-9 (n=85)* *>10 (n=29)*	1.56	1.32 - 1.79	833

We used a LASSO-regularized Cox proportional hazards model to perform a cross-validation

analysis with the survival outcomes of 833 cancers using baseline tumor expression profiles as

the predictors. Genes with a positive coefficient in >50% of the cross-validations were retained

to form an endocrine response gene set. Subsequently, we calculated a single sample gene set

enrichment score (GES)[15,16] for each sample using this gene set, and estimated the risk of death

on endocrine therapy using a Cox proportional hazards model. The resulting ENDORSE risk

estimates classified patients into three risk groups that predicted risk of adverse outcome (death)

on endocrine therapy. Additionally, we modeled and compared the survival outcomes using the

complete transcriptome and knowledge-based curated[17] and hallmark molecular signatures[18].

Stratifying METABRIC breast cancers based on ENDORSE scores

To demonstrate the practical utility of ENDORSE, we stratified the METABRIC ER+ breast

cancers based on estimated risk of death on endocrine therapy. Cancers with an ENDORSE risk

estimate or hazard ratio of ≥2 were classified into the high-risk category, cancers with an

estimate ≤1 in to low-risk, and all cancers with intermediate hazard ratios in to the medium-risk

category (Figure 1a). Analyses of the survival curves based on this classification shows

significant differences in the survival rates ($P=3.55\times10^{-24}$) of cancers stratified by the ENDORSE

risk estimates. In the METABRIC cohort, the majority of cancers were classified in the low or

medium risk class, while only 16% of the patients were classified in high-risk class (Figure 1b).

Next, to demonstrate the robustness of the ENDORSE risk estimates, we reanalyzed the Kaplan-

Meier survival curves of METABRIC patients in scenarios where either genes or samples were

missing from the training dataset (Figure 1c, d). We observed that even after reducing the

number of genes down to 10%, the log-rank test P-values were within the same order of

magnitude as the original stratification. Upon reduction of the sample size, the stratification

yielded significant P-values even when sub-sampled down to 10% of the original dataset.

We further compared the performance of the ENDORSE risk estimates with clinical covariates,

hormone receptor gene expression and other established breast cancer stratification models

including PAM50 intrinsic subtypes[6] and IntClust classes[12]. Univariate Cox models based on the

ENDORSE scores and risk estimates outperform all other variables evaluated. Additionally,

multivariate Cox-analysis with the ENDORSE signature along with hormone receptor

expression, PAM50 subtypes and IntClust classes demonstrates that ENDORSE captures unique

and significant proportion of the model variance over these predictors (Supplementary Tables 2-

4). Thus, the ENDORSE scores are non-redundant from existing molecular classifications and

augment risk stratification of ER+ breast cancers.

Supplementary Table 2: Multivariate ENDORSE Cox Models with PAM50 subtypes			
Variable	HR	95% C.I.	P-value
PAM50 claudin-low	1.13	0.515 - 2.46	0.77
PAM50 Her2	1.48	0.712 - 3.08	0.29
PAM50 LumA	0.864	0.444 - 1.68	0.67
PAM50 LumB	0.984	0.518 - 1.87	0.96
PAM50 NC	1.68	0.456 - 6.17	0.44
PAM50 Normal	1.38	0.652 - 2.92	0.4
ENDORSE Risk Groups: Medium Risk	2.37	1.79 - 3.13	1.4E-09
ENDORSE Risk Groups: High Risk	3.93	2.77 - 5.58	1.8E-14

Supplementary Table 3: Multivariate ENDORSE Cox Models with IntClust classes			
Variable	HR	95% C.I.	P-value
IntClust 10	0.421	0.175 - 1.02	0.054
IntClust 2	1.33	0.782 - 2.27	0.29
IntClust 3	0.875	0.546 - 1.4	0.58
IntClust 4 ER-	NA		
IntClust 4 ER+	1.38	0.872 - 2.17	0.17
IntClust 5	0.982	0.297 - 3.25	0.98
IntClust 6	1.61	0.98 - 2.64	0.06
IntClust 7	1.09	0.679 - 1.74	0.73
IntClust 8	1.03	0.668 - 1.58	0.9
IntClust 9	1.41	0.875 - 2.28	0.16
ENDORSE Risk Groups: Medium Risk	2.4	1.84 - 3.14	1.4E-10
ENDORSE Risk Groups: High Risk	4.19	3.05 - 5.77	1.1E-18

Supplementary Table 4: Multivariate ENDORSE Cox Models with HR gene expression			
Variable	HR	95% C.I.	P-value
ESR1 expression	0.953	0.874 - 1.04	0.28
PGR expression	0.854	0.757 - 0.964	0.01
ENDORSE Risk Groups: Medium Risk	2.39	1.83 - 3.12	1.9E-10
ENDORSE Risk Groups: High Risk	3.94	2.92 - 5.32	4.2E-19

Figure 1. Stratification of METABRIC ER+ breast cancers based on ENDORSE. A.
Kaplan-Meier survival curves and accompanying risk table of METABRIC ER+ breast cancers
stratified into low, medium and high-risk groups based on hazard ratios estimated using
ENDORSE scores. **B.** Histogram and cumulative density function plots showing frequency
distribution of samples based on estimated ENDORSE risk. **C and D.** A series of Kaplan-Meier
survival curves showing stratification of METABRIC ER+ breast cancers based on reduced
number of available genes (C) or samples (D) for calculating ENDORSE scores.

Validating ENDORSE performance in independent clinical datasets

To demonstrate the reliability and reproducibility of the ENDORSE risk estimate guided

stratification, we applied the Cox model coefficients obtained from the METABRIC models to

ER+ breast cancers in three independent clinical trial datasets. The first independent clinical trial

reported endocrine therapy treatment outcomes of 140 stage IV ER+ metastatic breast cancers,

78% of which had received prior endocrine therapy[19]. We calculated the ENDORSE gene set

enrichment scores from the transcriptomic data obtained from the metastases and stratified the

cancers based on METABRIC-derived ENDORSE coefficients (Figure 2a). Analysis of the

survival curves indicates accurate stratification of the cancers based on the risk estimates (P =

3×10^{-4}).

Next, we examined data from the NCT00265759 clinical trial which evaluated neoadjuvant

aromatase inhibitor (AI) treatment in Stage II or III ER+ breast cancers [20]. This study used 10%

Ki67 staining at the end of treatment (2-4 weeks) as the threshold for sensitive or resistant

cancers. We found that the ENDORSE risk estimates successfully stratified the cancers in

agreement with reported Ki67 % at both baseline (P=2.7×10^{-8}) and end of treatment (P=5.6×10^{-3})

(Figure 2b). The third independent clinical trial evaluated fulvestrant response in advanced

metastatic ER+ breast cancers previously treated with an antiestrogen (CONFIRM study)[21]. This

study classified tumors as resistant or sensitive based on a 10% Ki67 staining threshold. Again,

the stratification of these cancers based on ENDORSE risk estimates showed clear differences in

Ki67% across the risk groups (P = 1.6×10^{-9}), in addition to significant correlation with the

continuous risk score (P = 2.5×10^{-11}) (Figure 2c). Similar consistent patterns were observed with

the continuous risk scores. In both the CONFIRM and NCT00265759 trial, patients stratified in

low risk groups were consistently classified as sensitive based on the trial outcome data, while

high risk tumors were classified as resistant (Figure 2b, c).

In addition to the endocrine therapy trials in ER+ breast cancer, we also applied the ENDORSE

risk estimates to stratify ER- breast cancers or ER+ breast cancers on non-endocrine therapy as

negative controls. First, we compared ENDORSE risk estimates in a trial for neoadjuvant

trastuzumab and lapatinib in ER+/HER2+ or ER-/HER2+ breast cancers. In both molecular

subtypes, there was no significant difference in ENDORSE risk estimates between the partial

clinical response and residual disease groups (ER+ P = 0.465, ER- P = 0.824) (Figure 2d).

Similarly, stratification of the ER- cancers based on the estimated ENDORSE risk resulted in no

significant differences between the survival curves (P = 0.4) (Figure 2e). These results suggest

the ENDORSE risk estimate is not a general biomarker for aggressive breast cancers, but

specific to ER+ breast cancers on endocrine therapy.

Figure 2. Validation in independent clinical datasets. A. Kaplan-Meier survival curves of

ER+ breast cancer metastases stratified based on ENDORSE risk estimates, along with survival

risk table. **B and C.** Right panels show violin plots for ER+ breast cancers stratified based on

ENDORSE risk estimates (X-axis) and Ki67 % (Y-axis). The dotted line indicates a Ki67

staining level of 10%, a threshold used in both studies to classify cancers as sensitive or resistant

to therapy. The scatter plots in the left panel show correlation between continuous ENDORSE

risk estimates (X-axis) and Ki67% staining (Y-axis). Liner fit along with 95% confidence

intervals are also shown. **D.** Violin plots comparing ENDORSE risk scores in patients stratified

based on trial-reported clinical response. The left panel represents ER+/HER2+ cancers while the

right panel represents ER-/HER2+ cancers. **E.** Kaplan-Meier curves of ER- METABRIC breast

cancers stratified based on ENDORSE risk estimates.

Understanding the biology of high-risk cancers

To gain insights into the biology of high-risk cancers, we compared the gene set enrichment

scores, gene-level mutation and copy number alteration frequencies across the METABRIC

ENDORSE risk strata. We found that multiple pathways linked with functional loss of P53 and

RB were elevated, and among the most significant pathways in the high-risk cancers (Figure 3a,

Supplementary Figure 2). Concurrently, DNA-damage repair and cell cycle pathways that are

closely associated with p53 and Rb loss were also enriched (Figure 3a). Among gene signatures

that were enriched in the low/medium risk strata, we predominantly found pathways and

complexes associated with extracellular matrix interaction, including integrin, laminin,

hemidesmosome and basement membranes (Figure 3b, Supplementary Figure 1). To

demonstrate that the ENDORSE risk strata were not solely driven by a proliferative signal[22], we

calculated the meta-PCNA proliferation index and analyzed the multivariate Cox models

containing the proliferation index and the ENDORSE scores or risk groups as co-variates

(Supplementary Tables 5-6). This analysis reinforces the notion that the ENDORSE scores

capture information beyond mere proliferation.

Supplementary Table 5: Multivariate ENDORSE Cox Models with meta-PCNA			
Variable	**HR**	**95% C.I.**	**P-value**
meta-PCNA (Proliferation Index)	1.04	0.75 - 1.45	0.8
ENDORSE Risk Groups: Medium Risk	2.4	1.8 - 3.19	2.3E-09
ENDORSE Risk Groups: High Risk	3.99	2.78 - 5.74	7.2E-14

Supplementary Table 6: Multivariate ENDORSE Cox Models with meta-PCNA			
Variable	**HR**	**95% C.I.**	**P-value**
meta-PCNA (Proliferation Index)	0.978	0.703 - 1.36	0.89
ENDORSE Risk	1.88	1.64 - 2.16	5.7E-19

We next analyzed the somatic mutation and copy number profiles of the METABRIC cancers.

None of the copy number gain or losses were enriched in the high-risk tumors compared to

low/medium risk tumors. However, the somatic mutation frequencies for a few cancer-associated

genes were significant at the nominal threshold (Figure 3c, Supplementary Table 7). However,

only two genes, *TP53* (P = 6.6 x 10^{-5}) and *AKAP* (P = 3.3 x 10^{-4}), were significant at an FDR <

0.05 threshold. The functional impact of p53 mutation frequency in high-risk tumors was

corroborated by the enrichment of p53 loss of function-linked gene signatures (Figure 3a,

Supplementary Figure 2). Tumor sequencing efforts have shown high frequency of *AKAP*

mutations in metastatic lesions compared to primary tumors [23], corroborating the observed

enrichment in high-risk tumors.

Supplementary Table 7: Somatic mutation frequency					
Gene	Low Risk	Medium Risk	High Risk	Chi-square P-value	FDR
TP53	48	71	44	6.59E-05	0.001847
AKAP9	17	16	18	3.35E-04	0.009033
GATA3	74	59	11	2.91E-03	0.075576
CDH1	37	30	25	4.58E-03	0.114529
PIK3CA	255	152	65	2.87E-02	0.689714
AHNAK	35	32	15	7.26E-01	1
AHNAK2	90	66	26	4.66E-01	1
AKT1	13	22	8	8.39E-01	1
ARID1A	15	27	15	5.38E-02	1
BIRC6	12	26	10	5.03E-01	1
CBFB	44	15	10	7.95E-01	1
COL22A1	18	19	5	5.61E-01	1
DNAH11	25	38	14	7.64E-01	1
DNAH2	30	27	11	1.00E+00	1
DNAH5	30	20	10	1.00E+00	1
HERC2	19	29	12	5.37E-01	1
KMT2C	50	44	24	2.55E-01	1
LAMA2	19	19	8	1.00E+00	1
MAP3K1	80	44	18	2.43E-01	1
MLL2	30	23	9	8.24E-01	1
MUC16	63	60	31	1.96E-01	1
PDE4DIP	21	23	5	3.19E-01	1
RYR2	19	28	8	8.56E-01	1
SYNE1	50	34	22	2.38E-01	1
TBX3	28	22	6	3.22E-01	1
TG	20	15	11	2.20E-01	1
THADA	20	17	6	8.25E-01	1
USH2A	24	25	14	2.54E-01	1

a Pathways enriched in high risk tumors ER+ METABRIC tumors

b Pathways enriched in low/medium risk ER+ METABRIC tumors

c Mutation frequency in ER+ METABRIC breast cancers stratified by estimated risk

Figure 3: Biology of high-risk tumors. A-C. Violin plots comparing the single sample gene set enrichment scores of various pathways (Y-axis) in METABRIC tumors stratified by estimated

ENDORSE risk. The low and medium risk tumors were combined in one category for

comparison. **A.** Representative signatures for p53 loss, Rb loss, DNA damage repair and cell

cycle. **B.** Tumor-extra cellular matrix interaction pathways. **C.** Bar plots representing gene-level

mutation frequencies of various cancer-associated genes in low/medium vs. high-risk

METABRIC tumors. The p-values from Chi-square test are shown above the bars, with bold

letters indicating comparison significant at an FDR > 0.05 threshold.

Discussion

Gene expression biomarkers have been successfully integrated in clinical practice for prognosis and guiding treatment decisions in early-stage breast cancers, as evident from the success of the MINDACT and TAILORx studies[8,24]. However, gene expression biomarkers to guide treatment decisions in metastatic ER-positive breast cancer, either in the first-line or later line settings, are lacking. Thus, biomarkers for predicting optimal treatment of metastatic ER-positive breast cancers, particularly after progression on first-line aromatase inhibitor/CDK4/6 inhibitor combinations, could have a large impact on people with ER-positive breast cancer.

The development of new biomarkers based on patient transcriptome data is challenging. Since the number of available genes to train the statistical models tend to be much larger than the number of available samples ($p >> n$), it is quite easy to create complex prediction models that contain a large number of predictor variables. Often, such models perform very well in the training datasets, but the performance cannot be replicated in independent test datasets due to overfitting. A number of approaches have been proposed to address this issue. Broadly, these can be classified into unsupervised and supervised approaches. The unsupervised approach typically relies on grouping or clustering the samples into based on similarity of gene expression profiles, followed by analysis of association with survival outcomes [25]. Alternatively, a supervised approach is to perform dimensionality reduction prior to modelling the survival outcome or drug response using univariate or multivariate models [26].

In this study, we developed a biomarker for the prognosis of ER+ breast cancers using METABRIC data. We used a LASSO regularized Cox model for feature selection, effectively

reducing the dimensionality of the gene expression data. In addition, we adapted a pathway

signature approach in our framework, which further reduced the number of predictors down to a

single integrated variable as the final biomarker. We evaluated the reliability and robustness of

our biomarker through cross-validation analyses and simulations in the METABRIC ER+ cohort.

Our analyses revealed that the univariate ENDORSE biomarker consistently outperformed

multivariate models based on gene expression or literature-derived gene signatures (Figure 2).

Furthermore, we found that reducing the number of available genes to calculate the ENDORSE

risk estimates had a minimal impact on the ability of the biomarker to stratify METABRIC ER+

cancers. This distinct advantage of using the gene set enrichment scores (GES) over individual

genes or pathway predictors can be explained by the algorithm for GES calculation. The ssGSEA

method proposed by Barbie et al. replaces gene expression by their ranks, followed by

calculating the differences in the empirical distribution functions of the signature genes vs. all

other genes [16]. The rank-based method helps mitigate issues encountered due to batch effects and

differences in methods for transcriptome profiling, while the redundancy in the signature ensures

the scores remain consistent even in the case some genes are missing in the data. Consequently,

the ENDORSE biomarker was successful in stratifying ER+ breast cancers in multiple

independent validation datasets from diverse gene expression profiling platforms.

The ENDORSE biomarker stratified cancers based on estimated risk of death due to the disease

while on endocrine therapy. In addition to testing its potential to serve as a robust biomarker, we

explored the biology of the high-risk tumors for possible hints into their mechanism. We found

that high-risk tumors showed a consistent enrichment of pathways associated with loss of p53

and Rb, along with DNA damage repair and cell cycle progression. Mutations in the *TP53* have

long been associated with aggressiveness and chemotherapeutic resistance in hormone-receptor

negative breast cancers [27,28]. However, recent studies show that even though *TP53* are infrequent

in ER+ breast cancers, they have similar negative impact on patient outcome as hormone-

receptor negative breast cancers [29]. Similarly, inactivation of the tumor suppressor Rb has been

associated with therapeutic resistance in ER+ breast cancers [30,31]. A previous meta-analysis of

gene signatures by Venet et al. associated with breast cancer outcomes showed that most

signatures were redundant with proliferation [22]. While proliferation is indeed an important

feature of aggressive tumors, our analyses showed that proliferation signature did not encompass

the entirety of the signal from the ENDORSE biomarker.

To determine the clinical utility and clinical validity of a biomarker, it must be tested in a clinical

trial. According to the National Comprehensive Cancer Network (NCCN), options for metastatic

ER-positive, HER2-negative breast cancer after progression on aromatase inhibitor plus CDK4/6

inhibitor include fulvestrant (which requires the breast cancer to still be estrogen dependent) with

or withour alpelisib, exemestane (which also requires the breast cancer to still be estrogen

dependent) plus everolimus, or chemotherapy [32]. We will be testing the ENDORSE biomarker in

a clinical trial to assign patients to the regimens targeting estrogen vs chemotherapy.

Funding

Funding for this research was provided by the National Cancer Institute of the National Institutes

of Health through the U54 grant 1U54CA209978.

METHODS

Data retrieval, pre-processing and analysis

METABRIC gene expression, phenotypic and survival data were retrieved using cBioPortal for

cancer genomics [33]. Independent validation datasets used in this study were retrieved from the

NCBI GEO portal under the following accession IDs: GSE124647[19], GSE87411[20], GSE76040[21]

and GSE130788. For each gene expression dataset, we removed genes with zero variance and

summarized genes with multiple probes by mean expression. Each dataset was scaled, such that

the mean of each gene across the samples was zero with standard deviation equal to 1.

The analyses were performed in R 3.6.1, RStudio 1.2.1335.

Selecting samples for training models

The METABRIC cohort contained a total of 2509 samples. We constructed the training models

for ER+ cancers and endocrine therapy response by filtering the samples on the following

criteria: 1. The tumors were positive for estrogen receptor and negative for human epidermal

growth factor receptor 2 (ER+ and HER2-) in their immunohistochemistry profile, 2. If the

patient died, then disease was listed as cause of death, and 3. The patient did not receive

chemotherapy alone or in combination with endocrine therapy. After filtering based on these

criteria, we retained 833 samples with both gene expression and complete clinical data available.

Training features and model construction

We derived and compared four sets of features as potential predictors of long-term outcome of

ER+ METABRIC tumors, including curated gene set and hallmark gene set enrichment scores,

expression profiles of all genes and the ENDORSE gene signature (Figure 1). The curated gene

sets and hallmark gene sets (v7.0) were retrieved from MSigDb[17,18]. For each gene signature, the

gene set enrichment scores were calculated using the GSVA package for R[15] using the ssGSEA

method[16]. Next, the gene set enrichment scores (GES) or expression profiles of all genes, along

with age at diagnosis, were used as input features in LASSO-regularized Cox regression models,

with overall survival as the outcome variable[34]. The hazard function in the Cox model is defined

as:

$$h_i(t) = h_0(t)\exp\left(\beta x_i^T\right)$$

311   Where, X is a set of predictive features and h_0 is an arbitrary baseline hazard function. We

312   considered curated GES, hallmark GES and all genes each as individual sets of predictive

313   features in separate analyses. The coefficient (β) for each predictor in the model can be estimated

314   by maximizing the partial likelihood function $L(\beta)$, defined as:

$$L(\beta) = \prod_i \frac{\exp\left(\beta x_{j(i)}^T\right)}{\sum_{l \in R_i} \exp\left(\beta x_j^T\right)}$$

315   Where R_i is the set of indices of observations failing (events) at time t_i. In the LASSO Cox

316   model, the regularized coefficient is obtained by adding a penalty parameter λ to the log of the

317   likelihood function.

$$\hat{\beta} = \min -\frac{1}{N} l(\beta) + \lambda \, ||\beta||_1$$

Where, $l(\beta) = \log L(\beta)$. The λ penalty parameter was determined using 10-fold cross-validation

implemented in R package glmnet[36,37]. The optimal λ for the curated or hallmark GES and gene

expression feature sets were defined as the λ that minimized model deviance for each feature set.

We defined the preliminary endocrine resistance gene set (ENDORSE) using the features with

positive coefficients in the LASSO model for all genes, with optimal λ within one standard error

from the minimum model deviance. We further expanded the signature by including genes that

were positively correlated (Pearson's correlation > 0.75) with the selected features in the ER+

METABRIC samples. Next, we calculate the GES for each sample using GSVA and use the

GES as a predictive feature.

Repeated cross-validation, consensus ENDORSE signature and performance evaluation

To evaluate the performance of each set of predictive features and derive a consensus

ENDORSE model, we performed 10-fold cross-validation analysis repeated 50 times. In this

analysis, the features selected in the LASSO model from each set of predictors were evaluated in

multivariate Cox model. In each of the 50 repeats, the samples were split into 10 equal parts,

with one part serving as the test set and the remaining parts serving as the training set in each

fold of cross-validation. In each fold, the feature selection was performed for curated, hallmark

or ENDORSE GES and gene expression only using the training set. Then, the Cox model

coefficients were derived for each set of selected features using only the training set. To evaluate

performance, we applied the coefficients from the training model to the GES or gene expression

data from the test data. We reported the concordance indices of the Cox model fit in the training

and test data. In addition, we reported the correlation between the actual risk of event in the test

data and the predict risk of event based on the coefficients derived from the training data and

applied to the test data.

The consensus ENDORSE set was defined as the set of predictive features that were selected in

more 50% of the repeated cross-validations. This resulted in the selection of 63 genes that

comprised the consensus ENDORSE set for subsequent analysis involving risk or hazard ratio

(HR) estimation and predictions in independent datasets. After the consensus ENDORSE set was

defined, we obtained ENDORSE GES and Cox model coefficient for the complete ER+

METABRIC cohort.

$$h_i^{ENDORSE}(t) = h_0(t)\exp\left(\beta_E x_{E_i}^T\right)$$

Where, X_E is the set of ENDORSE GES and β_E are the associated coefficients. We then stratified

the tumors based on ENDORSE risk estimate by defining cancers with HR ≤ 1 as low-risk, HR ≥

2 as high-risk, and those with HR between 1 and 2 as medium risk.

METABRIC survival models and simulations

The ENDORSE model was evaluated in the METABRIC cohort by stratifying the ER+ cancers

based on estimated risk in the low, medium or high-risk categories. For comparisons with

clinical variables and other breast cancer classification approaches, including PAM50 and

IntClust, we reported the HR and p-values of the Cox model coefficients for these predictors in

univariate analyses. In addition, we reported the results of multivariate analyses that included the

PAM50 or IntClust classes along with the ENDORSE risk estimates.

We analyzed the survival curves of METABRIC ER+ tumors stratified based on ENDORSE risk

estimates using the log-rank test. To simulate the effects of random dropouts or missing data, we

sequentially reduced the total number of available genes to calculate the ENDORSE GES and

recalculated the risk estimate. Here, we randomly sampled and removed 10% of the genes in

each iteration until 90% of the genes were removed. In each iteration, we calculated the ability of

the recalculated risk estimate to stratify the cancers based on the difference in survival curves

using the log-rank test. Similarly, we simulated the impact of reduced sample set by sequentially

reducing the number of samples available for recalculating the ENDORSE scores by 10% in

each iteration and analyzed the difference in survival curves using the log-rank test.

Validation in independent clinical trial datasets

We evaluated the performance and ability of ENDORSE to predict treatment response in a set of

three independent clinical trials. These trials were selected based on criteria that the trial cohort

included any stage of ER+/HER2- breast cancers receiving endocrine therapy but did not receive

chemotherapy. In addition, the trials should have reported a clinical assessment of treatment

response or over survival outcomes, in addition to providing gene expression datasets. We found

three datasets the met the above criteria (GSE12647, GSE87411 and GSE76040). The first trial

(GSE12647) reported survival outcomes of 140 metastatic breast cancers on endocrine therapy.

For this dataset, we predicted the estimated risk of event based on the ENDORSE coefficients

derived from the METABRIC ER+ model. First, we calculated the ENDORSE GES for the test

samples using the baseline expression of the tumors. Next, we predicted the estimate risk of

event using the METABRIC-derived coefficients. We then evaluated the difference in survival

curves for the different strata using the log-rank test.

For the two subsequent datasets (GSE87411 and GSE76040), we again predict the estimated risk

based on ENDORSE GES from the baseline gene expression data. The NCT00265759 trial

evaluated neoadjuvant aromatase inhibitor response in a cohort of 109 stage II-III ER+ breast

cancers. This trial provided Ki67 data at both baseline and end of treatment and classified tumor

with Ki67 > 10% at end of treatment as non-responders. The CONFIRM trial evaluated 113

metastatic ER+ cancers on fulvestrant, and also reported Ki6 staining data at end of treatment.

Since both trials reported Ki67 as a determinant for treatment response, we compared the Ki67 %

across the ENDORSE risk strata or continuous risk scores as measure of performance evaluation.

In addition, we also evaluated the ENDORSE risk estimates in two negative control datasets. The

first negative control dataset (GSE130788) evaluated trastuzumab and lapatinib response in 62

ER+/HER+ or 48 ER-/HER+ breast cancers. The trial reported clinical response (partial clinical

response or residual disease) at the end of treatment. We predicted the estimated ENDORSE risk

scores using baseline gene expression and compared across the clinical response classes. As an

additional negative control, we also predicted and stratified the 429 ER- breast cancers in

METABRIC based on ENDORSE estimates. Then, we compared the difference in survival

curves using log-rank test.

Biological features associated with high-risk cancers

We assessed the features that may be important in explaining the biological difference between

high-risk cancers compared to low/medium risk cancers by comparing the curated and hallmark

enrichment scores, somatic mutation and copy number frequencies across the ER+ METABRIC.

For the comparison of GES, we performed t-tests for each signature and adjusted the p-values

using false discovery rate (FDR) or Benjamini-Hochberg method. In addition, we calculated the

effect size of the difference in means using Cohen's D method. An absolute Cohen's D > 0.8 was

considered as large effect.

The somatic mutations for the METABRIC ER+ cohort were summarized at the gene-level by

first removing all synonymous variants as non-consequential, then binarizing the gene by

mutation matrix based on presence of a mutation. Next, we filtered out genes with a sample

mutation frequency < 0.05. We then constructed a 2 x 2 contingency matrix for each gene and

the METABRIC ER+ tumor stratified based on low/medium risk or high risk. We calculated the

p-values for enrichment using the Chi-square test, followed by FDR adjustment. Similar to the

mutation analysis, we performed the Chi-square test on binary matrices of tumor gain or loss

across all genes, followed by FDR adjustment.

To compare the information captured by the ENDORSE scores with the proliferative meta-

PCNA signature [22], we calculated the proliferation index of the METABRIC ER+ tumors using

the R-package ProliferationIndex [37]. The package calculates the median expression of the meta-

PCNA genes, as described by Venet et al. [22]. Next, we performed multivariate Cox analysis of

the ER+ tumors with the proliferation index and either ENDORSE risk groups or continuous risk

estimates as covariates.

Data availability and code

All training and validation datasets used in this study are publicly available and listed under

"data retrieval, preprocessing and analysis". The sample code for reproducing the analyses in this

study are available at https://osf.io/bd3m7/?view_only=da4f860bd2474745880944fce1d433b1

References

1. Harvey, J. M., Clark, G. M., Osborne, C. K. & Allred, D. C. Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J. Clin. Oncol.* **17**, 1474–1481 (1999).

2. Rugo, H. S. *et al.* Endocrine Therapy for Hormone Receptor–Positive Metastatic Breast Cancer: American Society of Clinical Oncology Guideline. *JCO* **34**, 3069–3103 (2016).

3. Allison, K. H. *et al.* Estrogen and Progesterone Receptor Testing in Breast Cancer: ASCO/CAP Guideline Update. *JCO* **38**, 1346–1366 (2020).

4. Musgrove, E. A. & Sutherland, R. L. Biological determinants of endocrine resistance in breast cancer. *Nature Reviews Cancer* **9**, 631–643 (2009).

5. Spoerke, J. M. *et al.* Heterogeneity and clinical significance of ESR1 mutations in ER-positive metastatic breast cancer patients receiving fulvestrant. *Nature Communications* **7**, 1–10 (2016).

6. Parker, J. S. *et al.* Supervised Risk Predictor of Breast Cancer Based on Intrinsic Subtypes. *J Clin Oncol* **27**, 1160–1167 (2009).

7. Paik, S. *et al.* A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N. Engl. J. Med.* **351**, 2817–2826 (2004).

8. Cardoso, F. *et al.* 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *https://doi.org/10.1056/NEJMoa1602253* https://www.nejm.org/doi/10.1056/NEJMoa1602253 (2016) doi:10.1056/NEJMoa1602253.

9. Witten, D. M. & Tibshirani, R. Survival analysis with high-dimensional covariates. *Stat Methods Med Res* **19**, 29–51 (2010).

10.   Taylor, J. M. G., Ankerst, D. P. & Andridge, R. R. Validation of Biomarker-Based Risk

Prediction Models. *Clin Cancer Res* **14**, 5977–5983 (2008).

11.   Boutros, P. C. The path to routine use of genomic biomarkers in the cancer clinic.

Genome Res **25**, 1508–1513 (2015).

12.   Ali, H. R. *et al.* Genome-driven integrated classification of breast cancer validated in

over 7,500 samples. *Genome Biology* **15**, 431 (2014).

13.   Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours

reveals novel subgroups. *Nature* **486**, 346–352 (2012).

14.   Pereira, B. *et al.* The somatic mutation profiles of 2,433 breast cancers refine their

genomic and transcriptomic landscapes. *Nature Communications* **7**, 1–16 (2016).

15.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for

microarray and RNA-Seq data. *BMC Bioinformatics* **14**, 7 (2013).

16.   Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS -driven

cancers require TBK1. *Nature* **462**, 108–112 (2009).

17.   Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,

1739–1740 (2011).

18.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection.

Cell Systems **1**, 417–425 (2015).

19.   Sinn, B. V. *et al.* SETER/PR: a robust 18-gene predictor for sensitivity to endocrine

therapy for metastatic breast cancer. *NPJ Breast Cancer* **5**, 16 (2019).

20.   Ellis, M. J. *et al.* Ki67 Proliferation Index as a Tool for Chemotherapy Decisions During

and After Neoadjuvant Aromatase Inhibitor Treatment of Breast Cancer: Results From the

American College of Surgeons Oncology Group Z1031 Trial (Alliance). *J. Clin. Oncol.* **35**,

1061–1069 (2017).

21.   Jeselsohn, R. *et al.* TransCONFIRM: Identification of a Genetic Signature of Response to

Fulvestrant in Advanced Hormone Receptor-Positive Breast Cancer. *Clin. Cancer Res.* **22**,

5755–5764 (2016).

22.   Venet, D., Dumont, J. E. & Detours, V. Most Random Gene Expression Signatures Are

Significantly Associated with Breast Cancer Outcome. *PLOS Computational Biology* **7**,

e1002240 (2011).

23.   Kjällquist, U. *et al.* Exome sequencing of primary breast cancers with paired metastatic

lesions reveals metastasis-enriched mutations in the A-kinase anchoring protein family

(AKAPs). *BMC Cancer* **18**, 174 (2018).

24.   Sparano, J. A. *et al.* Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in

Breast Cancer. *New England Journal of Medicine* **379**, 111–121 (2018).

25.   Sotiriou, C. *et al.* Breast cancer classification and prognosis based on gene expression

profiles from a population-based study. *PNAS* **100**, 10393–10398 (2003).

26.   Paul, D., Bair, E., Hastie, T. & Tibshirani, R. "Preconditioning" for feature selection and

regression in high-dimensional problems. *Ann. Statist.* **36**, 1595–1618 (2008).

27.   Cattoretti, G., Rilke, F., Andreola, S., D'Amato, L. & Delia, D. P53 expression in breast

cancer. *International Journal of Cancer* **41**, 178–183 (1988).

28.   Elledge, R. M. *et al.* Prognostic significance of p53 gene alterations in node-negative

breast cancer. *Breast Cancer Res Tr* **26**, 225–235 (1993).

29.   Ungerleider, N. A. *et al.* Breast cancer survival predicted by TP53 mutation status differs

markedly depending on treatment. *Breast Cancer Research* **20**, 115 (2018).

30.   Bosco, E. E. *et al.* The retinoblastoma tumor suppressor modifies the therapeutic

response of breast cancer. *J Clin Invest* **117**, 218–228 (2007).

31.   Witkiewicz, A. K. & Knudsen, E. S. Retinoblastoma tumor suppressor pathway in breast

cancer: prognosis, precision medicine, and therapeutic interventions. *Breast Cancer Research*

16, 207 (2014).

32.   Gradishar, W. J. *et al.* Breast Cancer, Version 3.2020, NCCN Clinical Practice

Guidelines in Oncology. *Journal of the National Comprehensive Cancer Network* **18**, 452–

478 (2020).

33.   Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles using

the cBioPortal. *Sci Signal* **6**, pl1 (2013).

34.   Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *Journal of the Royal*

Statistical Society: Series B (Methodological) **58**, 267–288 (1996).

35.   Friedman, J. H., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear

Models via Coordinate Descent. *Journal of Statistical Software* **33**, 1–22 (2010).

36.   Simon, N., Friedman, J. H., Hastie, T. & Tibshirani, R. Regularization Paths for Cox's

Proportional Hazards Model via Coordinate Descent. *Journal of Statistical Software* **39**, 1–13

(2011).

37.   Ramaker, R. C. *et al.* RNA sequencing-based cell proliferation analysis across 19 cancers

identifies a subset of proliferation-informative cancers with a common survival signature.

Oncotarget **8**, 38668–38681 (2017).

a   Pathways enriched in high risk tumors ER+ METABRIC tumors

b   Pathways enriched in low/medium risk ER+ METABRIC tumors

c   Mutation frequency in ER+ METABRIC breast cancers stratified by estimated risk