

# **Title: Lifetime depression and age-related changes in body composition, cardiovascular function, grip strength and lung function: sex-specific analyses in the UK Biobank**

**Short title:** Age-related physiological changes in lifetime depression

**Authors:** Julian Mutz<sup>1\*</sup> and Cathryn M. Lewis<sup>1,2</sup>

## **Affiliations:**

1. Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, Greater London, SE5 8AF, United Kingdom.
2. Department of Medical and Molecular Genetics, Faculty of Life Sciences & Medicine, King's College London, London, Greater London, SE1 9RT, United Kingdom.

## **\*Corresponding author:**

Julian Mutz; Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London; Memory Lane, London, SE5 8AF, United Kingdom. Email:

[julian.mutz@gmail.com](mailto:julian.mutz@gmail.com).

**Keywords:** Ageing; Body composition; Cardiovascular function; Depression; Grip strength; Lung function

## Abstract

Individuals with depression, on average, die prematurely, have high levels of physical comorbidities and may experience accelerated biological ageing. A greater understanding of age-related changes in physiology could provide novel biological insights that may help inform strategies to mitigate excess mortality in depression. We used generalised additive models to examine age-related changes in 15 cardiovascular, body composition, grip strength and lung function measures, comparing males and females with a lifetime history of depression to healthy controls. The main dataset included 342,393 adults (mean age = 55.87 years, SD = 8.09; 52.61% females). We found statistically significant case-control differences for most physiological measures. There was some evidence that age-related changes in body composition, cardiovascular function, lung function and heel bone mineral density followed different trajectories in depression. These differences did not uniformly narrow or widen with age and differed by sex. For example, BMI in female cases was 1.1 kg/m<sup>2</sup> higher at age 40 and this difference narrowed to 0.4 kg/m<sup>2</sup> at age 70. In males, systolic blood pressure was 1 mmHg lower in depression cases at age 45 and this difference widened to 2.5 mmHg at age 65. These findings suggest that targeted screening for physiological function in middle-aged and older adults with depression is warranted to potentially mitigate excess mortality.

# 1 Introduction

2

3 Individuals with mental disorders, on average, die prematurely [1] and have high levels of physical comorbidities [2].

4 Although it is almost universally accepted that health deteriorates as part of the normal ageing process, there is

5 substantial variation in ageing trajectories [3]. Up to a third of older adults experience depression and in many cases

6 show greater levels of functional decline and cognitive impairment [4].

7

8 To better understand the heterogeneity in ageing trajectories, it would be useful to objectively assess whether groups or

9 individuals deviate from ‘normal’ age-related changes. To date, no single biological marker that fully captures the ageing

10 process has been discovered, although several candidate measures such as telomere attrition have been described [5]. The

11 measurement of blood-based biomarkers such as leukocyte telomere length is minimally invasive, relatively costly and

12 time-consuming, especially at population scale. Magnetic resonance imaging, for instance used to derive estimates of

13 brain age [6], is costly and less accessible. Physiological measures such as hand-grip strength can be easily obtained,

14 provide reliable information about age-related functional decline and predict morbidity and mortality risk [7-9].

15

16 Several studies have examined age-related variation in these measures [10-12]. However, less is known about the ageing

17 trajectories of individuals with depression and whether similar patterns are observed across different physiological

18 measures. A deeper understanding of age-related changes in physiology in individuals with depression might provide

19 insights into the premature mortality observed in mental disorders and could point towards potential targets for

20 prevention or intervention of accelerated ageing [13].

21

22 The aim of this study was to examine associations between age and multiple physiological measures, and to test for

23 differences in these measures between middle-aged and older adults with a lifetime history of depression and healthy

24 controls across age. Given major sex differences in physiology, age-related diseases and life expectancy [14], and in

25 prevalence of depression [15], analyses were conducted separately in males and females. In a separate study, we have

26 examined age-related changes in physiology in bipolar disorder [16].

27

28 Using data from up to ~342,000 UK Biobank participants aged 37-73, the following research questions were examined

29 for each physiological measure assessed during the baseline assessment: Are there average differences between males

30 and females with lifetime depression and healthy controls? Do age-related changes differ between males and females

31 with lifetime depression and healthy controls?

## 32 **Results**

33

### 34 **Study population**

35 Of the 502,521 UK Biobank participants, 392,467 (78.1%) individuals had complete data on all covariates. We yielded  
36 an analytical sample of 342,393 adults after excluding individuals who did not meet our inclusion criteria or whose  
37 lifetime depression status could not be determined. Sample sizes for physiological measures that were available in  
38 subsets of the UK Biobank population ranged from 120,843 to 224,805 participants ([Figure 1](#)).

39

### 40 **Sample characteristics**

41 The average age of participants in our main dataset was 55.87 years (SD = 8.09) and 52.61% of participants were  
42 female. About 22.25% of participants ( $n = 87,339$ ) met criteria for lifetime depression, 64.43% ( $n = 56,276$ ) of whom  
43 were female. Descriptive statistics and density plots of the physiological measures stratified by sex and lifetime  
44 depression status are presented in [Table 1](#) and [Supplement 5](#), respectively. Descriptive statistics of the covariates are  
45 presented in [Supplement 6](#).

46

### 47 **Sex differences**

48 We found moderate to large sex differences across all physiological measures. Female participants, on average, had  
49 lower hand-grip strength, systolic and diastolic blood pressure, BMI, fat-free mass, waist-hip ratio, peak expiratory flow,  
50 FEV<sub>1</sub>, forced vital capacity, heel bone mineral density and arterial stiffness than males. Their pulse rate, body fat  
51 percentage, fat mass and FEV<sub>1</sub>/FVC ratio were higher than in males ([Table 2](#)).

52

### 53 **Case-control differences**

54 We also found statistically significant differences between individuals with lifetime depression and healthy controls for  
55 most measures. Females with lifetime depression, on average, had lower hand-grip strength and lower blood pressure.  
56 Their pulse rate, BMI, body fat percentage, fat mass, fat-free mass, waist-hip ratio, peak expiratory flow, FEV<sub>1</sub>, forced  
57 vital capacity and FEV<sub>1</sub>/FVC ratio were higher compared to healthy controls. We did not find evidence of differences in  
58 heel bone mineral density or arterial stiffness between female cases and controls. Males with lifetime depression had  
59 lower hand-grip strength, blood pressure, peak expiratory flow, FEV<sub>1</sub>, forced vital capacity and heel bone mineral  
60 density, while their pulse rate, BMI, body fat percentage, fat mass, fat-free mass and waist-hip ratio were higher. We did  
61 not find evidence that the FEV<sub>1</sub>/FVC ratio or arterial stiffness differed between males with lifetime depression and  
62 healthy controls. The differences between individuals with lifetime depression and healthy controls were small, with the

63 largest effect size corresponding to a standardised mean difference of 0.16 (95% CI 0.15-0.17) for fat mass in females  
64 (Table 3).

65

#### 66 **Age-related physiological changes**

67 Results from the unadjusted GAMs suggested that hand-grip strength, fat-free mass, lung function and heel bone mineral  
68 density declined with age, while systolic blood pressure, body fat percentage, fat mass, waist-hip ratio and arterial  
69 stiffness increased with age. Non-linear associations with age were most evident for diastolic blood pressure, pulse rate  
70 and BMI. For example, diastolic blood pressure increased until about age 55 and decreased thereafter. BMI increased  
71 with age until participants were in their mid to late 50s and plateaued thereafter, except in healthy females where BMI  
72 increased until age 70 (Figure 2-3).

73

#### 74 **Case-control differences in age-related physiological changes**

75 Age-related changes in blood pressure, pulse rate, body composition and heel bone mineral density followed distinct  
76 trajectories in females with lifetime depression and healthy controls (Figure 4). There was also evidence of differences  
77 in age-related changes hand-in grip strength, blood pressure, pulse rate, lung function and heel bone mineral density  
78 between males with lifetime depression and healthy controls (Figure 5).

79

80 In females, we obtained similar results after adjustment for covariates, although differences between cases and controls  
81 were attenuated and there was greater overlap in the trajectories of pulse rate, lung function and heel bone mineral  
82 density. We found no evidence that age-related changes in arterial stiffness differed between female cases and controls  
83 (Supplement 7A). We also obtained similar results after adjustment for covariates in males, with some attenuation of  
84 differences between cases and controls (Supplement 7B). The formal statistical tests for differences in age-related  
85 changes in hand-grip strength and heel bone mineral density were no longer statistically significant, although age-related  
86 changes in forced vital capacity and FEV<sub>1</sub> differed between male cases and controls.

87

#### 88 **Sensitivity analyses**

89 Case-control numbers for the sensitivity analyses are presented in Supplement 8A. We retained between 35.15% to  
90 45.97% of cases and 100% of controls in the analyses in which cases included only those with lifetime depression  
91 according to at least two different measures. In the analyses in which we restricted our sample to participants with  
92 lifetime depression status assessed using the CIDI-SF, we retained between 24.69% to 40.67% of cases and between  
93 22.58% to 28.03% of controls.

94

95 We found similar results when we applied more stringent criteria for lifetime depression. Average differences between  
96 female cases and controls were slightly larger for all physiological measures. In males, differences were also larger,  
97 except that there was no evidence of differences in lung function between cases and controls ([Supplement 8B](#)). Most  
98 results were also similar when we restricted our analyses to individuals with lifetime depression status assessed using the  
99 CIDI-SF. In females, differences were slightly larger, except for hand-grip strength, and there was no evidence of  
100 differences in diastolic blood pressure. In males, there was no evidence that hand-grip strength or diastolic blood  
101 pressure differed between cases and controls, and differences in lung function measures were reversed (although there  
102 was no evidence of differences in peak expiratory flow or the FEV<sub>1</sub>/FVC ratio). Average differences in body  
103 composition, systolic blood pressure and pulse rate were slightly larger, while differences in heel bone mineral density  
104 were slightly smaller ([Supplement 8B](#)).

105

106 Differences in age-related changes in physiological measures between female cases and controls were similar to those  
107 observed in our main analyses ([Supplement 8C](#)). There was less evidence that age-related changes in systolic blood  
108 pressure differed between males with lifetime depression according to CIDI-SF and healthy controls. There was also  
109 some evidence that age-related changes in body composition followed slightly different trajectories in males with  
110 lifetime depression according to CIDI-SF, and there was more overlap in the lung function trajectories ([Supplement 8D](#)).

## 111 **Discussion**

112 We used data from the UK Biobank to examine differences in 15 physiological measures, between males and females,  
113 and between lifetime depression cases and healthy controls. The large sample size enabled us to highlight subtle  
114 differences between cases and controls, with heterogenous findings from previous studies that were often limited by  
115 small sample sizes. We demonstrate that case-control differences for several physiological measures vary by age,  
116 providing additional insights into the heterogeneity in ageing trajectories observed in individuals with a lifetime history  
117 of depression.

118

### 119 **Principal findings**

120 We observed moderate to large sex differences across all physiological measures. We also found small but statistically  
121 significant differences between lifetime depression cases and healthy controls for most measures. All physiological  
122 measures varied with age, and we observed several differences between the trajectories of depression cases and controls.  
123 Age-related changes in body composition differed between female cases and controls, while changes in cardiovascular  
124 function differed between cases and controls of both sexes. There was some evidence of differences in the heel bone  
125 mineral density trajectories of both sexes, although in males only prior to adjustment for covariates. We found some  
126 evidence that age-related changes in lung function differed between male cases and controls after adjustment for  
127 covariates. Differences between cases and controls did not uniformly narrow or widen with age. For example, BMI in  
128 female cases was 1.1 kg/m<sup>2</sup> higher at age 40 and this difference narrowed to 0.4 kg/m<sup>2</sup> at age 70. In male cases, systolic  
129 blood pressure was 1 mmHg lower at age 45 and this difference widened to 2.5 mmHg at age 65.

130

### 131 **Findings in the context of other studies**

132

#### 133 **Sex differences**

134 We confirm previous research on sex differences in hand-grip strength [10, 17-19], blood pressure [20], pulse rate [21,  
135 22], lung function [23-25], body fat percentage [26, 27], fat-free mass [26, 27] and waist-hip ratio [28]. Our finding that  
136 BMI was higher in males confirms findings from a German study [28], although studies in US and Chinese adults found  
137 no evidence of sex differences [26, 27]. The observation that fat mass was higher in females confirms findings from  
138 Sweden [29] and the US [26], although research in Chinese adults found no evidence of sex differences [27]. Males had  
139 higher heel bone mineral density, supporting previous studies of hand [30] and heel [31] bone mineral density. Our  
140 finding that arterial stiffness was higher in males supports a review which found greater arterial stiffness in males until at

141 least age 60 [32]. However, most studies included in another systematic review found no evidence of sex differences  
142 [33], and some studies found higher arterial stiffness in females [34-36].

143

#### 144 **Age-related physiological changes**

145 Our findings confirm previous research on age-related changes in hand-grip strength [10, 18, 19], blood pressure [12, 20,  
146 37, 38], fat mass [26, 29, 39], fat-free mass [26, 39, 40], waist-hip ratio [28], lung function [23-25, 41], bone mineral  
147 density [30, 42] and arterial stiffness [21, 34]. In males, we observed a small increase in pulse rate until age 55 and a  
148 decrease thereafter, similar to data from activity trackers in >90,000 US adults [22]. However, we found a slight decrease  
149 in pulse rate in females until their mid 50s and an increase thereafter. Although we observed some increase in BMI with  
150 age, comparable to results from Swedish cohorts [29], Chinese research found no evidence of age-related changes in  
151 BMI [27]. Our finding that body fat percentage increased with age confirms previous studies [29, 40], although we found  
152 limited evidence of a plateau or decrease before age 70 observed in previous research [26].

153

#### 154 **Sex differences in age-related physiological changes**

155 We confirm previous studies on sex differences in age-related changes in hand-grip strength [17, 18], blood pressure [12,  
156 20, 37], lung function [24, 41] and bone mineral density [30, 42]. In contrast to observations that pulse rate followed  
157 parallel ageing trajectories in both sexes [22], we found changes in pulse rate in females were mirrored by inverse  
158 changes in males.

159

160 Differences in BMI narrowed with age as BMI plateaued in males at age 50 and slightly decreased thereafter, while it  
161 increased in females until at least age 65. Greater age-related increase in BMI in females has been observed previously  
162 [29]. Body fat percentage followed similar trajectories with age in both sexes. However, research in Chinese adults  
163 suggested that body fat percentage plateaued in males over 40 and continued to increase in females [27], while results  
164 from the US showed that body fat percentage decreased in females and males over 55 and 65, respectively [26].  
165 Although Swedish data showed slightly greater age-related increase in fat mass in females [29], we found similar  
166 trajectories in both sexes. Our finding that fat mass plateaued earlier in females confirms research in US adults [26].  
167 However, research in Chinese adults found that fat mass remained stable in males from age 40 but continued to increase  
168 in females [27]. Our study did not support the finding that fat-free mass decreased earlier in females than in males [29].  
169 Fat-free mass decreased in both sexes, with steeper decline in males. Our finding that waist-hip ratio increased linearly  
170 with age in females confirms observations from China [43], but we did not find that waist-hip ratio plateaued in males at  
171 age 50 and decreased thereafter. Instead, we found that waist-hip ratio continued to increase until age 70. This increase

172 attenuated at old age, in line with German research [28]. The difference between the trajectories of males and females  
173 had an inverted U-shape, with the largest difference in waist-hip ratio between age 50 to 55.

174

175 We examined absolute changes in lung function. However, a review of longitudinal studies concluded that although there  
176 was a steeper decline in absolute lung function in males, evidence of sex differences in relative decline in lung function  
177 was limited [41]. Limited previous research examined sex differences in age-related changes in the FEV<sub>1</sub>/FVC ratio. We  
178 found that the difference in the FEV<sub>1</sub>/FVC ratio narrowed with age, with some variation at the lower and upper age  
179 range.

180

181 Finally, we found only limited evidence of a steeper age-related increase in arterial stiffness in males (in our study until  
182 the age of 50) as suggested by previous studies [21, 36], or that females over 65 had higher arterial stiffness than males  
183 [32]. Sex differences in arterial stiffness narrowed between age 60 and 70.

184

#### 185 **Case-control differences**

186 Our finding that hand-grip strength was lower in individuals with lifetime depression confirms previous research [44-47].  
187 Although several studies found stronger associations between hand-grip strength and depression in females [44, 48, 49]  
188 or no sex differences [45], we observed larger differences between male cases and controls. Results from the Northern  
189 Finland 1966 birth cohort suggested that low hand-grip strength was associated with increased depressive symptoms only  
190 in males [50], while in the Irish Longitudinal Study of Ageing higher hand-grip strength was associated with lower  
191 incident depression only in females [44].

192

193 The finding that cases had lower blood pressure confirms results from the Norwegian HUNT study [51, 52]. Although  
194 several studies found lower blood pressure in depression [51-54], other research found no evidence of differences in  
195 blood pressure [55] and a meta-analysis found higher incident hypertension in depression [56]. A 2-year follow-up study  
196 suggested that low blood pressure constitutes a risk factor for depression [57]. We confirmed previous analyses showing  
197 higher pulse rate in depression [58-60].

198

199 All body composition measures were elevated in cases. A meta-analysis of observational studies found an increased risk  
200 of obesity in depression [61], and findings from the 1958 British Birth Cohort found associations between depression and  
201 both underweight and obesity [62]. Previous studies of BMI and depression yielded mixed results, including higher BMI  
202 [63], lower BMI [64, 65], U-shaped association [66] or no association [67]. A Mendelian randomisation study suggested  
203 that BMI was a causal risk factor for depression but not vice versa [68]. Mixed results have also been observed for waist-

204 hip ratio, including lower risk of depressive symptoms in older females with high waist-hip ratio [65] and associations  
205 between depression and high waist-hip ratio [63, 69]. An analysis of UK Biobank participants found that females with  
206 probable depression had higher waist-hip ratio than males [70], although other research found higher waist-hip ratio in  
207 males with depression but not in females [71], or no evidence of an association with depressive symptoms [64]. Previous  
208 research found no evidence of associations between depression and total fat or lean mass, although central fat mass and  
209 lean mass were lower amongst older females with depression [72]. A Mendelian randomisation study suggested that fat  
210 mass, but not fat-free mass, was a causal risk factor for depression [68]. Although we found higher body fat percentage  
211 in cases, a recent analysis found associations between depression and higher body fat percentage only in females,  
212 although participants were aged 90 years [73].

213

214 In contrast to results from the Netherlands Study of Anxiety and Depression (NESDA) which suggested that peak  
215 expiratory flow was lower in females and higher in males with depression [74], we found higher peak expiratory flow in  
216 female cases. Male cases had lower peak expiratory flow, although not across all analyses. We found similar results for  
217 FEV<sub>1</sub>, forced vital capacity and FEV<sub>1</sub>/FVC ratio. Differences between male cases and controls were usually not observed  
218 in the sensitivity analyses, except that FEV<sub>1</sub> and forced vital capacity were higher in the subset of male cases classified  
219 according to CIDI-SF than in controls. Research in Germany found lower FEV<sub>1</sub> in depression but no evidence of  
220 differences in forced vital capacity or FEV<sub>1</sub>/FVC ratio [75]. Data from a community-based cohort in Korea suggested  
221 that FEV<sub>1</sub>, forced vital capacity and FEV<sub>1</sub>/FVC ratio were lower in depression, although only FEV<sub>1</sub> remained lower after  
222 adjustment for covariates [76]. However, a prospective study found that lower forced vital capacity and FEV<sub>0.75</sub> were  
223 associated with more depressive symptoms at follow-up in middle aged and older males [77].

224

225 We found that heel bone mineral density was lower in male cases, consistent with most previous studies [78]. There was  
226 some evidence that heel bone mineral density was higher in female cases in sensitivity analyses, contrary to most  
227 previous studies. A meta-analysis of 14 studies showed that spine and hip bone mineral density were lower in depression,  
228 especially in females [79]. One previous study suggested that heel bone mineral density was lower in female cases below  
229 the age of 40 but not in older females [80], although another study found lower heel bone mineral density also in  
230 postmenopausal females [81]. We did not find evidence that arterial stiffness differed between cases and controls. This  
231 finding contrasts with most previous studies that found higher arterial stiffness in depression [82, 83], including a recent  
232 UK Biobank study [84].

### 233 **Case-control differences in age-related physiological changes**

234 Hand-grip strength differences between female cases and controls were fairly stable across age, while generally  
235 narrowing with age in males. To our knowledge, the only study that examined changes in hand-grip strength in  
236 depression longitudinally found similar rates of decline in female cases and controls, in line with our findings. Contrary  
237 to our study, the authors found no evidence that hand-grip strength decreased in male cases [85].

238  
239 The HUNT study showed that prospective associations between depressive symptoms and changes in blood pressure  
240 were independent of age [51]. Findings from the Baltimore Longitudinal Study of Aging found associations between  
241 greater levels of depressive symptoms and lower diastolic blood pressure until age 45, and attenuated age-related decline  
242 thereafter in females but not in males [86]. We found that case-control differences in systolic blood pressure increased  
243 with age. There was no evidence of a difference in diastolic blood pressure until participants were in their early to mid  
244 50s, although diastolic blood pressure was lower in cases thereafter. This difference narrowed as participants approached  
245 age 70. To our knowledge, this is the first study to show that differences in pulse rate between cases and controls  
246 narrowed with age.

247  
248 This is also the first study to examine age-related changes in body composition in cases compared to controls. Across  
249 these measures, we found that differences in body composition narrowed with age in females, while generally remaining  
250 constant in males. There were some inconsistencies in the sensitivity analyses with more stringent case definition. These  
251 analyses suggested an increasing difference in body fat percentage, a flat inverted U-shape for fat mass and fat-free mass,  
252 and some variation in the difference smooth for waist-hip ratio.

253  
254 Results from the NESDA suggested a steeper decrease in peak expiratory flow in male cases during a 6-year follow-up,  
255 while rates of decline were similar in female cases and controls [85]. We also found some evidence of a steeper decrease  
256 in peak expiratory flow with age in male cases, although differences between female cases and controls remained stable.  
257 Rates of decline in the NESDA were similar in younger and older participants [85], concordant with our results showing  
258 an almost linear decrease in peak expiratory flow. While one previous study found a weak negative correlation between  
259 depressive symptoms and FEV<sub>1</sub> only in participants older than 50 [76], our study is, to our knowledge, the first to  
260 examine case-control differences in ageing trajectories across multiple lung function measures. We found that the  
261 difference in FEV<sub>1</sub> and forced vital capacity remained stable with age in females, with some variation in sensitivity  
262 analyses, while the difference widened with age in males, although not across all analyses. We did not find clear  
263 evidence of differences in the trajectories of FEV<sub>1</sub>/FVC ratio.

264

265 Differences in heel bone mineral density between cases and controls narrowed with age, approaching no difference in  
266 males in their late 60s and in females in their mid 50s, with some evidence of higher bone mineral density in older  
267 females. One previous study found lower heel bone mineral density in females with depression who were younger than  
268 40 but not in older females [80]. However, lower bone mineral density has also been found in postmenopausal females  
269 with depression [87]. A meta-analysis of 23 studies found lower bone mineral density in the spine, hip and forearm in  
270 premenopausal than in postmenopausal female cases compared to controls [88]. Our finding that differences in bone  
271 mineral density narrowed with age differs from a meta-analysis of five longitudinal studies which suggested that  
272 depression was associated with increased annual loss in hip and spine bone mineral density [79]. One study showed that  
273 elderly females with higher levels of depressive symptoms had an age-adjusted mean total hip bone mineral density  
274 decrease of 0.96% per year, compared to 0.69% in females with fewer symptoms [89]. Similar findings have been  
275 observed in males [90]. However, a prospective study of premenopausal females found no evidence of decreasing bone  
276 mineral density irrespective of depressive symptoms [91]. To our knowledge, this is the first study to examine  
277 differences in age-related changes in arterial stiffness between cases and controls. While we did not find clear evidence  
278 of different trajectories with age, there was some evidence that arterial stiffness was higher in female cases younger than  
279 age 50.

280  
281 Inconsistencies between studies could result from variation in sample characteristics, differences in assessment and  
282 definition of depression, or medication use. For example, previous research found that after adjustment for depression,  
283 antidepressant use was associated with lower spine bone mineral density in females but not in males [92], or that  
284 depression was associated with lower systolic blood pressure, while antidepressant use was associated with higher  
285 systolic and diastolic blood pressure [54]. Inconsistencies could also reflect differences in prevalence of specific  
286 symptoms. For example, somatic/vegetative symptoms but not mood/cognitive symptoms were previously associated  
287 with higher waist-hip ratio [69].

288  
289 Differences between cases and controls could reflect disease-related processes or poor underlying health status [93]  
290 occurring more frequently in cases. They could also result from biological mechanisms affecting depression and  
291 physiology (although a recent study did not find evidence of genetic correlations between depression and body  
292 composition [94]), or from treatment-related changes, for example lower bone mineral density due to antidepressant use  
293 [92]. Behavioural factors such as unhealthy lifestyles known to affect physiology (e.g., smoking and low levels of  
294 physical activity) might also contribute. It is likely that associations between depression and physiology are bi-directional  
295 in that depression is both cause and consequence of poor physiology [87]. Sex differences might be due to the relative  
296 frequency of different symptoms experienced by males and females or differences in prevalence of comorbidities.

297 **Weaknesses of the study**

298 Our research inevitably has limitations. Although we briefly remarked on this above, our analyses provide limited  
299 insights into the mechanism underpinning our results. The cross-sectional nature of our study presents additional  
300 uncertainty about whether these findings reflect changes in physiology due to ageing or cohort-specific differences in  
301 these measures and confounding factors. For example, previous research suggested that hand-grip strength is lower in  
302 younger cohorts [19], which could result in underestimating age-related decline in our study. Research on frailty indices  
303 provided evidence of stability across birth cohorts [95] as well as higher levels of frailty in more recent cohorts [96, 97].  
304 The finding that some physiological differences narrowed with age could result from selection bias leading to older  
305 participants with depression in our study being healthier, relative to their age group, than younger participants. Selection  
306 bias resulting in healthier older adults participating in the UK Biobank at higher rates could also result in underestimating  
307 age-related changes. Finally, some concerns remain about the validity and reliability of our definition of lifetime  
308 depression [98]. To maximise the sample size available for our main analyses, we classified cases using multiple data  
309 sources. However, these sources all have specific strengths and limitations, which have been discussed in detail  
310 elsewhere [98-100]. For example, mental disorders are likely underreported in linked health records. Self-reports might  
311 contain some level of misclassification because mental disorders are heterogenous in their presentation and have fuzzy  
312 boundaries, although participants were asked to only report diagnoses made by health professionals. To address these  
313 concerns, we conducted sensitivity analyses with more stringent case definition and found comparable results. Given that  
314 mental disorders fluctuate throughout the life course, we examined lifetime depression, which includes single episode  
315 and recurrent depression. Future studies could examine differences between single episode and recurrent depression as  
316 well as changes in physiology in relation to current psychiatric status at a more fine-scale timespan longitudinally.

317

318 **Generalisability**

319 The overall response rate of the UK Biobank was low (5.5%) and compared with non-responders, participants were  
320 older, more often female, of higher socio-economic status and reported fewer health conditions compared with data from  
321 a nationally representative survey [101]. An empirical investigation comparing the UK Biobank with data from 18 cohort  
322 studies with conventional response rates found the direction of associations to be similar, although with differences in  
323 magnitude [102]. There are additional concerns about the representativeness of individuals with mental disorders because  
324 participation in research can be influenced by selection bias due to mental health [103]. Our findings might not  
325 generalise to younger and older populations and conclusions are limited to adults between the ages of 37 to 73.

326 Additional studies including children and adolescents as well as the elderly are needed.

327 **Implications**

328 The primary aim of this study was to examine differences in age-related physiological changes between individuals with  
329 lifetime depression and healthy controls. Understanding age-related changes in physiology in individuals with depression  
330 is of public health importance, given that variation in these measures is linked to morbidity, mortality and other  
331 functional outcomes. The differences observed between cases and controls suggest that targeted screening for  
332 physiological function in middle-aged and older adults with depression is warranted to potentially mitigate excess  
333 mortality. Future studies aimed at further disentangling aetiology are needed to inform preventative and treatment  
334 strategies. Interventions such as resistance training might help improve physiological health and reduce depressive  
335 symptoms [104], although multiple factors associated with health status should be considered [105].

## 336 **Methods**

### 337 **Study population**

338 The UK Biobank is a prospective study of >500,000 UK residents aged 37–73 at baseline, recruited between 2006–2010.  
339 Details of the study rationale and design have been reported elsewhere [106]. Briefly, individuals registered with the UK  
340 National Health Service (NHS) and living within a 25-mile (~40 km) radius of one of 22 assessment centres were invited  
341 to participate (9,238,453 postal invitations sent). Participants provided information on their sociodemographic  
342 characteristics, health behaviours and medical history. They also underwent physical examination including height,  
343 weight and blood pressure measurement, and had blood and urine samples taken. Linked hospital inpatient records are  
344 available for most participants and primary care records are currently available for approximately half of participants. A  
345 subset of 157,366 out of 339,092 invited participants (46%) completed an online follow-up mental health questionnaire  
346 (MHQ) between 2016-2017, covering 31% of all participants.

347

### 348 **Exposures**

349 Age at baseline assessment was the primary explanatory variable.

350

351 Lifetime depression was assessed as part of the MHQ using a modified version of the depression module of the  
352 Composite International Diagnostic Interview Short Form (CIDI-SF) and defined according to DSM-5 criteria for major  
353 depressive disorder ([Supplement 1](#)) [107]. To achieve maximum coverage of the UK Biobank study, we also included in  
354 our definition of lifetime depression individuals who had reported during the nurse-led interview at baseline that a doctor  
355 had told them that they had depression (field 20002), participants who reported in response to a single-item question on  
356 the MHQ that a professional (doctors, nurse or person with specialist training such as psychologist or therapist) had  
357 diagnosed them with depression (field 20544), participants with a hospital inpatient record containing an ICD-10 code  
358 for depression (F32-F33; [Supplement 2](#)), participants with a primary care record containing a Read v2 or CTV3 code for  
359 depression (see [108] for diagnostic codes and data extraction procedures), and those who were classified as individuals  
360 with probable depression according to Smith et al. [109] based on additional questions that were introduced during the  
361 later stages of the baseline assessment ([Supplement 3](#)). We excluded individuals with any record of bipolar disorder or  
362 psychosis.

363

364 Healthy controls were defined as individuals who did not meet our criteria for lifetime depression and did not have a  
365 record of other mental disorders: (i) no “schizophrenia”, “mania/bipolar disorder/manic depression”, “anxiety/panic  
366 attacks”, “obsessive compulsive disorder”, “anorexia/bulimia/other eating disorder”, “post-traumatic stress disorder”

367 reported during the nurse-led interview at baseline (field 20002); (ii) no mental disorders reported in response to the  
368 single-item question on the MHQ (field 20544); (iii) no self-reported current psychotropic medication use at baseline  
369 ([Supplement 4](#)) [100]; (iv) no linked hospital inpatient record that contained any ICD-10 Chapter V code except organic  
370 causes or substance use (F20-F99); (v) no primary care record containing a diagnostic code for mental disorders [108];  
371 (vi) not classified as individual with probable bipolar disorder according to Smith et al. [109]; (vi) no Patient Health  
372 Questionnaire-9 (PHQ-9) sum score of  $\geq 5$ , which was assessed as part of the MHQ.

373

## 374 **Outcomes**

375 We examined 15 continuous physiological measures that were obtained at the baseline assessment, including maximal  
376 hand-grip strength, systolic and diastolic resting blood pressure, resting pulse rate, body mass index (BMI), waist-hip  
377 ratio, fat mass, fat-free mass, body fat percentage, peak expiratory flow, forced vital capacity (FVC), forced expiratory  
378 volume in one second (FEV<sub>1</sub>), FVC/FEV<sub>1</sub> ratio, heel bone mineral density and arterial stiffness.

379

## 380 **Physiological measures**

381 All physiological measures were collected by certified healthcare technicians or nurses using a direct data entry system.  
382 Participants were asked to remove any outer garments, shoes, socks or tights. The assessment lasted about 10-15  
383 minutes.

384

## 385 **Hand-grip strength**

386 Hand-grip strength in whole kilogram force units was measured using a Jamar J00105 hydraulic hand dynamometer  
387 (measurement range 0-90 kg). Participants were asked to sit upright in a chair and place their forearms on armrests. The  
388 dynamometer handle was set to the second incremental slot or, in participants with very large hands, moved to the third  
389 slot. Participants kept their elbow adjacent to their side and bent to a 90° angle with their thumb facing upward. A  
390 maximal score was obtained from each participant's right and left hand. We used the maximal grip strength of the  
391 participant's self-reported dominant hand. If no data on handedness were available, we used the highest value of both  
392 hands. This variable has been used previously in UK Biobank research [110], although other studies have used the  
393 highest value of left and right hand [111] or calculated the average grip strength from both hands [9]. We used absolute  
394 units because these are the simplest to use in risk assessment. A previous UK Biobank study found no evidence of  
395 differences in mortality or disease incidence prediction when grip strength was expressed in absolute terms compared to  
396 when expressed relative to anthropometric traits (height, weight, fat-free mass and BMI) [9].

### 397 **Body composition**

398 Weight measurements were obtained with a Tanita BC-418 MA body composition analyser, or, in individuals with  
399 pacemaker or females who reported that they were or might be pregnant, using a manual scale. Standing height was  
400 measured using a Seca 202 height measure. Body mass index (BMI) was calculated as weight divided by standing height  
401 squared ( $\text{kg/m}^2$ ). Waist and hip circumference in cm were measured using a Wessex non-stretchable sprung tape. Waist-  
402 to-hip ratio was calculated by dividing waist circumference by hip circumference. Whole body fat mass and fat-free mass  
403 in kg and body fat percentage (measurement range 1-75%) were estimated by electrical bio-impedance.

404

### 405 **Pulmonary function**

406 Volumetric measures of lung function were quantified using breath spirometry with a Vitalograph Pneumotrac 6800  
407 following standard procedures. Participants were asked to record two to three blows, each lasting for at least six seconds,  
408 within a period of approximately six minutes. The reproducibility of the first two blows was automatically compared  
409 and, if acceptable (defined as a  $\leq 5\%$  difference in forced vital capacity (FVC) and forced expiratory volume in one  
410 second ( $\text{FEV}_1$ )), a third blow was not required. FVC in litres describes the maximum amount of air that can be exhaled  
411 when blowing out as fast as possible after a deep breath.  $\text{FEV}_1$  in litres describes the amount of air that can be exhaled in  
412 one second when blowing out as fast as possible. We used the derived best measure for both FVC and  $\text{FEV}_1$  which was  
413 the maximum value from reproducible spirograms, in line with previous research [112]. Given that these measures are  
414 affected by several factors unrelated to pulmonary function (e.g., effort and body size), we also calculated the ratio of  
415  $\text{FEV}_1$  to FVC. Peak expiratory flow (PEF) in litres per minute represents a person's maximum speed of expiration. PEF  
416 is determined by physiological factors such as lung volume and elasticity or expiratory muscle strength. It is used for  
417 monitoring asthma and diagnosing chronic obstructive pulmonary disease. We calculated the average of all available  
418 readings. Spirometry was not performed in participants who confirmed or were unsure that they had any of the following  
419 contra-indications: chest infection in the last month (i.e., influenza, bronchitis, severe cold or pneumonia); history of a  
420 detached retina; heart attack or surgery to eyes, chest or abdomen in the last three months; history of a collapsed lung;  
421 pregnancy in the 1st or 3rd trimester; currently on medication for tuberculosis.

422

### 423 **Heel bone mineral density**

424 Heel bone mineral density was estimated by quantitative ultrasound assessment of the calcaneus using a Sahara Clinical  
425 Bone Sonometer. Participants were asked to sit with their back straight and had their left heel measured first.  
426 Measurement of both heels was performed only during later stages of the baseline assessment. Measures of speed of  
427 sound in metres per second and broadband ultrasound attenuation (decibels/megahertz) were combined into a

428 quantitative ultrasound index. From this an estimate of heel bone mineral density in grams per cm<sup>2</sup> was derived based on  
429 the assumption that sound waves travel differently through denser bones.

430

### 431 **Cardiovascular measures**

432 Blood pressure, pulse rate and arterial stiffness (pulse wave velocity) were measured by trained nurses during a separate  
433 stage of the baseline assessment.

434

### 435 **Blood pressure**

436 Seated systolic and diastolic resting blood pressure in millimetres of mercury (mmHg) was measured twice using an  
437 Omron 705 IT digital blood pressure monitor (measurement range 0-255 mmHg) following standard procedures.

438 Participants were asked to loosen or remove any restrictive clothing and put their arm on a desktop. Measurements were  
439 taken from the left upper arm or, if impractical, from the right arm. If there was a problem with the measurement, it was  
440 repeated. Three different cuff sizes were available and a Seca tape was used to determine the circumference of the  
441 midpoint of the upper arm. If the largest cuff size was too small or the digital blood pressure monitor did not produce a  
442 reading, a manual sphygmomanometer was used in conjunction with a stethoscope. A second measurement was taken at  
443 least one minute after the first measure, following the same procedure. We used an average of the two readings to reduce  
444 potential measurement error.

445

### 446 **Pulse rate**

447 Resting pulse rate in beats per minute was recorded during the blood pressure measurements using the Omron 705 IT  
448 device or, exceptionally, a manual sphygmomanometer. We used an average of the two readings to reduce potential  
449 measurement error.

450

### 451 **Arterial stiffness**

452 Arterial stiffness (vascular reactivity) is an independent predictor of cardiovascular risk and mortality that can be  
453 measured rapidly, inexpensively, and without special training [113]. Resting pulse wave velocity was measured non-  
454 invasively using finger photoplethysmography with a PulseTrace PCA2 infra-red sensor. The pulse waveform was  
455 recorded over a period of 10-15 seconds, with the sensor clipped to the end of the index finger of the non-dominant hand  
456 while the participant was sitting. If the waveform did not fill at least 2/3 of the display of the device, or did not stabilize  
457 within one minute, the measurement was repeated on a larger finger or on the thumb. The shape of the waveform is  
458 directly related to the time it takes for the pulse wave to travel through the arterial tree in the lower body and to be  
459 reflected back to the finger. The time between the peaks of the waveform (the pulse wave peak-to-peak time, i.e., the

460 difference between the peak values of direct and reflected components) was divided by the participant's height to obtain  
461 the arterial stiffness index in metres per second. A higher score on the index represents stiffer arteries. The method has  
462 been externally validated and is highly correlated with the gold-standard carotid-femoral pulse wave velocity [114].

463

#### 464 **Exclusion criteria**

465 Participants for whom their genetic sex, inferred from the relative intensity of biological markers on the Y and X  
466 chromosomes, and self-reported sex did not match were excluded. Participants with missing data for any covariates or  
467 who responded "do not know" or "prefer not to answer" were excluded from all analyses.

468

#### 469 **Covariates**

470 Covariates included ethnicity (White, Asian, Black, Chinese, Mixed-race or other), gross annual household income  
471 (<£18,000, £18,000–30,999, £31,000–51,999, £52,000–100,000 or >£100,000), physical activity (number of days per  
472 week spent walking, engaging in moderate-intensity physical activity and engaging in vigorous-intensity physical  
473 activity for  $\geq 10$  minutes continuously), smoking status (never, former or current), alcohol intake frequency (never,  
474 special occasions only, one to three times a month, once or twice a week, three or four times a week or daily or almost  
475 daily), sleep duration (hours per day) and, for cardiovascular measures, current use of antihypertensive medications at  
476 baseline (yes/no). See our previous publication [105] for further details on these variables.

477

#### 478 **Statistical analyses**

479 All analyses, except the sensitivity analyses, were pre-specified prior to inspection of the data (preregistration:  
480 [osf.io/pc76g](https://osf.io/pc76g)) and algorithms were tested on simulated data. Statistical analyses were conducted using R (version 3.6.0).

481

482 Sample characteristics were summarised using means and standard deviations or counts and percentages. We also present  
483 the number of individuals who met our criteria for lifetime depression and healthy control. Kernel density plots of all  
484 physiological measures stratified by lifetime depression case status are presented separately for males and females.

485 Differences between the sexes and between cases and controls were estimated using standardised mean differences ( $\pm$   
486 95% confidence intervals).

487

488 We examined the relationship between each physiological measure and age using generalised additive models (GAMs)  
489 with the 'mgcv' package in R [115]. Each measure was modelled against a penalised regression spline function of age  
490 with separate smooths for individuals with lifetime depression and healthy controls. GAMs are flexible modelling  
491 approaches that allow for the relationship between an outcome variable and a continuous exposure to be represented by a

492 non-linear smooth curve while controlling for covariates. They are particularly useful if a linear model does not capture  
493 key aspects of the relationship between variables and attempt to achieve maximum goodness-of-fit while maintaining  
494 parsimony of the fitted curve to minimize overfitting. Smoothing parameters were selected using the restricted maximum  
495 likelihood method and we used the default option of ten basis functions to represent smooth terms in each model.

496

497 Two models were fit for males and females separately:

498

- 499 • Unadjusted model: physiological measure ~ lifetime depression + s(age, by lifetime depression).
- 500 • Adjusted model: physiological measure ~ lifetime depression + s(age, by lifetime depression) + covariates  
501 (ethnicity, household income, physical activity, smoking status, alcohol intake frequency and sleep duration).

502

503 where s(age, by lifetime depression) represents the smooth function for age, stratified by lifetime depression status. For  
504 cardiovascular measures, the adjusted model additionally included self-reported antihypertensive medication use.

505

506 To formally test whether the relationship between physiological measures and age differed between individuals with  
507 lifetime depression and healthy controls, we also fit, in separate analyses, models that included a reference smooth for  
508 healthy controls and a difference smooth for individuals with lifetime depression compared to healthy controls. For these  
509 analyses, lifetime depression status was coded as an ordered factor in R. If the difference smooth differs from zero,  
510 individuals with lifetime depression and healthy controls follow a different trend with age.

511

512 Adjusted *p*-values were calculated using the `p.adjust()` command in R to account for multiple testing across each set of  
513 analyses of the 15 physiological measures. Two methods were used: (1) Bonferroni and (2) Benjamini & Hochberg  
514 [116], all two-tailed, with  $\alpha = .05$  and false discovery rate of 5%, respectively.

515

516 To address potential concerns about the validity and reliability of our case definition, we conducted sensitivity analyses  
517 in which we restricted the samples to (i) individuals with lifetime depression according to at least two different data  
518 sources and (ii) individuals with lifetime depression status assessed using the CIDI-SF. These sensitivity analyses were  
519 not preregistered.

520 **Authorship contributions**

521 CML acquired the studentship funding, interpreted the findings and critically reviewed the manuscript. JM conceived the  
522 idea of the study, acquired the data, carried out the statistical analysis, interpreted the findings, wrote the manuscript and  
523 revised the manuscript for final submission. Both authors read and approved the final manuscript. JM had full access to  
524 all data used in this study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

526 **Acknowledgments**

527 This research has been conducted using data from UK Biobank, a major biomedical database. This project made use of  
528 time on Rosalind HPC, funded by Guy's & St Thomas' Hospital NHS Trust Biomedical Research Centre (GSTT-BRC),  
529 South London & Maudsley NHS Trust Biomedical Research Centre (SLAM-BRC), and Faculty of Natural Mathematics  
530 & Science (NMS) at King's College London.

532 **Conflicts of Interest and Funding**

533 JM receives studentship funding from the Biotechnology and Biological Sciences Research Council (BBSRC) (ref:  
534 2050702) and Eli Lilly and Company Limited. CML is part-funded by the National Institute for Health Research (NIHR)  
535 Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College  
536 London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department  
537 of Health and Social Care. CML is a member of the Scientific Advisory Board of Myriad Neuroscience.

539 **Ethics**

540 Ethical approval for the UK Biobank study has been granted by the National Information Governance Board for Health  
541 and Social Care and the NHS North West Multicentre Research Ethics Committee (11/NW/0382). No project-specific  
542 ethical approval is needed. Data access permission has been granted under UK Biobank application 45514.

544 **Data sharing statement**

545 The data used are available to all bona fide researchers for health-related research that is in the public interest, subject to  
546 an application process and approval criteria. Study materials are publicly available online at <http://www.ukbiobank.ac.uk>.

548 **Supplementary material**

549 Supplementary information is available online.

## 550 **References**

- 551 1. Plana-Ripoll O, Weyer N, Momen NC, Christensen MK, Iburg KM, Laursen TM, McGrath JJ. Changes over  
552 time in the differential mortality gap in individuals with mental disorders. *JAMA Psychiatry*. 2020; 77: 648-50.
- 553 2. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ, Walters EE, Wang PS. The  
554 epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*.  
555 2003; 289: 3095-105.
- 556 3. Daskalopoulou C, Koukounari A, Wu Y-T, Terrera GM, Caballero FF, de la Fuente J, Tyrovolas S,  
557 Panagiotakos DB, Prince M, Prina M. Healthy ageing trajectories and lifestyle behaviour: the Mexican Health and Aging  
558 Study. *Scientific Reports*. 2019; 9: 1-10.
- 559 4. Wilkinson P, Ruane C, Tempest K. Depression in older adults. *The BMJ*. 2018; 363: k4922.
- 560 5. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013; 153: 1194-  
561 217.
- 562 6. Cole JH, Franke K. Predicting age using neuroimaging: innovative brain ageing biomarkers. *Trends in*  
563 *Neurosciences*. 2017; 40: 681-90.
- 564 7. Veronese N, Stubbs B, Fontana L, Trevisan C, Bolzetta F, Rui MD, Sartori L, Musacchio E, Zambon S, Maggi  
565 S, Perissinotto E, Corti MC, Crepaldi G, et al. A comparison of objective physical performance tests and future  
566 mortality in the elderly people. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*. 2017; 72:  
567 362-8.
- 568 8. Celis-Morales CA, Welsh P, Lyall DM, Steell L, Petermann F, Anderson J, Iliodromiti S, Sillars A, Graham N,  
569 Mackay DF, Pell JP, Gill, JMR, Sattar N, et al. Associations of grip strength with cardiovascular, respiratory, and cancer  
570 outcomes and all cause mortality: prospective cohort study of half a million UK Biobank participants. *The BMJ*. 2018;  
571 361: k1651.
- 572 9. Ho FK, Celis-Morales CA, Petermann-Rocha F, Sillars A, Welsh P, Welsh C, Anderson J, Lyall DM, Mackay  
573 DF, Sattar N, Gill, JMR, Pell JP, Gray SR. The association of grip strength with health outcomes does not differ if grip  
574 strength is used in absolute or relative terms: a prospective cohort study. *Age and Ageing*. 2019; 48: 684-91.
- 575 10. Dodds RM, Syddall HE, Cooper R, Benzeval M, Deary IJ, Dennison EM, Der G, Gale CR, Inskip HM, Jagger  
576 C, Kirkwood TB, Lawlor DA, Robinson SM, et al. Grip strength across the life course: normative data from twelve  
577 British studies. *PloS One*. 2014; 9: e113637.
- 578 11. Cheng S, Xanthakis V, Sullivan LM, Vasan RS. Blood pressure tracking over the adult life course: patterns and  
579 correlates in the Framingham heart study. *Hypertension*. 2012; 60: 1393-9.

- 580 12. Ji H, Kim A, Ebinger JE, Niiranen TJ, Claggett BL, Merz CNB, Cheng S. Sex differences in blood pressure  
581 trajectories over the life course. *JAMA Cardiology*. 2020; 5: 19-26.
- 582 13. Wolkowitz OM. Accelerated biological aging in serious mental disorders. *World Psychiatry*. 2018; 17: 144-5.
- 583 14. Austad SN, Bartke A. Sex differences in longevity and in responses to anti-aging interventions: a mini-review.  
584 *Gerontology*. 2016; 62: 40-6.
- 585 15. McCarthy MM, Nugent BM, Lenz KM. Neuroimmunology and neuroepigenetics in the establishment of sex  
586 differences in the brain. *Nature Reviews Neuroscience*. 2017; 18: 471-84.
- 587 16. Mutz J, Young AH, Lewis CM. Age-related changes in physiology in individuals with bipolar disorder.  
588 medRxiv. 2021.
- 589 17. Cooper R, Hardy R, Sayer AA, Ben-Shlomo Y, Birnie K, Cooper C, Craig L, Deary IJ, Demakakos P, Gallacher  
590 J, McNeill G, Martin RM, Starr JM, et al. Age and gender differences in physical capability levels from mid-life  
591 onwards: the harmonisation and meta-analysis of data from eight UK cohort studies. *PloS One*. 2011; 6: e27899.
- 592 18. Keevil VL, Hayat S, Dalzell N, Moore S, Bhaniani A, Luben R, Wareham NJ, Khaw K-T. The physical  
593 capability of community-based men and women from a British cohort: the European Prospective Investigation into  
594 Cancer (EPIC)-Norfolk study. *BMC Geriatrics*. 2013; 13: 93.
- 595 19. Beller J, Miething A, Regidor E, Lostao L, Epping J, Geyer S. Trends in grip strength: Age, period, and cohort  
596 effects on grip strength in older adults from Germany, Sweden, and Spain. *SSM-Population Health*. 2019; 9: 100456.
- 597 20. Burt VL, Whelton P, Roccella EJ, Brown C, Cutler JA, Higgins M, Horan MJ, Labarthe D. Prevalence of  
598 hypertension in the US adult population: results from the Third National Health and Nutrition Examination Survey,  
599 1988-1991. *Hypertension*. 1995; 25: 305-13.
- 600 21. AlGhatrif M, Strait JB, Morrell CH, Canepa M, Wright J, Elango P, Scuteri A, Najjar SS, Ferrucci L, Lakatta  
601 EG. Longitudinal trajectories of arterial stiffness and the role of blood pressure: the Baltimore Longitudinal Study of  
602 Aging. *Hypertension*. 2013; 62: 934-41.
- 603 22. Quer G, Gouda P, Galarnyk M, Topol EJ, Steinhubl SR. Inter-and intraindividual variability in daily resting  
604 heart rate and its associations with age, sex, sleep, BMI, and time of year: retrospective, longitudinal cohort study of  
605 92,457 adults. *Plos One*. 2020; 15: e0227709.
- 606 23. Liao SY, Lin X, Christiani DC. Occupational exposures and longitudinal lung function decline. *American*  
607 *Journal of Industrial Medicine*. 2015; 58: 14-20.
- 608 24. Nunn A, Gregg I. New regression equations for predicting peak expiratory flow in adults. *The BMJ*. 1989; 298:  
609 1068-70.

- 610 25. van Oostrom SH, Engelfriet PM, Verschuren WM, Schipper M, Wouters IM, Boezen M, Smit HA, Kerstjens  
611 HA, Picavet HSJ. Aging-related trajectories of lung function in the general population—The Doetinchem Cohort Study.  
612 PloS One. 2018; 13: e0197250.
- 613 26. Li C, Ford ES, Zhao G, Balluz LS, Giles WH. Estimates of body composition with dual-energy X-ray  
614 absorptiometry in adults. The American Journal of Clinical Nutrition. 2009; 90: 1457-65.
- 615 27. He X, Li Z, Tang X, Zhang L, Wang L, He Y, Jin T, Yuan D. Age-and sex-related differences in body  
616 composition in healthy subjects aged 18 to 82 years. Medicine. 2018; 97: e11152.
- 617 28. Frenzel A, Binder H, Walter N, Wirkner K, Loeffler M, Loeffler-Wirth H. The aging human body shape. NPJ  
618 Aging and Mechanisms of Disease. 2020; 6: 1-15.
- 619 29. Larsson I, Lissner L, Samuelson G, Fors H, Lantz H, Näslund I, Carlsson L, Sjöström L, Bosaeus I. Body  
620 composition through adult life: Swedish reference data on body composition. European Journal of Clinical Nutrition.  
621 2015; 69: 837-42.
- 622 30. Malkin I, Bigman G, Matias R, Kalichman L, Seibel M, Kobylansky E, Livshits G. Age-related changes of  
623 bone strength phenotypes: observational follow-up study of hand bone mineral density. Archives of Osteoporosis. 2006;  
624 1: 59-68.
- 625 31. Ross RD, Shah RC, Leurgans SE, Buchman AS, Bennett DA. Association of heel bone mineral density with  
626 incident disability and mortality in community-dwelling older adults. JBMR Plus. 2020; 4: e10390.
- 627 32. Merz AA, Cheng S. Sex differences in cardiovascular ageing. Heart. 2016; 102: 825-31.
- 628 33. Cecelja M, Chowienczyk P. Dissociation of aortic pulse wave velocity with risk factors for cardiovascular  
629 disease other than hypertension: a systematic review. Hypertension. 2009; 54: 1328-36.
- 630 34. Mitchell GF, Parise H, Benjamin EJ, Larson MG, Keyes MJ, Vita JA, Vasan RS, Levy D. Changes in arterial  
631 stiffness and wave reflection with advancing age in healthy men and women: the Framingham Heart Study.  
632 Hypertension. 2004; 43: 1239-45.
- 633 35. Russo C, Jin Z, Palmieri V, Homma S, Rundek T, Elkind MS, Sacco RL, Di Tullio MR. Arterial stiffness and  
634 wave reflection: sex differences and relationship with left ventricular diastolic function. Hypertension. 2012; 60: 362-8.
- 635 36. Coutinho T, Borlaug BA, Pellikka PA, Turner ST, Kullo IJ. Sex differences in arterial stiffness and ventricular-  
636 arterial interactions. Journal of the American College of Cardiology. 2013; 61: 96-103.
- 637 37. Franklin SS, Gustin IV W, Wong ND, Larson MG, Weber MA, Kannel WB, Levy D. Hemodynamic patterns of  
638 age-related changes in blood pressure: the Framingham Heart Study. Circulation. 1997; 96: 308-15.
- 639 38. Wolf-Maier K, Cooper RS, Banegas JR, Giampaoli S, Hense H-W, Joffres M, Kastarinen M, Poulter N,  
640 Primatesta P, Rodríguez-Artalejo F. Hypertension prevalence and blood pressure levels in 6 European countries, Canada,  
641 and the United States. JAMA. 2003; 289: 2363-9.

- 642 39. Kyle UG, Genton L, Slosman DO, Pichard C. Fat-free and fat mass percentiles in 5225 healthy subjects aged 15  
643 to 98 years. *Nutrition*. 2001; 17: 534-41.
- 644 40. Jackson AS, Janssen I, Sui X, Church TS, Blair SN. Longitudinal changes in body composition associated with  
645 healthy ageing: men, aged 20–96 years. *British Journal of Nutrition*. 2012; 107: 1085-91.
- 646 41. Thomas ET, Guppy M, Straus SE, Bell KJ, Glasziou P. Rate of normal lung function decline in ageing adults: a  
647 systematic review of prospective cohort studies. *BMJ open*. 2019; 9: e028150.
- 648 42. Malkin I, Karasik D, Livshits G, Kobylansky E. Modelling of age-related bone loss using cross-sectional data.  
649 *Annals of Human Biology*. 2002; 29: 256-70.
- 650 43. Qian X, Su C, Zhang B, Qin G, Wang H, Wu Z. Changes in distributions of waist circumference, waist-to-hip  
651 ratio and waist-to-height ratio over an 18-year period among Chinese adults: a longitudinal study using quantile  
652 regression. *BMC Public Health*. 2019; 19: 700.
- 653 44. McDowell CP, Gordon BR, Herring MP. Sex-related differences in the association between grip strength and  
654 depression: results from the Irish Longitudinal Study on Ageing. *Experimental Gerontology*. 2018; 104: 147-52.
- 655 45. Ashdown-Franks G, Stubbs B, Koyanagi A, Schuch F, Firth J, Veronese N, Vancampfort D. Handgrip strength  
656 and depression among 34,129 adults aged 50 years and older in six low-and middle-income countries. *Journal of*  
657 *Affective Disorders*. 2019; 243: 448-54.
- 658 46. Han K-M, Chang J, Yoon H-K, Ko Y-H, Ham B-J, Kim Y-K, Han C. Relationships between hand-grip strength,  
659 socioeconomic status, and depressive symptoms in community-dwelling older adults. *Journal of Affective Disorders*.  
660 2019; 252: 263-70.
- 661 47. Volaklis K, Mamadjanov T, Meisinger C, Linseisen J. Association between muscular strength and depressive  
662 symptoms. *Wiener Klinische Wochenschrift*. 2019; 131: 255-64.
- 663 48. Veronese N, Stubbs B, Trevisan C, Bolzetta F, De Rui M, Solmi M, Sartori L, Musacchio E, Zambon S,  
664 Perissinotto E, Baggio G, Crepaldi G, Manzato E, et al. Poor physical performance predicts future onset of depression in  
665 elderly people: Progetto Veneto Anziani Longitudinal Study. *Physical Therapy*. 2017; 97: 659-68.
- 666 49. Smith L, White S, Stubbs B, Hu L, Veronese N, Vancampfort D, Hamer M, Gardner B, Yang L. Depressive  
667 symptoms, handgrip strength, and weight status in US older adults. *Journal of Affective Disorders*. 2018; 238: 305-10.
- 668 50. Suija K, Timonen M, Suviola M, Jokelainen J, Järvelin M-R, Tammelin T. The association between physical  
669 fitness and depressive symptoms among young adults: results of the Northern Finland 1966 birth cohort study. *BMC*  
670 *Public Health*. 2013; 13: 535.
- 671 51. Hildrum B, Mykletun A, Holmen J, Dahl AA. Effect of anxiety and depression on blood pressure: 11-year  
672 longitudinal population study. *The British Journal of Psychiatry*. 2008; 193: 108-13.

- 673 52. Hildrum B, Mykletun A, Stordal E, Bjelland I, Dahl AA, Holmen J. Association of low blood pressure with  
674 anxiety and depression: the Nord-Trøndelag Health Study. *Journal of Epidemiology & Community Health*. 2007; 61: 53-  
675 8.
- 676 53. Tikhonoff V, Hardy R, Deanfield J, Friberg P, Kuh D, Muniz G, Pariente CM, Hotopf M, Richards M.  
677 Symptoms of anxiety and depression across adulthood and blood pressure in late middle age: the 1946 British birth  
678 cohort. *Journal of Hypertension*. 2014; 32: 1590-99.
- 679 54. Licht CM, De Geus EJ, Seldenrijk A, Van Hout HP, Zitman FG, Van Dyck R, Penninx BW. Depression is  
680 associated with decreased blood pressure, but antidepressant use increases the risk for hypertension. *Hypertension*. 2009;  
681 53: 631-8.
- 682 55. Shinn EH, Poston WSC, Kimball KT, St. Jeor ST, Foreyt JP. Blood pressure and symptoms of depression and  
683 anxiety: a prospective study. *American Journal of Hypertension*. 2001; 14: 660-4.
- 684 56. Meng L, Chen D, Yang Y, Zheng Y, Hui R. Depression increases the risk of hypertension incidence: a meta-  
685 analysis of prospective cohort studies. *Journal of Hypertension*. 2012; 30: 842-51.
- 686 57. Paterniti S, Verdier-Taillefer M-H, Geneste C, Bisslerbe J-C, Alperovitch A. Low blood pressure and risk of  
687 depression in the elderly: a prospective community-based study. *The British Journal of Psychiatry*. 2000; 176: 464-7.
- 688 58. Gould CE, Beaudreau SA. Association between depression and anxiety on blood pressure dysregulation and  
689 pulse in the Health and Retirement Study. *International Journal of Geriatric Psychiatry*. 2013; 28: 1045-53.
- 690 59. Lechin F, van der Dijs B, Orozco B, Lechin ME, Báez S, Lechin AE, Rada I, Acosta E, Arocha L, Jiménez V.  
691 Plasma neurotransmitters, blood pressure, and heart rate during supine-resting, orthostasis, and moderate exercise  
692 conditions in major depressed patients. *Biological Psychiatry*. 1995; 38: 166-73.
- 693 60. Kemp AH, Quintana DS, Quinn CR, Hopkinson P, Harris AW. Major depressive disorder with melancholia  
694 displays robust alterations in resting state heart rate and its variability: implications for future morbidity and mortality.  
695 *Frontiers in Psychology*. 2014; 5: 1387.
- 696 61. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, Zitman FG. Overweight, obesity, and  
697 depression: a systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry*. 2010; 67:  
698 220-9.
- 699 62. Geoffroy M-C, Li L, Power C. Depressive symptoms and body mass index: co-morbidity and direction of  
700 association in a British birth cohort followed over 50 years. *Psychological Medicine*. 2014; 44: 2641-52.
- 701 63. Rivenes AC, Harvey SB, Mykletun A. The relationship between abdominal fat, obesity, and common mental  
702 disorders: results from the HUNT study. *Journal of Psychosomatic Research*. 2009; 66: 269-75.
- 703 64. Ho RC, Niti M, Kua EH, Ng TP. Body mass index, waist circumference, waist-hip ratio and depressive  
704 symptoms in Chinese elderly: a population-based study. *International Journal of Geriatric Psychiatry*. 2008; 23: 401-8.

- 705 65. Zhi T, Wang Q, Liu Z, Zhu Y, Wang Y, Shi R, Wang Z, Chu X, Wang X, Jiang X. Body mass index, waist  
706 circumference and waist-hip ratio are associated with depressive symptoms in older Chinese women: results from the  
707 Rugao Longevity and Ageing Study (RuLAS). *Aging & Mental Health*. 2017; 21: 518-23.
- 708 66. De Wit LM, Van Straten A, Van Hertem M, Penninx BW, Cuijpers P. Depression and body mass index, a u-  
709 shaped association. *BMC Public Health*. 2009; 9: 14.
- 710 67. Gale C, Sayer AA, Cooper C, Dennison E, Starr J, Whalley L, Gallacher JE, Ben-Shlomo Y, Kuh D, Hardy R,  
711 Craig L, Deary IJ. Factors associated with symptoms of anxiety and depression in five cohorts of community-based older  
712 people: the HALCyon (Healthy Ageing across the Life Course) Programme. *Psychological Medicine*. 2011; 41: 2057-73.
- 713 68. Speed MS, Jepsen OH, Børghlum AD, Speed D, Østergaard SD. Investigating the association between body fat  
714 and depression via Mendelian randomization. *Translational Psychiatry*. 2019; 9: 1-9.
- 715 69. Baldofski S, Mauche N, Dogan-Sander E, Bot M, Brouwer I, Paans N, Cabout M, Gili M, van Grootheest G,  
716 Hegerl U, Owens M, Roca M, Visser M, et al. Depressive symptom clusters in relation to body weight status: results  
717 from two large European multi-center studies. *Frontiers in Psychiatry*. 2019; 10: 858.
- 718 70. Ul-Haq Z, Smith DJ, Nicholl BI, Cullen B, Martin D, Gill JM, Evans J, Roberts B, Deary IJ, Gallacher J,  
719 Hotopf M, Craddock N, Mackay DF, et al. Gender differences in the association between adiposity and probable major  
720 depression: a cross-sectional study of 140,564 UK Biobank participants. *BMC Psychiatry*. 2014; 14: 153.
- 721 71. Wiltink J, Michal M, Wild PS, Zwiener I, Blettner M, Münzel T, Schulz A, Kirschner Y, Beutel ME.  
722 Associations between depression and different measures of obesity (BMI, WC, WHtR, WHR). *BMC Psychiatry*. 2013;  
723 13: 223.
- 724 72. Mezuk B, Golden SH, Eaton WW, Lee HB. Depression and body composition among older adults. *Aging &*  
725 *Mental Health*. 2012; 16: 167-72.
- 726 73. Giovannini S, Macchi C, Liperoti R, Laudisio A, Coraci D, Loreti C, Vannetti F, Onder G, Padua L, Bonaccorsi  
727 G. Association of body fat with health-related quality of life and depression in nonagenarians: the Mugello Study.  
728 *Journal of the American Medical Directors Association*. 2019; 20: 564-8.
- 729 74. van Milligen BA, Lamers F, Guus T, Smit JH, Penninx BW. Objective physical functioning in patients with  
730 depressive and/or anxiety disorders. *Journal of Affective Disorders*. 2011; 131: 193-9.
- 731 75. Spitzer C, Gläser S, Grabe HJ, Ewert R, Barnow S, Felix SB, Freyberger HJ, Völzke H, Koch B, Schäper C.  
732 Mental health problems, obstructive lung disease and lung function: findings from the general population. *Journal of*  
733 *Psychosomatic Research*. 2011; 71: 174-9.
- 734 76. Park Y, Jung JY, Kim YS, Chung KS, Song JH, Kim SY, Kim EY, Kang YA, Park MS, Chang J, Leem AY.  
735 Relationship between depression and lung function in the general population in Korea: a retrospective cross-sectional  
736 study. *International Journal of Chronic Obstructive Pulmonary Disease*. 2018; 13: 2207-13.

- 737 77. Giltay EJ, Nissinen A, Giampaoli S, Zitman FG, Kromhout D. Low respiratory function increases the risk of  
738 depressive symptoms in later life in men. *Psychosomatic Medicine*. 2010; 72: 53-60.
- 739 78. Rosenblat J, Gregory J, Carvalho A, McIntyre R. Depression and disturbed bone metabolism: a narrative review  
740 of the epidemiological findings and postulated mechanisms. *Current Molecular Medicine*. 2016; 16: 165-78.
- 741 79. Wu Q, Magnus J, Liu J, Bencaz A, Hentz J. Depression and low bone mineral density: a meta-analysis of  
742 epidemiologic studies. *Osteoporosis International*. 2009; 20: 1309-20.
- 743 80. Williams LJ, Pasco JA, Jacka FN, Hodge JM, Kotowicz MA, Berk M. Quantitative Heel Ultrasound (QUS)  
744 measures of bone quality in association with mood and anxiety disorders. *Journal of Affective Disorders*. 2013; 146:  
745 395-400.
- 746 81. Atteritano M, Lasco A, Mazzaferro S, Macri I, Catalano A, Santangelo A, Bagnato G, Bagnato G, Frisina N.  
747 Bone mineral density, quantitative ultrasound parameters and bone metabolism in postmenopausal women with  
748 depression. *Internal and Emergency Medicine*. 2013; 8: 485-91.
- 749 82. Peng L, Bi S, Liu X, Long T, Zhao Y, Li F, Yang T, Zhang C. Association between depressive symptoms and  
750 arterial stiffness: a cross-sectional study in the general Chinese population. *BMJ Open*. 2020; 10: e033408.
- 751 83. Seldenrijk A, van Hout HP, van Marwijk HW, de Groot E, Gort J, Rustemeijer C, Diamant M, Penninx BW.  
752 Depression, anxiety, and arterial stiffness. *Biological Psychiatry*. 2011; 69: 795-803.
- 753 84. Dregan A, Rayner L, Davis KA, Bakolis I, de la Torre JA, Das-Munshi J, Hatch SL, Stewart R, Hotopf M.  
754 Associations between depression, arterial stiffness, and metabolic syndrome among adults in the UK Biobank Population  
755 Study: a mediation analysis. *JAMA Psychiatry*. 2020; 77: 598-606.
- 756 85. Lever-van Milligen BA, Lamers F, Smit JH, Penninx BW. Six-year trajectory of objective physical function in  
757 persons with depressive and anxiety disorders. *Depression and Anxiety*. 2017; 34: 188-97.
- 758 86. Shah MT, Zonderman AB, Waldstein SR. Sex and age differences in the relation of depressive symptoms with  
759 blood pressure. *American Journal of Hypertension*. 2013; 26: 1413-20.
- 760 87. Erez HB, Weller A, Vaisman N, Kreitler S. The relationship of depression, anxiety and stress with low bone  
761 mineral density in post-menopausal women. *Archives of Osteoporosis*. 2012; 7: 247-55.
- 762 88. Yirmiya R, Bab I. Major depression is a risk factor for low bone mineral density: a meta-analysis. *Biological*  
763 *Psychiatry*. 2009; 66: 423-32.
- 764 89. Diem SJ, Blackwell TL, Stone KL, Yaffe K, Cauley JA, Whooley MA, Ensrud KE. Depressive symptoms and  
765 rates of bone loss at the hip in older women. *Journal of the American Geriatrics Society*. 2007; 55: 824-31.
- 766 90. Diem S, Harrison S, Haney E, Cauley J, Stone K, Orwoll E, Ensrud K. Depressive symptoms and rates of bone  
767 loss at the hip in older men. *Osteoporosis International*. 2013; 24: 111-9.

- 768 91. Cizza G, Mistry S, Nguyen VT, Eskandari F, Martinez P, Torvik S, Reynolds JC, Gold PW, Sinai N, Csako G.  
769 Do premenopausal women with major depression have low bone mineral density? A 36-month prospective study. *PLoS*  
770 *One*. 2012; 7: e40894.
- 771 92. Mezuk B, Eaton WW, Golden SH, Wand G, Lee HB. Depression, antidepressants, and bone mineral density in a  
772 population-based cohort. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2008; 63:  
773 1410-5.
- 774 93. Pinto E. Blood pressure and ageing. *Postgraduate Medical Journal*. 2007; 83: 109-14.
- 775 94. Hübel C, Gaspar HA, Coleman JR, Hanscombe KB, Purves K, Prokopenko I, Graff M, Ngwa JS, Workalemahu  
776 T, O'Reilly PF, Bulik CM, Breen G. Genetic correlations of psychiatric traits with body composition and glycemic traits  
777 are sex- and age-dependent. *Nature Communications*. 2019; 10: 1-12.
- 778 95. Bäckman K, Joas E, Falk H, Mitnitski A, Rockwood K, Skoog I. Changes in the lethality of frailty over 30  
779 years: evidence from two cohorts of 70-year-olds in Gothenburg Sweden. *The Journals of Gerontology: Series A*. 2017;  
780 72: 945-50.
- 781 96. Yu R, Wong M, Chong K, Chang B, Lum C, Auyeung T, Lee J, Lee R, Woo J. Trajectories of frailty among  
782 Chinese older people in Hong Kong between 2001 and 2012: an age-period-cohort analysis. *Age and Ageing*. 2018; 47:  
783 254-61.
- 784 97. Yang Y, Lee LC. Dynamics and heterogeneity in the process of human frailty and aging: evidence from the US  
785 older adult population. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2010; 65: 246-55.
- 786 98. Glanville KP, Coleman JR, Howard DM, Pain O, Hanscombe KB, Jermy B, Arathimos R, Hübel C, Breen G,  
787 O'Reilly PF, Lewis CM. Multiple measures of depression to enhance validity of Major Depressive Disorder in the UK  
788 Biobank. *BJPsych Open*. 2021; 7: e44.
- 789 99. Davis K, Hotopf M. Mental health phenotyping in UK Biobank. *Progress in Neurology and Psychiatry*. 2019;  
790 23: 4-7.
- 791 100. Davis KA, Cullen B, Adams M, Brailean A, Breen G, Coleman JR, Dregan A, Gaspar HA, Hübel C, Lee W,  
792 McIntosh AM, Nolan J, Pearsall R, et al. Indicators of mental disorders in UK Biobank—A comparison of approaches.  
793 *International Journal of Methods in Psychiatric Research*. 2019; 28: e1796.
- 794 101. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, Collins R, Allen NE. Comparison of  
795 sociodemographic and health-related characteristics of UK Biobank participants with those of the general population.  
796 *American Journal of Epidemiology*. 2017; 186: 1026-34.
- 797 102. Batty GD, Gale CR, Kivimäki M, Deary IJ, Belle S. Comparison of risk factor associations in UK Biobank  
798 against representative, general population based studies with conventional response rates: prospective cohort study and  
799 individual participant meta-analysis. *The BMJ*. 2020; 368: m131.

- 800 103. Knudsen AK, Hotopf M, Skogen JC, Øverland S, Mykletun A. The health status of nonparticipants in a  
801 population-based health study: the Hordaland Health Study. *American Journal of Epidemiology*. 2010; 172: 1306-14.
- 802 104. Ansai JH, Rebelatto JR. Effect of two physical exercise protocols on cognition and depressive symptoms in  
803 oldest-old people: a randomized controlled trial. *Geriatrics & Gerontology International*. 2015; 15: 1127-34.
- 804 105. Mutz J, Roscoe CJ, Lewis CM. Exploring health in the UK Biobank: associations with sociodemographic  
805 characteristics, psychosocial factors, lifestyle and environmental exposures. *medRxiv*. 2021.
- 806 106. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic D, Delaneau O,  
807 O'Connell J, Cortes A, Welsh S, Young A, et al. The UK Biobank resource with deep phenotyping and genomic data.  
808 *Nature*. 2018; 562: 203-9.
- 809 107. Davis KA, Coleman JR, Adams M, Allen N, Breen G, Cullen B, Dickens C, Fox E, Graham N, Holliday J,  
810 Howard LM, John A, Lee W, et al. Mental health in UK Biobank—development, implementation and results from an  
811 online questionnaire completed by 157 366 participants: a reanalysis. *BJPsych Open*. 2020; 6: E18.
- 812 108. Fabbri C, Hagenaars SP, John C, Williams AT, Shrine N, Moles L, Hanscombe KB, Serretti A, Shepherd DJ,  
813 Free RS, Wain LV, Tobin MD, Lewis CM. Genetic and clinical characteristics of treatment-resistant depression using  
814 primary care records in two UK cohorts. *Molecular Psychiatry*. 2021.
- 815 109. Smith DJ, Nicholl BI, Cullen B, Martin D, Ul-Haq Z, Evans J, Gill JM, Roberts B, Gallacher J, Mackay D,  
816 Hotopf M, Deary I, Craddock N, et al. Prevalence and characteristics of probable major depression and bipolar disorder  
817 within UK Biobank: cross-sectional study of 172,751 participants. *PloS One*. 2013; 8: e75362.
- 818 110. Firth JA, Smith L, Sarris J, Vancampfort D, Schuch F, Carvalho AF, Solmi M, Yung AR, Stubbs B, Firth J.  
819 Handgrip strength is associated with hippocampal volume and white matter hyperintensities in major depression and  
820 healthy controls: a UK Biobank study. *Psychosomatic Medicine*. 2020; 82: 39-46.
- 821 111. Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older  
822 adults and its association with multimorbidity and mortality: a prospective analysis of 493 737 UK Biobank participants.  
823 *The Lancet Public Health*. 2018; 3: e323-e32.
- 824 112. Doiron D, de Hoogh K, Probst-Hensch N, Fortier I, Cai Y, De Matteis S, Hansell AL. Air pollution, lung  
825 function and COPD: results from the population-based UK Biobank study. *European Respiratory Journal*. 2019; 54:  
826 1802140.
- 827 113. Said MA, Eppinga RN, Lipsic E, Verweij N, van der Harst P. Relationship of arterial stiffness index and pulse  
828 pressure with cardiovascular disease and mortality. *Journal of the American Heart Association*. 2018; 7: e007621.
- 829 114. Millasseau SC, Kelly R, Ritter J, Chowienczyk P. Determination of age-related increases in large artery  
830 stiffness by digital pulse contour analysis. *Clinical Science*. 2002; 103: 371-7.
- 831 115. Wood SN. Generalized additive models: an introduction with R. 2017.

- 832 116. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple  
833 testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. 1995; 57: 289-300.

## 834 **Figure legends**

835

836 **Figure 1.** Flowchart of study population. The main dataset included hand-grip strength, blood pressure, pulse rate and  
837 measures of body composition.

838

839 **Figure 2.** Generalised additive models of age-related changes in physiological measures in females with lifetime  
840 depression and healthy controls. The solid lines represent physiological measures against smoothing functions of age.  
841 The shaded areas correspond to approximate 95% confidence intervals ( $\pm 2 \times$  standard error). FEV<sub>1</sub> = forced expiratory  
842 volume in one second; FVC = forced vital capacity.

843

844 **Figure 3.** Generalised additive models of age-related changes in physiological measures in males with lifetime  
845 depression and healthy controls. The solid lines represent physiological measures against smoothing functions of age.  
846 The shaded areas correspond to approximate 95% confidence intervals ( $\pm 2 \times$  standard error). FEV<sub>1</sub> = forced expiratory  
847 volume in one second; FVC = forced vital capacity.

848

849 **Figure 4.** Difference smooths comparing age-related changes in physiological measures of females with lifetime  
850 depression to healthy controls. The shaded areas correspond to approximate 95% confidence intervals ( $\pm 2 \times$  standard  
851 error). Negative values on the y-axes correspond to lower values in females with lifetime depression compared to healthy  
852 controls. The horizontal lines represent no difference between female cases and controls. FEV<sub>1</sub> = forced expiratory  
853 volume in one second; FVC = forced vital capacity; Bonf. = Bonferroni; BH = Benjamini & Hochberg.

854

855 **Figure 5.** Difference smooths comparing age-related changes in physiological measures of males with lifetime  
856 depression to healthy controls. The shaded areas correspond to approximate 95% confidence intervals ( $\pm 2 \times$  standard  
857 error). Negative values on the y-axes correspond to lower values in males with lifetime depression compared to healthy  
858 controls. The horizontal lines represent no difference between male cases and controls. FEV<sub>1</sub> = forced expiratory volume  
859 in one second; FVC = forced vital capacity; Bonf. = Bonferroni; BH = Benjamini & Hochberg.

860 **Tables**  
861

**Table 1.** Physiological measures

	Female		Male		
	Overall (N=502521)	Healthy control (N=123842)	Depression (N=56276)	Healthy control (N=131212)	Depression (N=31063)
<b>Hand-grip strength</b>					
Mean (SD)	31.77 (11.34)	25.10 (6.46)	24.73 (6.67)	41.34 (9.09)	40.37 (9.37)
Missing	3203				
<b>Systolic blood pressure</b>					
Mean (SD)	137.87 (18.65)	135.07 (19.20)	132.32 (18.28)	141.11 (17.38)	139.02 (16.78)
Missing	1325				
<b>Diastolic blood pressure</b>					
Mean (SD)	82.26 (10.15)	80.63 (9.98)	80.26 (9.93)	84.19 (9.96)	83.78 (10.02)
Missing	1323				
<b>Pulse rate</b>					
Mean (SD)	69.42 (11.26)	69.83 (10.38)	70.05 (10.52)	67.74 (11.59)	68.70 (11.99)
Missing	1323				
<b>Body mass index</b>					
Mean (SD)	27.43 (4.80)	26.63 (4.86)	27.35 (5.39)	27.62 (4.01)	28.08 (4.45)
Missing	3105				
<b>Body fat percentage</b>					
Mean (SD)	31.45 (8.55)	35.98 (6.77)	36.87 (7.02)	24.93 (5.63)	25.55 (5.88)
Missing	10408				
<b>Fat mass</b>					
Mean (SD)	24.86 (9.57)	26.12 (9.55)	27.67 (10.60)	21.85 (7.87)	22.88 (8.71)
Missing	10973				
<b>Fat-free mass</b>					
Mean (SD)	53.22 (11.50)	44.43 (4.85)	45.05 (5.17)	63.74 (7.61)	64.25 (7.93)
Missing	10176				
<b>Waist-hip ratio</b>					
Mean (SD)	0.87 (0.09)	0.81 (0.07)	0.82 (0.07)	0.93 (0.06)	0.94 (0.07)
Missing	2265				
		N=43133	N=19146	N=52359	N=12177
<b>Peak expiratory flow</b>					
Mean (SD)	407.82 (130.78)	355.08 (76.97)	359.35 (76.98)	511.67 (115.38)	509.05 (117.30)
Missing	168566				
<b>Forced expiratory volume 1s</b>					
Mean (SD)	2.85 (0.78)	2.50 (0.53)	2.53 (0.53)	3.45 (0.73)	3.43 (0.75)
Missing	149197				
<b>Forced vital capacity</b>					
Mean (SD)	3.78 (0.98)	3.28 (0.64)	3.31 (0.64)	4.59 (0.88)	4.57 (0.90)
Missing	149197				
<b>FEV<sub>1</sub> / FVC</b>					
Mean (SD)	0.75 (0.07)	0.76 (0.06)	0.76 (0.06)	0.75 (0.07)	0.75 (0.07)
Missing	149197				
		N=85994	N=31736	N=89790	N=17285
<b>Heel bone mineral density</b>					
Mean (SD)	0.54 (0.14)	0.52 (0.12)	0.52 (0.12)	0.58 (0.15)	0.57 (0.15)
Missing	180831				
		N=38592	N=25073	N=42889	N=14289
<b>Arterial stiffness</b>					
Mean (SD)	9.34 (4.05)	8.70 (4.41)	8.71 (3.27)	9.91 (4.27)	9.97 (3.30)
Missing	332721				

Note: SD = standard deviation; FEV<sub>1</sub> = forced expiratory volume in one second; FVC = forced vital capacity.

Table 2. Sex differences in physiological measures (females compared to males)

Variable	SMD	95% CI	
Hand-grip strength	-2.051	-2.060	-2.043
Systolic blood pressure	-0.358	-0.364	-0.351
Diastolic blood pressure	-0.361	-0.368	-0.354
Pulse rate	0.179	0.172	0.186
Body mass index	-0.185	-0.191	-0.178
Body fat percentage	1.771	1.763	1.779
Fat mass	0.502	0.496	0.509
Fat-free mass	-3.006	-3.016	-2.997
Waist-hip ratio	-1.775	-1.782	-1.767
Peak expiratory flow	-1.569	-1.582	-1.557
Forced expiratory volume 1s	-1.460	-1.473	-1.448
Forced vital capacity	-1.674	-1.687	-1.661
FEV <sub>1</sub> / FVC	0.165	0.154	0.176
Heel bone mineral density	-0.397	-0.405	-0.388
Arterial stiffness	-0.304	-0.315	-0.292

*Note:* SMD = standardised mean difference; CI = confidence interval; FEV<sub>1</sub> = forced expiratory volume in one second; FVC = forced vital capacity. All Bonferroni-adjusted *p*-values for Welch's *t*-test < 0.001. Negative values correspond to lower values in females.

Table 3. Differences in physiological measures between individuals with lifetime depression and healthy controls

Variable	Female					Male				
	SMD	95% CI		<i>p</i> <sub>Bonf.</sub>	<i>p</i> <sub>BH</sub>	SMD	95% CI		<i>p</i> <sub>Bonf.</sub>	<i>p</i> <sub>BH</sub>
Hand-grip strength	-0.056	-0.066	-0.046	<0.001	<0.001	-0.106	-0.118	-0.093	<0.001	<0.001
Systolic blood pressure	-0.145	-0.155	-0.135	<0.001	<0.001	-0.121	-0.133	-0.108	<0.001	<0.001
Diastolic blood pressure	-0.037	-0.047	-0.028	<0.001	<0.001	-0.041	-0.053	-0.028	<0.001	<0.001
Pulse rate	0.022	0.012	0.032	<0.001	<0.001	0.083	0.070	0.095	<0.001	<0.001
Body mass index	0.144	0.134	0.154	<0.001	<0.001	0.112	0.100	0.124	<0.001	<0.001
Body fat percentage	0.129	0.119	0.139	<0.001	<0.001	0.111	0.098	0.123	<0.001	<0.001
Fat mass	0.156	0.146	0.166	<0.001	<0.001	0.129	0.117	0.141	<0.001	<0.001
Fat-free mass	0.124	0.114	0.134	<0.001	<0.001	0.066	0.053	0.078	<0.001	<0.001
Waist-hip ratio	0.090	0.080	0.100	<0.001	<0.001	0.138	0.126	0.151	<0.001	<0.001
Peak expiratory flow	0.055	0.038	0.072	<0.001	<0.001	-0.023	-0.042	-0.003	0.382	0.032
Forced expiratory volume 1s	0.051	0.034	0.068	<0.001	<0.001	-0.024	-0.044	-0.005	0.241	0.022
Forced vital capacity	0.046	0.029	0.063	<0.001	<0.001	-0.022	-0.042	-0.003	0.421	0.032
FEV <sub>1</sub> / FVC	0.018	0.001	0.035	0.569	0.044	-0.011	-0.031	0.009	>0.999	0.276
Heel bone mineral density	0.013	-0.0001	0.026	0.789	0.056	-0.077	-0.093	-0.061	<0.001	<0.001
Arterial stiffness	0.004	-0.012	0.020	>0.999	0.571	0.015	-0.004	0.034	>0.999	0.078

Note: SMD = standardised mean difference; CI = confidence interval; Bonf. = Bonferroni; BH = Benjamini & Hochberg; FEV<sub>1</sub> = forced expiratory volume in one second; FVC = forced vital capacity. *P*-values for Welch's t-test. Negative values correspond to lower values in depression cases.









