

Social predictors of food insecurity during the stay-at-home order due to the COVID-19 pandemic in Peru. Results from a cross-sectional web-based survey

Jorge L. Cañari-Casaño^{1,2}, Omaira Cochachin-Henostroza³, Oliver A. Elorreaga^{1,2,4}, Gandy Dolores-Maldonado⁵, Anthony Aquino-Ramírez³, Sindy Huaman-Gil^{5,6}, Juan P. Giribaldi-Sierralta^{5,6}, Juan Pablo Aparco^{7,8}, Daniel A. Antiporta⁹, Mary E. Penny¹⁰

AFILIATION INSTITUTIONAL

1. Clima, Latin American Center of Excellence for Climate Change and Health, Universidad Peruana Cayetano Heredia.
2. Emerge, Emerging Diseases and Climate Change Research Unit, School of Public Health and Administration, Universidad Peruana Cayetano Heredia.
3. Grupo de Investigación Nutobes, Universidad Nacional Mayor de San Marcos, Lima, Perú
4. GMINIS Research Group, Pontificia Universidad Católica del Perú
5. Escuela Profesional de Nutrición, Facultad de Medicina, Universidad Nacional Mayor de San Marcos, Lima, Perú
6. Sociedad Científica de Estudiantes de Nutrición, Universidad Nacional Mayor de San Marcos, Lima, Perú
7. Centro Nacional de Alimentación y Nutrición, Instituto Nacional de Salud, Lima, Perú
8. Universidad San Ignacio de Loyola, Carrera de Nutrición y Dietética, Lima, Perú.
9. Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA.
10. Nutrition Research Institute, Lima, Peru.

SUMMARY

Background

Stay-at-home orders and social distancing have been implemented as the primary tools to reduce the spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). However, this approach has indirectly lead to the unemployment of 2·3 million Peruvians, in Lima, Perú alone. As a result, the risk of food insecurity may have increased, especially in low-income families who rely on a daily wage. This study estimates the prevalence of moderate or severe food insecurity (MSFI) and identifies the associated factors that explain this outcome during the stay-at-home order.

Methods

A cross-sectional web-based survey, with non-probabilistic sampling, was conducted between May 18 and June 30, 2020, during the stay-at-home order in Peru. We used social media advertisements on Facebook to reach 18-59-year-olds living in Peru. MSFI was assessed using the Food Insecurity Experience Scale (FIES). Rasch model methodology requirements were considered, and factors associated with MSFI were selected using stepwise forward selection. A Poisson generalized linear model (Poisson GLM), with log link function, was employed to estimate adjusted prevalence ratios (aPR).

Findings

This analysis is based on 1846 replies. The prevalence of MSFI was 23·2%, and FIES proved to be an acceptable instrument with reliability 0·72 and infit 0·8-1·3. People more likely to experience MSFI were those with low income (less than 255 US\$/month) in the pre-pandemic period (aPR 3·77; 95%CI, 1·98-7·16), those whose income was significantly reduced during the pandemic period (aPR 2·27; 95%CI, 1·55-3·31), and those whose savings ran out in less than 21 days (aPR 1·86; 95%CI, 1·43-2·42). Likewise, heads of households (aPR 1·20; 95%CI, 1·00-1·44) and those with probable SARS-CoV2 cases as relatives (aPR 1·29; 95%CI, 1·05-1·58) were at an increased risk of MSFI. Additionally, those who perceived losing weight during the pandemic (aPR 1·21; 95%CI, 1·01-1·45), and increases in processed foods prices (aPR 1·31; 95%CI, 1·08-1·59), and eating less minimally processed food (aPR 1·82; 95%CI, 1·48-2·24) were more likely to experience MSFI.

Interpretation

People most at risk of MSFI were those in a critical economic situation before and during the pandemic. Social protection policies should be reinforced to prevent or mitigate these adverse effects.

Funding

None.

INTRODUCTION

By the end of January 2020, the world had experienced over 100 million SARS-CoV2 cases, of whom approximately 2.3 million have died.⁽¹⁾ Peru has had around 1 million cases and more than 41,000 have died from SARS-CoV2.⁽²⁾ Moreover, Peru is one of the countries with the highest SARS-CoV2 mortality rate in the world.⁽³⁾ After declaring a global health emergency,⁽⁴⁾ the World Health Organization (WHO) recommended social distancing and closing public and private services, among other measures to contain the spread of SARS-CoV2.

Two principal measures adopted by many countries were stay-at-home orders and border closures (i.e., national lockdowns). In Peru, these were implemented from March 16 to June 30, 2020. People were only allowed to go out for essential activities, for instance to buy food and medicine. However, due to the precarious labour market in Peru, where informal labour constitutes 50% of all labour,⁽⁵⁾ and pre-pandemic poverty the purchasing power of Peruvian families was severely affected by the lockdown. The Peruvian government tried to help this vulnerable population by giving social aid (i.e., financial stimulus to families, early withdrawal from the private retirement pension, and food donations) to face the stay-at-home orders' side effects. It has been suggested that this was not enough due to inefficient distribution and the already critical situation exacerbated by COVID-19 pandemic.⁽⁶⁾

stay-at-home orders could affect food insecurity due to the resulting economic crisis, massive loss of jobs, high food prices, and growing demand for medical care.⁽⁷⁾ These results may generate short- and medium-term adverse health effects.⁽⁷⁾ Recent research on the short-term impacts of stay-at-home orders supports this idea. One of the first studies in Peru reported a considerable income reduction after stay-at-home orders in 37% of the participants,⁽⁸⁾ and over 2.3 million lost jobs in Lima alone.⁽⁹⁾ Similar occurrences have been reported outside Peru.⁽¹¹⁻¹⁰⁾ Furthermore, higher food prices, changes in diets, increased depression, anxiety and physical violence against women have all been associated with stay-at-home orders.^(11,12,13)

Recent studies have also reported an increase in food insecurity after implementation of stay-at-home orders,⁽¹⁴⁻¹⁶⁾ which worsens as the crisis intensifies.⁽¹¹⁾ These results are alarming, especially considering that the World Food Program estimated the number of people experiencing food insecurity will increase to 265 million by the end of 2020,⁽¹⁷⁾ accompanied by potential negative effects on health and household relations.⁽⁷⁾

While MSFI was estimated at 29.9% between 2014-2016 in Peru,⁽¹⁸⁾ little data exist regarding how stay-at-orders have affected MSFI. The present study aimed to estimate the prevalence of MSFI and determine the factors associated with MSFI in our study population.

Research in context

Evidence before this study

Stay-at-home orders established to stop the spread of the COVID-19 pandemic have caused profound adverse effects in low- and middle-income countries such as Peru, specifically in economically vulnerable populations, who are most at risk for food insecurity. We searched PubMed servers up to Jan 20, 2020, for peer-reviewed articles published using the terms “(food insecurity OR food security) AND (COVID-19 OR SARS-CoV2) OR (stay-at-home order OR Quarantine OR lockdown)”, without data or language restrictions. Our search found few studies that have examined the direct or indirect impact of stay-at-home orders on moderate-severe food insecurity (MSFI). However, the available evidence is scarce in low- and middle-income settings, especially Latin American countries such as Peru.

Added value of this study

To our knowledge, this study provides one of the first pieces of evidence about the prevalence of MSFI, measured using the Food Insecurity Experience Scale (FIES), and its associated factors during the stay-at-home order in Peru. Our findings agree with the growing scientific literature reporting that those most at risk of MSFI occupy tenuous economic positions before and during the pandemic.

Implications of all the available evidence

It is urgent to implement social protection policies that aim to strengthen real-time systems focusing on identifying and providing social assistance to vulnerable populations most at risk of MSFI. These systems should consider that non-poor households can quickly descend into poverty due to the inability of day laborers to work during stay-at-home orders. It is also essential to identify appropriate social aid strategies (money transfers, food donations, communal pots of food, etc.) according to the vulnerable population's characteristics and state of the pandemic. These recommendations may mitigate the adverse effects of stay-at-home orders on food MSFI.

METHODS

Study design and participants

In this cross-sectional study an online survey administered through the Qualtrics platform was used to estimate the prevalence of and factors associated with MSFI. The survey was administered to Peruvian adults, 18 to 59 years old who had internet and social media (Facebook) access by mobile devices or computers.

Procedures

The survey was accessible between May 18 and June 30 during the stay-at-home order. The dissemination of this study was done by: (1) paid advertisements from the “Colegio de Nutricionistas del Perú” official social media site; (2) social media posts by public and private universities and Peruvian research centres, and (3) by dissemination of the online survey link by participants through their personal social media profile or through their WhatsApp groups and contacts.

Participants who clicked on the link were redirected to a Qualtrics survey platform to read more about the study and consent before starting the survey. The survey could be answered more than once, and if this happened, participants had to mention why they were repeating the survey. Participants who completed the study were invited to download educational material on “Healthy Eating” and asked to share the survey link through their social media and WhatsApp contacts.

The online survey collected sociodemographic information (i.e., place of residence, sex, age, marital status, nationality, educational level, employment status, whether they suffered from a chronic disease or disability, SARS-CoV2 symptoms or any relative that tested positive for SARS-CoV2 in participant environment); FIES; household information (i.e., number of household members, average monthly income, economic condition during stay-at-home order, emergency savings, government financial support, water connection and drainage); perception about food prices (i.e., prices relative to pre-pandemic period: “lower price”, “same price”, “higher price”, “do not know/prefer not to answer”); perception about food consumption (i.e., “eat less”, “eat the same”, “eat more”, “do not eat/prefer not to answer”); perception about places to buy and waiting time, and perception about body weight status before and during the pandemic using body shape images were included in the questionnaire. To validate the survey questionnaire, a pilot study was conducted by phone call to 23 people of different age groups.

Food Insecurity Measurement

We used the Food Insecurity Experience Scale (FIES) at the individual level to determine the prevalence of MSFI. The FIES is an eight-item questionnaire with dichotomous responses (i.e., Yes or No) about experiences with food insecurity at the household or individual level.

The FIES was developed by Food and Agriculture Organization’s (FAO) Voices of the Hungry (VoH) project and was validated in 151 countries with Gallup World Poll (GWP) data in 2014.⁽¹⁹⁻²¹⁾ Several studies have used FIES to determine food insecurity in different fields and context.⁽²²⁻²⁸⁾

We adapted the FIES questionnaire to the Peruvian context taking into consideration a study by Vargas and Penny,⁽²⁹⁾ and a pilot study (n=23) where content and understanding were evaluated. Minor modifications were done to the formulation of the questions and the whole questionnaire was translated into Spanish (Table 1).

To analyze and determine MSFI, we only considered people who responded positively (value: 1) or negatively (value: 0) to all questions on the FIES. Since the FIES scale fulfilled all Rasch model assumptions (reliability and infit), a raw score was obtained (i.e., sum of affirmative responses). Individuals with raw scores between 4-8 were classified as MSFI and those with scores of 0-3 were classified as not having MSFI.⁽³⁰⁾ Similar cut off points were used to determine MSFI in other studies.^(22, 26, 31, 32)

Table 1: Food Insecurity Experiences Scale (FIES)¹

Order	Questions	Items	Possible answers
Q1	During the last month of quarantine, was there a time when, because of lack of money or other resources, you were worried you would not have enough food to eat?	WORRIED	1= Yes 2= No 3= Don’t Know 4=Prefer not to answer
Q2	During the last month of quarantine, was there a time when, because of lack of money or other resources, you could not eat healthy and nutritious food? For example: meat, fish, fruits, or vegetables.	HEALTHY	1= Yes 2= No 3= Don’t Know 4=Prefer not to answer

Q3	During the last month of quarantine, was there a time when, because of lack of money or other resources, you had to eat only a few kinds of foods?	FEWFOODS	1= Yes 2= No 3= Don't Know 4=Prefer not to answer
Q4	During the last month of quarantine, was there a time when, because of lack of money or other resources, you had to reduce the number of meals per day because you did not have enough food? For example, skip breakfast, lunch, or dinner?	SKIPPED	1= Yes 2= No 3= Don't Know 4=Prefer not to answer
Q5	During the last month of quarantine, was there a time when, because of lack of money or other resources, you had to eat less than you thought you should eat?	ATELESS	1= Yes 2= No 3= Don't Know 4=Prefer not to answer
Q6	During the last month of quarantine, was there a time when, because of lack of money or other resources, your food ran out in your household?	RANOUT	1= Yes 2= No 3= Don't Know 4=Prefer not to answer
Q7	During the last month of quarantine, was there a time when, because of lack of money or other resources, you went to sleep hungry because there was not enough food?	HUNGRY	1= Yes 2= No 3= Don't Know 4=Prefer not to answer
Q8	During the last month of quarantine, was there a time when, because of lack of money or other resources, you went without eating for a whole day?	WHLDAY	1= Yes 2= No 3= Don't Know 4=Prefer not to answer

¹ The Spanish version of the FIES is provided in the supplementary material.

Covariates

Sociodemographic variables including age, sex, place of residence, educational level, if respondent was a head of household, and number of household members were considered. Household income level before and during the pandemic were related to an individual's financial status and use and duration of savings during the stay-at-home order. Variables related to social assistance included support given by the government or any institution. Variables used to understand changes since the emergency began, included changes in location where food was bought and principal food groups consumed (i.e., healthy or minimally processed food, culinary ingredients, processed food, and ultra-processed foods). Body weight perceptions before and during the pandemic were evaluated using the Stunkard scale.⁽³³⁾ Finally, variables related to COVID-19 pandemic were also considered; specifically, participants reporting a SARS-CoV2 diagnosis, loss of smell/taste, or whose household members had or had had SARS-CoV2.

Data Analysis

Rash model assumptions for the FIES scale were evaluated using the RM.weight package in R statistical software,^(21, 34) where performance of an eight-item questionnaire was assessed utilizing infit statistic and the ability to distinguish different levels of food insecurity in participants was evaluated with the reliability of FIES scale. After verifying the fulfilment of the FIES scale's assumptions, food insecurity was determined using the raw score (the sum of affirmative responses). Descriptive analysis of categorical variables and bivariate analysis between MSFI and other predictor variables were performed by proportions and chi-square test, respectively. Factors associated with the predictor variables and MSFI were determined using forward stepwise selection, starting with the most significant predictor variable. We used Poisson generalized linear models (Poisson GLMs) with log link function, to estimate adjusted prevalence ratios (aPR) fitted at the departmental level. Data analysis was performed using Stata 15.0 (Stata Corp., College Station, Texas – United States) and software program R studio (Version 4.0.4; R Foundation for Statistical Computing, Vienna, Austria).

Ethical Issues

The study was approved by the human ethics committee of “Instituto de Investigación Nutricional” (CIEI-IIN), Lima – Peru, N° 394-2020/CIEI-IIN.

RESULTS

A total of 2643 people answered the study's informed consent, and 116 declined to participate in the study. Fifty-two participants were excluded because we could not determine if they had taken the survey more than once or they retook the survey for an inadequate reason and 629 participants because they did not answer all the questions on the FIES scale. The final sample considered for the analysis was 1846 participants (Figure 1).

Figure 1: Flow diagram




Table 2 shows the infit statistic and the Rasch model parameters. Infit values were acceptable, and Rasch model reliability in the FIES scale was 0.72 (data not shown in the table) which was also acceptable (greater than 0.7).

Table 2: Evaluation of the assumptions of the Rash model of the FIES scale.

Item	Item_sev.*	St.err.†	Infit‡	Outfit
WORRIED	-2.37100181	0.0804195	1.2714576	2.2715038
FEWFOOD	-1.99919744	0.0785325	0.8542389	0.7456257
HEALTHY	-0.72783574	0.0818627	1.0229524	0.9484038
ATELESS	-0.8382067	0.08112009	0.8300268	0.7194007
SKIPPED	0.02735437	0.0891645	0.9363935	0.8611874
RUNOUT	0.69432414	0.09915824	1.1374597	1.2012408
HUNGRY	1.31138247	0.1125294	0.8408805	0.7309963
WHLDAY	3.90325585	0.25963247	1.0598114	0.7132612

*Item_sev: Item severity. †St.err: Standard error. ‡Infit: Infit statistic. Acceptable values are in the range of 0.7 - 0.13.

Most of the participants were women (74.9%) with a technical or university education (85.3%). About a third were heads of household (34.2%) and lived-in provinces other than Metropolitan Lima (32.2%). Regarding economic factors, a low proportion of people (12.4%) received less than the minimum wage in Peru (less than 255 US\$/month) before the pandemic and only 23.4% continued receiving income during the stay-at-home orders. On the other hand, over half of participants (56.7%) had emergency savings left to address the pandemic, while 16% and 9.5% of people ran out of savings in 22-35 days and less than 21 days since the start of the stay-at-home order, respectively.

As for factors related explicitly to the stay-at-home order, we found 17% of people had relatives with probable cases of SARS-CoV2, 30% stayed at home without working but had worked in the pre-pandemic period, 38.1% were working during the stay at home period, and 26.8% were not working before or during the stay at home order. 71.6% reported that the process to buy food took more time than before, and only 23.4% mentioned they received some support (include the government and private support) during the COVID-19 pandemic. In relation to body weight perceptions, 24.9% people perceived losing weight, 32.1% maintained their weight, and 30.7% gained weight. Regarding natural or minimally processed food consumption, 30% of participants mentioned that they ate less of this food group, and 39% ate more. It was found that 23.3% of participants experienced MSFI (Table 3).

Table 3. General characteristics of the population studied.

Characteristics	N (n=1846)	(%)
Sociodemographic factors		
Sex		
Female	1383	(74.9%)
Male	451	(24.4%)
Missing/prefer not to answer	12	(0.7%)
Age		
18-25	723	(39.2%)
26-35	516	(30.0%)
36-59	607	(32.9%)
Level of education		
Higher university education	1353	(73.3%)
Higher technical education	222	(12.0%)
Up to secondary education	250	(13.5%)

Missing/prefer not to answer	21 (1.1%)
Head of household	
No	1167 (63.2%)
Yes	631 (34.2%)
Missing/prefer not to answer	48 (2.6%)
Place of residence	
Province of Metropolitan Lima	1249 (67.7%)
Other provinces or departments	594 (32.2%)
Missing/prefer not to answer	3 (0.2%)
Economic factors	
Household income pre-pandemic period (US\$/month)	
More than 1788	181 (9.8%)
767-1787	452 (24.5%)
256-766	644 (34.9%)
Less than 255	228 (12.4%)
Missing/prefer not to answer	341 (18.5%)
Household income during COVID-19 pandemic	
Still receiving income	432 (23.4%)
Income decreased slightly	542 (29.4%)
Income decreased considerably or ceased	748 (40.5%)
Missing/prefer not to answer	124 (6.7%)
Duration of savings to address the COVID-19 pandemic (grouped in days)	
Still have savings	1047 (56.7%)
It ran out between 22 to 35 days	296 (16.0%)
It ran out in less than 21 days	175 (9.5%)
Missing/prefer not to answer	328 (17.8%)
Factors related to the COVID-19 pandemic	
People with relatives with probable cases of SARS-CoV2	
Without probable cases of SARS-CoV2	1529 (82.8%)
With probable cases of SARS-CoV2	314 (17.0%)
Missing/prefer not to answer	3 (0.2%)
The main situation during the COVID-19 pandemic	
Staying at home without working	555 (30.1%)
Usually does not work	494 (26.8%)
Working in this period	704 (38.1%)
Missing/prefer not to answer	93 (5.0%)
Waiting time to purchase of most food	
It is slower, takes longer to do the shopping	1321 (71.6%)
It is the same as the pre-pandemic time	210 (11.4%)
It is faster. It takes less time	78 (4.2%)
Missing/prefer not to answer	237 (12.8%)
Change in the usual place of purchase of most food	

Changed the place of purchase	875 (47.4%)
Did not change the place of purchase	736 (39.9%)
Missing/prefer not to answer	235 (12.7%)
Perception of food price of processed foods during COVID-19 pandemic	
Lowered or maintained	564 (30.6%)
Price increased	1026 (55.6%)
Missing/prefer not to answer	256 (13.9%)
Social support in the COVID-19 pandemic	
Received some support during the COVID-19 pandemic	
Yes	432 (23.4%)
No	1326 (71.8%)
Missing/prefer not to answer	88 (4.8%)
Perception of body condition	
Perception of weight loss during the COVID-19 pandemic	
Maintained or gained weight	1159 (62.8%)
Lost weight	459 (24.9%)
Missing/prefer not to answer	228 (12.4%)
Food patterns and food insecurity	
Perception of consumption of minimally processed foods during the COVID-19 pandemic	
Ate the same or more	1089 (59.0%)
Ate less	553 (30.0%)
Missing/prefer not to answer	204 (11.1%)
Moderate-Severe Food insecurity (MSFI)	
No	1418 (76.8%)
Yes	428 (23.2%)

Table 4. Bivariate analysis of potential factors associated with moderate or severe food insecurity.

Variables	Moderate or Severe Food Insecurity		P*
	No (total=1418) n (%)	Yes (total=428) n (%)	
Sociodemographic factors			
Head of household			0.018
No	914 (78.3%)	253 (21.7%)	
Yes	463 (73.4%)	168 (26.6%)	
Economic factors			
Household income pre-pandemic period (US\$/month)			< 0.001
More than 1788	168 (92.8%)	13 (7.2%)	
767-1787	403 (89.2%)	49 (10.8%)	
256-766	461 (71.6%)	183 (28.4%)	
Less than 255	113 (49.6%)	115 (50.4%)	

Household income during COVID-19 pandemic			< 0.001
Still receiving income	395 (91.4%)	37 (8.6%)	
Income decreased slightly	457 (84.3%)	85 (15.7%)	
Income decreased considerably or ceased	466 (62.3%)	282 (37.7%)	
Duration of savings to address the COVID-19 pandemic (grouped in days)			< 0.001
Still have savings	908 (86.7%)	139 (13.3%)	
It ran out between 22 to 35 days	161 (54.4%)	135 (45.6%)	
It ran out in less than 21 days	85 (48.6%)	90 (51.4%)	
Factors related to the COVID-19 pandemic			
People with relatives with probable cases of SARS-CoV2			< 0.001
Without probable cases of SARS-CoV2	1200 (78.5%)	329 (21.5%)	
With probable cases of SARS-CoV2	216 (68.6%)	98 (31.2%)	
The main situation during the Covid-19 pandemic			< 0.001
Staying at home without working	385 (69.4%)	170 (30.6%)	
Usually does not work	384 (77.7%)	110 (22.3%)	
Working in this period	581 (82.5%)	123 (17.5%)	
Waiting time to purchase of most food			0.017
It is slower, takes longer to do the shopping	1000 (75.7%)	321 (24.3%)	
It is the same as the pre-pandemic time	173 (82.4%)	37 (17.6%)	
It is faster. It takes less time	67 (85.9%)	11 (14.1%)	
Change in the usual place of purchase of most food			0.013
Changed the place of purchase	656 (75.0%)	219 (25.0%)	
Did not change the place of purchase	590 (80.2%)	146 (19.8%)	
Perception of food price of processed foods during the COVID-19 pandemic			< 0.001
Lowered or maintained	465 (82.5%)	99 (17.6%)	
Price increased	757 (73.8%)	269 (26.2%)	
Social support in the COVID-19 pandemic			
Received some support during the COVID-19 pandemic			< 0.001
Yes	284 (65.7%)	148 (34.3%)	
No	1066 (80.4%)	260 (19.6%)	
Perception of body condition			
Perception of weight loss during the COVID-19 pandemic			< 0.001
Maintained or gained weight	929 (80.2%)	230 (19.8%)	
Lost weight	318 (69.3%)	141 (30.7%)	
Food patterns and food insecurity			
Perception of consumption of minimally processed foods during the COVID-19 pandemic			< 0.001
Ate the same or more	944 (86.7%)	145 (13.3%)	
Ate less	323 (58.4%)	230 (41.6%)	

* Chi-square test.

In bivariate analysis between MSFI and principal predictor variables, most were found to be significant, except for participant sex and age (table 4). The factors associated with MSFI are shown in Table 5. Among sociodemographic factors, participants who were heads of households had a prevalence of MSFI up to 1.2 times higher compared to participants who were not (aPR

1.20, CI95%: 1.00-1.44). In relation to economic factors, people whose households had an average monthly income of < 255 US\$/month and between 256 to 766 US\$/month in the pre-pandemic period had a prevalence of MSFI up to 3.8 (aPR 3.77 CI95%: 1.98-7.16) and 2.8 (aPR 2.80 CI95%: 1.50-5.23) times higher, respectively compared to households with an average monthly income of more than 1788 US\$/month. People whose savings to address the emergency ran out in between 22 to 35 days and less than 21 days since the start of the stay-at-home order, had a prevalence of MSFI up to 1.84 (aPR 1.84, CI95%: 1.46-2.31) and 1.86 (aPR 1.86, CI95%: 1.43-2.42) times higher, respectively, compared to people with savings left to address the emergency.

On the other hand, participants with relatives who were probable cases of SARS-CoV2 had a prevalence of MSFI up to 1.29 (aPR 1.29, CI95%: 1.05-1.58) times higher relative to those who did not. Participants who reported a loss of weight during the pandemic period had a prevalence of MSFI 1.21 (aPR 1.21, CI95%: 1.01-1.45) times higher than those who maintained or gained weight. Lastly, participants who reported consuming less natural or minimally processed food had a prevalence of MSFI 1.82 (aPR 1.82, CI95%: 1.48-2.24) times higher than those who ate the same or more of that food group. Participants who reported increases in processed foods prices had a prevalence of MSFI 1.31 (aPR 1.31; 95%CI, 1.08-1.59) times higher than those who reported that prices were maintained or decreased.

Table 5. Prevalence ratios of factors associated with moderate-severe food insecurity. Peru

Variables	Multivariate regression analysis.		
	aPR*	(95% CI)	p
Sociodemographic factors			
Head of household			
No	Ref.		
Yes	1.20	(1.00 - 1.44)	0.047
Economic factors			
Household income pre-pandemic period (US\$/month)			
More than 1788	Ref.		
767-1787	1.47	(0.75 – 2.86)	0.259
256-766	2.80	(1.50 - 5.23)	0.001
Less than 255	3.77	(1.98 - 7.16)	< 0.001
Household income during COVID-19 pandemic			
Still receiving income	Ref.		
Income decreased slightly	1.54	(1.02 - 2.31)	0.039
Income decreased considerably or ceased	2.27	(1.55 - 3.31)	< 0.001
Duration of savings to address the COVID-19 pandemic (grouped in days)			
Still have savings	Ref.		
It ran out between 22 to 35 days	1.84	(1.46 - 2.31)	< 0.001
It ran out in less than 21 days	1.86	(1.43 - 2.42)	< 0.001
Factors related to the COVID-19 pandemic			
People with relatives with probable cases of SARS-CoV2			
Without probable cases of SARS-CoV2	Ref.		

With probable cases of SARS-CoV2	1.29	(1.05 - 1.58)	0.015
Perception of food price of processed foods during COVID-19 pandemic			
Lowered or maintained	Ref.		
Price increased	1.31	(1.08 - 1.59)	0.007
Perception of body condition			
Perception of weight loss during the COVID-19 pandemic			
Maintained or gained weight	Ref.		
Lost weight	1.21	(1.01 - 1.45)	0.034
Food patterns and food insecurity			
Perception of consumption of minimally processed foods during the COVID-19 pandemic			
Ate the same or more	Ref.		
Ate less	1.82	(1.48 - 2.24)	< 0.001

aPR: Prevalence ratios adjusted. Multivariate regression analysis fitting by department level considering the constitutional province of Callao as a department.

DISCUSSION

The present study found a prevalence of MSFI of 23.2%. Factors associated with it were having an income below the minimum living wage in the pre-pandemic period, having experienced a substantial reduction in income during the pandemic, ran out of savings in the first 21 days of the pandemic period, being a head of household, having relatives who were probable cases of SARS-CoV2, perceiving weight loss during pandemic period, reporting that they ate less natural or minimally processed food, and having perceived an increase in prices of processed foods.

In Peru in 2014, the reported prevalence of MSFI was 29.9%;⁽³⁵⁾ however, our study found a prevalence of MSFI of 23.2% in the first 30-45 days of stay-at-home order. Although these results are not comparable with MSFI initial values, the prevalence of MSFI in this study may be underestimated because of some particular characteristics of the population analysed. For example, 73.3% had a university education, only 12.4% reported having monthly income less than 255 US\$/month before the pandemic, and participants required internet access and a social network account to access the survey, attributes that low-income people generally do not have and they are precisely the population most at risk of suffering food insecurity.^(22, 26) Other studies that have evaluated Food Insecurity in the context of the COVID-19 pandemic using FIES have reported higher MSFI prevalence. In Jordan, the prevalence of MSFI was 59.2%,⁽³⁶⁾ in Kenya and Uganda MSFI increased from 50% to 88% and from 43% to 87%, from before to during the pandemic, respectively.⁽¹⁵⁾ Our study measures individual MSFI; however, there is already evidence of the effect stay-at-home order on household MSFI. In Mexico, it was reported that the prevalence of household MSFI increased from 24.2% in 2018 to 30.2% in June 2020.⁽³¹⁾ Vermont also reported is an increase of MSFI from 18.3% pre-pandemic to 24.4% during the pandemic.⁽¹⁴⁾ In rural Bangladesh the prevalence of MSFI increased from 8.3% pre-pandemic to 51.8% during the pandemic.⁽¹¹⁾

Our study is one of the first investigations into MSFI during the stay-at-home order due to COVID-19 pandemic in Peru. We found that up to 40.5% of participants experienced a considerable reduction in their income, similar results were reported by the Young Lives study in Peru with 37% of households affected by a salary cut or suspension without payment.⁽⁸⁾ These high percentages of income losses could be related to the fact that in Peru, 50% of income came

from informal employment in the pre-pandemic period.⁽⁵⁾ This could explain our results, i.e., those who had lower income (compared to high-income people) before the pandemic, had less capacity to save money to face emergency situations, and had substantial income reduction during the pandemic, are shown to be more likely to have MSFI. These findings are consistent with other studies, for example, in Jordan it was reported that people who had a monthly income per capita below the poverty line were more likely to have moderate (OR: 5.33; 95% CI: 4.44–6.40) and severe food insecurity and severe food insecurity (OR: 6.87; 95% CI: 5.542–8.512).⁽³⁶⁾ Similarly, other studies that explored determinants of food insecurity during the COVID pandemic reported a higher risk of MSFI in low-income people,⁽³⁷⁾ people who lost their job,^(14, 38) or people with substantial income loss.⁽³⁹⁾

The Peruvian government implemented social aid programs aimed at poor populations to help them face the stay-at-home order; this aid consisted mainly of direct payment of 760 Peruvian soles (208.5 US\$), early withdrawal from the private retirement pension, and food donations. In our study, we found that most respondents reported not having been part of this beneficiary population. For example, only 15.7% received some type of support from the government and of these only 25.6% was allocated to people with pre-pandemic monthly income below the minimum living wage (less than 255 US\$) and 55.6% to people with monthly income from 256 to 766 US\$ (data not shown in tables). We do not have information on the percentage of the vulnerable population that did not receive government aid to face the stay-at-home order in Peru. However, it has been reported that the data source used to identify vulnerable households was designed for regular contexts and does not allow for the identification of families that abruptly entered poverty due to the mandatory stay-at-home order.⁽⁵⁾ Problems with receiving money from social assistance were also reported, because nearly 14 million adult Peruvians (59.8% of the total adult population) did not have a bank account.⁽⁴⁰⁾ For this reason, our results invite further analysis and discussion on the targeting and delivery of government social aid to vulnerable populations during the pandemic.

On the other hand, the high food price perception during the pandemic (55.6% participants reported an increase in the price of processed food in our study) may have limited the capacity to buy or access food during the pandemic. Loss of jobs^(11, 41) and increase in food prices⁽¹²⁾ are some of the side effects of stay-at-home orders reported by other studies.

In this study we found that people who had reported consuming less minimally processed food were more likely to experience MSFI, this may be due to reduced access to food in terms of quantity and quality, increased food price or having some fear of SARS-CoV2 infection in markets because products are not packaged.⁽⁴²⁾

It is possible that one of the reasons heads of households were more likely to suffer from MSFI could be stress and anxiety related to getting food and having to prioritize feeding their family over themselves.⁽⁷⁾ This result could be related to the perception of losing weight that, in our study, was associated with MSFI.

Our findings demonstrate a greater chance of experiencing MSFI in participants with relatives who were probable cases of SARS-CoV2. Even though there is no evidence about a direct relationship between MSFI and SARS-CoV2, we suggest that there could be an indirect relationship, as household economic resources could be allocated primarily to treatment or care of the infected household member instead of providing food at home.

The study has some limitations. Because it is a cross-sectional study; it cannot establish causal relationships. There might be bias as participants were not randomly selected, thus the results shown are only generalizable to our study population.

It is important to discuss how the sample was obtained. This study employed a web-based survey where participants were not randomly selected. Ideally, we expected to have a sample with sociodemographic characteristics similar to the Peruvian national population (21.2% and 20.8% within the poorest and poor quintiles, respectively; 28.8% residing in the province of Metropolitan Lima; 60% of women between 15-49 years old having at least secondary education and around 40% having higher education).^(43, 44) However, our sample consisted mainly of women (74.9%) and people who residing in the province of Metropolitan Lima (67.7%); with only 12.9% of those women have at least secondary level education, 87.1% have higher or university education (data not shown) and with 12.4% of households reporting an average monthly income of less than 255 US\$/month in pre-pandemic period (similar to the poorest quintile of the Peruvian national population). Considering these discrepancies between the general Peruvian population and our study sample, we can mention that this study's sample has a similar distribution to Peruvian population that live in Metropolitan Lima, with an upper-middle income socioeconomic status and it is not generalizable to the whole Peru. However, we showed important evidence on MSFI in the context of the stay-at-home order due to the COVID-19 pandemic in Peru.

In conclusion, people who went through MSFI were economically vulnerable both before and during stay-at-home order due to COVID-19 pandemic in Peru. Factors most strongly associated with MSFI included being a head of household, consuming less minimally processed food, and perceiving losing weight. It is necessary to implement effective social assistance policies to prevent or mitigate these adverse effects.

REFERENCES

1. Ensheng Dong HD, Lauren Gardner. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet*. 2020;20:533-4.
2. MINSA. Sala situacional COVID-19 Perú: Ministerio de Salud 2020 [Available from: https://covid19.minsa.gob.pe/sala_situacional.asp].
3. Worldometer. COVID-19 Coronavirus Pandemic. Reported Cases and Deaths by Country, Territory, or Conveyance 2020 [Available from: https://www.worldometers.info/coronavirus/?utm_campaign=homeAdvegas1?].
4. Sohrabi C AZ, O'Neill N, et al. World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19) *Int J Surg*. 2020;2020(76):71-6.
5. Jaramillo M ÑH. COVID-19 y shock externo Impactos económicos y opciones de política en el Perú. Serie de documentos de política pública PNUD América Latina y el Caribe 2020.
6. GRADE. The COVID-19 Pandemic and Food Insecurity in Peru, by Eduardo Zegarra: Group for the Analysis of Development 2020 [Available from: <https://www.grade.org.pe/en/novedades/the-covid-19-pandemic-and-food-insecurity-in-peru-by-eduardo-zegarra/>].
7. Anna L SW, Kartika P, Hilary S. A conceptual model for understanding the rapid COVID-19–related increase in food insecurity and its impact on health and healthcare. *Am J Clin Nutr*. 2020;00(1–8).
8. Lives Y. Listening to Young Lives at Work in Peru: First Call. 2020 Agosto 2020.: GRADE, IIN; 2020 [updated Agosto 2020. Available from: <https://www.younglives.org.uk/content/listening-young-lives-work-peru-first-call>].
9. INEI. Nota de prensa INEI: Empleo en Lima Metropolitana disminuyó en 2 millones 318 mil 300 personas durante el trimestre marzo-abril-mayo del presente año 2020 [updated 15/06/2020 Available from: <https://www.inei.gob.pe/prensa/noticias/empleo-en-lima-metropolitana-disminuyo-en-2-millones-318-mil-300-personas-durante-el-trimestre-marzo-abril-mayo-del-presente-ano-12253/>].
10. Parker Kim MJ, Brown Anna. About Half of Lower-Income Americans Report Household Job or Wage Loss Due to COVID-19 2020 [Available from: <https://www.pewsocialtrends.org/2020/04/21/about-half-of-lower-income-americans-report-household-job-or-wage-loss-due-to-covid-19/>].
11. Hamadani JD HM, Baldi AJ, et al. Immediate impact of stay-at-home orders to control COVID-19 transmission on socioeconomic conditions, food insecurity, mental health, and intimate partner violence in Bangladeshi women and their families: an interrupted time series. *Lancet Glob Health*. 2020;S2214-109X(20):30366-1.
12. Akter S. The impact of COVID-19 related 'stay-at-home' restrictions on food prices in Europe: findings from a preliminary analysis. *Food Secur*. 2020:1-7.
13. Bracale R VC. Changes in food choice following restrictive measures due to Covid-19. *Nutr Metab Cardiovasc Dis*. 2020;30(9):1423-6.
14. Niles MT, Bertmann F, Belarmino EH, Wentworth T, Biehl E, Neff RA. The Early Food Insecurity Impacts of COVID-19. *Nutrients*. 2020;12(7):2096.
15. Kansime MK, Tambo JA, Mugambi I, Bundi M, Kara A, Owuor C. COVID-19 implications on household income and food security in Kenya and Uganda: Findings from a rapid assessment. *World development*. 2021;137:105199-.
16. Dondi A, Candela E, Morigi F, Lenzi J, Pierantoni L, Lanari M. Parents' Perception of Food Insecurity and of Its Effects on Their Children in Italy Six Months after the COVID-19 Pandemic Outbreak. *Nutrients*. 2021;13(1):121.
17. IFPRI. 2020 Global report on food crises. Joint analysis for better decisions 2020 [Available from: <https://www.ifpri.org/publication/2020-global-report-food-crises-joint-analysis-better-decisions>].

18. FAO I, UNICEF, WFP, WHO. The State of Food Security and Nutrition in the World 2020. Transforming food systems for affordable healthy diets Rome2020 [Available from: <https://doi.org/10.4060/ca9692en>].
19. FAO. Methods for estimating comparable prevalence rates of food insecurity experienced by adults throughout the world Rome: FAO; 2016 [Available from: <http://www.fao.org/3/a-i4830e.pdf>].
20. Ballard TJ KA, Cafiero C, Schmidhuber J. Better measurement of food insecurity in the context of enhancing nutrition. *Ernahrungs Umschau*. 2014;61(2):38-41.
21. Carlo Cafiero SV, Mark Nord. Food security measurement in a global context: The food insecurity experience. 2018;116:146-52.
22. Smith MD, Rabbitt MP, Coleman- Jensen A. Who are the World's Food Insecure? New Evidence from the Food and Agriculture Organization's Food Insecurity Experience Scale. *World Development*. 2017;93:402-12.
23. Sinclair K, Ahmadigheidari D, Dallmann D, Miller M, Melgar-Quiñonez H. Rural women: Most likely to experience food insecurity and poor health in low- and middle-income countries. *Global Food Security*. 2019;23:104-15.
24. Frongillo E NH, Smith M, Coleman-Jensen A. Food Insecurity Is More Strongly Associated with Poor Subjective Well-Being in More-Developed Countries than in Less-Developed Countries. *The Journal of Nutrition*. 2018;149(2):330-5.
25. Grimaccia E NA. Food Insecurity Individual Experience: A Comparison of Economic and Social Characteristics of the Most Vulnerable Groups in the World. *Social Indicators Research*. 2019;143(1):391-410.
26. Wambogo EA, Ghattas H, Leonard KL, Sahyoun NR. Validity of the Food Insecurity Experience Scale for Use in Sub-Saharan Africa and Characteristics of Food-Insecure Individuals. *Current Developments in Nutrition*. 2018;2(9).
27. Karpyn A HG, Knowles Z. Validity of the Food Insecurity Experience Scale and Prevalence of Food Insecurity in the Bahamas. 2020.
28. Asfahani Fea. Food Insecurity and Subjective Wellbeing Among Arab Youth Living in Varying Contexts of Political Instability. *Journal of Adolescent Health*. 2018; 64.
29. Vargas S PM. Measuring food insecurity and hunger in Peru: A qualitative and quantitative analysis of an adapted version of the USDA's Food Insecurity and Hunger Module. *Public Health Nutrition*. 2010;13(10):1488-97.
30. Ballard TJ, Kepple, A.W. & Cafiero, C. The food insecurity experience scale: development of a global standard for monitoring hunger worldwide. Technical Paper Rome: FAO; 2013 [Available from: http://www.fao.org/fileadmin/templates/ess/voh/FIES_Technical_Paper_v1-1.pdf].
31. Gaitán-Rossi P, Vilar-Compte M, Teruel G, Pérez-Escamilla R. Food insecurity measurement and prevalence estimates during the COVID-19 pandemic in a repeated cross-sectional survey in Mexico. *Public health nutrition*. 2021;24(3):412-21.
32. Depa J GF, Muller A, Eleraky L, Hilzendegen C, Stroebele-Benschop N. Prevalence of food insecurity among food bank users in Germany and its association with population characteristics. *Preventive medicine reports*. 2018;9:96-101.
33. Stunkard AJ, Sørensen T, Schulsinger F. Use of the Danish Adoption Register for the study of obesity and thinness. *Research publications - Association for Research in Nervous and Mental Disease*. 1983;60:115-20.
34. RM.weights: Weighted Rasch Modeling and Extensions using Conditional Maximum Likelihood [Internet]. 2018. Available from: <https://cran.r-project.org/web/packages/RM.weights/index.html>.
35. FAO F, OMS, PMA, UNICEF. El estado de la seguridad alimentaria y la nutrición en el mundo 2019. Protegerse frente a la desaceleración y el debilitamiento de la economía Roma: FAO; 2019 [
36. Elsahoryi N, Al-Sayyed H, Odeh M, McGrattan A, Hammad F. Effect of Covid-19 on food security: A cross-sectional survey. *Clin Nutr ESPEN*. 2020;40:171-8.

37. Manfrinato CV, Marino A, Condé VF, Franco MdCP, Stedefeldt E, Tomita LY. High prevalence of food insecurity, the adverse impact of COVID-19 in Brazilian favela. *Public Health Nutrition*. 2020;1-14.
38. Owens MR, Brito-Silva F, Kirkland T, Moore CE, Davis KE, Patterson MA, et al. Prevalence and Social Determinants of Food Insecurity among College Students during the COVID-19 Pandemic. *Nutrients*. 2020;12(9).
39. Kent K, Murray S, Penrose B, Auckland S, Visentin D, Godrich S, et al. Prevalence and Socio-Demographic Predictors of Food Insecurity in Australia during the COVID-19 Pandemic. *Nutrients*. 2020;12(9).
40. Peruano E. MEF: Todos los peruanos tendrán una cuenta bancaria al 28 de julio del 2021 Lima2020 [Available from: <https://elperuano.pe/noticia/102330-mef-todos-los-peruanos-tendran-una-cuenta-bancaria-al-28-de-julio-del-2021>].
41. Ahmed F IA, Pakrashi D, Rahman T, Abu S. Determinants and Dynamics of Food Insecurity During COVID-19 2020 [updated September 2. Available from: <http://users.monash.edu/~asaduli/pub/foodsec-covid.pdf>].
42. Scarmozzino F VF. Covid-19 and the Subsequent Lockdown Modified Dietary Habits of Almost Half the Population in an Italian Sample. *Foods*. 2020;9(5):675.
43. INEI. National Institute of Statistics and Informatic. Demographic and Family Health Survey (DHS) 2019 [Available from: <https://proyectos.inei.gob.pe/endes/>].
44. National Institute of Statistics and Informatics. Peru: Population Projections, by Department, Province and District, 2018-2020". 2020 [Available from: https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1715/].

AUTHORSHIP CONTRIBUTIONS

OCH, MP and JLC led the design of the study. JLC, OCH, SH, GD, AA, and JPG contributed to designing and validating the instrument and collecting of data. JLC and OAE carried out the data analysis. DAA, JPA and MP were responsible for the interpretation of the results. JLC contributed to the initial drafting of the manuscript based on comments from DAA, MP, and JPA. All authors contributed to the interpretation of data, critical revision of the manuscript and approved the final version.

DECLARATION OF INTERESTS

We declare no competing interests.

DATA SHARING

Requests of survey data will be considered for research purposes, while does not conflict with other requests by the study authors. Requests should be made to Jorge L. Cañari-Casaño (jorge.canari@outlook.com).

ACKNOWLEDGMENTS

We want to thank Rikolto in Latin America and the Belgian cooperation for their contribution in designing the advertising campaign on social media. We also thank Colegio de Nutricionistas del Perú, Universidad Nacional Mayor de San Marcos and all the public Peruvians universities that sharing at not cost our survey online from their official social media pages.

JLC is a doctoral student studying an Epidemiological Research Doctorate at Universidad Peruana Cayetano Heredia under FONDECYT/CIENCIACTIVA scholarship EF033-235-2015 and supported by training grant D43 TW007393 awarded by the Fogarty International Center of the US National Institutes of Health.

We also express our acknowledgement to Hilary Creed for her comments on our protocol of study and John Neseemann, for his review of the English version of our manuscript.

CORRESPONDENT AUTHOR

Jorge Luis Cañari Casaño

Email: jorge.canari@outlook.com