Title

Global impact and cost-effectiveness of one-dose versus two-dose human papillomavirus vaccination schedules: a comparative modelling analysis

Kiesha Prem[*], Yoon Hong Choi[†], Élodie Bénard[†], Emily A Burger[†], Liza Hadley, Jean-François Laprise, Mary Caroline Regan, Mélanie Drolet, Stephen Sy, Kaja Abbas, Allison Portnoy, Jane J Kim, Marc Brisson, Mark Jit[*]

[*]Joint Corresponding Author
[†]Joint Second Author; contributed equally

Authors and Affiliations

Kiesha Prem
Joint Corresponding Author
Degree: PhD
Affiliations: Department of Infectious Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, United Kingdom; Saw Swee Hock School of Public Health, National University of Singapore and National University Health System, Singapore
Email: kiesha.prem@lshtm.ac.uk

Yoon Hong Choi
Joint Second Author
Degree: PhD
Affiliations: Modelling and Economics Unit, Data, Analytics and Surveillance, UK Health Security Agency, London, United Kingdom
Email: yoon.choi@ukhsa.gov.uk

Élodie Bénard
Joint Second Author
Degree: MSc
Affiliations: Centre de recherche du CHU de Québec - Université Laval, Québec, QC, Canada
Email: elodie.benard.1@ulaval.ca

Emily A Burger
Joint Second Author
Degree: PhD
Affiliations: Center for Health Decision Science, Harvard T.H. Chan School of Public Health, Boston, MA, USA; Department of Health Management and Health Economics, University of Oslo, Oslo, Norway
Email: eburger@hsph.harvard.edu

Liza Hadley
Degree: MMath

Affiliations: Department of Infectious Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, United Kingdom; Disease Dynamics Unit, Department of Veterinary Medicine, University of Cambridge, Cambridge, United Kingdom
Email: lh667@cam.ac.uk

Jean-François Laprise
Degree: PhD
Affiliations: Centre de recherche du CHU de Québec - Université Laval, Québec, QC, Canada
Email: jean-francois.laprise@crchudequebec.ulaval.ca

Mary Caroline Regan
Degree: BA
Affiliations: Center for Health Decision Science, Harvard T.H. Chan School of Public Health, Boston, MA, USA
Email: mregan@hsph.harvard.edu

Mélanie Drolet
Degree: PhD
Affiliations: Centre de recherche du CHU de Québec - Université Laval, Québec, QC, Canada
Email: melanie.drolet@crchudequebec.ulaval.ca

Stephen Sy
Degree: MS
Affiliations: Center for Health Decision Science, Harvard T.H. Chan School of Public Health, Boston, MA, USA
Email: ssy@hsph.harvard.edu

Kaja Abbas
Degree: PhD
Affiliations: Department of Infectious Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, United Kingdom
Email: kaja.abbas@lshtm.ac.uk

Allison Portnoy
Degree: ScD
Affiliations: Center for Health Decision Science, Harvard T.H. Chan School of Public Health, Boston, MA, USA
Email: aportnoy@mail.harvard.edu

Prof Jane J Kim
Degree: PhD
Affiliations: Center for Health Decision Science, Harvard T.H. Chan School of Public Health, Boston, MA, USA
Email: jkim@hsph.harvard.edu

Prof Marc Brisson
Degree: PhD
Affiliations: Centre de recherche du CHU de Québec - Université Laval, Québec, QC, Canada; Department of Social and Preventive Medicine, Université Laval, Québec, QC, Canada; MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, London, UK

95   Email: marc.brisson@crchudequebec.ulaval.ca
96
Prof Mark Jit
Joint Corresponding Author
Degree: PhD
Affiliations: Department of Infectious Disease Epidemiology, Faculty of Epidemiology and Population
Health, London School of Hygiene & Tropical Medicine, London, United Kingdom; Modelling and
Economics Unit, National Infection Service, Public Health England, London, United Kingdom; School
of Public Health, University of Hong Kong, Hong Kong Special Administrative Region, China
Email: mark.jit@lshtm.ac.uk

Abstract

Background

To eliminate cervical cancer as a public health problem, the World Health Organization currently recommends routine vaccination of adolescent girls with two doses of the human papillomavirus (HPV) vaccine before sexual initiation. However, many countries have yet to implement HPV vaccination because of financial or logistical barriers to delivering two doses outside the infant immunisation programme.

Methods

Using three independent HPV transmission models, we estimated the long-term health benefits and cost-effectiveness of one-dose versus two-dose HPV vaccination, in 188 countries, assuming that one dose of the vaccine gives either a shorter duration of full protection (20 or 30 years) or lifelong protection but lower vaccine efficacy (e.g., 80%) compared to two doses. We simulated routine vaccination with the 9-valent HPV vaccine in 10-year-old girls at 80% coverage for the years 2021–2120, with a one-year catch-up campaign of 11–14-year-old girls at 80% coverage in the first year of the programme.

Results

Over the years 2021–2120, one-dose vaccination at 80% coverage was projected to avert 112.9 million (range of medians: 75.8–176.2) and 148.0 million (111.6–187.6) cervical cancer cases assuming one dose of the vaccine confers 20 and 30 years of protection, respectively. Should one dose of the vaccine provide lifelong protection at 80% vaccine efficacy, 155.2 million (143.7–170.3) cervical cancer cases could be prevented. Around 65 to 889 additional girls would need to be vaccinated with the second dose to prevent one cervical cancer case, depending on the epidemiological profiles of the country. Across all income groups, the threshold cost for the second dose was low: from 0.85 (0.07–3.82) USD in low-income countries to 18.08 (-3.62–85.64) USD in high-income countries, assuming one-dose confers 30-year protection.

<u>Conclusions</u>

Results were consistent across the three independent models and suggest that one-dose

vaccination has similar health benefits to a two-dose programme while simplifying vaccine

delivery, reducing costs, and alleviating vaccine supply constraints. The second dose may be

cost-effective if there is a shorter duration of protection from one dose, cheaper vaccine

and vaccination delivery strategies, and high burden of cervical cancer.

Background

Cervical cancer is the fourth leading cause of cancer mortality among women globally with an estimated 570 000 new cases and 311 000 deaths in 2018, with the majority of deaths occurring in low- and middle-income countries (LMICs) (1). Persistent infection with high-risk genotypes of human papillomavirus (HPV) is a necessary precursor of cervical cancer.

Primary prevention of cervical cancer is available with four highly efficacious prophylactic vaccines—two 2-valent, one 4-valent, one 9-valent—that are currently licensed for protection against HPV infection (2–5). All protect against the two most carcinogenic HPV types, 16 and 18, which are responsible for 70% of cervical cancer cases globally (6–8). Some additionally protect against HPV types 6 and 11, which do not cause cancer but are responsible for most cases of anogenital warts, and against other high-risk types such as HPV 31, 33, 45, 52, and 58 (either directly or through cross-protection), which have been linked to a further 20% of cervical cancer cases (6–8).

Multiple analyses including the global Papillomavirus Rapid Interface for Modelling and Economics (PRIME) model developed in collaboration with the World Health Organization (WHO) (9,10) have found HPV vaccination to be cost-effective in almost all countries. The HPV vaccines were initially administered as a three-dose regimen over six months. In 2014, the WHO Strategic Advisory Group of Experts on Immunization reviewed the evidence for dose reduction and recommended a two-dose regimen for individuals below 15 years of age (11). With the availability of vaccines and screening tests that allow detection of both high-risk HPV types and neoplasias that are precursors to cervical cancer, the Secretary-General

of WHO has called for global elimination of cervical cancer as a public health problem, i.e.,

achieving the measurable global targets set by WHO (12). Current WHO guidelines

recommend that all countries vaccinate females aged 9–14 years against HPV (13).

Although some of these vaccines have been licensed for more than a decade, LMICs with

the highest incidence of cervical cancer are disproportionately less likely to introduce the

HPV vaccine into their routine immunisation programmes (9,14–16). High vaccine

procurement and delivery costs coupled with logistical constraints surrounding the delivery

of a two-dose regimen outside the infant vaccination schedule has hampered vaccine

introduction and uptake (17). Despite the financial support of Gavi, the Vaccine Alliance,

many LMICs have yet to introduce HPV vaccines into their routine programmes (18,19).

Since 2017, constrained supply of the 4-valent and 9-valent HPV vaccines has further

delayed vaccine introductions in many countries (20,21). Moreover, physical distancing

measures such as school closures and national lockdowns in response to the current COVID-

19 pandemic (22) have caused eligible populations to miss doses of HPV vaccine (21).

These financial, logistical, and supply constraints have motivated research into one-dose

vaccination schedules. If proven effective, one-dose HPV vaccination would simplify vaccine

delivery and lower costs of national vaccination programmes (18,23). It could also expedite

the introduction of HPV vaccines into national immunisation schedules for LMICs,

potentially protecting many more females against cervical cancer (19).

Evidence is emerging from immunogenicity trials, post-hoc analyses of efficacy trials, and

post-licensure observational studies to suggest that one dose of the HPV vaccine may

provide a high level of protection against incident and persistent HPV infections.

A systematic review of participants in six clinical trials who received only one dose of HPV

vaccination, because they did not complete their allocated schedules, suggests that this

schedule may be as effective as two doses in preventing HPV infection in up to seven years

of follow-up (24). However, evidence on the non-inferior efficacy of a single-dose schedule

from participants randomised to receive one dose has yet to emerge (expected in 2025).

Furthermore, antibody titres in immunogenicity trials were lower than in those receiving

two or three doses. While inferior antibody titres may not necessarily translate to inferior

protection, at this point, there is still uncertainty about the efficacy and durability of one-

dose vaccination.

Additionally, in the event that one-dose vaccination protection is slightly worse than two or

three doses, populations may still be almost as well protected through indirect (herd)

protection. Such effects can be examined using HPV transmission dynamic models. To date,

model-based analyses set in the United Kingdom (UK) (25), the United States (US), and

Uganda (26,27) suggest that one-dose schedules would be cost-effective and would prevent

almost as many cancers as two-dose or three-dose schedules if one dose confers at least 20

years of protection or has at least 80% efficacy against HPV 16/18 infection.

In this paper, we compare the impact and cost-effectiveness of one-dose versus two-dose

vaccination in 188 countries, assuming that one dose of the vaccine gives either shorter

duration of protection or lower vaccine efficacy compared to two doses. We use a hybrid

approach: firstly, we consider the age-specific impact that HPV vaccines may have using the

results of multiple independent HPV transmission dynamic models, and secondly,

extrapolate these effects to the remaining countries in the world using data on population

demographics and cervical cancer burden synthesised in a single model (PRIME).

Methods

To assess the extent to which one-dose HPV vaccination schedules will provide similar

protection and be cost-effective compared to two doses, we compared the impact of three

different strategies: (1) no HPV vaccination; (2) a one-dose HPV vaccination schedule in

which we assume that one dose of the HPV vaccine confers either 20 or 30 years of full

protection or 80% vaccine efficacy (VE) over the lifetime; and (3) a two-dose HPV

vaccination schedule in which two doses of the vaccine would provide lifetime protection at

100% VE. The minimum duration of protection in the waning scenarios for one-dose reflects

the availability of over 10 years of data from various studies—ESCUDDO trial (28,29), IARC

India post-randomisation analysis (30)—that do not show any evidence of waning of either

clinical or immunological protection (31).

Fig 1 provides an overview of the data sources and key steps of the modelling framework

described in the following sections. We synthesised the long-term population-wide impact

of HPV vaccination on cervical cancer incidence by age and time predicted by three

published transmission dynamic models: (i) the Public Health England (PHE) model, a

compartmental dynamic model set in the UK (32); (ii) the HPV-ADVISE model, an individual-

based dynamic model set in Uganda, Nigeria, India, Vietnam (27,33), and Canada (34,35);

and (iii) the Harvard model, a hybrid model that links two individual-based models, set in

the US, Uganda, El Salvador, and Nicaragua (14,36). In total, we combined results from 10

model-country scenarios. The models have been extensively reviewed and used to inform

vaccine policy (including by the UK's Joint Committee on Vaccination and Immunisation (37),

the World Health Organization's Strategic Advisory Group of Experts on Immunization

(11,38,39) and the US Advisory Committee on Immunization Practice (40–43). The models

stratify population by age, gender, and sexual activity-based risk group, as well as screening

behaviour-based risk group in the HPV-ADVISE and Harvard models. They capture HPV

natural history and disease, as well as HPV transmission as informed by country-specific

sexual behaviour surveys. More details about the models can be found in the

Supplementary Materials. For the scenarios where one dose confers a shorter duration of

protection (i.e., 20 or 30 years), we assume 100% VE, as suggested by clinical trial

populations (24,28,30,31). We modelled routine annual vaccination with the 9-valent

vaccine in 10-year-old girls to begin in 2021 and run uninterrupted until 2120. We also

included catch-up vaccination of girls aged 11–14 years in the first year of the programme.

Throughout, vaccine coverage was assumed to be 80%. In sensitivity analyses, we

investigated the impact of a one-dose vaccination schedule with a bivalent vaccine

(**Supplementary Materials**).

Fig 1. Overview of the data sources and the key steps of the modelling. To compare the impact and
cost-effectiveness of one-dose versus two-dose vaccination in 188 countries, we adopted a hybrid
approach. First, we synthesised the age-specific impact of HPV vaccines of three published
transmission dynamic models—PHE, HPV-ADVISE, Harvard—from 11 model-country settings.
Second, we derived the primary impact of vaccination using a static model (PRIME). Third, we
extrapolated the primary and secondary effects to the remaining countries in the world. Fourth, we
measured and compared population-level impact (e.g., cervical cancers averted, number of females
needed to be vaccinated, threshold costs of the first and second dose of the vaccine) for three

vaccine strategies: no HPV vaccination (the counterfactual); a one-dose HPV vaccination schedule in
which we assume that one dose of the vaccine provides either a shorter duration of protection (20
or 30 years) or lower vaccine efficacy (i.e., 80%) compared to two doses; and a two-dose HPV
vaccination schedule in which two doses of the vaccine provides lifetime protection.
263

Using PRIME, we then estimated the primary impact of a two-dose vaccination schedule,

without herd effects and waning immunity, in 188 countries. Full details of PRIME, including

model equations and updates, are available at (9,10). As PRIME is a static model, it cannot

estimate herd effects, nor can it capture the effect of waning vaccine immunity. Here, we

introduced a novel method which compares results from PRIME and the three dynamic

models—PHE, HPV-ADVISE, Harvard—set in nine countries—UK, US, Canada, Nigeria,

Uganda, India, Vietnam, El Salvador, and Nicaragua. We calculated the difference between

cervical cancer incidence predicted by PRIME and each of the dynamic models to derive the

secondary effects of vaccination, which is a combination of waning immunity (20/30-year

duration vs lifetime protection and lower vaccine efficacy) and herd effects at every age and

time-point. We then calculated the ratio of secondary to primary vaccine impact. By

assuming that the primary impact of a vaccine (i.e., vaccine with lifetime protection and no

herd effects) is different in every country as estimated by PRIME, we extrapolated the ratio

(secondary to primary) to other countries to project the secondary effects of vaccination,

using a similar approach as two comparative modelling analyses conducted by the WHO's

Cervical Cancer Elimination Modelling Consortium (14,44). A meta-analysis by Drolet and

colleagues showed a significant decrease in the prevalence of HPV 16 and 18 among women

aged 20–24 years (risk ratio [RR] 0.34, 95% CI 0.23–0.49) and 25–29 years (RR 0.63, 95% CI

0.41–0.97) (45). As most of the women in these age groups were unvaccinated, the meta-

analysis found evidence of similar herd effects more than four years after the introduction

of HPV vaccination.

Uncertainty in predictions was captured by generating multiple simulations from the three

dynamic models representing different plausible parameter sets. For the PHE model, 100

runs were simulated from the best-fitting parameter sets to capture uncertainty in the

duration of infection, duration of natural immunity, screening accuracy, the progression of

cervical cancer, age-specific prevalence, and the number of sexual partners. For HPV-

ADVISE, 1000 runs were simulated from 50 parameter sets that simultaneously fit country-

specific behavioural and epidemiological data. These 50 parameter sets illustrate the

uncertainty in sexual behaviour, HPV transmission, the natural history of HPV-related

diseases, and screening. For the Harvard model that reflect two sexual behaviour settings

(low- and high-HPV prevalence), 50 best-fitting dynamic transmission model parameter sets,

capturing variations in genotype- and sex-specific transmission probability, and genotype-

and sex-specific natural immunity, were propagated through four cervical carcinogenesis

models that have been previously calibrated (i.e., fit) to the US, Uganda, El Salvador, or

Nicaragua (14,36).

<u>Effectiveness and cost-effectiveness measures</u>

For each country, we estimated the number of cervical cancer cases, deaths, and disability-

adjusted life years (DALYs)—caused by HPV 16, 18, 31, 33, 45, 52, and 58—occurring under

each scenario by age and time since vaccination in females born in the years 2011–2110

(Section 1.8 **Supplementary materials**). We then compared the impact of a one-dose

schedule (giving 20/30 years protection or lifelong protection but at 80% initial VE) with no

vaccination, and a two-dose schedule (giving lifetime protection at 100% VE) with a one-

dose schedule. We calculated the number of females needed to vaccinate with one dose,

and the number of females needed to give an additional (i.e., second) dose, to avert one

cervical cancer case, death, or DALY. We also projected the threshold cost to pay for the

first and second dose of vaccine, which is the maximum that could be paid for the first dose

(compared to no vaccination) and second dose (compared to one dose only) for the

incremental cost-effectiveness ratio to remain below country-specific gross domestic

product (GDP) per capita (in 2017 USD). We used the GDP per capita estimates by the World

Bank (46), but also considered a lower threshold , i.e., 30–40% and 60–65% of GDP per

capita in low-income and middle- to high-income countries (47,48). The time horizon of the

analysis was from 2021 to 2120; we accrued all health benefits of vaccination up to the end

of the routine vaccination programme (i.e., the year 2120) or age 100 of all vaccinated

cohorts, whichever came first. Using modelled results from the 10 model-country pairs, we

projected the outcome measures in 188 countries and aggregated the results by World Bank

income groups. After projecting the various measures of effectiveness and cost-

effectiveness under the several vaccination scenarios, we compared the outcomes

generated with results from the 10 model-country pairs. After projecting the various

measures of effectiveness and cost-effectiveness under the vaccination scenarios, we

compared the outcomes generated with results from the 10 model-country scenarios. We

presented the results, aggregated by World Bank income groups (details in the

Supplementary Materials), as the median (and 80% uncertainty intervals (UI)) from each of

the model-country predictions. Both health outcomes and costs were discounted at 0% and

3% per year (49).

Results

In 188 countries over the years 2021–2120, the models projected that routine annual

vaccination of 10-year-old girls (plus a one-year catch-up campaign of girls aged 11–14

years) with one-dose of the 9-valent HPV vaccine at 80% coverage would avert 112.9 million

(range of medians: 75.8–176.2) and 148.0 million (111.6–187.6) cervical cancer cases should

one dose of the vaccine confer 20 and 30 years of protection, respectively (**Fig 2**; with the

equivalent cumulative and discounted benefits figures in the **Supplementary Materials**).

Under a scenario of one dose of the vaccine providing lifelong protection at 80% initial VE,

the models predicted that 155.2 million (143.7–170.3) cervical cancer cases would be

prevented (**Fig 2**). A one-dose schedule conferring 20 years of protection would avert 65.0%

(range of medians: 45.8–91.8%) of the cases averted by the vaccination schedule providing

lifelong protection at 100% VE (**Fig 3**). However, if the duration of protection increases to 30

years, a one-dose schedule would avert more cases at 87.9% (range of medians: 59.6–100%)

of the cases averted by the vaccination schedule providing lifelong protection at 100% VE

(**Fig 3**). Similarly, for the scenario where one dose of the vaccine provides lifelong protection

but at lower VE (of 80%), most of the cases (88.3% (range of medians: 86.2–96.7%)) can still

be averted (**Fig 3**).

Fig 2. Cervical cancers averted by routine one-dose HPV vaccination by country income groups.
The lines represent the median projections of the 10 model-country settings: the PHE model in
black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. The
grey area corresponds to the additional cases averted in the vaccinated cohort after the 100 years of
routine vaccination. Cancers averted were discounted at 0%. Only cervical cancer caused by HPV 16,
18, 31, 33, 45, 52, and 58, which could be averted by the 9-valent HPV vaccine, were considered.

Fig 3. Cervical cancers averted by routine one-dose HPV vaccination as a proportion of cervical
cancers averted by routine HPV vaccination programmes conferring lifelong protection at 100%
vaccine efficacy. The median percentage (intervals: 10–90th percentile) of cancers averted by a
one-dose schedule compared to a two-dose programme of the 10 model-country settings: the PHE
model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in
blue. Health outcomes were discounted at 0%. Only cervical cancer caused by HPV 16, 18, 31, 33, 45,
52, and 58, which could be averted by the 9-valent HPV vaccine, were considered.

Due to large disparities in age-standardised cervical cancer incidence across country income

groups in 2021, the number of cases averted by routine vaccination programmes is higher in

low-income countries (31.9 million (range of medians: 21.2–48.6), if one-dose confers 20

years of protection) than in high-income countries (4.8 million (range of medians: 3.5–6.9)).

More cervical cancers could be averted if one dose of the vaccine confers a longer duration

of protection, i.e., at 30 years or lifelong but lowered VE. Assuming waning of protection at

20 years (on average) after vaccination, the PHE model parameterised with data from the

UK projected that a one-dose schedule could avert 91.8% (80%UI 76.2–99.8%) of the cases

averted by a two-schedule vaccination schedule. However, the HPV-ADVISE and Harvard

models, mostly parameterised with data from LMICs, projected that 61.0% (range of

medians: 45.8–87.6%) could be averted (**Fig 3**).

The models consistently projected that fewer girls need to be vaccinated with the first dose

to prevent one cervical cancer case in low-income countries (32 (range of medians: 23–51))

than middle-income (43 (range of medians: 31–69)) and high-income countries (83 (range of

medians: 67–122)) if one-dose confers 20 years of protection (**Fig 4A–C**). However,

variations across models were observed for the projections of the number of girls needed to

be vaccinated with the second dose to prevent one cervical cancer case. Compared to the

HPV-ADVISE and Harvard models, the PHE model projected that more girls need to be

vaccinated with the second dose to avert one cervical cancer case when the protection from

one dose of the vaccine wanes 20 years after vaccination (**Fig 4A–D**; 889 (80%UI 93–27 700)

girls if one dose confers 20 years of protection). However, if one-dose confers lifelong

protection but at lowered VE, the differences between the PHE, HPV-ADVISE and Harvard

models decrease. When we discounted health outcomes, the model predicted that more

girls need to be vaccinated to avert one cervical cancer case (**Fig 4E–H**).

389
Fig 4. Number of girls needed to be vaccinated with the first and second dose to avert one
additional cervical cancer case by income group. The lines represent the median projections of the
10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the
Harvard model-country pairs in blue. The grey area corresponds to the additional cases averted in
the vaccinated cohort after the 100 years of routine vaccination. Health outcomes were discounted
at 3% (panels A–D) and 0% (panels E–H).

Across all income groups, the threshold (i.e., maximum) cost for the second dose to remain

cost-effective was low—from 0.85 (range of medians: 0.07–3.82) USD in low-income

countries to 18.08 (range of medians: -3.62–85.64) USD in high-income countries if one-

dose confers 30 year protection to—as few additional cancers would be averted with a

longer duration of protection (≥30 years) or higher VE (>80%). With a higher GDP per capita,

middle- and high-income countries have a higher threshold cost (**Fig 5**). However, if one-

dose confers ≤20 years of protection, the threshold cost for the second dose to remain cost-

effective is borderline at 3.24 (range of medians: 0.34–5.35) USD in low-income countries

and 62.93 (range of medians: 12.63–117.45) USD in high-income countries, suggesting that

duration of protection remains the main driver of uncertainty.

407
Fig 5. Threshold cost to pay for the first and second dose of vaccine by country income groups. The
threshold cost is the maximum that could be paid for the first dose (compared to no vaccination)

and second dose (compared to one dose only) for the incremental cost-effectiveness ratio to remain
below the cost-effectiveness threshold. Two cost-effectiveness thresholds are presented: a lower
threshold as suggested by Jit (2020) in panels A–D and country gross domestic product (GDP) per
capita (in 2017 USD) in panels E–H. The lower cost-effectiveness threshold presented in panels A–D
is 30–40% and 60–65% of GDP per capita in low-income and middle- to high-income countries,
respectively. Cost and health outcomes were discounted at 3% and 0%, respectively.

Discussion

In this study, three independent transmission dynamic models projected consistent results

suggesting that routine one-dose HPV vaccine programmes at 80% coverage worldwide

could provide a high level of population protection and be cost-effective. We considered

three assumptions of the one-dose schedule: one dose of the HPV vaccine confers either 20

or 30 years of protection at full VE, or lifelong protection but at 80% VE. Across all

assumptions, one-dose schedules provide large population impacts on cervical cancer, while

the difference in population impact of the one-dose versus two-dose vaccination schedule is

small if one dose confers ≥30 years of protection or lifelong protection but at 80% VE. This

underscores the significant potential public health impact of the one-dose vaccination

schedule if vaccine uptake is high across all countries (21).

Although trials (28–30) and post-randomisation analyses (30) suggest that the duration of

protection of one dose of the vaccine is more than ten years, it is uncertain how long

vaccinated individuals will remain protected and how the vaccine would wane beyond the

first decade. The threshold duration of protection for a one-dose schedule to avert the

majority of vaccine-preventable cancers is associated with the ages at which individuals are

vaccinated and reach peak sexual activity, which varies between countries. If one dose of

the vaccine confers ≤20 years of protection, giving the second dose may have a larger health

impact, especially in settings where HPV transmission persists decades after vaccination.

However, if one dose provides a longer duration of protection (≥30 years), administering the

second dose will bring about few health gains at potentially high costs. Hence, decisions on

offering the second dose should account for the duration of protection provided by the first

dose, whether it covers the peak years of sexual activity and HPV transmission, and the

costs of delivering the additional dose. The second dose becomes more cost-effective if the

protection from one dose is less than 20 years, the costs of the vaccine and delivering it are

lower than current reported costs, and/or the local burden of cervical cancer is high.

Our comparison of one- and two-dose vaccination schedules is motivated by several

advantages of a one-dose schedule. Firstly, many LMICs have yet to implement national HPV

vaccination programmes because of the challenges of delivering two vaccine doses to

adolescent females (17). Compared to two-dose HPV vaccination, a one-dose HPV

vaccination schedule would be cheaper and easier to implement (e.g., no follow-up of

vaccinated individuals would be required), potentially enabling more LMICs to introduce

HPV vaccine into national immunisation schedules (21,23). More recently, HPV vaccine

implementation in LMICs has been delayed due to constraints in HPV vaccine supply (20,21).

Our model-based analysis predicts that routinely vaccinating 10-year-old girls at 80%

coverage in LICs could result in four times (population-adjusted) more cervical cases averted

than in high-income countries. Under our one-dose assumptions, routine one-dose HPV

vaccination programmes could protect up to 155 million females against cervical cancer

globally over the years 2021–2120.

Secondly, the COVID-19 pandemic has disrupted several routine immunisation programmes

(50–52), including HPV vaccination (21,53). Abbas and colleagues predicted that the benefits

of resuming routine childhood immunisation services outweigh the risk of being infected

with COVID-19 during the vaccination visits (52), reinforcing WHO's call for all countries to

continue routine immunisation services safely (54). With physical distancing measures such

as school closures and national lockdowns being implemented in many countries to cope

with the COVID-19 pandemic (22), health officials grapple with reconfiguring school-based

HPV vaccine delivery (53–55). Compared to the two-dose vaccination schedule, a one-dose

schedule would further minimise interactions between vaccinees and health workers,

simplifying vaccine delivery while also decreasing SARS-CoV-2 exposure.

The lack of country-specific behavioural, virological, and clinical data in many countries

limits fitting transmission dynamic models individually to most countries (56). However, in

this comparative modelling study, we synthesised results from three published dynamic

models based in nine countries, covering high-, middle- and low-income settings across

three continents and a wide variety of epidemiological characteristics for HPV transmission

and cervical cancer (14,56). Our approach provides a common framework using PRIME for

population demographics, cervical cancer burden, and impact/cost-effectiveness

calculations, while varying representation of HPV transmission and cervical cancer natural

history across the three dynamic models. We then extrapolated the age- and time-

dependent ratio of the secondary to primary impacts of vaccine strategies to other

countries. While there may be considerable uncertainty around extrapolating this ratio to

another country, the use of 10 model-country pairs lends confidence that we are likely to

have captured the range of possible outcomes for most countries. More precise estimates

would require fitting these models to additional specific countries, for which calibration

data are not available (56,57).

Our model projections of vaccine impact also involve other sources of uncertainty that we

did not explicitly quantify. The PRIME model uses country-specific cervical cancer burden

from the Global Cancer Incidence, Mortality and Prevalence (GLOBOCAN) database (58),

which may underestimate the full burden of HPV-related disease, and thus vaccine impact,

in LMICs (14). In this study, we only assessed the effect of HPV vaccination on cervical

cancers. If we also accounted for the vaccine impact on other HPV-related cancers, we

would anticipate a greater value of HPV vaccination programmes (26,59). However, the

paucity of data on the efficacy of one dose on non-cervical cancers complicates the analysis

evaluating their vaccine impact. Because the health gains from the second dose are small,

any minor variations in gains will amplify the variability in the number needed to vaccinate

with the additional dose. Finally, we project the impact of HPV vaccination on cervical

cancers over the next century. Over the past decades, we have witnessed substantial

demographic (60) and behavioural changes (61,62) with extraordinary improvements in

public health (63). In 2020, the COVID-19 pandemic has caused substantial disruptions to

population demography (64) and sexual behaviour (65), with uncertainty around the longer-

term consequences of such disruption. Moreover, over the next century, we expect to see

continued advancements in pre-cancer screening and treatment services, which will further

decrease cervical cancer incidence. Such uncertainties in life expectancy, population, and

economic forecasts have significant implications for our predictions.

Conclusion

Under the scenarios where a single HPV vaccine dose confers more than 30 years of protection or 80% efficacy with lifelong protection, routine one-dose HPV vaccination provides the majority of health benefits to the two-dose programme while simplifying vaccine delivery, reducing costs, and circumventing vaccine supply constraints. The second dose may be cost-effective if there is a shorter duration of protection from one dose, cheaper vaccine and vaccination delivery strategies, and high burden of cervical cancer. These results are fairly consistent when projected from three independent transmission dynamic models used in nine countries. The outcomes of our comparative modelling analysis contribute to the extensive evidence base, including emerging evidence from the single-dose HPV vaccine trials and observational studies, which would be beneficial to policymakers when they consider HPV vaccination in their populations.

References

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin. 2018;

2. Garland SM, Hernandez-Avila M, Wheeler CM, Perez G, Harper DM, Leodolter S, et al. Quadrivalent Vaccine against Human Papillomavirus to Prevent Anogenital Diseases. N Engl J Med [Internet]. 2007 May 10 [cited 2020 Oct 30];356(19):1928–43. Available from: www.nejm.org

3. Paavonen J, Naud P, Salmerón J, Wheeler CM, Chow SN, Apter D, et al. Efficacy of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine against cervical infection and precancer caused by oncogenic HPV types (PATRICIA): final analysis of a double-blind, randomised study in young women. Lancet. 2009 Jul 25;374(9686):301–14.

4. Joura EA, Giuliano AR, Iversen O-E, Bouchard C, Mao C, Mehlsen J, et al. A 9-Valent HPV Vaccine against Infection and Intraepithelial Neoplasia in Women. N Engl J Med [Internet]. 2015 Feb 19 [cited 2020 Oct 30];372(8):711–23. Available from: https://www.nejm.org/doi/full/10.1056/NEJMoa1405044

5. Qiao YL, Wu T, Li RC, Hu YM, Wei LH, Li CG, et al. Efficacy, Safety, and Immunogenicity of an Escherichia coli-Produced Bivalent Human Papillomavirus Vaccine: An Interim Analysis of a Randomized Clinical Trial. J Natl Cancer Inst [Internet]. 2020 Feb 1 [cited 2020 Nov 4];112(2):145–53. Available from: https://academic.oup.com/jnci/article/112/2/145/5488952

6. de Sanjose S, Quint WGV, Alemany L, Geraets DT, Klaustermeier JE, Lloveras B, et al. Human papillomavirus genotype attribution in invasive cervical cancer: a retrospective cross-sectional worldwide study. Lancet Oncol. 2010;

7. Serrano B, Alemany L, Tous S, Bruni L, Clifford GM, Weiss T, et al. Potential impact of a nine-valent vaccine in human papillomavirus related cervical disease. Infect Agent Cancer [Internet]. 2012 Dec 29 [cited 2020 Oct 30];7(1):1–13. Available from: https://link.springer.com/articles/10.1186/1750-9378-7-38

8. de Martel C, Georges D, Bray F, Ferlay J, Clifford GM. Global burden of cancer attributable to infections in 2018: a worldwide incidence analysis. Lancet Glob Heal [Internet]. 2020 Feb 1 [cited 2020 Oct 30];8(2):e180–90. Available from: www.thelancet.com/lancetgh

9. Jit M, Brisson M, Portnoy A, Hutubessy R. Cost-effectiveness of female human papillomavirus vaccination in 179 countries: A PRIME modelling study. Lancet Glob Heal [Internet]. 2014 [cited 2020 Oct 30];2(7):e406–14. Available from: https://pubmed.ncbi.nlm.nih.gov/25103394/

10. Abbas KM, van Zandvoort K, Brisson M, Jit M. Effects of updated demography, disability weights, and cervical cancer burden on estimates of human papillomavirus vaccination impact at the global, regional, and national levels: a PRIME modelling study. Lancet Glob Heal [Internet]. 2020 Apr 1 [cited 2020 Oct 30];8(4):e536–44. Available from: http://gco.iarc.fr

11. World Health Organization. Human papillomavirus vaccines: WHO position paper, October 2014. Wkly Epidemiol Rec Relev épidémiologique Hebd. 2014;89(43):465–91.

12. World Health Organization. WHO | WHO leads the way towards the elimination of cervical cancer as a public health concern. World Health Organization; 2018.

564   13.   Simms KT, Steinberg J, Caruana M, Smith MA, Lew J Bin, Soerjomataram I, et al.
565   Impact of scaled up human papillomavirus vaccination and cervical screening and the
566   potential for global elimination of cervical cancer in 181 countries, 2020–99: a
567   modelling study. Lancet Oncol. 2019;

568   14.   Brisson M, Kim JJ, Canfell K, Drolet M, Gingras G, Burger EA, et al. Impact of HPV
569   vaccination and cervical screening on cervical cancer elimination: a comparative
570   modelling analysis in 78 low-income and lower-middle-income countries. Lancet
571   [Internet]. 2020 Feb 22 [cited 2020 Oct 30];395(10224):575–90. Available from:
572   https://doi.org/10.1016/

573   15.   PATH. Global HPV Vaccine Introduction Overview: projected and current national
574   introductions, demonstration/pilot projects, gender-neutral vaccination programs,
575   and global HPV vaccine introduction maps (2006–2022). [Internet]. 2020 May [cited
576   2020 Oct 30]. Available from: https://www.path.org/resources/global-hpv-vaccine-
577   introduction-overview/

578   16.   WHO. Immunization, vaccines and biologicals: data, statistics and graphics. WHO
579   [Internet]. 2020 [cited 2020 Oct 30]; Available from:
580   http://www.who.int/immunization/monitoring_surveillance/data/en/

581   17.   Gallagher KE, LaMontagne DS, Watson-Jones D. Status of HPV vaccine introduction
582   and barriers to country uptake. Vaccine. 2018 Aug 6;36(32):4761–7.

583   18.   LaMontagne DS, Bloem PJN, Brotherton JML, Gallagher KE, Badiane O, Ndiaye C.
584   Progress in HPV vaccination in low- and lower-middle-income countries. Int J Gynecol
585   Obstet [Internet]. 2017 Jul 1 [cited 2020 Oct 30];138:7–14. Available from:
586   http://doi.wiley.com/10.1002/ijgo.12186

587   19.   Kreimer AR, Cernuschi T, Rees H, Saslow D, Porras C, Schiller J. Prioritisation of the
588   human papillomavirus vaccine in a time of constrained supply. Vol. 4, The Lancet
589   Child and Adolescent Health. Elsevier B.V.; 2020. p. 349–51.

590   20.   World Health Organization. Global market study: HPV vaccines. World Health
591   Organization; 2019.

592   21.   World Health Organization. Meeting of the Strategic Advisory Group of Experts on
593   Immunization, April 2022: conclusions and recommendations – Réunion du Groupe
594   stratégique consultatif d'experts sur la vaccination, avril 2022: conclusions et
595   recommandations. Wkly Epidemiol Rec [Internet]. 2022 [cited 2022 Jun
596   20];97(24):261–76. Available from: https://apps.who.int/iris/handle/10665/356580

597   22.   Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and
598   management practices during coronavirus outbreaks including COVID-19: a rapid
599   systematic review. Lancet Child Adolesc Heal [Internet]. 2020 [cited 2020 Apr
600   22];0(0). Available from:
601   https://linkinghub.elsevier.com/retrieve/pii/S235246422030095X

602   23.   Barnabas R V., Brown ER, Onono MA, Bukusi EA, Njoroge B, Winer RL, et al. Efficacy of
603   Single-Dose Human Papillomavirus Vaccination among Young African Women. NEJM
604   Evid [Internet]. 2022 Apr 11 [cited 2022 Jun 20];1(5). Available from:
605   https://evidence.nejm.org/doi/abs/10.1056/EVIDoa2100056

606   24.   Whitworth HS, Gallagher KE, Howard N, Mounier-Jack S, Mbwanji G, Kreimer AR, et
607   al. Efficacy and immunogenicity of a single dose of human papillomavirus vaccine
608   compared to no vaccination or standard three and two-dose vaccination regimens: A
609   systematic review of evidence from clinical trials. Vol. 38, Vaccine. Elsevier Ltd; 2020.
610   p. 1302–14.

611   25.   Jit M, Brisson M, Laprise JF, Choi YH. Comparison of two dose and three dose human
612   papillomavirus vaccine schedules: Cost effectiveness analysis based on transmission
613   model. BMJ. 2015 Jan 7;350. Available from:
614   http://www.bmj.com/content/350/bmj.g7584?tab=related#datasupp
615   26.   Burger EA, Campos NG, Sy S, Regan C, Kim JJ. Health and economic benefits of single-
616   dose HPV vaccination in a Gavi-eligible country. Vaccine. 2018 Aug 6;36(32):4823–9.
617   27.   Drolet M, Laprise JF, Martin D, Jit M, Bénard É, Gingras G, et al. Optimal human
618   papillomavirus (HPV) vaccination strategies to prevent cervical cancer in low- and
619   middle-income countries in the context of limited resources: A mathematical
620   modeling analysis. Lancet Infect Dis. 2021;
621   28.   Porras C, Sampson JN, Herrero R, Gail MH, Cortés B, Hildesheim A, et al. Rationale
622   and design of a double-blind randomized non-inferiority clinical trial to evaluate one
623   or two doses of vaccine against human papillomavirus including an epidemiologic
624   survey to estimate vaccine efficacy: The Costa Rica ESCUDDO trial. Vaccine. 2022 Jan
625   3;40(1):76–88.
626   29.   ClinicalTrials.gov. Comparing One or Two Doses of the Human Papillomavirus Vaccine
627   for the Prevention of Human Papillomavirus Infection, ESCUDDO Study:
628   NCT03180034 [Internet]. [cited 2022 Feb 3]. Available from:
629   https://clinicaltrials.gov/ct2/show/NCT03180034
630   30.   Basu P, Malvi SG, Joshi S, Bhatla N, Muwonge R, Lucas E, et al. Vaccine efficacy
631   against persistent human papillomavirus (HPV) 16/18 infection at 10 years after one,
632   two, and three doses of quadrivalent HPV vaccine in girls in India: a multicentre,
633   prospective, cohort study. Lancet Oncol. 2021 Nov 1;22(11):1518–29.
634   31.   Single-dose-HPV-Vaccine-Evaluation-Consortium. Review of the current published
635   evidence on single-dose HPV vaccination 3rd edition 2020. 2020.
636   32.   Choi YH, Jit M, Gay N, Cox A, Garnett GP, Edmunds WJ. Transmission dynamic
637   modelling of the impact of human papillomavirus vaccination in the United Kingdom.
638   Vaccine [Internet]. 2010 May 28 [cited 2020 Oct 30];28(24):4091–102. Available from:
639   https://pubmed.ncbi.nlm.nih.gov/19909831/
640   33.   Brisson M, Laprise JF, Martin D, Drolet M, Van De Velde N, Boily MC. Technical
641   Appendix HPV-ADVISE LMIC. 2017.
642   34.   Van De Velde N, Boily MC, Drolet M, Franco EL, Mayrand MH, Kliewer E V., et al.
643   Population-level impact of the bivalent, quadrivalent, and nonavalent human
644   papillomavirus vaccines: A model-based analysis. J Natl Cancer Inst [Internet]. 2012
645   Nov 21 [cited 2020 Oct 30];104(22):1712–23. Available from: http://www.marc-
646   brisson.net/HPVadvise.pdf
647   35.   Brisson M, Van De Velde N, Drolet M, Laprise JF, Boily MC. Technical Appendix HPV-
648   ADVISE Canada [Internet]. 2012. Available from: http://www.marc-
649   brisson.net/HPVadvise.pdf
650   36.   Burger EA, Smith MA, Killen J, Sy S, Simms KT, Canfell K, et al. Projected time to
651   elimination of cervical cancer in the USA: a comparative modelling study. Lancet
652   Public Heal [Internet]. 2020 Apr 1 [cited 2021 Jan 12];5(4):e213–22. Available from:
653   www.thelancet.com/
654   37.   Hall AJ. The United Kingdom Joint Committee on Vaccination and Immunisation.
655   Vaccine. 2010 Apr 19;28(SUPPL. 1):A54–7.
656   38.   World Health Organization. Human papillomavirus vaccines: WHO position paper,
657   May 2017. Weekly Epidemiological Record= Relevé épidémiologique hebdomadaire

[Internet]. 2017; Available from:
https://www.who.int/immunization/policy/position_papers/hpv/en/

39. World Health Organization. Meeting of the Strategic Advisory Group of Experts on
immunization, October 2019 – conclusions and recommendations. Wkly Epidemiol
Rec Relev épidémiologique Hebd [Internet]. 2019 [cited 2020 Nov 3];49(47):541–560.
Available from: http://www.who.int/wer/2019/wer9447/en/

40. Laprise JF, Chesson HW, Markowitz LE, Drolet M, Martin D, Bénard É, et al.
Effectiveness and cost-effectiveness of human papillomavirus vaccination through
age 45 years in the United States. Ann Intern Med. 2020 Jan 7;172(1):22–9.

41. Brisson M, Laprise JF, Chesson HW, Drolet M, Malagón T, Boily MC, et al. Health and
Economic Impact of Switching from a 4-Valent to a 9-Valent HPV Vaccination Program
in the United States. J Natl Cancer Inst. 2016 Jan 1 [cited 2021 Jan 12];108(1).
Available from: https://pubmed.ncbi.nlm.nih.gov/26438574/

42. Chesson HW, Laprise JF, Brisson M, Markowitz LE. Impact and Cost-effectiveness of 3
Doses of 9-Valent Human Papillomavirus (HPV) vaccine among US females previously
vaccinated with 4-valent hpv vaccine. J Infect Dis. 2016 Jun 1;213(11):1694–700.
Available from: /pmc/articles/PMC4857476/?report=abstract

43. Laprise JF, Markowitz LE, Chesson HW, Drolet M, Brisson M. Comparison of 2-dose
and 3-dose 9-valent human papillomavirus vaccine schedules in the United States: A
cost-effectiveness analysis. J Infect Dis. 2016 Sep 1;214(5):685–8.

44. Canfell K, Kim JJ, Brisson M, Keane A, Simms KT, Caruana M, et al. Mortality impact of
achieving WHO cervical cancer elimination targets: a comparative modelling analysis
in 78 low-income and lower-middle-income countries. Lancet. 2020 Feb
22;395(10224):591–603. Available from:
http://www.thelancet.com/article/S0140673620301574/fulltext

45. Drolet M, Bénard É, Pérez N, Brisson M, Ali H, Boily MC, et al. Population-level impact
and herd effects following the introduction of human papillomavirus vaccination
programmes: updated systematic review and meta-analysis. Lancet. 2019 Aug
10;394(10197):497–509. Available from:
http://www.thelancet.com.libproxy1.nus.edu.sg/article/S0140673619302983/fulltext

46. World Bank. GDP per capita (current US$) | Data. Available from:
https://data.worldbank.org/indicator/NY.GDP.PCAP.CD

47. Jit M. Informing Global Cost-Effectiveness Thresholds Using Country Investment
Decisions: Human Papillomavirus Vaccine Introductions in 2006-2018. Value Heal.
2020 Oct 16;

48. Ochalek J, Abbas K, Claxton K, Jit M, Lomas J. Assessing the value of human
papillomavirus vaccination in Gavi-eligible low-income and middle-income countries.
BMJ Glob Heal. 2020 Oct 20;5(10):e003006. Available from:
https://gh.bmj.com/lookup/doi/10.1136/bmjgh-2020-003006

49. World Health Organization. WHO Guide on Standardization of Economic Evaluations
of Immunization Programmes. WHO. 2019 Oct; Available from:
http://www.who.int/immunization/documents/who_ivb_19.10/en/

50. World Health Organization. Pulse survey on continuity of essential health services
during the COVID-19 pandemic: interim report, 27 August 2020. Available from:
https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-
2020.1

51. Nelson R. COVID-19 disrupts vaccine delivery. Lancet Infect Dis. 2020 May

1;20(5):546. Available from: www.thelancet.com/infection

52.   Abbas K, Procter SR, van Zandvoort K, Clark A, Funk S, Mengistu T, et al. Routine
childhood immunisation during the COVID-19 pandemic in Africa: a benefit–risk
analysis of health benefits versus excess risk of SARS-CoV-2 infection. Lancet Glob
Heal. 2020 Oct 1;8(10):e1264–72. Available from: www.thelancet.com/lancetgh

53.   UNICEF Supply Division. Human papillomavirus (HPV) vaccine: supply and demand
update | UNICEF Supply Division. Available from:
https://www.unicef.org/supply/reports/human-papillomavirus-hpv-vaccine-supply-
and-demand-update

54.   World Health Organization. WHO and UNICEF warn of a decline in vaccinations during
COVID-19. Available from: https://www.who.int/news/item/15-07-2020-who-and-
unicef-warn-of-a-decline-in-vaccinations-during-covid-19

55.   Suwantika AA, Boersma C, Postma MJ. The potential impact of COVID-19 pandemic
on the immunization performance in Indonesia. Expert Rev Vaccines. 2020 Aug
2;19(8):687–90. Available from:
https://www.tandfonline.com/doi/full/10.1080/14760584.2020.1800461

56.   Jit M, Levin C, Brisson M, Levin A, Resch S, Berkhof J, et al. Economic analyses to
support decisions about HPV vaccination in low- and middle-income countries: A
consensus report and guide for analysts. Vol. 11, BMC Medicine. BioMed Central;
2013. p. 23. Available from:
http://bmcmedicine.biomedcentral.com/articles/10.1186/1741-7015-11-23

57.   Kim JJ, Brisson M, Edmunds WJ, Goldie SJ. Modeling Cervical Cancer Prevention in
Developed Countries. Vaccine. 2008 Aug 19;26(SUPPL. 10):K76. Available from:
/pmc/articles/PMC2769256/?report=abstract

58.   International Agency for Research on Cancer. Global Cancer Observatory. Available
from: https://gco.iarc.fr/

59.   Gillison ML, Chaturvedi AK, Lowy DR. HPV prophylactic vaccines and the potential
prevention of noncervical cancers in both men and women. Vol. 113, Cancer. NIH
Public Access; 2008. p. 3036–46. Available from:
/pmc/articles/PMC6264789/?report=abstract

60.   United Nations Department of Economic and Social Affairs Population Division. World
Population Prospects. 2019. Available from: https://population.un.org/wpp/

61.   Baussano I, Lazzarato F, Brisson M, Franceschi S. Human papillomavirus vaccination at
a time of changing sexual behavior. Emerg Infect Dis. 2016 Jan 1;22(1):18–23.
Available from: /pmc/articles/PMC4696692/?report=abstract

62.   Wellings K, Collumbien M, Slaymaker E, Singh S, Hodges Z, Patel D, et al. Sexual
behaviour in context: a global perspective. Vol. 368, Lancet. Elsevier Limited; 2006. p.
1706–28. Available from: https://pubmed.ncbi.nlm.nih.gov/17098090/

63.   Cutler D, Miller G. The role of public health improvements in health advances: The
twentieth-century United States. Demography. 2005;42(1):1–22. Available from:
https://link-springer-com.ez.lshtm.ac.uk/article/10.1353/dem.2005.0002

64.   Goldstein JR, Lee RD. Demographic perspectives on the mortality of COVID-19 and
other epidemics. Proc Natl Acad Sci U S A. 2020 Sep 8;117(36):22035–41. Available
from: www.pnas.org/cgi/doi/10.1073/pnas.2006392117

65.   Ibarra FP, Mehrad M, Di Mauro M, Peraza Godoy MF, Cruz EG, Nilforoushzadeh MA,
et al. Impact of the COVID-19 pandemic on the sexual behavior of the population. The
vision of the east and the west. Vol. 46, International Braz J Urol. Brazilian Society of

Urology; 2020. p. 104–12. Available from: http://orcid.org/0000-0003-4687-7353
753
754
755

Abbreviations

DALY   Disability-adjusted life year
GDP   Gross domestic product
HPV   Human papillomavirus
LMICs Low- and middle-income countries
PHE   Public Health England
PRIME Papillomavirus Rapid Interface for Modelling and Economics
RR   Risk ratios
UI   Uncertainty intervals
UK   United Kingdom
US   United States
USD   United States Dollar
VE   Vaccine efficacy
WHO   World Health Organization

Funding

Financial support for this project was provided by PATH on behalf of the Single-Dose HPV Vaccine Evaluation Consortium which includes Harvard University (Harvard), London School of Hygiene & Tropical Medicine (LSHTM), PATH, US National Cancer Institute (NCI), University of British Columbia, Canada (UBC), CHU de Québec-Université Laval, Quebec (CHU), University of Witwatersrand Reproductive Health and HIV Institute (Wits RHI), US Centers for Disease Control and Prevention (CDC), and the World Health Organization (WHO). The work was also part funded by the Bill & Melinda Gates Foundation (OPP1157270) and the Fonds de recherche du Québec - Santé (FRQS) Research Scholars award (to MB), and a Foundation scheme grant from the Canadian Institutes of Health Research (CIHR; grant number FDN-143283). This research was also enabled in part by support provided by Compute Canada (www.computecanada.ca).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' Contributions

KP and MJ conceptualised the study, curated and analysed the data, interpreted the findings, and drafted manuscript. YHC, EB, EAB, LH, and JFL curated and analysed the data, interpreted the findings, and were major contributors in writing the manuscript. JJK and MB interpreted the findings and were major contributors in writing the manuscript. MCR, MD, SS, KA, and AP analysed the data, interpreted the findings, and reviewed the manuscript. All authors read and approved the final manuscript.

Declaration of interests

783 We declare no competing interests.

Data availability

785 All analysis codes are available at https://github.com/kieshaprem/hpv-1-dose.

Acknowledgement

787 We thank members of the Single-Dose HPV Vaccine Evaluation Consortium for comments
788 and helpful discussion on this work.

789
790
791

Figure captions

Fig 1. Overview of the data sources and the key steps of the modelling. To compare the impact and cost-effectiveness of one-dose versus two-dose vaccination in 188 countries, we adopted a hybrid approach. First, we synthesised the age-specific impact of HPV vaccines of three published transmission dynamic models—PHE, HPV-ADVISE, Harvard—from 10 model-country settings. Second, we derived the primary impact of vaccination using a static model (PRIME). Third, we extrapolated the primary and secondary effects to the remaining countries in the world. Fourth, we measured and compared population-level impact (e.g., cervical cancers averted, number of females needed to be vaccinated, threshold costs of the first and second dose of the vaccine) for three vaccine strategies: no HPV vaccination (the counterfactual); a one-dose HPV vaccination schedule in which we assume that one dose of the vaccine provides either a shorter duration of protection (20 or 30 years) or lower vaccine efficacy (i.e., 80%) compared to two doses; and a two-dose HPV vaccination schedule in which two doses of the vaccine provides lifetime protection.

Fig 2. Cervical cancers averted by routine one-dose HPV vaccination by country income groups. The lines represent the median projections of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. The grey area corresponds to the additional cases averted in the vaccinated cohort after the 100 years of routine vaccination. Cancers averted were discounted at 0%. Only cervical cancer caused by HPV 16, 18, 31, 33, 45, 52, and 58, which could be averted by the 9-valent HPV vaccine, were considered.

Fig 3. Cervical cancers averted by routine one-dose HPV vaccination as a proportion of cervical cancers averted by routine HPV vaccination programmes conferring lifelong protection at 100% vaccine efficacy. The median percentage (intervals: 10–90th percentile) of cancers averted by a one-dose schedule compared to a two-dose programme of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. Health outcomes were discounted at 0%. Only cervical cancer caused by HPV 16, 18, 31, 33, 45, 52, and 58, which could be averted by the 9-valent HPV vaccine, were considered.

Fig 4. Number of girls needed to be vaccinated with the first and second dose to avert one additional cervical cancer case by income group. The lines represent the median projections of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. The grey area corresponds to the additional cases averted in the vaccinated cohort after the 100 years of routine vaccination. Health outcomes were discounted at 3% (panels A–D) and 0% (panels E–H).

Fig 5. Threshold cost to pay for the first and second dose of vaccine by country income groups. The threshold cost is the maximum that could be paid for the first dose (compared to no vaccination) and second dose (compared to one dose only) for the incremental cost-effectiveness ratio to remain below the cost-effectiveness threshold. Two cost-effectiveness thresholds are presented: a lower threshold as suggested by Jit (2020) in panels A–D and country gross domestic product (GDP) per capita (in 2017 USD) in panels E–H. The lower cost-effectiveness threshold presented in panels A–D is 30–40% and 60–65% of GDP per capita in low-income and middle- to high-income countries, respectively. Cost and health outcomes were discounted at 3% and 0%, respectively.

 Figures

Figure 1. Overview of the data sources and the key steps of the modelling. To compare the impact and cost-effectiveness of one-dose versus two-dose vaccination in 188 countries, we adopted a hybrid approach. First, we synthesised the age-specific impact of HPV vaccines of three published transmission dynamic models—PHE, HPV-ADVISE, Harvard—from 10 model-country settings. Second, we derived the primary impact of vaccination using a static model (PRIME). Third, we extrapolated the primary and secondary effects to the remaining countries in the world. Fourth, we measured and compared population-level impact (e.g., cervical cancers averted, number of females needed to be vaccinated, threshold costs of the first and second dose of the vaccine) for three vaccine strategies: no HPV vaccination (the counterfactual); a one-dose HPV vaccination schedule in which we assume that one dose of the vaccine provides either a shorter duration of protection (20 or 30 years) or lower vaccine efficacy (i.e., 80%) compared to two doses; and a two-dose HPV vaccination schedule in which two doses of the vaccine provides lifetime protection.

Figure 2. Cervical cancers averted by routine one-dose HPV vaccination by country income groups. The lines represent the median projections of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. The grey area corresponds to the additional cases averted in the vaccinated cohort after the 100 years of routine vaccination. Cancers averted (health outcomes) were discounted at 0%. Only cervical cancer caused by HPV 16, 18, 31, 33, 45, 52 and 58, which could be averted by the 9-valent HPV vaccine, were considered.

840
841

Protection from 1 dose

Figure 3. Cervical cancers averted by routine one-dose HPV vaccination as a proportion of cervical cancers averted by routine HPV vaccination programmes conferring lifelong protection at 100% vaccine efficacy. The median percentage (intervals: 10–90th percentile) of cancers averted by a one-dose schedule compared to a two-dose programme of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. Health outcomes were discounted at 0%. Only cervical cancer caused by HPV 16, 18, 31, 33, 45, 52 and 58, which could be averted by the 9-valent HPV vaccine, were considered.

842
843
844

Figure 4. Number of girls needed to be vaccinated with the first and second dose to avert one additional cervical cancer case by income group. The lines represent the median projections of the 10 model-country settings: the PHE model in black, HPV-ADVISE model-country pairs in red, and the Harvard model-country pairs in blue. The grey area corresponds to the additional cases averted in the vaccinated cohort after the 100 years of routine vaccination. Health outcomes were discounted at 3% (panels A–D) and 0% (panels E–H).

Figure 5. Threshold cost to pay for the first and second dose of vaccine by country income groups. The threshold cost is the maximum that could be paid for the first dose (compared to no vaccination) and second dose (compared to one dose only) for the incremental cost-effectiveness ratio to remain below the cost-effectiveness threshold. Two cost-effectiveness thresholds are presented: a lower threshold as suggested by Jit (2020) in panels A–D and country gross domestic product (GDP) per capita (in 2017 USD) in panels E–H. The lower cost-effectiveness threshold presented in panels A–D is 30–40% and 60–65% of GDP per capita in low-income and middle- to high-income countries, respectively. Cost and health outcomes were discounted at 3% and 0%, respectively.

Protection from 1 dose

Protection from 1 dose