

1 **Lockdown and non-COVID-19 deaths: Cause-specific**  
2 **mortality during the first wave of the 2020 pandemic in**  
3 **Norway. A population-based register study**

4 Guttorm Raknes<sup>1,2\*</sup>

5 Marianne Sørli Strøm<sup>1,3</sup>

6 Gerhard Sulo<sup>4,5</sup>

7 Simon Øverland<sup>4</sup>

8 Mathieu Roelants<sup>6</sup>

9 Petur Benedikt Juliusson<sup>1,7</sup>

10

11 \*Corresponding author

12 [guttorm.raknes@fhi.no](mailto:guttorm.raknes@fhi.no)

13

14 Address:

15 Department of Health Registry Research and Development

16 Norwegian Institute of Public Health

17 Postboks 973 Sentrum

18 NO-5808 Bergen

19 Norway

20 Tel: +47 476 30 220

21

22 Word count: 5375

23

24 Affiliations

- 25 1) Department of Health Registry Research and Development, Norwegian Institute of Public Health,  
26 Bergen, Norway.
- 27 2) Raknes Research, Ulset, Norway.
- 28 3) Institute of Global Health and Primary Care, Faculty of Medicine, University of Bergen, Bergen,  
29 Norway.
- 30 4) Centre for Disease Burden, Division of Mental and Physical Health, Norwegian Institute of Public  
31 Health, Bergen, Norway.
- 32 5) Oral Health Centre of Expertise in Western Norway-Hordaland, Bergen, Norway.
- 33 6) Environment and Health, Department of Public Health and Primary Care, KU Leuven – University of  
34 Leuven, Leuven, Belgium
- 35 7) Department of Clinical Science, Faculty of Medicine, University of Bergen, Bergen, Norway.
- 36

37 **KEYWORDS**

38 COVID-19, Pandemic, Cause of Death, Mortality, Register Study, Norway

39

40 **ABSTRACT**

41 **Objective:** To explore the potential impact of the first wave of COVID-19 pandemic on all cause and  
42 cause-specific mortality in Norway.

43 **Design** Population based register study.

44 **Setting** The Norwegian cause of Death Registry and the National Population Register of Norway.

45 **Participants** All recorded deaths in Norway during March to May from 2010 to 2020.

46 **Main outcome measures** Rate (per 100 000) of all-cause mortality and causes of death in the EU  
47 Shortlist for Causes of Death March to May 2020. The rates were age-standardised and adjusted to a  
48 100% register coverage and compared with a 95% prediction interval (PI) based on corresponding  
49 rates for 2010-2019.

50 **Results** 113 710 deaths were included, of which 10 226 from 2020. We did not observe any deviation  
51 from predicted total mortality. There were fewer than predicted deaths from chronic lower  
52 respiratory diseases excluding asthma (11.4, 95% PI 11.8 to 15.2) and from other non-ischemic, non-  
53 rheumatic heart diseases (13.9, 95% PI 14.5 to 20.2). The death rates were higher than predicted for  
54 Alzheimer's disease (7.3, 95% PI 5.5 to 7.3) and diabetes mellitus (4.1, 95% PI 2.1 to 3.4).

55 **Conclusions** There was no significant difference in the frequency of the major causes of death in the  
56 first wave of the 2020 COVID-19 pandemic in Norway. An increase in diabetes mellitus deaths and  
57 reduced mortality due to some heart and lung conditions may be linked to infection control  
58 measures.

59 **DECLARATIONS**

60 **Acknowledgements**

61 We would like to thank Olaug Margrethe Askeland and Gunhild Forland Slungård for preparing the  
62 data files at short notice.

63 **Competing interests**

64 All authors have completed the Unified Competing Interest form (available on request from the  
65 corresponding author) and declare: no support from any organisation for the submitted work; no  
66 financial relationships with any organisations that might have an interest in the submitted work in  
67 the previous three years, no other relationships or activities that could appear to have influenced the  
68 submitted work.

69 **Funding**

70 This study was funded by the Norwegian Institute for Public Health, to which all authors are  
71 affiliated. There was no external funding.

72 **Copyright/licensing**

73 The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of  
74 all authors, an exclusive non exclusive licence on a worldwide basis to the BMJ Publishing Group Ltd  
75 to permit this article (if accepted) to be published in BMJ editions and any other BMJ PGL products  
76 and sublicences such use and exploit all subsidiary rights, as set out in our licence.

77 **Author contributions**

78 All authors contributed substantially to the conception and design of the study, to the interpretation  
79 of data, to critical revision of the manuscript's intellectual content, and approved the final version.  
80 GR (the guarantor) performed the analyses and drafted the manuscript.

81 The authors agree to be accountable for all aspects of the work in ensuring that questions related to  
82 the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### 83 **Transparency declaration**

84 The lead author (guarantor of manuscript) affirms that this manuscript is an honest, accurate, and  
85 transparent account of the study being reported; that no important aspects of the study have been  
86 omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have  
87 been explained.

### 88 **Patient and public involvement statement**

89 The study was partly initiated due to questions from the press and public. Patients and the public  
90 were not involved in the development and conduct of the study or interpretation or presentation of  
91 the results. It is not possible for either researchers or the general population to influence which  
92 parameters are registered in the Norwegian Cause of Death Registry. All data are freely available on  
93 request.

### 94 **Data sharing statement**

95 All analyses were performed on aggregated data, we did not have access to directly or indirectly  
96 person identifiable data at any time.

97 The main analyses are based on cause of death data that are available from the Norwegian Cause of  
98 Death Registry (<http://statistikkbank.fhi.no/dar/>), a public, open access repository (in Norwegian  
99 only). The repository is updated regularly, and late incoming death certificates provide a gradually  
100 improving coverage. The analyses in this study are based on data retrieved on December 11, 2020;  
101 these data are found in in Supplementary table 3.

102 Data on the number of death certificates submitted electronically are not publicly available, but  
103 these will be made available on request. Weekly death statistics from the National Population  
104 Register can be accessed via Statistics Norway's webpages ([https://www.ssb.no/befolkning/artikler-  
105 og-publikasjoner/her-finner-du-ukentlige-tall-pa-antall-dode](https://www.ssb.no/befolkning/artikler-og-publikasjoner/her-finner-du-ukentlige-tall-pa-antall-dode)).

106 **Reporting guideline**

107 A filled in STROBE checklist for cross sectional studies is submitted. We are aware that this is not  
108 intended for population based register studies, but most items in the STROBE checklist are relevant.

109 **Protocol**

110 Not applicable

111 **Patient consent**

112 Not applicable.

113 The article has not been submitted to any other journal previously.

114

115 **INTRODUCTION**

116 The first two Norwegians deaths linked to COVID-19 in Norway were recorded on March 12<sup>th</sup> 2020

117 .[1] The same day the Norwegian government imposed the most comprehensive infection control

118 measures since World War II (Text box1) .[2]

119 Total mortality figures indicate no excess mortality in Norway in spring 2020 .[3] However, total

120 mortality figures do not reflect potential changes in the cause composition of deaths or specific

121 changes for certain causes.

122 There are concerns that the lockdown and social distancing measures may cause unintended

123 negative consequences for public health and mortality .[4] Some of the specific concerns relate to

124 patients potentially avoiding contact with health services, even for potentially critical symptoms such

125 as chest pain or signs of stroke ,[5] or increased suicide rates due to worsening of mental illness in

126 the context of social distancing .[6] On the other hand, social distancing may contribute to fewer

127 infectious diseases in general, such as pneumonia and influenza. The general health of patients with

128 chronic conditions such as heart disease, dementia, and diabetes, may also be more stable in the

129 absence of acute infections, leading to lower hospitalization rates and maybe mortality.

130 In this study, we used data from the Norwegian Cause of Death Registry to explore whether cause-

---

**TEXT BOX1: MEASURES IMPOSED BY THE NORWEGIAN GOVERNMENT ON  
MARCH 12, 2020 TO LIMIT THE SPREAD OF THE SARS-COV-2 VIRUS**

---

Schools, universities, and kindergartens, shops, and services considered non-essential were shut down

Mandatory 14-day quarantine for all persons entering Norway from countries with a high transmission rate, or after exposure to potential contagion

People were told to isolate themselves and, if possible, to get tested at the slightest sign of respiratory infection

General calls for hand hygiene, to keep a distance of at least two meters in the public space, to work from home whenever possible, and to avoid using public transport

131 specific mortality in Norway during the first wave of the COVID-19 pandemic (March to May 2020)  
132 was different from that observed during the same months in the preceding years.

## 133 **METHODS**

134 We conducted an observational, population-based register study.

135 Data were obtained from the Norwegian Cause of Death Registry and included information from all  
136 death certificates received by December 11<sup>th</sup>, 2020.

137 The study included all deaths occurring from March 1st until May 31st, 2020, and all deaths from  
138 March 1<sup>st</sup> to May 31<sup>st</sup> in the years 2010-2019 as reference. The data included Norwegian citizens who  
139 died abroad, but not foreign citizens who died in Norway.

140 Main outcomes were mortality rates (per 100 000 inhabitants) for all items on the 2012 EU Shortlist  
141 for Causes of Death .[7] Analyses were performed for all-cause mortality, deaths caused by diseases  
142 and external causes, the 17 level-1 causes (including external causes), on all 35 level-2 causes, and all  
143 31 level-3 causes. We present results of analyses for all level 2 and 3 causes of death from the EU  
144 shortlist when the observed age-standardised rates were outside the 95% prediction interval. Special  
145 attention was given to suicide rates, as it was feared that this rate would increase during the  
146 lockdown. For respiratory diseases, we calculated rates both including and excluding COVID-19.

147 The cause of death is reported by a physician to the Norwegian Cause of Death Registry and the  
148 Population Register by completing and mailing an international death certificate (1993 version) [8] or  
149 online by submitting an electronic death certificate [9].

150 The ICD-10 code of the cause of death in the register is determined from the death certificate by an  
151 automated coding system (IRIS) that is based on the Automatic Classification of Medical Entry  
152 (ACME) software .[10, 11] When needed, the cause of death is manually evaluated by trained staff  
153 from the Norwegian Cause of Death Registry. The term “Cause of Death” used throughout this  
154 manuscript is equivalent to the “underlying cause of death” as defined by WHO .[12]

155 **Bias**

156 Although it is mandatory by law to report all deaths to the Norwegian Cause of Death Registry, not all  
157 deaths are captured. In some cases, no death certificate is submitted, or some of the involved parties  
158 may fail to forward the document to the register. Approximately two percent of death certificates  
159 are missing in a typical year. Also, a portion of death certificates is submitted with a delay of several  
160 months. Since we retrieved the data several months in advance of the annual publication of the  
161 official Norwegian cause of death statistics, bias due to a lower coverage of late incoming certificates  
162 is a concern for the 2020 data.

163 Efforts were made to reduce number of missing causes of death in the 2020 data. The introduction of  
164 the digital death certificate contributed to speeding up the logistics, as the register receives  
165 electronic certificates within 24 hours. Since the cause of death is more often missing for persons  
166 who died abroad, and we also analysed the data excluding these.

167 We were aware of an increase in the number of deaths at home (outside hospitals and nursing  
168 homes), probably secondary to the infection control measures. This could lead to bias, for example  
169 due to a larger proportion of death certificates submitted by a general practitioner. Electronic death  
170 certificates were available before, during and after the first wave of the pandemic in 2020, and  
171 differences between the electronic and paper registration forms could lead to systematic differences  
172 in the reporting of cause of death. Before 2020, death certificates were almost exclusively paper-  
173 based.

174 The pandemic itself may also have influenced how doctors determined causes of death on the death  
175 certificate. For example, COVID-19 may have changed the level of precision for respiratory causes of  
176 death.

177 **Statistical methods**

178 Age-standardised death rates for the period March-May 2020 were compared with those of the  
179 same period during years 2010-2019. Age-standardised mortality rates were computed by the direct

180 standardization method, using 5-year age strata and the European Standard Population of 2013 as  
181 standard population .[13] The ‘at risk’ population was defined as the Norwegian population  
182 according to Statistics Norway on January 1<sup>st</sup> of each year .[14]

183 The age-standardised rates were adjusted to 100% coverage for all years to reduce the impact of  
184 missing data when comparing different years. We adjusted each cause of death group, subgroup or  
185 diagnose with a proportion of missing data, corresponding to the proportion the outcome made up  
186 of the total non-missing data for each year. The death rates used for the analyses are thus age-  
187 standardised, adjusted data, but raw rates not adjusted for missing data also represented in figures.

188 The age-standardised rates were adjusted to 100% coverage for all years to reduce the impact of  
189 missing data when comparing different years. We adjusted the rate of each cause of death by  
190 multiplying the total number of deaths in a given year with the proportion of deaths due to known  
191 causes. The death rates used for the analyses are thus age-standardised, adjusted for missing data,  
192 but raw rates not adjusted for missing data also represented in tables and figures.

193 Observed rates of all cause or disease specific causes of mortality in March – May 2020 were  
194 compared with projections based on the same causes during the same months in 2010-2019.  
195 Projections were estimated with linear regression and reported as a 95% prediction interval. Rates  
196 outside this interval were considered as a statistically significant change. A Durbin-Watson test was  
197 used to detect autocorrelation among the annual observations. + summary of findings from this test.

198 The number of cases with missing cause of death was determined by comparing the number of  
199 available death certificates with the total number of deaths according to the National Population  
200 Register. We assumed a proportional distribution of causes of death among the missing death  
201 certificates. To assess whether missing or delayed death certificates introduced bias, we examined  
202 the distribution of age and sex of the deceased and the location of death (home or hospital, nursing  
203 home). A similar comparison was made for causes of deaths based on electronic vs. paper-based  
204 death certificates.

205 **Ethics**

206 All analyses were performed on aggregated data, we did not have access to directly or indirectly  
207 person identifiable data at any time. Updated data are publicly available from the data repository of  
208 the Norwegian Cause of Death Registry .[15]

209 It is statutory that all deaths that include Norwegian citizens must be registered with the cause of  
210 death in the Norwegian Cause of Death Registry .[16] Prior consent to the storage of personal data in  
211 the register is not obtained, and it is not possible to opt out. Approval from the ethics committee or  
212 privacy ombudsman for research was neither required nor expedient.

213 **RESULTS**

214 In total, 113 710 deaths were identified in the months March, April, May since 2010. There were 216  
215 cases of COVID-19 as listed cause of death.

216 Overall, the cause of death was missing in 2 286 cases (2.0%). In 2020 the cause of death was  
217 unknown in 568 cases (5.6%), including 131 (out of a total of 133) deaths that occurred abroad.

218 Patient characteristics for deaths in data are presented in Table 1.

219

220 **Table 1.** Characteristics of persons dying during March-May, 2010-2020.

	2010-2019		2020	
<b>Mean age (<math>\pm</math> SD)</b>	78.7	( $\pm$ 15.2)	78.7	( $\pm$ 14.8)
<b>Median age (IQR)</b>	82	(17)	82	(17)
<b>N (%) females</b>	52 910	(51.1)	4 909	(50.8)
<b>N (%) external cause</b>	6 394	(6.3)	598	(6.2)
<b>N (%) death at home</b>	14 214	(14.0)	1 554	(16.1)
<b>N all deaths</b>	103 484		10 226	

221

222 Rates of all causes, diseases, and level-1 EU shortlist causes of death are presented in Table 2 while  
223 rates of all (level 1 to 3) EU shortlist causes of death are presented in Supplementary table 1.  
224 Observations for all causes, diseases, and external causes are presented in Figure 1, and for the most  
225 frequent level 1 causes (cancer, cardiovascular diseases, and respiratory diseases with and without  
226 COVID-19) in Figure 2. Causes of death for which the observed age-standardised rate fell outside the  
227 prediction interval are presented in Table 3 and in Figure 3. Suicide rates are found in Figure 4. There  
228 was a lower than predicted rate for other heart diseases (-19.7%) and other chronic lower respiratory  
229 diseases (-13.6%) (Table 3, Figure 2). There was a higher-than-expected rate for diabetes mellitus  
230 (49.9%) and Alzheimer's disease (14.9%) (Table 3, Figure 3). Among the rarer causes of death there  
231 was a higher-than-predicted rate for oesophageal cancer, non-malignant neoplasms and certain  
232 conditions originating in the perinatal period. The suicide rate did not deviate significantly from the  
233 projection based on 2010-2019 (Figure 3). No significant autocorrelation was observed.

234 Characteristics of persons with or without known cause of death, death at home and death  
235 elsewhere, and cause of deaths reported electronically or with a paper-based death certificate are  
236 presented in Supplementary table 2. Age and sex distribution were almost identical in cases with or  
237 without known cause of death, and in digital and paper-based death certificates. Among deaths at  
238 home, median age was nine years lower, there was a lower proportion females and external causes  
239 of death were more frequent.

240

241 **Table 2.** March to May death rates for all chapters in the EU Shortlist for Causes of Death. Death rates are adjusted for missing causes of death, unadjusted  
 242 rates are also shown. Predicted death rates for 2020 with 95% prediction intervals. All rates are age-standardised. \*Respiratory diseases also presented with  
 243 COVID-19 (officially not in EU shortlist chapter 8).

EU shortlist chapter	Underlying cause of death	2010-2019	2020				
		Average death rate	N	Predicted death rate	Unadjusted rate	Death rate	95% prediction interval
	All causes	238.53	1 0226	214.86	214.38	214.38	(202.42 to 227.30)
	Diseases	224.48	9 280	201.12	190.48	200.96	(188.93 to 213.31)
<b>1</b>	Infectious and parasitic diseases	5.69	219	5.52	4.64	4.89	(4.43 to 6.61)
<b>2</b>	Neoplasms	65.25	2 695	58.46	56.13	59.25	(55.94 to 60.98)
<b>3</b>	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism	0.83	36	0.61	0.76	0.80	(0.31 to 0.92)
<b>4</b>	Endocrine, nutritional, and metabolic diseases	5.75	295	5.22	6.18	6.52	(4.24 to 6.20)
<b>5</b>	Mental and behavioural disorders	14.76	775	18.29	16.55	17.45	(15.5 to 21.07)
<b>6</b>	Diseases of the nervous system and the sense organs	11.23	625	13.07	13.22	13.94	(11.78 to 14.35)
<b>7</b>	Diseases of the circulatory system	70.31	2 182	50.53	46.03	48.55	(46.18 to 54.89)
<b>8</b>	Diseases of the respiratory system (excl. COVID-19)	26.49	1 005	27.53	21.29	22.45	(22.95 to 32.11)
*	Diseases of the respiratory system (incl. COVID-19)	26.49	1 221	27.53	25.93	27.34	(22.95 to 32.11)
*	COVID-19	0.00	216	0.00	4.64	4.89	(0.00 to 0.00)
<b>9</b>	Diseases of the digestive system	7.45	302	6.98	6.34	6.69	(5.92 to 8.03)
<b>10</b>	Diseases of the skin and subcutaneous tissue	0.60	27	0.79	0.58	0.61	(0.48 to 1.10)
<b>11</b>	Diseases of the musculoskeletal system/connective tissue	1.62	76	1.97	1.60	1.69	(1.49 to 2.45)
<b>12</b>	Diseases of the genitourinary system	4.99	191	4.37	4.04	4.26	(3.04 to 5.71)
<b>13</b>	Complications of pregnancy, childbirth and puerperium	0.01	0	0.00	0.00	0.00	(-0.02 to 0.03)
<b>14</b>	Certain conditions originating in the perinatal period	0.35	22	0.18	0.40	0.43	(0.00 to 0.36)
<b>15</b>	Congenital malformations and chromosomal abnormalities	0.63	37	0.65	0.71	0.75	(0.44 to 0.85)
<b>16</b>	Symptoms, signs, ill-defined causes	8.53	357	6.94	7.38	7.79	(4.06 to 9.82)
<b>17</b>	External causes of morbidity and mortality	13.99	598	13.63	12.07	12.76	(11.08 to 16.18)

244

245

246 **Table 3: Observed and predicted mortality from** causes of death where the death rate of March to May 2020 deviated statistically significantly from the  
 247 95% prediction interval. Sorted descending after number of deaths. Death rates are adjusted for missing causes of death, unadjusted rates are also shown.  
 248 All rates are age-standardised.

EU Shortlist chapter	Underlying cause of death	2010-19	2020				
		Average death rate	N	Predicted death rate	Unadjusted death rate	Death rate	95% prediction interval
7.2	Other heart diseases	20.16	622	17.33	13.20	13.92	(14.50 to 20.17)
8.3.2	<i>Chronic lower respiratory diseases (excl. Asthma)</i>	13.08	509	13.48	10.79	11.38	(11.80 to 15.15)
6.2	Alzheimer's disease	5.23	323	6.36	6.94	7.31	(5.47 to 7.26)
4.1	Diabetes mellitus	3.64	185	2.76	3.92	4.13	(2.10 to 3.42)
2.2	Non-malignant neoplasms (benign and uncertain)	1.52	87	1.37	1.84	1.94	(0.80 to 1.93)
2.1.2	<i>Malignant neoplasm of oesophagus</i>	1.22	74	1.14	1.51	1.60	(0.72 to 1.56)
<b>14</b>	<b>Certain conditions originating in the perinatal period</b>	<b>0.35</b>	<b>22</b>	<b>0.18</b>	<b>0.40</b>	<b>0.43</b>	<b>(0.00 to 0.36)</b>

249

250

## 251 **DISCUSSION**

### 252 **Principal findings**

253 In this analysis of mortality during the first wave of the COVID-19 pandemic in Norway we observed  
254 no excess mortality and no substantial changes in the rates for neoplasms, cardiovascular disease,  
255 suicide, or other external causes. There was a lower than predicted death rate for respiratory  
256 diseases, which is probably attributable to lower mortality from chronic lung diseases (excluding  
257 asthma). The mortality rate was also lower than predicted for other non-ischemic heart diseases  
258 (excluding valvular and rheumatic heart conditions) and for Alzheimer’s disease. The observed death  
259 rate of diabetes mellitus was 45% higher than predicted. Within neoplasms, we observed higher than  
260 expected rates for oesophageal cancer and benign neoplasms.

### 261 **Strengths and imitations**

262 The Cause of Death Registry represent the main source of information for monitoring mortality. The  
263 Registry is based on established international coding practices that offer comparability and can be  
264 used to examine trends over time. The Norwegian Cause of Death Registry covers all deaths in  
265 Norway, and since it is possible to verify and validate its content using other sources, it is regarded as  
266 highly reliable .[17] As in the other Nordic countries, the Norwegian society is characterized by a high  
267 degree of trust between citizens and in the authorities. As a result, there is evidence of a high degree  
268 of compliance with governmental measures during the pandemic, which has probably contributed to  
269 low coronavirus transmission and death rates during the pandemic .[18] There is thus reason to  
270 believe that Norway is well suited for studies on the effects of population-based infection control  
271 measures.

272 Although death registers typically cover the whole population, they are prone to bias due to  
273 incomplete information on individual cases. Our main concern was bias due to loss or late incoming  
274 death certificates. However, as shown in Supplementary table 2, descriptive characteristics indicate  
275 that the group of patients with or without known cause of death was not systematically different.

276 The proportion of deaths at home from March to May 2020 was 15% higher than in the same months  
277 in 2010-2019. Both age and the proportion of females were considerably lower among deaths at  
278 home in 2020 (Supplementary table 2), but this was also true in 2010-19 (data not shown). The  
279 absolute increase in deaths at home (200 -250 estimated deaths) should not be sufficient to skew the  
280 findings significantly. Differences in patient characteristics from electronic and paper-based death  
281 certificates were also marginal, and thus unlikely to be a significant source of bias.

282 Doctors often have limited information about the patient when recording the cause of death on a  
283 death certificate, and a subjective assessment may occur. There may have been some systematic  
284 changes in the diagnostic practice and determination of the cause of death during the pandemic,  
285 especially when it comes to respiratory diseases. A more frequent use of chest x- rays may have  
286 biased the rates of the different respiratory causes of death. These effects are difficult to quantify,  
287 but we do not believe that the resulting bias is clinically relevant for most causes of death.

288 Another limitation is the large and diverse number of outcomes. We examined 83 EU shortlist causes  
289 of deaths without correcting for multiple testing. Even under normal circumstances it is therefore  
290 likely that some rates would fall outside the expected prediction interval. We could have used wider  
291 prediction intervals to adjust for multiple comparisons and avoid false positive findings, but this  
292 would at the same time increase the probability of a false negative finding. A comparison of the  
293 causes of death in Table 3 with a 99% prediction interval, shows that only diabetes mellitus remains  
294 significantly outside the predicted range. This means that great care should be taken when  
295 interpreting the results.

## 296 **Strengths & weaknesses in relation to other studies & key differences**

297 We have been unable to identify other studies with similar focus. The SARS-COV-2 virus was only  
298 identified in December 2019, and access to high-quality data with sufficient detail in such a short  
299 time is limited. Some studies reported on all-cause mortality ,[3,19] in relation to COVID-19 and  
300 infection control measures, but data on specific causes of death is usually lacking.

301

302 **Possible mechanisms & explanations for findings**

303 Only 216 Norwegians died from COVID-19 in March to May 2020, and the infection rate was also very  
304 low. Reports from the Norwegian National Institute of Public Health even showed a lower-than-  
305 expected mortality in some of the weeks in the first wave of the pandemic ,[20] indicating that  
306 misclassification of COVID-19 deaths was almost non-existent. It is therefore unlikely that SARS-COV-  
307 2 infections had a significant impact on the mortality of other causes in Norway in this period.

308 The unexpected low death rate from chronic lung diseases could be a consequence of the anti-  
309 pandemic measures. For instance, exacerbations from chronic obstructive pulmonary disease (COPD)  
310 are often caused by respiratory infections. Social distancing and improved hygiene may have been  
311 protective, even for non-respiratory diseases. The acute myocardial infarction death rate was the  
312 lowest since recordings started in 1970, but this fits into a decades long downward trend. In contrast,  
313 the number of deaths due to Alzheimer's disease has increased since the early 1990s. The observed  
314 mortality rate for Alzheimer's disease in 2020 was slightly higher than the upper limit of the  
315 prediction interval. Many Alzheimer's patients live in nursing homes where infection control  
316 measures were extra strict. There have been concerns that this patient group may have problems  
317 complying with the infection control measures, and that basic needs of vulnerable patients have  
318 been set aside .[21]

319 The most unexpected and striking finding was the large and significant increase in diabetes mellitus  
320 as cause of death (4.1 deaths per 100 000). The increase was 45% compared to the predicted rate,  
321 which is equivalent to an excess of 74 deaths in March to May 2020. One explanation could be that  
322 some diabetes patients inappropriately avoided contact with healthcare professionals, which may  
323 have contributed to poor blood glucose control. A recent survey among diabetes nurses across  
324 Europe confirmed that there are significant physical and psychological problems in their patient

325 populations during COVID-19, and that clinical diabetes services have been disrupted .[22] We were  
326 not able to differentiate between type 1 and type 2 diabetes.

327 The observed increase in deaths from oesophageal cancer, benign neoplasms and perinatal  
328 conditions should be followed closely in the time ahead. However, we find it unlikely that these  
329 findings are associated with infection control measures because there is no obvious underlying  
330 mechanism. These causes are also rare, and therefore more prone to become outliers in a study  
331 with many outcomes. The perinatal causes were 140% higher than predicted, but there were only 22  
332 deaths. Deaths due to oesophageal cancer and non-malignant neoplasms were only marginally  
333 higher than the upper limit of the 95% prediction interval.

334 We did not find an increased suicide rate, but the observation window of our study is limited to the  
335 first three months of the pandemic in Norway. It is possible that negative effects of the pandemic  
336 and social distancing measures will have more impact over time. So far, COVID-19 is coded separately  
337 in the cause of death register, and not as an EU shortlist chapter 8 respiratory disease. As seen in  
338 Figure 2, respiratory disease as cause of death is within the 95% prediction interval when including  
339 COVID-19.

#### 340 **Potential implications for clinicians or policymakers**

341 This study did not reveal an alarming increase in the causes of death that could suggest unacceptable  
342 negative effects of the infection control measures on public health or mortality, including suicide.

343 The clinical relevance of the observed changes in cause of death rate is uncertain. Diabetes mellitus  
344 may be a notable exception, and extra efforts should be made to monitor diabetes mortality and  
345 maybe even improve diabetes care throughout the pandemic.

#### 346 **Unanswered questions and future research**

347 The current pandemic is caused by a virus that was only discovered in December 2019. We have  
348 currently only examined mortality during the first wave in Norway, but similar continuous monitoring

349 is warranted until the population is sufficiently vaccinated. Results from this study highlight the  
350 importance to closely monitor morbidity and mortality due to diabetes mellitus in particular.

## 351 REFERENCES

- 352 1. Verdens Gang. Coronaviruset.  
353 [https://www.vg.no/spesial/2020/corona/?utm\\_source=corona-](https://www.vg.no/spesial/2020/corona/?utm_source=corona-)  
354 [widget&utm\\_term=d7#norway](https://www.vg.no/spesial/2020/corona/?utm_source=corona-widget&utm_term=d7#norway) (accessed January 2021).
- 355 2. The Norwegian Directorate of Health. Helsedirektoratet har vedtatt omfattende tiltak for å  
356 hindre spredning av Covid-19. [https://www.helsedirektoratet.no/nyheter/helsedirektoratet-](https://www.helsedirektoratet.no/nyheter/helsedirektoratet-har-vedtatt-omfattende-tiltak-for-a-hindre-spredning-av-covid-19)  
357 [har-vedtatt-omfattende-tiltak-for-a-hindre-spredning-av-covid-19](https://www.helsedirektoratet.no/nyheter/helsedirektoratet-har-vedtatt-omfattende-tiltak-for-a-hindre-spredning-av-covid-19) (Published 12.03.2020)  
358 (accessed January 2021).
- 359 3. Kontis V, Bennett JE, Rashid T et al. Magnitude, demographics and dynamics of the effect of  
360 the first wave of the COVID-19 pandemic on all-cause mortality in 21 industrialized countries.  
361 *Nat Med* 2020;26:1919–28. doi:10.1038/s41591-020-1112-0.
- 362 4. Bavli I, Sutton B, Galea S. Harms of public health interventions against covid-19 must not be  
363 ignored. *BMJ* 2020 2;371:m4074. doi: 10.1136/bmj.m4074.
- 364 5. Feral-Pierssens AL, Claret PG, Chouihed T. Collateral damage of the COVID-19 outbreak:  
365 expression of concern. *Eur J Emerg Med* 2020;27:233-4. doi:  
366 10.1097/MEJ.0000000000000717.
- 367 6. Lennon JC. What lies ahead: Elevated concerns for the ongoing suicide pandemic. *Psychol*  
368 *Trauma* 2020;12:S118-9. doi: 10.1037/tra0000741.
- 369 7. Eurostat. European Shortlist for Causes of Death, 2012.  
370 [https://ec.europa.eu/eurostat/ramon/nomenclatures/index.cfm?TargetUrl=LST\\_NOM\\_DTL\\_](https://ec.europa.eu/eurostat/ramon/nomenclatures/index.cfm?TargetUrl=LST_NOM_DTL_)  
371 [LINEAR&IntCurrentPage=2&StrNom=COD\\_2012](https://ec.europa.eu/eurostat/ramon/nomenclatures/index.cfm?TargetUrl=LST_NOM_DTL_LINEAR&IntCurrentPage=2&StrNom=COD_2012) (accessed January 2021).
- 372 8. World Health Organization. The International Form of Medical Certificate of Cause of Death.  
373 [https://crvsgateway.info/The-International-Form-of-Medical-Certificate-of-Cause-of-](https://crvsgateway.info/The-International-Form-of-Medical-Certificate-of-Cause-of-Death~356)  
374 [Death~356](https://crvsgateway.info/The-International-Form-of-Medical-Certificate-of-Cause-of-Death~356) (Accessed January 2021).

- 375 9. Norwegian Institute of Public Health. Elektronisk dødsmelding.  
376 <https://www.fhi.no/hn/helseregistre-og-registre/dodsarsaksregisteret/elektronisk->  
377 [dodsmelding/](https://www.fhi.no/hn/helseregistre-og-registre/dodsarsaksregisteret/elektronisk-dodsmelding/) (accessed January 2021).
- 378 10. Iris Institute. About Iris Institute. [https://www.dimdi.de/dynamic/en/classifications/iris-](https://www.dimdi.de/dynamic/en/classifications/iris-institute/)  
379 [institute/](https://www.dimdi.de/dynamic/en/classifications/iris-institute/) (accessed January 2021).
- 380 11. Lu TH. Using ACME (Automatic Classification of Medical Entry) software to monitor and  
381 improve the quality of cause of death statistics. *J Epidemiol Community Health* 2003;57:470-  
382 1. doi: 10.1136/jech.57.6.470.
- 383 12. World Health Organization. 4.1.2 Underlying cause of death. In: World Health Organization.  
384 International statistical classification of diseases and related health problems: instruction  
385 manual (Vol. 2). World Health Organization. P23.
- 386 13. European Commission. Eurostat. Revision of the European Standard Population: report of  
387 Eurostat's task force: 2013 edition. LU: : Publications Office 2013. doi:10.2785/11470.
- 388 14. Norway. 07459: Population, by sex and one-year age groups (M) 1986 - 2020  
389 <https://www.ssb.no/en/statbank/table/07459/> (accessed January 2021).
- 390 15. Norwegian Institute of Public Health. Dødsårsaksregisteret - statistikkbank  
391 <http://statistikkbank.fhi.no/dar/> (accessed February 2021).
- 392 16. FOR-2001-12-21-1476. Forskrift om innsamling og behandling av opplysninger i  
393 Dødsårsaksregisteret (Dødsårsaksregisterforskriften) (2001).  
394 <https://lovdata.no/dokument/SF/forskrift/2001-12-21-1476> (accessed January 2021).
- 395 17. Phillips DE, Lozano R, Naghavi M et al. A composite metric for assessing data on mortality  
396 and causes of death: the vital statistics performance index. *Popul Health Metr* 2014;12:14.  
397 doi: 10.1186/1478-7954-12-14.
- 398 18. Conyon MJ, He L, Thomsen S. Lockdowns and COVID-19 Deaths in Scandinavia. *SSRN Journal*  
399 2020. doi:10.2139/ssrn.3616969 [Published Online First: 2020].

- 400 19. Nørgaard SK, Vestergaard LS, Nielsen J et al. Real-time monitoring shows substantial excess  
401 all-cause mortality during second wave of COVID-19 in Europe, October to December 2020.  
402 Euro Surveill 2021;26. doi: 10.2807/1560-7917.ES.2021.26.1.2002023.
- 403 20. Norwegian Institute of Public Health. [Total mortality monitoring. Report week 4/2021.]  
404 [https://www.fhi.no/globalassets/dokumenterfiler/influenza/influensoovervaking/0421--  
405 overvaking-av-totaldodelighet-2021-uke-4.pdf](https://www.fhi.no/globalassets/dokumenterfiler/influenza/influensoovervaking/0421--overvaking-av-totaldodelighet-2021-uke-4.pdf) (In Norwegian) (Accessed February 2021).
- 406 21. Neerland BE. The pandemic of the vulnerable. Tidsskr Nor Lægeforen 2020 Published online  
407 first: 2020 doi: 10.4045/tidsskr.20.0583.
- 408 22. Forde R, Arente L, Ausili D et al. The impact of the COVID-19 pandemic on people with  
409 diabetes and diabetes services: A pan-European survey of diabetes specialist nurses  
410 undertaken by the Foundation of European Nurses in Diabetes survey consortium. Diabet  
411 Med. doi:10.1111/dme.14498 [Published Online First: 30 December 2020].

412

## 413 Figure legends

414

415 **Figure 1:** Observed and predicted death rates (linear) with 95% confidence bands for all-  
416 cause mortality (panel a), total mortality excluding external causes (panel b) and mortality  
417 due to external causes (panel c).

418 **Figure 2:** Observed and predicted death rates (linear) with 95% confidence bands for  
419 respiratory diseases (panel a), neoplasms (panel b), and cardiovascular diseases causes  
420 (panel c). For 2020, mortality from respiratory diseases is presented both with and without COVID-  
421 19.

422 **Figure 3:** Observed and predicted death rates (linear) with 95% confidence bands for causes of death  
423 with higher of lower rates than expected.

424 **Figure 4:** Observed and predicted death rate (linear) with 95% confidence bands for suicide and  
425 intentional self-harm.


## 426 SUPPLEMENTARY MATERIAL

427 **Supplementary table 1:** Observed and predicted mortality rates with 95% prediction intervals for all  
428 analysed items in the EU Shortlist for Causes of Death and deaths from COVID-19. Diseases and level  
429 1 (EU shortlist main chapters) in bold. Level 3 in italics. Rates (per 100 000) are age-standardised.  
430 Predictions are based on age-standardised rates adjusted to 100% coverage, adjusting for cases with  
431 unknown cause of death. \*Respiratory diseases also presented with COVID-19 (officially not in EU  
432 shortlist chapter 8).


433 **Supplementary table 2:** Characteristics of persons (2020) with or without known cause of death,  
434 death at home and death elsewhere, and cause of deaths reported digitally or with a paper-based  
435 death certificate. Supplementary file 3: Underlying data. Cause specific deaths March to May 2010 to  
436 2020 with rates recorded in the Norwegian Cause of Death Registry by 11 December 2020. The  
437 causes have been grouped according to the EU Shortlist for Causes of Disease. Level 1 corresponds to  
438 the 17 main chapter

439 **Supplementary table 3:** Underlying data. Cause specific deaths March to May 2010 to 2020 (column  
440 E to O) with rates (per 100000, column P to Z) recorded in the Norwegian Cause of Death Registry by  
441 11 December 2020. The causes have been grouped according to the EU Shortlist for Causes of  
442 Disease. Level 1 corresponds to the 17 main chapters. \*Respiratory diseases also presented with  
443 COVID-19 (officially not in EU shortlist chapter 8).


### All causes




### Diseases



### External causes of morbidity and mortality



### Diseases of the respiratory system




**a**

### Neoplasms




**b**

### Diseases of the circulatory system




**c**

### Alzheimer's disease




### Other heart diseases




● Death rate    × Unadjusted rate    ..... 95% prediction band    — 95% confidence band    — Regression 2010-2019


### Other chronic lower respiratory diseases



### Diabetes mellitus



# Suicide and intentional self-harm



o Death rate    x Unadjusted rate    ..... 95% prediction band    — 95% confidence band    — Regression 2010-2019