

Correcting excess mortality for pandemic-associated population decreases.

Jeremy Samuel Faust, MD MS, Brigham and Women's Hospital Department of Emergency Medicine, Emergency Medicine, Harvard Medical School (corresponding author).
Chengan Du, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation.
Shu-Xia Li, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation.
Zhenqiu Lin, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation
Harlan M. Krumholz, MD SM, Yale School of Medicine, Center for Outcomes Research and Evaluation.

Abstract

Objectives: We identify a correction for excess mortality that takes the sudden unexpected changes in the size of the United States population into account.

Design: This is a weekly cross-sectional analysis of all-cause mortality since week 5, 2020. We describe and apply a simple correction that takes population changes into account in order to provide corrected weekly estimates of expected deaths for 2020 and 2021.

Setting: The United States.

Participants: All United States residents.

Interventions: The covid-19 pandemic.

Main outcome measures: Expected and excess mortality for the United States during the covid-19 period.

Results

As of week 53, 2020 (ending January 2, 2021), approximately >10,200 more excess deaths have occurred in the United States than could be detected if expected deaths projections were not amended to reflect population decreases during 2020. The figure is projected to rise to >12,600 (>600 weekly) by week 5, 2021. Assuming recent excess mortality and pandemic-associated visa reductions continue until the earliest time herd immunity could be approached resulting from a combination of infections and vaccinations (week 17, 2021), if point estimates of expected deaths are not corrected, expected deaths will be overestimated (and therefore potential excess mortality underestimated) by ~43,000 during 2021, or >53,300 since the outbreak of the pandemic measurement period (beginning week 5, 2020). By late December 2021, weekly expected death differences are projected to approach 1,000 per week.

Conclusions

Current models measuring excess mortality should be revised immediately so that public health officials do not lose the ability to detect ongoing excess mortality as the population changes continue to compound, lowering the number of weekly expected deaths. A similar approach should be used in the middle and late phases of all future pandemics.

Correcting excess mortality for pandemic-associated population decreases.

Jeremy Samuel Faust, MD MS, Brigham and Women's Hospital Department of Emergency Medicine, Emergency Medicine, Harvard Medical School (corresponding author).

Chengang Du, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation.

Shu-Xia Li, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation.

Zhenqiu Lin, PhD, Yale-New Haven Hospital, Center for Outcomes Research and Evaluation

Harlan M. Krumholz, MD SM, Yale School of Medicine, Center for Outcomes Research and Evaluation.

Introduction

In a pandemic, the ability to detect excess mortality is paramount to policy decision-making. Detection of excess mortality relies on accurate mortality expectations. Unfortunately, covid-19 has corrupted critical assumptions upon which usual predictive models rely. Specifically, there are currently fewer permanent United States residents than projected before the pandemic, owing to excess mortality and decreases in inward migration.

Current Centers for Disease Control and Prevention and other published mortality models do not take real-time population changes into account, assuming that established population trends have not deviated substantially, and therefore minimally affect the accuracy of expected death projections.¹⁻⁴ Sudden changes in the United States population size during 2020 renders that assumption inadequate.

Excess mortality will be underestimated if expected death projections are not lowered to account for these population changes. The cumulative effect of this substantially blunts the ability to detect excess mortality during periods in which mortality rates only modestly exceed projected expectations. Moreover, the insidious compounding of these effects continuously decreases the population denominator, thereby further evading detection.

Here, we describe a simple correction procedure to improve the estimate of excess mortality during the 2020 covid-19 pandemic period and show its effect. We then conservatively project the anticipated magnitude of effect that is likely to occur during 2021 unless the technique we describe is widely adopted. This approach has widespread applicability to future excess death estimates when population sizes are in flux.

Methods

We performed a cross-sectional analysis of weekly excess deaths in the United States during the covid-19 pandemic period.

First, we determined excess deaths, defined as the difference between observed and expected deaths, using a “standard” approach.^{5,6} For granularity and accuracy, the United States population was divided into 6 age brackets and further divided by sex for a total of 12 demographic groups as follows: 0-24 female, 0-24 male, 25-44 female, 25-44 male, 45-64 female, 45-64 male, 65-74 female, 65-74 male, 75-84 female, 75-84 male, ≥ 85 female, and ≥ 85 male.

For observed mortality we used weekly figures provided by the National Center for Health Statistics at the United States Centers for Disease Control and Prevention. For weeks 6-53 2020 (with week 53 ending January 2nd, 2021, as per the CDC), published weighted figures were used in order to minimize known lag effects.⁷

To calculate standard expected deaths, we considered ongoing changes due to normal population growth and aging on a weekly basis, as elsewhere.⁵ Specifically, we applied seasonal autoregressive integrated moving average (ARIMA) to project the weekly population for all 12 groups for 2020-2021 using the data from the US Centers for Disease Control and Prevention for 2014-2019.^{8,9} The seasonal period was chosen at 52 weeks to project weekly death expectations for 2020 and 2021 for all 12 groups. We then summed the estimated contributions from the groups to obtain a point estimate of the total number of expected weekly deaths for the United States from week 6 of 2020 through Week 52 of 2021. Both 80% and 95% confidence interval estimates were determined by combining the confidence interval from the 12 groups, assuming independence among groups (see Tables S4-S16).

Next, for each of the 12 groups, we calculated “corrected” weekly estimates for expected deaths taking pandemic-associated changes in population into effect. To do this, weekly all-cause mortality incident rates using point estimates calculated via the standard approach above were first determined. Then, for each week, we calculated a revised estimated population for each of the 12 groups by subtracting the cumulative number of excess deaths during the covid-19 pandemic to that point and the cumulative decrease in permanent resident visas granted to persons arriving from abroad (owing to official policies halting such movements) for that group from the corresponding original population projection.⁹

The decrease in permanent resident visas granted during 2020 was determined by subtracting the month-over-month difference between 2020 and 2019 data, as reported by the United States Department of State (Table S1; see Supplemental Appendix Notes).¹⁰ Monthly figures were divided by 4 to create weekly estimates. We then apportioned the resulting difference among the 12 demographic groups according to the demographic breakdown of visas granted in 2019 (Table S2).¹¹ Finally, to calculate the corrected expected number of weekly deaths, the previously determined all-cause mortality incident rate was multiplied by the revised population estimate (i.e. the corrected denominator).

The mathematical description of this procedure is provided as follows:

We assume there were 53 weeks during 2020. For week $6 < i \leq 53$, we defined the original projected population for each of the 12 demographics (denoted by k , where $1 \leq k \leq 12$) in week i of 2020 to be $n_{i,k}$; the original point estimate for all-cause deaths to be $\hat{m}_{i,k}$; the sum of all previous weeks’ excess death to be $P_{i-1,k}$ (starting from week 6, 2020); the sum of all previous weeks’ decreases in inward immigration to be $Q_{i-1,k}$ (starting from week 6, 2020). Therefore, the corrected expected deaths for week i (since week 6, 2020) after correction ($\hat{m}_{i,k_{adj}}$) satisfies:

Equation 1:

$$\hat{m}_{i,k_adj} = (n_{i,k} - P_{i-1,k} - Q_{i-1,k}) \times \frac{\hat{m}_{i,k}}{n_{i,k}}$$

The above equation also holds for the weeks in 2021, where $P_{i-1,k}$ and $Q_{i-1,k}$ denote the sum of all previous weeks' excess death and the sum of all previous weeks' decreases in inward immigration since week 6, 2020.

In order to project expected deaths for 2021, we assumed that pandemic-related population changes would continue through week 17 and cease at that point (the week ending May 1, 2021). For simplicity, excess deaths for week 1-17 of 2021 were estimated using the provisional mean excess mortality for each of the 12 demographics reported for weeks 49-52, 2020 (see Tables S4-S16, orange text), though these data remain incomplete and CDC models suggest this assumption will underestimate mean excess mortality during this time.¹² To estimate ongoing decreases in permanent visas for these weeks, we conservatively assumed a decrease of 70% for weeks 1-17 of 2021, although the mean during the pandemic period through December 2020 was actually >80% (Supplemental Appendix Notes, Table S3, and Tables S4-S16, orange text).^{9,10} Expected and excess deaths were reported to the hundreds place in the text and to the ones place in tables and figures.

This study was not subject to institutional review because it relied solely upon data available to the public.

All the analysis was done through R version 4.0.2 and Microsoft Excel version 16.44.

Results

Our study included 3,079,937 all-cause deaths in the United States from week 5, 2020 through week 53 of 2020, based on preliminary CDC/NCHS data as of February 4, 2021.^{72/10/2021 1:03:00 PM} Absent the covid-19 pandemic, we estimated that the United States population would have increased from approximately 329,525,900 in week 5, 2020 to 331,381,200 by week 53, 2020 residents during the study period (Table 1). Based on standard calculations that take this growth into account, approximately 438,200 excess deaths occurred during that period (lower 95% confidence boundary 282,400) based on preliminary data. However, as of week 53 2020, we estimate that the United States population was actually 330,624,000 persons (757,200 fewer than originally projected), owing to the noted pandemic-associated excess mortality and a measured decrease in the number new permanent visa-granted residents arriving to the United States (approximately 316,300 persons) since February 2020 (Table 1, Table S1). As a result of this denominator change, as of week 53, 2020 we measured that approximately >10,200 more excess deaths occurred in the United States in 2020 than detectable using standard uncorrected approaches (Figure 1).

Due to the compounding effect, the number of missed excess deaths will rise to >12,600 (>600 weekly) by week 5, 2021 (Figure 1-2). Assuming recent excess mortality and pandemic-associated visa reductions continue until the earliest feasible time herd

immunity could be approached resulting from a combination of infections and vaccinations (week 17, 2021), if point estimates of expected deaths are not corrected to reflect these changes, we project expected deaths will be overestimated (and therefore potential excess mortality underestimated) by ~43,000 during 2021 (Figure 2), or >53,300 since the outbreak of the pandemic measurement period (beginning week 5, 2021). By late December 2021, weekly expected death differences are projected to approach 1,000 per week (~61,200 expected deaths using the standard calculation versus ~60,200 using the population-corrected calculation (Figure 3).

Discussion

We find that excess mortality in the United States has been underestimated during the covid-19 pandemic because current models in use do not consider changes in population that have occurred during the crisis. Here, we show that by taking into account such population changes weekly, the number of expected deaths decreases, meaning that excess mortality is greater than would otherwise be detectable.

We also show that this effect is compounding over time such that these quantitative differences will soon translate into an important qualitative reversal. If the measured number of deaths in a coming month is similar to the originally expected deaths estimate, observers will incorrectly conclude that excess mortality is no longer occurring. However, in comparison to the corrected estimate, excess deaths would, in fact, be occurring (Figure 3, the growing area between blue and gray lines). Such a failing to detect ongoing excess death could lead to premature policy changes such as the early curtailing of non-pharmacologic interventions. Moreover, after the pandemic subsides, failure to make the corrections identified here would eventually and misleadingly seem to appear to epidemiologists and public health officials as *deficit mortality* (i.e., measuring fewer deaths than expected). This approach would inaccurately depict a scenario in which unadjusted post-pandemic death rates appear to be lower than at any time in history, when they are not. In addition, because much of the excess mortality that has occurred during the covid-19 pandemic has been in persons with medical comorbidities, the surviving post-pandemic population is likely to collectively have a lower crude all-cause mortality rate than the pre-pandemic population because a smaller proportion of them will have significant comorbidities. Therefore, even the corrected estimates of expected mortality in the future we provide may *still* be overestimated. In fact, our findings obliquely points to the need for a systemic post-pandemic population risk recalibration so that public health professionals can continue to track disease-specific mortality for virtually every cause of death accurately.

A strength of our correction procedure for calculating excess mortality is that it relies upon real data for 2020 (week 5, 2020 to week 53, 2020). A weakness is that we must currently rely upon incomplete provisional data. Therefore, reporting lags render the figures presented to be lower boundary estimates of the effect we have unveiled for 2020, which also artificially lowers our 2021 projections.

An acknowledged weakness in our 2021 projections is that it was necessary to estimate the magnitude of excess mortality and the decrease in permanent visas for weeks 1-17 of

2021. In addition, the assumption that all such pandemic-associated changes would continue until May 1, 2021, and then cease suddenly was arbitrary. However, we believe our approach for this period was conservative; we used average counts from weeks 49-52 of 2020, the most recent for which adequate data is currently available, although the subsequent weeks (January 2021) corresponded to the deadliest weeks of the pandemic in the United States yet recorded.¹³ Thus, our projections for 2021 are almost certain to underestimate the true magnitude of the effect described here. In addition, the changes due to immigration described herein also represent lower boundary estimates, as we considered only permanent visas granted from persons arriving directly from abroad (See Supplemental Appendix Notes). Based on prior years and government travel data (Table S3), approximately twice as many fewer permanent residents likely arrived to the United States during the pandemic period than would be represented by CDC mortality data.

To our knowledge, this report is the first to describe the underestimation of excess mortality resulting from real-time changes in the population during a pandemic or outbreak of any kind. During the covid-19 pandemic, most of the effect we describe resulted from excess mortality rather than immigration. However, in a future pandemic with higher mortality rates among young and middle-aged adults (who comprise a majority of inward migrants, see Table S2), that balance could change. This means that more inclusive and more definitive real-time reporting on the decreases in inward migration might be more important in accurately measuring this effect in future pandemics. In nations with more outward migration than inward migration, the effects described here could have the opposite effect. However, these potential eventualities are built into our procedure (See Equation 1).

We propose that in the middle and late phases of the covid-19 pandemic—and future pandemics that similarly cause substantial and measurable changes in the population size—the correction methodology described here will help officials more accurately detect excess mortality later in the pandemic. Failing to adopt the approach described here is almost certain to lead to the premature declaration that excess mortality has ceased to occur both now and in any similar instances in the future. That the problem we identify here has not been considered in the past is a result of the fact that there has never been an event with this magnitude of excess mortality during an era in which real-time state, national, and even international data became available to epidemiologists and public officials with such a short time lag and on an ongoing basis.

Excess mortality remains one of the most reliable metrics for measuring a pandemic's effect, as it is agnostic to cause of death. However, to leverage excess death data properly, the ability to detect it later in pandemics must be maintained. Future research will include updated data and assess whether the effects described herein can be reliably applied at the city or state level, where denominators are smaller. In addition, the correction methodology we have identified may be improved upon by further by considering excess mortality in certain risk pools such as among persons living in long term care facilities, or in certain high-risk areas. To ensure our policy decisions are based on the best available evidence, officials should update United States and global expected death estimates immediately.

Acknowledgements:

We wish to thank Lauren M. Rossen, PhD, MS (National Center for Health Statistics), for facilitating the public release of National Center for Health Statistics data used for this study, and Julia Gelatt, PhD (Migration Policy Institute) for expertise and help identifying migration data. Neither individual received compensation for their contributions.

Patient and public involvement statement: It was not appropriate or possible to involve patients or the public in the design, or conduct, or reporting, or dissemination plans of our research.

Contributor statement: Dr. Faust conceived of the project, designed it, and wrote the manuscript in consultation with Dr. Krumholz. Dr. Du, Li, and Lin provided the statistical modeling framework and planning. Dr. Du performed the primary statistical modeling. analysis. All authors were involved in the analysis of the findings and contributed to the manuscript creation process either via direct writing or editing.

Funding: There was no specific funding for this investigation.

References

1. Woolf SH, Chapman DA, Sabo RT, Weinberger DM, Hill L. Excess Deaths From COVID-19 and Other Causes, March-April 2020. *JAMA*. 2020;324(5):510. doi:10.1001/jama.2020.11787
2. Weinberger DM, Chen J, Cohen T, et al. Estimation of Excess Deaths Associated With the COVID-19 Pandemic in the United States, March to May 2020. *JAMA Intern Med*. 2020;180(10):1336. doi:10.1001/jamainternmed.2020.3391
3. Faust JS, Krumholz HM, Dickerson KL, Lin Z, Gilman C, Walensky RP. *Mortality among Adults Ages 25-44 in the United States During the COVID-19 Pandemic*. *Epidemiology*; 2020. doi:10.1101/2020.10.21.20217174
4. Rossen LM. Excess Deaths Associated with COVID-19, by Age and Race and Ethnicity – United States, January 26–October 3, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69. doi:10.15585/mmwr.mm6942e2
5. Shiels MS, Almeida JS, García-Closas M, Albert PS, Freedman ND, de González AB. Impact of Population Growth and Aging on Estimates of Excess U.S. Deaths During the COVID-19 Pandemic, March to August 2020. *Ann Intern Med*. Published online December 15, 2020. doi:10.7326/M20-7385
6. Faust JS, Krumholz HM, Du C, et al. All-Cause Excess Mortality and COVID-19–Related Mortality Among US Adults Aged 25-44 Years, March-July 2020. *JAMA*. Published online December 16, 2020. doi:10.1001/jama.2020.24243

7. National Center for Health Statistics O. AH Excess Deaths by Sex, Age, and Race | Data | Centers for Disease Control and Prevention. Accessed February 7, 2021. <https://data.cdc.gov/NCHS/AH-Excess-Deaths-by-Sex-Age-and-Race/m74n-4hbs>
8. US Centers for Disease Control and Prevention. Underlying Cause of Death, 1999-2019, D100F141. Accessed January 4, 2021. <https://wonder.cdc.gov/controller/saved/D76/D100F141>
9. Federal Register. Declaring a National Emergency Concerning the Novel Coronavirus Disease (COVID-19) Outbreak. Accessed January 4, 2021. https://webcache.googleusercontent.com/search?q=cache:C_yPvWLV1JwJ:https://www.federalregister.gov/documents/2020/03/18/2020-05794/declaring-a-national-emergency-concerning-the-novel-coronavirus-disease-covid-19-outbreak+&cd=1&hl=en&ct=clnk&gl=us
10. US Department of State. Monthly Immigrant Visa Issuance Statistics. Accessed January 8, 2021. <https://travel.state.gov/content/travel/en/legal/visa-lawo/visa-statistics/immigrant-visa-statistics/monthly-immigrant-visa-issuances.html>
11. US Department of Homeland Security. Table 8. Persons Obtaining Lawful Permanent Resident Status by Sex, Age, Marital Status, and Occupation: Fiscal Year 2019. Department of Homeland Security. Published September 20, 2020. Accessed January 4, 2021. <https://www.dhs.gov/immigration-statistics/yearbook/2019/table8>
12. US Centers for Disease Control and Prevention. COVID-19 Forecasts: Deaths. Centers for Disease Control and Prevention. Published February 11, 2020. Accessed January 4, 2021. <https://www.cdc.gov/coronavirus/2019-ncov/covid-data/forecasting-us.html>
13. Totals for the US. The COVID Tracking Project. Accessed February 8, 2021. <https://covidtracking.com/data/national>

Supplemental Appendix: See attached pdf.

Table 1. Weekly population, calculated expected, observed, excess deaths, and inward migration related population changes, United States, Week 5, 2020 through Week 52, 2021.

Week ending	Week #	Weekly population	Expected deaths (point estimate)	Lo.80	Hi.80	Lo.95	Hi.95	Observed deaths (weighted)	Excess deaths	Excess deaths (cumulative)	Decrease in usual inward migration	Decrease in usual inward migration (cumulative)	Corrected population (original-excess deaths-decreased migration)
2/8/20	Week 6	329,525,873	59,403	58,799	60,007	58,479	60,327	59,369	-34	-34	-737	-737	329,525,873
2/15/20	Week 7	329,566,071	58,895	58,185	59,606	57,650	60,141	58,783	-112	-146	-737	-1,474	329,566,842
2/22/20	Week 8	329,606,270	58,199	57,375	59,023	56,568	59,830	58,866	667	521	-737	-2,211	329,607,153
2/29/20	Week 9	329,646,468	57,708	56,801	58,616	55,719	59,698	59,274	1,566	2,087	-737	-2,948	329,646,684
3/7/20	Week 10	329,686,667	58,422	57,456	59,387	56,624	60,219	59,643	1,221	3,308	3,309	361	329,685,317
3/14/20	Week 11	329,726,865	57,858	56,849	58,868	55,844	59,872	58,637	779	4,087	3,309	3,670	329,720,249
3/21/20	Week 12	329,767,063	57,296	56,255	58,336	55,088	59,504	59,171	1,875	5,962	3,309	6,978	329,759,668
3/28/20	Week 13	329,807,262	56,723	55,660	57,786	54,291	59,154	62,972	6,249	12,211	3,309	10,287	329,794,683
4/4/20	Week 14	329,847,460	56,865	55,786	57,943	54,354	59,376	72,235	15,370	27,581	9,263	19,550	329,825,323
4/11/20	Week 15	329,887,659	56,261	55,172	57,351	53,518	59,005	79,020	22,759	50,340	9,263	28,813	329,840,888
4/18/20	Week 16	329,927,857	55,303	54,206	56,400	52,213	58,392	76,699	21,396	71,736	9,263	38,076	329,849,065
4/25/20	Week 17	329,968,056	54,721	53,619	55,823	51,354	58,088	73,831	19,110	90,846	9,263	47,339	329,858,604
5/2/20	Week 18	330,008,254	54,814	53,708	55,920	51,451	58,177	69,236	14,422	105,268	9,674	57,013	329,870,430
5/9/20	Week 19	330,048,453	53,785	52,677	54,893	49,986	57,584	66,767	12,982	118,250	9,674	66,687	329,886,532
5/16/20	Week 20	330,088,651	53,398	52,288	54,508	49,512	57,284	64,429	11,031	129,281	9,674	76,361	329,904,075
5/23/20	Week 21	330,128,849	53,533	52,422	54,644	49,691	57,375	61,564	8,031	137,312	9,674	86,035	329,923,568
5/30/20	Week 22	330,169,048	52,934	51,822	54,046	48,803	57,065	59,613	6,679	143,991	9,674	95,709	329,946,062
6/6/20	Week 23	330,209,246	53,592	52,479	54,704	49,691	57,492	58,828	5,236	149,227	9,241	104,950	329,969,907
6/13/20	Week 24	330,249,445	53,116	52,002	54,229	49,070	57,162	57,963	4,847	154,074	9,241	114,190	329,995,629
6/20/20	Week 25	330,289,643	52,791	51,677	53,905	48,636	56,945	57,921	5,130	159,204	9,241	123,431	330,021,740
6/27/20	Week 26	330,329,842	52,638	51,524	53,752	48,402	56,875	58,423	5,785	164,989	9,241	132,671	330,047,567
7/4/20	Week 27	330,370,040	52,997	51,883	54,112	48,803	57,192	59,744	6,747	171,736	8,789	141,460	330,072,741
7/11/20	Week 28	330,408,930	52,693	51,578	53,807	48,441	56,944	61,835	9,142	180,878	8,789	150,249	330,096,095
7/18/20	Week 29	330,447,820	52,286	51,171	53,401	47,902	56,670	63,070	10,784	191,662	8,789	159,038	330,117,054
7/25/20	Week 30	330,486,710	52,067	50,952	53,183	47,593	56,542	64,127	12,060	203,722	8,789	167,827	330,136,371
8/1/20	Week 31	330,525,600	52,223	51,107	53,338	47,782	56,663	64,098	11,875	215,597	8,789	176,616	330,154,412
8/8/20	Week 32	330,564,490	52,430	51,315	53,546	48,077	56,783	63,730	11,300	226,897	7,998	184,614	330,172,638
8/15/20	Week 33	330,603,380	51,855	50,739	52,971	47,313	56,398	63,820	11,965	238,862	7,998	192,611	330,192,231
8/22/20	Week 34	330,642,270	51,965	50,849	53,081	47,417	56,512	62,683	10,718	249,580	7,998	200,609	330,211,159
8/29/20	Week 35	330,681,160	52,069	50,953	53,185	47,591	56,547	61,214	9,145	258,725	7,998	208,606	330,231,333
9/5/20	Week 36	330,720,051	52,197	51,081	53,313	47,796	56,598	60,274	8,077	266,802	6,145	214,751	330,253,080
9/12/20	Week 37	330,758,941	52,580	51,463	53,696	48,328	56,831	59,662	7,082	273,885	6,145	220,896	330,277,748
9/19/20	Week 38	330,797,831	52,856	51,739	53,973	48,741	56,971	59,649	6,793	280,678	6,145	227,040	330,303,411
9/26/20	Week 39	330,836,721	53,226	52,109	54,343	49,215	57,237	60,376	7,150	287,828	6,145	233,185	330,329,363
10/3/20	Week 40	330,875,611	53,686	52,569	54,804	49,884	57,489	59,083	5,397	293,225	7,820	241,005	330,354,959
10/10/20	Week 41	330,914,501	53,808	52,690	54,925	50,049	57,567	60,344	6,536	299,761	7,820	248,825	330,380,632
10/17/20	Week 42	330,953,391	54,925	53,808	56,043	51,598	58,253	58,865	3,940	303,700	7,820	256,644	330,405,166
10/24/20	Week 43	330,992,281	55,156	54,038	56,274	51,886	58,427	60,450	5,294	308,994	7,820	264,464	330,432,297
10/31/20	Week 44	331,031,171	54,989	53,871	56,107	51,721	58,258	61,637	6,648	315,642	7,820	272,284	330,458,074
11/7/20	Week 45	331,070,061	55,968	54,849	57,086	52,994	58,942	65,143	9,175	324,817	6,327	278,611	330,482,496
11/14/20	Week 46	331,108,951	56,593	55,475	57,712	53,864	59,322	66,589	9,996	334,813	6,327	284,938	330,505,884
11/21/20	Week 47	331,147,841	56,823	55,704	57,941	54,196	59,449	69,005	12,182	346,995	6,327	291,265	330,528,451
11/28/20	Week 48	331,186,731	56,639	55,521	57,758	53,975	59,303	70,466	13,827	360,822	6,327	297,592	330,548,832
12/5/20	Week 49	331,225,621	57,905	56,786	59,024	55,661	60,148	74,310	16,405	377,227	7,222	304,814	330,567,569
12/12/20	Week 50	331,264,511	58,347	57,227	59,466	56,375	60,318	77,924	19,577	396,804	7,358	312,172	330,582,831
12/19/20	Week 51	331,303,401	58,310	57,191	59,430	56,263	60,358	76,769	18,459	415,263	7,358	319,530	330,594,786
12/26/20	Week 52	331,342,291	58,593	57,473	59,712	56,622	60,563	74,019	15,426	430,689	7,358	326,888	330,607,859
1/2/21	Week 53	331,381,182	60,280	59,160	61,400	58,865	61,696	67,807	7,527	438,216	7,358	334,247	330,623,965

Figure 1. Cumulative difference between corrected and uncorrected expected deaths, United States, week 5, 2020 through week 52, 2021.




Figure 2. Weekly difference between corrected and uncorrected expected deaths, United States, week 5, 2020 through week 52, 2021.




Figure 3. Observed versus uncorrected expected and corrected expected deaths, United States, week 5, 2020 through week 52, 2021.

