Treatment of irritant contact dermatitis in healthcare settings during the COVID19 pandemic: The emollient Dermol 500 exhibits virucidal activity against influenza A virus and SARS-CoV-2.

Christine T. Styles[1], Michael Vanden Oever[1], Jonathan Brown[1], Sweta Rai[2], Sarah Walsh[2], Finola M. Ryan[3,4], Wendy S. Barclay[1] and Rachel S. Edgar[1]*

[1]Department of Infectious Diseases, Imperial College London, St Mary's Campus, Norfolk Place, London W2 1PG, UK.
[2]Department of Dermatology, King's College Hospital NHS Foundation Trust, Denmark Hill, London SE5 9RS.
[3] Department of Occupational Health & Wellbeing, Jennie Lee House, King's College Hospital NHS Foundation Trust, London SE5 8AD, UK.
[4]Department of Targeted Intervention, University College London, Gower Street, London WC1E 6BT, UK.
*Correspondence to Rachel S. Edgar; rachel.edgar@imperial.ac.uk

KEY MESSAGES

1. What is already known about this subject?
   - Work-related contact dermatitis is a prominent issue among healthcare workers, and likely exacerbated by the enhanced hand hygiene and personal protective equipment required to control infection during the COVID19 pandemic.
   - The antimicrobial lotion Dermol 500 is frequently prescribed as an emollient and soap substitute to help prevent and treat dermatitis, but its use during the COVID19 pandemic was not advised as its capacity to inactivate viruses was unknown.
2. What are the new findings?
   - Increased incidence of irritant contact dermatitis was recorded amongst healthcare workers at King's College Hospital NHS Foundation Trust in 2020 compared to 2019.
   - Dermol 500 lotion and its antimicrobial components, benzalkonium chloride (BAK) and chlorohexidine dihydrochloride (CD), exhibit virucidal activity against influenza A virus and SARS-CoV-2, the virus responsible for COVID19 pandemic.
3. How might this impact policy or clinical practice in the foreseeable future?
   - Our results demonstrate that Dermol 500 can be safely used as a soap substitute to treat work-related contact dermatitis in clinical care settings during the COVID19 pandemic.
   - Employers can meet their obligations under COSHH to eliminate workplace exposure to a harmful substance and substitute with an alternative product for hand hygiene.

ABSTRACT

Objectives: To investigate whether the antimicrobial emollient Dermol 500 and its active components, benzalkonium chloride (BAK) and chlorohexidine dihydrochloride (CD), exhibit virucidal activity thus informing whether Dermol 500 is a suitable soap substitute for use during the COVID19 pandemic, to combat the increased incidence of work-related contact dermatitis in clinical settings that we report here.

Methods: Inactivation of influenza A virus and SARS-CoV-2 by Dermol 500 and the independent and combined virucidal activity of the Dermol 500 components BAK and CD was assessed by influenza A virus and SARS-CoV-2 infectivity assays. Viruses were treated with concentrations of BAK and CD comparable to Dermol 500, and lower, and infectivity of the viruses assessed by titration.

Results: Dermol 500 exhibits comparable virucidal activity to alcohol-based sanitisers against influenza A virus and SARS-CoV-2. In addition, the Dermol 500 components BAK and CD exhibit independent and synergistic virucidal activity against influenza A virus and SARS-CoV-2, the causative agent of COVID19.

Conclusions: The synergistic virucidal activity of the Dermol 500 components BAK and CD makes Dermol 500 suitable as a soap substitute to treat and prevent work-related contact dermatitis in healthcare settings.

IAV: Influenza A virus
SARS-CoV-2:   Severe acute respiratory syndrome coronavirus 2
PPE: Personal protective equipment
COSHH: Control of Substances Hazardous to Health Regulations
RIDDOR: Reporting of Injuries, Diseases and Dangerous Occurrences Regulations
BAK: Benzalkonium chloride
CD: Chlorohexidine dihydrochloride
RT: Room temperature
SFM: Serum-free medium
CCV: Canine coronavirus
HSV1:   Herpes simplex virus 1
HIV: Human immunodeficiency virus
CMV: Cytomegalovirus
RSV: Respiratory syncytial virus
SARS-CoV-1:   Severe acute respiratory syndrome coronavirus 1

INTRODUCTION

Hand hygiene is imperative in healthcare settings and hand washing with soap or use of alcohol-based hand sanitisers are advised worldwide to assist infection control[1]. However, such hand hygiene measures and use of personal protective equipment (PPE) such as gloves and masks can precipitate irritant contact dermatitis[2]. This skin condition is characterised by damage to the stratum corneum layer of the epidermis, resulting in cracked, dry and itchy skin with impaired barrier function, and advances to severe irritation and inflammation[3]. Following the development of dermatitis there is an elevated risk of bacterial colonisation and subsequent transmission to other healthcare workers or patients.

Work-related contact dermatitis in healthcare professionals has been exacerbated by the infection control measures required during the COVID19 pandemic. 84.6% of surveyed Wuhan healthcare workers reported adverse skin reactions attributed to increased hand washing frequency[4]. Since March 2020, we have observed a sharp increase in NHS staff presenting with irritant hand dermatitis (Figure 1)[5]. Between 26th March – 6th May, 468 staff within King's College Hospital NHS Foundation Trust presenting to a staff dermatology clinic were diagnosed with irritant hand dermatitis. For comparison, only 8 staff had this diagnosis recorded over the same time period in 2019 at the occupational health department of the hospital. A 9.5 fold increase in irritant contact dermatitis diagnoses has been observed overall in 2020 compared to 2019 (Figure 1C), with significant increases in both occupational health walk-in cases and those diagnosed following management referral (Figures 1A and B).

Although dermatitis is a prominent issue for healthcare workers, it is manageable with appropriate interventions[3]. Under UK health and safety law, legislation details that employers have a duty of care and statutory legal obligation to employees to minimise and avoid health problems caused by work, including development of skin conditions such as work-related dermatitis[6]. 'The Control of Substances Hazardous to Health Regulations, (COSHH) 2002, (as amended)' requires employers to identify substances that may be hazardous to health, assess risks and implement adequate control measures to control exposure and protect health[7]. Indeed, if the contact dermatitis is considered solely caused by work, it is reportable under the 'Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (RIDDOR) 2013'[8][9]. Prior to the COVID19 pandemic, the use of soap substitutes and emollients was advised to treat work-related contact dermatitis in healthcare workers by the Royal College of Physicians and other medical bodies[2], meeting the COSHH requirement to eliminate exposure to the hazardous substance and substitute with an alternative. Notably, Dermol 500 lotion is usually prescribed as an antimicrobial emollient and soap substitute to aid skin conditions such as dermatitis[10]. However, due to a lack of evidence regarding whether Dermol 500 can inactivate virus particles, its use in clinical settings was discouraged during the COVID19 pandemic[11].

Figure 1. Comparative analysis of staff diagnosed with irritant contact dermatitis at King's College Hospital NHS Foundation Trust during 2019 and 2020.

Irritant contact dermatitis diagnoses following (A) walk-in presentation to occupational health, (B) management referral to occupational health, and (C) presentation at a staff dermatology clinic established by Drs. Rai and Walsh between 26th March – 6th May 2020 to address a surge in staff with skin conditions.

Dermol 500 contains the antimicrobial components benzalkonium chloride (BAK) and chlorohexidine dihydrochloride (CD)[10], which both have broad and well established antimicrobial activity[12][13]. This includes activity against the bacterial genuses *Chlamydia*[14], *Pseudomonas*[15], *Streptococcus*[16] and *Staphlococcus*[17], amongst others. BAK has been independently shown to exhibit virucidal activity against various enveloped viruses, including Canine coronavirus (CCV)[18], herpes simplex virus 1 (HSV1)[19], human immunodeficiency virus (HIV)[19] cytomegalovirus (CMV)[14] and respiratory syncytial virus (RSV)[14]. In addition, surface disinfectants where BAK is the principle active component have been shown to inactivate the SARS-CoV-1 coronavirus[20]. Virucidal activity has been demonstrated for the parental compound of CD (chlorohexidine), with alternative chlorohexidine salt form derivatives chlorohexidine gluconate and chlorohexidine acetate inactivating HIV and HSV-1[19][21].

Synergistic or cumulative virucidal activity of BAK and CD within the context and concentration found in Dermol 500 lotion has not previously been reported. The purpose of this study was to assess the virucidal activity of Dermol 500 and its antimicrobial components BAK and CD against influenza A virus (IAV) and SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), the virus responsible for COVID19 disease.

METHODS

Cell culture

Madin-Darby Canine Kidney (MDCK) cells and VeroE6 cells were routinely cultured at 37°C with 5% CO_2 in Dulbecco's modified Eagle's Medium (DMEM; Gibco-Life Technologies) supplemented with 10% FCS, 1% penicillin/streptomycin and either Glutamax (MDCK) or NEAA (Vero E6)(Gibco-Life Technologies).

Influenza infectivity assay

The influenza A virus used in this study was strain A/PR8/8/34 (H1N1), henceforth referred to as IAV. The virus was propagated in confluent MDCK flasks in the presence of TPCK-treated trypsin at 1 μg/ml (Worthingtons Bioscience) and serum-free DMEM containing 1% penicillin/streptomycin and 1% Glutamax (Gibco Life-Technologies)(SFM). MDCK cells were seeded in 6 well plates 24 to 72h prior to influenza infectivity assays.

Due to their viscosity, Dermol 500 lotion and commercially available alcohol-based sanitisers were diluted 1:1 to 50% concentration in SFM to allow for accurate pipetting. Virus was incubated with Dermol 500/alcohol-based sanitisers at the final concentrations described, or with SFM alone as a control. The alcohol-based hand sanitisers assessed were: 1) Cuticura Original Hand Sanitiser, 2) Lifebuoy Hand Hygiene Sanitiser 3) Hygiene Vision Europe Ltd Hand Hygiene Sanitiser. Alternatively, virus was incubated with BAK and CD at the concentrations described (Sigma Aldrich; CD suspended in DMSO), culture medium alone or DMSO vehicle control (Supplementary Figure 1).

Virus was incubated with the treatment for the stated times at either room temperature (RT) or 32°C (average skin temperature). Following incubation, SFM was added to the virus/treatment samples to a 10^{-1} dilution then subject to 10-fold serial dilutions. The limit of detection was therefore 10^1 virus particles/mL. Due to adverse effects of BAK and CD on cultured *in vitro* cell lines the limit of detection for BAK and CD treatments between 0.05 - 0.1% was 10^2 virus particles/mL without filtration. Where noted, IAV was filtered to remove BAK/CD post-incubation using 15 mLlAmicon Ultra-15 centrifugal filter with a 50,000 molecular weight cut off (UFC905008, Merck) and centrifugation (4000 x g, 10 min). Filtered virus was resuspended in SFM and serially diluted as above.

Infectious virus was titred via standard plaque assay. Briefly, confluent MDCK cells were incubated with 1ml of virus dilutions for 1 hour at 37°C. Virus dilutions were aspirated and cells overlaid with 3ml SFM containing 0.8% Avicel® (FMC BioPolymer), 0.14% BSA and TPCK-treated trypsin at 1 μg/ml (Worthingtons Bioscience). After 3 days, infected cells were fixed in PBS with 8-10% formalin, stained with 0.1% toulidine blue and viral plaque forming units (PFU) assessed.

SARS-CoV-2 TCID50 infectivity assay

The SARS-CoV-2 strain used in this study was SARS-CoV-2/England/IC19 and is henceforth referred to as 'SARS-CoV-2'[22]. Vero E6 cells were seeded into 96-well plates in supplemented DMEM for confluence the next day. SARS-CoV-2 was incubated with either Dermol 500 or alcohol-based sanitiser diluted as for IAV above. Alternatively, BAK and CD diluted to 0.2% in assay diluent (DMEM, 0.3% BSA, 1X NEAA, 1X penicillin/streptomycin) were mixed with an equal volume of SARS-CoV-2 to a final concentration of 0.1% BAK and 0.1% CD. Virus was also mixed with assay diluent in the absence of Dermol 500, alcohol-based sanitisers or CD/BAK. After stated incubation period, neat sample was added to the first column of confluent Vero E6 cells and Log_{10}/half-Log_{10} dilution series immediately performed across the plate in assay diluent. Four technical replicates were performed for each sample. Plates were incubated for 5 days at 37°C before adding crystal violet stain (0.1% w/v) to live cells. Wells were scored for either an intact, stained cell sheet or the absence of cells due to virus-induced cytopathic effect. For each condition, the Spearman-Karber method was used to calculate the 50% tissue culture infectious dose (TCID50) of the virus.

Statistical analysis

All data was analysed using GraphPad Prism and presented as mean ± standard deviation (mean±SD). For each treatment condition, at least 2 independent biological replicates were performed.

185 **RESULTS**
186

187 **Dermol 500 exhibits virucidal activity against IAV and SARS-CoV-2 comparable to alcohol-based hand**
188 **sanitisers.**
189

To assess whether Dermol 500 emollient exhibits virucidal activity, we examined the capacity of IAV and
SARS-CoV-2 to infect cells following incubation with Dermol 500 (Figure 2). Dermol 500 lotion was diluted in
cell culture medium to allow for accurate pipetting, therefore could only be assessed at maximum 25% of its
original concentration. Even at this dilution, we observed a significant reduction in the amount of infectious
IAV after 30min incubation at skin temperature 32°C (~100-fold reduction in virus particles; Figure 2A), and
a similar decrease in SARS-CoV-2 infectivity after only 5min (Figure 2B). In parallel, we assessed the virucidal
activity of 3 commercially available alcohol-based hand sanitisers diluted equivalently. Importantly, Dermol
500 performed as well or better than these alcohol-based sanitisers at decreasing viral infectivity. The
virucidal effect of Dermol 500 was concentration-dependent, performing less well at a 12.5% dilution
(Supplementary Figure 2), indicating increased efficacy against viruses when Dermol 500 is used at its
intended concentration. In addition, increased incubation time correlated with enhanced reduction in
infectious virus.

Figure 2. Dermol 500 lotion exhibits virucidal activity against IAV and SARS-CoV-2.

(A) IAV was incubated with Dermol 500 or alcohol-based hand sanitisers at 25% final
concentration, or medium-only control, for 5–120min at 32°C. Virus infectivity was assessed
by plaque assay (n=2; mean±SD). 2-way ANOVA [treatment x time]: treatment ****$P<0.0001$,
time $P>0.05$; multiple comparisons versus control: *$P<0.05$, **$P<0.01$, ***$P<0.001$. (B) SARS-
CoV-2 was incubated with Dermol 500 or alcohol-based hand sanitisers at 25% final
concentration for 5–60min, or assay diluent control for 60min, at 32°C. Virus infectivity was
assessed by TCID50 assay (n=2; mean±SD). 2-way ANOVA [treatment x time]: treatment
****$P<0.0001$, time **$P<0.0032$; multiple comparisons versus control: *$P<0.05$, **$P<0.01$.

The antimicrobial agents benzalkonium chloride (BAK) and chlorohexidine dihydrochloride (CD) rapidly inactivate IAV and SARS-CoV-2 at the concentrations found in Dermol 500 lotion.

BAK and CD are considered the active antimicrobial agents in Dermol 500[10], and alone have been reported to exhibit some virucidal activity[14] [18-21]. We tested their ability to inactivate IAV and SARS-CoV-2 in combination, to determine whether they contribute to the virucidal activity of Dermol 500 lotion at their intended concentration (Figure 3). Within 30 seconds of incubation with 0.1% BAK/0.1% CD the level of infectious IAV decreased 100,000-fold, and by 3 minutes infectious virus had reached undetectable levels (Figure 3A). For SARS-CoV-2, no virus-induced cytopathic effect was observed for samples incubated with 0.1% BAK/0.1% CD across all incubation time frames (10^2 TCID50/mL detection limit) indicating a >10,000-fold reduction in virus infectivity within 0.5min of contact with BAK/CD, below the limit of detection. We conclude that BAK and CD together are extremely potent virucidal agents against influenza and coronaviruses at the concentrations found in Dermol 500 lotion.

Figure 3. Benzalkonium chloride (BAK) and chlorohexidine dihydrochloride (CD) exhibit rapid virucidal activity against IAV and SARS-CoV 2 at the concentration in Dermol 500.

(A) IAV was incubated at 32°C with 0.1% BAK + 0.1% CD, or medium-only control, for 1-5min. IAV infectivity was assessed by plaque assay directly (unfiltered; 10^2 PFU/mL detection limit; n=2; mean±SD), or parallel samples were subject to filtration to remove BAK/CD prior to plaque assay (filtered; 10^1 PFU/mL detection limit; n=2; mean±SD). 2-way ANOVA [treatment x time]: unfiltered treatment **$P=0.0031$, time $P>0.05$; filtered treatment ***$P=0.0005$, time $P>0.05$. (B) SARS-CoV-2 was incubated at RT with 0.1% BAK + 0.1% CD for either 0.5min (n=2), 10min (n=2) or 30min (n=4), or assay diluent control for 30min (n=4)(mean±SD). Virus infectivity was assessed by TCID50 assay. No virus-induced cytopathic effect was observed for samples incubated with 0.1% BAK/0.1% CD across all incubation time frames (10^2 TCID50/mL detection limit). 2-tailed Mann-Whitney test: **$P=0.002$.

The antimicrobial agents BAK and CD independently inactivate IAV with different efficiencies, and act synergistically to disrupt viral infectivity.

We next investigated the virucidal activity of decreasing concentrations of BAK and CD, both in combination and independently to further characterise their efficacy (Figure 4). Within a 30 second time frame, we observed a significant decrease in IAV infectivity with the lowest concentration of BAK tested (0.0125%), indicating that amounts could be reduced in future virucidal formulations. CD alone showed no significant decrease in infectious virus over this time frame, but showed synergistic interaction with BAK by enhancing its efficacy against IAV in a time- and temperature-dependent manner (Figure 4; Supplementary Figures 3 and 4). At skin temperature, CD starts to exert an independent virucidal effect at higher concentrations by 5 minutes, and by 30 minutes the lowest CD concentration has reduced IAV infectivity by 100-1000 fold. 0.0125% BAK/0.1% CD was the most efficient combination we tested, with IAV infectivity below detectable levels within 30 seconds (Supplementary Figure 4).

Figure 4. Viral inactivation by BAK and CD at concentrations less than 0.1%.

0.5min incubation time

IAV was incubated at 32°C with decreasing concentrations of BAK, CD or in combination. Infectivity was assessed by plaque assay (n≥2; mean±SD). 2-way ANOVA[treatment x %]: 0.5min treatment *P<0.0279, % ****P<0.0001; 5mi n treatment P>0.05, % ***P=0.002; 30min treatment P>0.05, % ****P<0.0001; multiple comparisons versus 0%: *P<0.05, **P<0.01, ****P<0.0001.

5min incubation time

30min incubation time

231
232
233 **DISCUSSION AND CONCLUSIONS**
234
Any employment that requires hand hygiene, exposure to irritants and PPE such as masks and gloves poses
a risk of work-related skin disease, with irritant contact dermatitis the most common occupational
inflammatory skin disease in Europe[23]. With the prominent worldwide focus on hand hygiene and PPE during
the COVID19 pandemic, increased occurrence of work-related contact dermatitis will likely extend beyond
clinical settings into commercial employment industries, particularly other key workers. Occupational
dermatitis is a reportable disease under UK health and safety leglislation[6 8], and employers are legally obliged
to minimise exposure to harmful substances and avoid health problems including skin conditions. The
provision of emollients and soap substitutes, such as Dermol 500 lotion, are advised for treatment and
prevention of dermatitis[3], but not recommended currently due to a lack of evidence regarding their virucidal
efficacy[11]. This study clearly demonstrates Dermol 500, and its antimicrobial components≥BAK and CD, exhibit
robust virucidal activity against IAV and SARS-CoV-2, enabling its use against work-related contact dermatitis
during the COVID19 pandemic.
247
Synonymous with the cumulative activity of BAK and CD against bacteria[15], these compounds act
synergistically to inactivate two different enveloped viruses, IAV and SARS-CoV-2. Extrapolating these results,
we predict that Dermol 500 will be effective against other enveloped viruses, especially as both BAK and CD
have been shown to independently inactivate the enveloped viruses HSV-1 and HIV[19 21]. Indeed, BAK has been
shown to inactivate a broad range of enveloped viruses including CCV[18], CMV[14], RSV[14] and SARS-CoV-1[20].
Interestingly, 0.1% BAK is a component of many commercial nasal sprays, although listed as a preservative
rather than an active ingredient[24].
255
256 The mechanisms underpinning BAK and CD viral inactivation are poorly understood, and beyond the scope
257 of this paper, but both have been commonly used antimicrobials since the 1950's. They are known to disrupt
258 the physical integrity and biochemical functions of the lipid membranes of both gram-negative and gram-
259 positive bacteria[12 14 25-27]. The synergistic activity of BAK and CD indicates they may inactivate viruses via

complementary mechanisms. Beyond disrupting lipid viral envelopes, BAK alone has been shown to inactivate various non-enveloped viruses including human coxsackie virus[19], human adenoviruses[28] and enteroviruses[14]. However, BAK virucidal activity is less potent against non-enveloped viruses and requires longer incubation times to achieve viral inactivation[14]. BAK and CD virucidal activity could also encompass denaturation of viral glycoproteins required for attachment and entry. Indeed, chlorohexidine antimicrobial activity has been associated with inhibition of membrane bound proteins as well as coagulation and denaturation of cytoplasmic components, and may therefore access and denature viral proteins, especially in the presence of BAK[12 29]. The virucidal activity of these compounds and Dermol 500 lotion against non-enveloped viruses requires further study, but we predict they will be at least as efficacious as alcohol-based sanitisers, which inefficiently inactivate the majority of non-enveloped viruses[1 30].

To conclude, this study confirms Dermol 500 lotion and its constituents inactivate SARS-CoV-2 and the common respiratory virus influenza A. This broad virucidal and antimicrobial activity of Dermol 500 emollient makes it a safe and efficacious soap substitute in healthcare settings to combat the adverse effects of frequent hand washing and PPE. Its use can prevent and treat work-related irritant contact dermatitis, helping curb the large increases in this skin condition seen during this pandemic. In turn, this will decrease instances of staff sickness absence, or restriction from clinical work due to skin disease, enhance the safe and timely delivery of clinical care, and benefit staff wellbeing.

Contributors
CTS designed the study, performed laboratory experiments, analysed the data, wrote and edited the manuscript. MVO and JB performed laboratory experiments and analysed the data. SR and SW provided clinical data and edited the manuscript. FMR proposed the research question, provided clinical data and edited the manuscript. WSB edited the manuscript and acquired funding. RSE designed the study, analysed the data, wrote and edited the manuscript, and acquired funding. All authors reviewed and approved the final manuscript.

Funding
This research was supported by a Wellcome Trust Sir Henry Dale Fellowship (208790/Z/17/Z) awarded to Rachel S. Edgar, funding awarded to Wendy Barclay by the Oak Foundation, and conducted as part of the G2P consortium funded by UKRI.

Competing interests
None declared.

Patient consent for publication
Not required.

Ethics approval
Not required

Provenance and peer review
Not commissioned; pending external peer review.

Data availability statement
Data is available on request.

Acknowledgments
We'd like to thank Dr Lucia Batty, Dr Lisa Curran and Mr James Kesson (King's College Hospital NHS Foundation Trust, London) for their assistance during the preparation of this manuscript, and Dr John O'Neill (MRC Laboratory of Molecular Biology, Cambridge) for helpful discussion.

REFERENCES

1. . WHO Guidelines on Hand Hygiene in Health Care: First Global Patient Safety Challenge Clean Care Is Safer Care. Geneva2009.
2. Smedley J, Williams S, Peel P, et al. Management of occupational dermatitis in healthcare workers: a systematic review. *Occup Environ Med* 2012;69(4):276-79. doi: 10.1136/oemed-2011-100315
3. Smedley J, Group OD, Group BD. Concise guidance: diagnosis, management and prevention of occupational contact dermatitis. *Clin Med (Lond)* 2010;10(5):487-90. doi: 10.7861/clinmedicine.10-5-487 [published Online First: 2010/12/02]
4. Lin P, Zhu S, Huang Y, et al. Adverse skin reactions among healthcare workers during the coronavirus disease 2019 outbreak: a survey in Wuhan and its surrounding regions. *Br J Dermatol* 2020;183(1):190-92. doi: 10.1111/bjd.19089 [published Online First: 2020/04/08]
5. Walsh SC, S. . Introduction of a hand dermatitis clinic to reduce occupational dermatoses during the covid-19 pandemic. *medRxiv 2021020421250767* 2021 doi: https://doi.org/10.1101/2021.02.04.21250767 [published Online First: 08/02/2021]
6. Wrigglesworth F, Earl B. A guide to the Health and Safety at Work Act. Rev. ed. London: Industrial Society, 1978 1974.
7. Great Britain H, Safety Executive ap. Control of substances hazardous to health : the Control of Substances Hazardous to Health Regulations 2002 (as amended) : approved code of practice and guidance. Sixth edition. ed.
8. Great Britain H, Safety E. A guide to the Reporting of Injuries, Diseases and Dangerous Occurrences Regulations 1995, RIDDOR 95. Sudbury: HSE Books 1996.
9. Waclawski E. Disease reporting after the Reporting of Injuries, Diseases, and Dangerous Occurrence Regulations (1995) (RIDDOR) is revised. *Occup Med (Lond)* 2013;63(3):168-9. doi: 10.1093/occmed/kqs220 [published Online First: 2013/04/09]
10. Whitefield M. Effectiveness of a new antimicrobial emollient in the management of eczema/dermatitis. *J Dermatol Treat* 1998;9(2):103-09. doi: Doi 10.3109/09546639809161381
11. British Society for Cutaneous Allergy BAoD. Update on occupational skin disease during the coronavirus pandemic: British Society for Cutaneous Allergy & British Association of Dermatologists; 2020 [updated 20/04/20. Available from: https://www.bad.org.uk/shared/get-file.ashx?itemtype=document&id=66882020.
12. McDonnell G, Russell AD. Antiseptics and disinfectants: Activity, action, and resistance. *Clin Microbiol Rev* 1999;12(1):147-+. doi: Doi 10.1128/Cmr.12.1.147
13. Pereira BMP, Tagkopoulos I. Benzalkonium Chlorides: Uses, Regulatory Status, and Microbial Resistance. *Appl Environ Microb* 2019;85(13) doi: UNSP e00377-19
10.1128/AEM.00377-19
14. Belec L, Tevi-Benissan C, Bianchi A, et al. In vitro inactivation of Chlamydia trachomatis and of a panel of DNA (HSV-2, CMV, adenovirus, BK virus) and RNA (RSV, enterovirus) viruses by the spermicide benzalkonium chloride. *J Antimicrob Chemother* 2000;46(5):685-93. doi: 10.1093/jac/46.5.685 [published Online First: 2000/11/04]
15. Richards RM, Mcbride RJ. Enhancement of Benzalkonium Chloride and Chlorhexidine Acetate Activity against Pseudomonas-Aeruginosa by Aromatic Alcohols. *J Pharm Sci-Us* 1973;62(12):2035-37. doi: DOI 10.1002/jps.2600621232
16. Mosca A, Russo F, Miragliotta G. In vitro antimicrobial activity of benzalkonium chloride against clinical isolates of Streptococcus agalactiae. *J Antimicrob Chemoth* 2006;57(3):566-68. doi: 10.1093/jac/dki474

17. Castillo JA, Clapes P, Infante MR, et al. Comparative study of the antimicrobial activity of
bis(Nalpha-caproyl-L-arginine)-1,3-propanediamine dihydrochloride and chlorhexidine
dihydrochloride against Staphylococcus aureus and Escherichia coli. *J Antimicrob Chemother*
2006;57(4):691-8. doi: 10.1093/jac/dkl012 [published Online First: 2006/02/10]

18. Saknimit M, Inatsuki I, Sugiyama Y, et al. Virucidal Efficacy of Physicochemical Treatments
against Coronaviruses and Parvoviruses of Laboratory-Animals. *Exp Anim Tokyo*
1988;37(3):341-45. doi: DOI 10.1538/expanim1978.37.3_341

19. Wood A, Payne D. The action of three antiseptics/disinfectants against enveloped and non-
enveloped viruses. *J Hosp Infect* 1998;38(4):283-95. doi: Doi 10.1016/S0195-6701(98)90077-
9

20. Rabenau HF, Kampf G, Cinatl J, et al. Efficacy of various disinfectants against SARS coronavirus.
J Hosp Infect 2005;61(2):107-11. doi: 10.1016/j.jhin.2004.12.023

21. Baqui AAMA, Kelley JI, Jabra-Rizk MA, et al. In vitro effect of oral antiseptics on human
immunodeficiency virus-1 and herpes simplex virus type 1. *J Clin Periodontol* 2001;28(7):610-
16. doi: DOI 10.1034/j.1600-051x.2001.028007610.x

22. McKay PF, Hu K, Blakney AK, et al. Self-amplifying RNA SARS-CoV-2 lipid nanoparticle vaccine
candidate induces high neutralizing antibody titers in mice. *Nat Commun* 2020;11(1):3523.
doi: 10.1038/s41467-020-17409-9 [published Online First: 2020/07/11]

23. Diepgen TL. Occupational skin-disease data in Europe. *Int Arch Occup Environ Health*
2003;76(5):331-8. doi: 10.1007/s00420-002-0418-1 [published Online First: 2003/04/12]

24. Ho CY, Wu MC, Lan MY, et al. In vitro effects of preservatives in nasal sprays on human nasal
epithelial cells. *Am J Rhinol* 2008;22(2):125-9. doi: 10.2500/ajr.2008.22.3154 [published
Online First: 2008/04/18]

25. Barrettbee K, Newboult L, Edwards S. The Membrane Destabilizing Action of the Antibacterial
Agent Chlorhexidine. *Fems Microbiol Lett* 1994;119(1-2):249-53.

26. Cheung HY, Wong MMK, Cheung SH, et al. Differential Actions of Chlorhexidine on the Cell Wall
of Bacillus subtilis and Escherichia coli. *Plos One* 2012;7(5) doi: ARTN e36659
10.1371/journal.pone.0036659

27. Gilbert P, Moore LE. Cationic antiseptics: diversity of action under a common epithet. *J Appl*
Microbiol 2005;99(4):703-15. doi: 10.1111/j.1365-2672.2005.02664.x

28. Romanowski EG, Yates KA, Shanks RMQ, et al. Benzalkonium Chloride Demonstrates
Concentration-Dependent Antiviral Activity Against Adenovirus In Vitro. *J Ocul Pharmacol Th*
2019;35(5):311-14. doi: 10.1089/jop.2018.0145

29. Denyer SP. Mechanisms of action of antibacterial biocides. *Int Biodeter Biodegr* 1995;36(3-
4):227-45. doi: Doi 10.1016/0964-8305(96)00015-7

30. Golin AP, Choi D, Ghahary A. Hand sanitizers: A review of ingredients, mechanisms of action,
modes of delivery, and efficacy against coronaviruses. *Am J Infect Control* 2020 doi:
10.1016/j.ajic.2020.06.182 [published Online First: 2020/06/23]
400