Genome-wide association studies of 27 accelerometry-derived physical activity

measurements identified novel loci and genetic mechanisms

Guanghao Qi,[1][†] Diptavo Dutta,[1][†] Andrew Leroux,[1,2] Debashree Ray,[1,3] John Muschelli,[1] Ciprian

Crainiceanu[1] and Nilanjan Chatterjee[1,4]*

[1]Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

21205, USA

[2]Department of Biostatistics and Informatics, University of Colorado, Aurora, CO 80045, USA

[3]Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore,

MD 21205, USA

[4]Department of Oncology, Johns Hopkins School of Medicine, Baltimore, MD 21205, USA

18 **This work was supported by an R01 grant from the National Human Genome Research**

19 **Institute [1 R01 HG010480-01].**

21 **†These authors contributed equally to this work**

22 ***Correspondence to Nilanjan Chatterjee (nilanjan@jhu.edu)**

Abstract

Physical inactivity (PA) is an important risk factor for a wide range of diseases. Previous genome-wide association studies (GWAS), based on self-reported data or a small number of phenotypes derived from accelerometry, have identified a limited number of genetic loci associated with habitual PA and provided evidence for involvement of central nervous system in mediating genetic effects. In this study, we derived 27 PA phenotypes from wrist accelerometry data obtained from 88,411 UK Biobank study participants. Single-variant association analysis based on mixed-effects models and transcriptome-wide association studies (TWAS) together identified 5 novel loci that were not detected by previous studies of PA, sleep duration and self-reported chronotype. For both novel and previously known loci, we discovered associations with novel phenotypes including active-to-sedentary transition probability, light-intensity PA, activity during different times of the day and proxy phenotypes to sleep and circadian patterns. Follow-up studies including TWAS, colocalization, tissue-specific heritability enrichment, gene-set enrichment and genetic correlation analyses indicated the role of the blood and immune system in modulating the genetic effects and a secondary role of the digestive and endocrine systems. Our findings provided important insights into the genetic architecture of PA and its underlying mechanisms.

Key words: Genome-wide association studies, physical activity, novel loci, blood and immune mechanisms

Introduction

Regular physical activity (PA) is associated with lower risk of a wide range of diseases, including

cancer, diabetes, cardiovascular disease(Kyu et al., 2016), Alzheimer's disease(Rovio et al.,

2005), as well as mortality(Smirnova et al., 2019; Leroux et al., 2019). However, studies have

indicated that large majority of US adults and adolescents are insufficiently active(Piercy et al.,

2018), and thus PA interventions have great potential to improve public health. PA was shown

to have a substantial genetic component, and understanding its genetic mechanism can inform

the design of individualized interventions(Lightfoot et al., 2018; Moore-Harrison & Lightfoot,

2010). For example, people who are genetically pre-disposed to low PA may benefit more from

early and more frequent guidance.

A number of previous genome-wide association studies (GWAS) on physical activity have relied

on self-reported phenotypes, which are subject to perception and recall error(De Moor et al.,

2009; Hara et al., 2018; Kim et al., 2014; Klimentidis et al., 2018). Recently, wearable devices

have been used extensively to collect physical activity data objectively and continuously for

multiple days. To date, there have been two GWAS based on acceleromtery-derived activity

phenotypes. Both studies used data from the UK Biobank study(Doherty et al., 2017; Bycroft et

al., 2018) but only focused on a few summaries of these high-density PA measurements. One

study considered two accelerometry-derived phenotypes (average acceleration and fraction

accelerations > 425 milli-gravities) and identified 3 loci associated with PA(Klimentidis et al.,

2018). A second study used a machine learning approach to extract PA phenotypes, including

overall activity, sleep duration, sedentary time, walking and moderate intensity activity(Doherty

et al., 2018). This study identified 14 loci associated with PA and found that the central nervous

system (CNS) plays an essential role in modulating the genetic effects on PA. However, both

studies used a small number of phenotypes, which may not capture the complexity of PA

patterns.

Recent studies suggest that in addition to the total volume of activity, other PA summaries may

be strongly associated with human health and mortality risk. For example, the transition

between active and sedentary states was strongly associated with measures of health and

mortality(Leroux et al., 2019; Schrack et al., 2019). PA relative amplitude, a proxy for sleep

quality and circadian rhythm, was strongly associated with mental health(Rock, Goodwin,

Harmer, & Wulff, 2014a). Moderate-to-vigorous PA (MVPA) and light intensity PA (LIPA) have

also been reported to be associated with health(McGregor, Palarea-Albaladejo, Dall,

Stamatakis, & Chastin, 2019; Young & Haskell, 2018). Thus, there is increasing evidence that

objectively measured PA in the free-living environment is a highly complex phenotype that

requires a large number of summaries that provide complementary information. Understanding

the genetic mechanisms behind these summaries is critical for understanding the genetic

regulation of activity behavior and informing targeted interventions.

In this paper, we conducted genome-wide association analysis using 27 accelerometry-derived

PA measurements from UK Biobank data(Bycroft et al., 2018; Doherty et al., 2017). The

phenotypes cover a wide range of features including volumes of activity, activity during

different times of the day, active to sedentary transition probabilities, principal components

and proxies for circadian rhythm (**Table 1**). We conducted GWAS using a mixed-model-based

method, fastGWA(Jiang et al., 2019), to identify variants associated with the above phenotypes.

We also conducted transcriptome-wide association studies (TWAS)(Gusev et al., 2016) across

48 tissues to identify genes and tissues harboring the associations. We further conducted

colocalization (Giambartolomei et al., 2014), tissue-specific heritability enrichment(Finucane et

al., 2018; Finucane et al., 2015), gene-set enrichment(Watanabe, Taskesen, van Bochoven, &

Posthuma, 2017) and genetic correlation(Bulik-Sullivan et al., 2015a) analyses to further reveal

the underlying biological mechanisms. We identified 5 novel loci associated with PA and

showed that, in addition to the CNS, blood and immune related mechanisms could play an

important role in modulating the genetic effects on activity, and digestive and endocrine tissues

could play a secondary role.

Material and Methods

Study Cohort and Physical Activity Phenotypes

The UK Biobank study consists of ~500,000 individuals in the United Kingdom with

comprehensive genotype and phenotype data(Bycroft et al., 2018). We used a subset of the

103,712 individuals who were invited and agreed to participate in the accelerometry sub-study

where participants wore a wrist-worn accelerometer for up to 7 days(Doherty et al., 2017).

Accelerometry data from participants are available at multiple resolutions. Here, the individual-

specific set of accelerometry-based phenotypes was derived from the five-second level

acceleration data provided by the UK Biobank team. We further compressed the data by

averaging the 5-second level data within every minute. Individuals were screened for poor

quality data using indictors provided by the UK Biobank. In addition, we required individuals to

have at least 3 days (12am-12am) of sufficient wear time defined as estimated wear time

greater than 95% of the day (>= 1368 minutes). Our inclusion criteria for this analysis closely

mirrors that described in a related paper from our group(Leroux et al., 2020) with the exception

that we did not exclude participants younger than 50 at the time of accelerometer wear or

based on missing demographic and lifestyle data and instead excluded individuals based on

ancestry and genotype data (see subsection *Genotype Data* below).

Physical activity phenotypes were all calculated at the day level and then averaged within study

participants across days to obtain one measure for each phenotype and study participant. This

led to in 31 PA phenotypes for 93,745 study participants that covered a wide spectrum of

information. The phenotypes are described in **Table 1** and briefly summarizes as follows: 1)

total volume of activity (total acceleration (TA), total log acceleration (TLA)); 2) activity during

12 disjoint two-hour windows of the day (TLA 12am-2am, TLA 2am-4am, …, TLA 10pm-12am);

3) duration of sedentary state (ST), LIPA and MVPA; 4) principal components of the log-

transformed minute-level activity profiles (PC1-6); 5) active-to-sedentary transition probability

(ASTP) and sedentary-to-active transition probability (SATP); 6) proxy phenotypes for circadian

patterns, including dynamic activity ratio estimate (DARE), activity during the most active 10

hours (M10) and least active 5 hours (L5) of the day, timing of M10 and L5, and PA relative

amplitude ($\frac{M10-L5}{M10+L5}$). They included most of the phenotypes used in the previous PA association

studies as well (See **Table 1** for details)(Doherty et al., 2018; Klimentidis et al., 2018). The exact

procedure for deriving study participant-specific phenotypes is described in detail in the

supplemental material of the related paper from our group(Leroux et al., 2020). The

phenotypes were inverse-normal transformed

$$INT(y_i) = \ \Phi^{-1}\left[\frac{rank(y_i)-k}{n-2k+1}\right], \quad k = 3/8,$$

where the transformed variables have mean 0 and variance 1.

Removing Highly Correlated Phenotypes

Some of the initial 31 PA phenotypes were highly correlated (**Figure S1**). To avoid counting

similar phenotypes multiple times, if two phenotypes had correlation > 0.8 we removed one of

them. First, we removed total acceleration (TA), duration of sedentary state (ST), PC1 and M10

due to their high correlation with total log acceleration (TLA). TLA was retained as the main

metric for the total volume of activity. Since two previous GWASs used TA as the main metric

for the volume of activity (Doherty et al., 2018; Klimentidis et al., 2018), we chose TLA instead

of TA to avoid repetition with existing work. In addition, the original distribution of TA is highly

skewed, which may not be completely addressed by the INT. In total, 4 phenotypes were

removed and 27 PA phenotypes were retained for the association analysis.

Genotype Data

The imputed genotype data for ~93 million variants, using UK10K, 1000 Genomes (Phase 3) and

Haplotype Reference Consortium as reference panel, provided by UK Biobank were used and

merged with the PA phenotype data. Following the Neale lab UK Biobank GWAS pipeline

(https://github.com/Nealelab/UK_Biobank_GWAS/blob/master/imputed-v2-

[gwas/README.md](https://github.com/Nealelab/UK_Biobank_GWAS/blob/master/imputed-v2-gwas/README.md)), we excluded study participants according to the following criteria: 1) non-

white ancestry; 2) putative sex chromosome aneuploidy; 3) an excessive number of relatives

(more than 10 putative third-degree relatives in the kinship table); 4) sample was not in the

input for phasing of chr1-chr22. After applying these exclusion criteria the sample was further

reduced to 88,411 study particiants for downstream analysis.

We conducted variant quality control to ensure that genetic variants with poor genotyping

quality do not affect the results. Specifically, variants that satisfy any of the following criteria

were removed: 1) imputation INFO score < 0.8; 2) MAF < 0.01; 3) Hardy-Weinberg Equilibrium

(HWE) p-value < 1×10^{-6}; 4) missing in more than 10% study participants. After the filtering,

8,951,705 variants remained for downstream analysis, of which 8,067,228 (90.1%) were single

nucleiotide polymorphisms (SNPs) and the rest (9.9%) were insertion-deletions (INDELs).

Association Analysis

We used a fast mixed-effects model method, fastGWA(Jiang et al., 2019), for genome-wide

association analysis. Like other mixed-effects model methods, fastGWA allows the inclusion of

related and unrelated individuals but improves computational efficiency by incorporating a

sparse genetic relationship matrix (GRM). The GRM measures the genetic similarity between

individuals and each element is the correlation of genotypes between a pair of individuals. We

constructed the GRM using linkage disequilibrium (LD)-pruned variants that had MAF > 5% and

were present in HapMap3 (LD-pruning was done in PLINK using the following set up as

recommended in Jiang et al (Jiang et al., 2019): window size = 1000Kb, step-size = 100 and r^2 =

0.9). We further computed a sparse-GRM at sparsity level 0.05 to capture the genetic

relatedness between the closely related individuals only and reduced others to zero. We used

the Haseman-Elston regression to estimate the variance of the random effects as an

intermediate step of fastGWA. This approach is orders of magnitude faster than the previous

state-of-the-art, BOLT-LMM(Loh et al., 2015; Loh, Kichaev, Gazal, Schoech, & Price, 2018).

Models were adjusted for age, sex and the first 20 genetic principal components as covariates.

Because the PA phenotypes are correlated, principal component analysis (PCA) was conducted

on the phenotypes to estimate the number of independent phenotypes before setting the

GWAS significance threshold. At least 19 phenotype PCs were needed to explain 99% percent of

the PA phenotypic variance (**Figure S2**). Variants with p-value below the threshold

$5 \times 10^{-8}/19 = 2.63 \times 10^{-9}$ were declared to be statistically significant, which accounted for

the number of independent phenotypes. LD clumping was conducted based on the minimum p-

value across phenotypes. The requirements for the lead SNPs of different loci were to have

$r^2 < 0.1$ and be at least > 500kb apart.

A locus was defined as novel if its lead variant is >500kb from the lead variant of any known loci

discovered by the following GWASs on PA, sleep, and circadian rhythm: (1) Doherty et al study

on a smaller set of accelerometry-derived PA phenotypes(Doherty et al., 2018); (2) Klimentidis

et al study on self-reported and accelerometry-derived PA(Klimentidis et al., 2018); (3) Dashti et

al study on self-reported sleep duration(Dashti et al., 2019); (4) Jones et al study on circadian

rhythm(Jones et al., 2019). Considering the diversity of PA phenotypes, we further ensure these

loci do not have associations with other related traits not listed above, by searching for the lead

variants in Open Targets Genetics (OTG, https://genetics.opentargets.org/). A locus remains

novel if the lead variant: 1) is not associated at $p < 5 \times 10^{-8}$ with traits whose names include

the following keywords: accelerometry, physical, exercise, sleep, nap, circadian and

chronotype; 2) and is not in LD with previously reported GWAS lead variants for these traits

$(r^2 > 0.5)$.

Transcriptome-wide association studies (TWAS) were conducted using the FUSION R

program(Gusev et al., 2016) with reference models generated from 48 tissues of GTEx v7(GTEx

et al., 2017). TWAS analysis was limited to the 18 traits with at least one genome-wide

significant variant ($p < 2.63 \times 10^{-9}$). Multiple testing due to the large number of tissue-trait

combinations (48*18=864) was addressed by a two-stage adjustment approach: 1) for each

variant, the Benjamini-Hochberg (BH) adjustment was applied across all tissue-trait pairs; 2)

each variant with BH-adjusted p-value 2.5×10^{-6} was then identified (accounting for 20,000

protein-coding genes). Since there can be multiple genes in close proximity to each other, to

identify independent loci detected by TWAS analysis, genes were clustered based on significant

associations. A clumping approach was used, which selected the gene with the smallest

minimum p-value across tissue-trait pairs and removed the other genes with a transcription

start site (TSS) within 1Mb of the lead gene TSS. The process continued by identifying the gene

with the next smallest minimum p-value and iterating. The only exception was when the lead

gene of the cluster was not a protein-coding gene (e.g., pseudogene, lncRNA) and a protein-

coding gene was in the cluster. In this case the protein-coding gene with the smallest minimum

p-value was identified as the lead gene. This led to independent gene clusters at genomic loci

which were least 1Mb apart, i.e., none of the lead gene TSS is within the cis region of another

lead gene.

Enrichment Analysis

Stratified LD score regression(Finucane et al., 2015; Finucane et al., 2018) was used to identify

the tissues and genomic annotations enriched by the heritability for PA. For tissue specific

analysis, chromatin-based annotations were used as derived from the ENCODE and Roadmap

data(ENCODE, 2012; Roadmap et al., 2015) by Finucane et al(Finucane et al., 2018). The

annotations were based on narrow peaks of DNase I hypersensitivity site (DHS) and five

activating histone marks (H3K27ac, H3K4me3, H3K4me1, H3K9ac and H3K36me3) observed for

111 tissues or cell types, resulting in a total of 489 annotations. Stratified LD score regression

computes the heritability attributed to each annotation and computes a coefficient and a p-

value that characterize enrichment.

241 In a separate analysis, the enrichment of TWAS signals was evaluated among the genes that

242 have been reported to be associated to different traits, using FUMA(Watanabe et al., 2017;

243 Watanabe, Umićević Mirkov, de Leeuw, van den Heuvel, & Posthuma, 2019). For a given PA

244 trait, we defined a gene-set as the genes that were significant at an exome-wide level ($p <$

245 2.5×10^{-6}) and investigated whether these genes overlapped with the genes that have been

246 mapped to genome-wide significant variants for different traits as reported in GWAS

247 catalog(Buniello et al., 2019). The collection of such genes have been detailed in Molecular

248 Signatures Database (MSigDB)(Liberzon et al., 2011). We used FUMA to compute the

249 proportion of genes related to other diseases and traits that were also identified by our TWAS

250 analysis and computed enrichment p-values using the Fisher's exact test.

252 *Colocalization Analysis*

254 For each susceptibility locus of PA (**Table 2**), colocalization analysis was conducted between its

255 most significantly associated phenotype and eQTL effects on gene expression in 48 tissues in

256 GTEx v7(GTEx et al., 2017). SNPs within +-200kb radius of the lead SNP were used and genes

257 that had at least one significant eQTL (q-value < 0.05) in the region were considered. Analysis

258 was conducted using the R package COLOC (Giambartolomei et al., 2014) and GWAS and eQTL

259 effects were identified as being colocalized if PP4 > 0.8.

261 *Heritability and Genetic Correlation Analysis*

Heritability of activity phenotypes was estimated using Haseman-Elston regression as an

intermediate output of fastGWA(Jiang et al., 2019). Our fastGWA analysis computed sparse

GRM at sparsity level 0.05 as recommended by the fastGWA paper (see "Association analysis").

However, this cutoff may miss the subtle relatedness in the sample and affect heritability

estimate. As a sensitivity analysis, we re-estimated the heritability using a lower sparsity

threshold at 0.02 to capture more subtle relatedness. The genetic correlation between 18 PA

traits and 238 complex traits and diseases was estimated using LD score regression(Bulik-

Sullivan et al., 2015a) implemented in LD Hub(Zheng et al., 2017). In particular, we focused on

four broad groups of traits and diseases (A) cholesterol levels (B) anthropometric traits (C)

autoimmune disease and (D) miscellaneous traits including psychiatric, neurological, cognitive

and personality traits. For each trait and within each category, we applied a false discovery rate

correction to the p-values corresponding to the genetic correlation estimated using LD score

regression, to account for multiple testing. Any genetic correlation with FDR-adjusted p-value

less that 10% were declared as significant.

Results

Genetic Loci Associated with Physical Activity

Single-variant genome-wide association analysis identified a total of 16 independent loci,

including three novel ones compared to previous studies (**Table 2 and Figure 1**). All three novel

loci were discovered on chromosome 3: the locus indexed by rs3836464 was associated with

ASTP; the locus indexed by rs9818758 was associated with relative amplitude, which is a proxy

sleep behavior and circadian rhythm(Rock, Goodwin, Harmer, & Wulff, 2014b); the locus

indexed by indel 3:131647162_TA_T (no rsid available) was associated with TLA 2am-4am

which is a proxy phenotype for activity during sleep. LIPA appeared to be associated with other

SNPs near 3:131647162_TA_T but not the lead variant itself, indicating multiple independent

signals at the same locus (**Figure 1**). Nearest coding genes for the novel loci include *SEC13*,

USP4, and *CPNE4*.

Our analysis also identified novel phenotypes for several known loci (**Table 2**). The strongest

signal was seen for the locus indexed by rs113851554 which is associated with multiple sleep

and circadian rhythm proxy phenotypes including TLA 12am-2am ($p = 6.7 \times 10^{-37}$), TLA 2am-

4am ($p = 7.9 \times 10^{-39}$), average log acceleration during the least active 5 hours of the day (L5,

$p = 1.3 \times 10^{-33}$), timing of L5 ($p = 5.4 \times 10^{-22}$) and PA relative amplitude ($p = 6.9 \times 10^{-15}$).

This locus was previously identified to be associated with accelerometry-derived sleep duration

in UK Biobank(Doherty et al., 2018). Among other known loci, 5 were only discovered in the

GWAS of self-reported circadian rhythm(Jones et al., 2019) but not in the other studies

considered (**Table 2**, last column). In our analysis, the loci indexed by rs1144566, rs9369062 and

rs12927162 were associated with sleep proxy phenotypes including timing of L5, TLA 12am-

2am, TLA 2am-4am and TLA 10pm-12am. Three other loci, indexed by rs301799, rs2909950 and

rs12717867, were associated with TLA 6pm-8pm and LIPA, respectively.

In addition to the phenotypic associations above, other variants in some of the loci captured

associations that are not reflected by the lead variants. In particular, variants in high LD with

rs2138543 are associated with a wide range of phenotypes including LIPA, MVPA, activity

during two-hour windows, and a number of proxy phenotypes for circadian patterns (**Table S1**).

Transcriptome-Wide Association Study and Colocalization Analysis

We performed transcriptome-wide association studies (TWAS) (Gamazon et al., 2015; Gusev et

al., 2016) for each PA trait based on gene expression data across 48 tissues available through

GTEx (version 7)(GTEx et al., 2017). Our analysis identified 15 loci (**Table 3, Figure 2, Table S2**)

with significant association in at least one trait-tissue pair analysis after correcting for multiple

testing (Benjamini-Hochberg corrected p-value $< 2.5 \times 10^{-6}$, see **Methods**). We identified two

novel loci. One of them was indexed by *RN7SKP16*, whose higher expression in brain putamen

basal ganglia is genetically associated with lower level of MVPA. Another was indexed by

pseudogene *PDXDC2P (16q22.1)*, whose higher expression in esophagus mucosa and EBV

transformed lymphocytes appeared to be genetically associated with lower level of TLA 6am-

8am (**Figure 2**). See **Table S2** for details of these associations. These loci was not previously

reported by any prior GWAS and were not close to any of the 3 novel regions detected by our

single variants analysis (**Table 3**).

The TWAS analysis also identified novel PA phenotypes, potential target genes and underlying

tissues for many of the known loci or novel loci detected through single variant analysis (**Table**

3). Consistent with a previous study (Doherty et al., 2018), the TWAS analysis showed that

genetic association for PA traits often points towards involvement of CNS (**Table 3**). Further, our

analysis indicates consistent involvement of blood and immune, digestive and endocrine

systems in modulating the genetic effects on PA. See **Table S2** for a complete list of associated

tissues. Among the 15 loci significant in TWAS analysis, the lead genes of 4 loci were

significantly associated with PA phenotypes via the blood and immune tissues. For example, the

genetically predicted expression of *PBX3* and *KANSL1* in the blood and immune tissues (whole

blood, EBV-transformed lymphocytes) were each associated with 3 PA phenotypes. The genes

associated with PA via blood and immune tissues were also associated via digestive (esophagus

mucosa, small intestine - terminal ileum, colon – sigmoid, etc) and/or endocrine (thyroid,

pituitary) tissues, but only one of them overlapped with the 9 genes that were associated with

PA phenotypes through the CNS (**Table 3**). Another locus, represented by *C3orf62*, were

associated with PA relative amplitude only via the digestive and endocrine tissues (esophagus

mucosa, colon – sigmoid, thyroid) but not the blood/immune tissues or CNS. These findings

suggested two potential pathways for the genetic regulation of PA: a primary pathway involving

the CNS (brain in particular) and a secondary pathway involving the blood/immune system and,

potentially, the digestive and endocrine systems. The actual biological processes involved in the

pathways are beyond the scope of this paper and may be worth future investigation.

Several genes that were found to be significantly associated to specific PA traits in our TWAS

analysis, were also found to be highly overlapping with genes that were previously reported to

be associated with various traits and diseases including but not limited to neuropsychiatric

diseases, behavioral traits, anthropometric traits and autoimmune diseases (**Figure S3**). For

example, we found that the genes associated with TLA across different tissues, are enriched for

genes that have been associated with neuroticism, bipolar disorder, Parkinson's disease,

cognitive function and several others indicating the putative involvement of the CNS in the

genetic mechanism of TLA. Additionally, the genes associated with relative amplitude

overlapped highly with those associated with several autoimmune diseases like inflammatory

bowel disease, ulcerative colitis in addition to different behavioral and cognitive traits (**Figure**

S3). These results further supported the possible involvement of both CNS as well as the blood

and immune system in the genetic mechanism of PA traits.

We performed a colocalization analysis to gain further insights on the tissue specific activity of

the significant genetic loci. Among the 16 loci significantly associated with PA, 9 loci colocalized

with the eQTL signals for at least one gene and one tissue with a colocalization probability

(PP4) > 0.8 (**Table S3**). Colocalization occurred in a similar set of tissues as those that harbored

the TWAS associations (**Table 3 and Table S3**), namely the CNS, blood and immune (whole

blood, spleen, EBV transformed lymphocytes), digestive (esophagus, colon) and endocrine

(thyroid, testis, adrenal gland) tissues, and also in a number of cardiovascular tissues that were

not highlighted by TWAS. Among the 15 lead genes for TWAS significant loci, the eQTL signal of

4 genes (*RERE, C3orf62, PBX3* and *RP11-396F22.1*) colocalized with PA GWAS signal in at least

one tissue. Colocalization also occurred in two other secondary genes (*RP5-1115A15.1* and

CASC10).

Analysis of Heritability and Co-Heritability

Our fastGWA analysis estimated genome-wide heritability of PA phenotypes as an intermediate

output. The estimates appeared to be dependent on the sparsity level of the genetic

relationship matrix (**Figure S4**). We chose the results under the lower cutoff (0.02) since it

captured more subtle relatedness and should give more accurate heritability estimates. The

estimates of heritability varied across different PA phenotypes. A number of traits were

estimated to have higher heritability than others, including TLA (0.15), TLA 6pm-8pm (0.15),

MVPA (0.14) (**Figure S4**). Afternoon and pre-sleep evening activity (TLA 4pm to 12am) appeared

to be more heritable than morning activity (TLA 2am to 12pm). As could be expected,

phenotypes with higher heritability tend to have a higher average χ^2 statistic for genetic

associations, and a QQ plot which deviate further from the null line (**Figure S5**). The magnitude

of heritability estimates was generally consistent with previous studies, which reported 10-20%

heritability for PA traits (Doherty et al., 2018; Klimentidis et al., 2018). We also notice that

heritability estimated using restricted maximum likelihood (REML) tended to be slightly higher

Haseman-Elston regression estimates (**Figure S4**).

We further used stratified LD-score regression for partitioning heritability by functional

annotations of genome(Finucane et al., 2015; Finucane et al., 2018). Consistent with TWAS

findings, this analysis also indicated possible role for blood and immune system in addition to

CNS for genetic regulation of PA (**Figure 3**). In particular, heritabilities for both TLA and LIPA

were enriched for DNase I hypersensitivity sites (DHS) in primary B cells from peripheral blood

and that for TLA 12pm-2pm were enriched for H3K27ac in spleen. We also found potential

enrichment in other traits, though they were not significant after FDR adjustment. For example,

for TLA 8am-10am, MVPA, and ASTP the heritability enrichment in active chromatin regions of

blood/immune tissues were all close to being statistically significant (**Figure S6**).

We further used LD score regression(Bulik-Sullivan et al., 2015a; Bulik-Sullivan et al., 2015b) to

explore genetic correlation between PA phenotypes and four broad groups of complex traits

and diseases (**Figure S7**). Genetic correlations were identified (FDR < 10%) between PA

phenotypes and: (1) neurological, psychiatric and cognitive traits, including Alzheimer's disease

(AD), attention-deficit hyperactivity disorder (ADHD), depressive symptoms, intelligence, and

neo-conscientiousness; (2) auto-immune diseases, with the strongest correlation for multiple

sclerosis and weaker correlations for Crohn's disease and primary billary cirrhosis; (3) obesity-

related anthropometric traits and (4) cholesterol levels. Most PA traits have negative genetic

correlation with obesity-related traits and triglycerides, and positive genetic correlation with

HDL cholesterol. The directions of genetic correlation with the other two categories of traits are

mixed (**Figure S7**). These results broadly supported our previous results indicating the role of

CNS and blood/immune related mechanisms in the genetics of PA traits.

Discussion

In summary, our study provided novel insights to genetic architecture of physical activity

through genome-wide association analysis of an extensive set of accelerometry based PA

phenotypes, derived in the UK biobank study, and a series of follow-up genomic analyses. We

identified a total of six novel loci, most of which were associated with PA phenotypes not

considered in previous studies (Dashti et al., 2019; Doherty et al., 2018; Jones et al., 2019;

Klimentidis et al., 2018). Our analysis also identified novel phenotypes associated with the

known loci. Further, we provided multiple independent lines of evidence that genetic

mechanisms for association for PA involve the blood and immune system.

Compared to the 15 loci identified by the two previous GWASs on accelerometry-based PA

(Doherty et al., 2018; Klimentidis et al., 2018), the novel loci we discovered have increased the

number of PA susceptibility loci by 33%. Most of the novel loci were connected to the

expression of genes, pseudogenes or long non-coding RNAs (lncRNA, **Table 2 and 3**), and

C3orf62 was also supported by evidence of colocalization (**Table S3**). The novel locus indexed by

rs9818758 overlaps with the TWAS locus index by *C3orf62*. Though it was unclear how *C3orf62*

is involved in PA, two secondary genes in the locus, *ARIH2* and *DAG1* (**Table 3**), appeared to be

involved in the following biological processes: *ARIH2* was found to be essential for

embryogenesis by regulating the immune system(Lin et al., 2013); *DAG1* was found to play a

role in the regeneration of skeletal muscles(Cohn et al., 2002). Both processes appeared

relevant for PA. We argue that the two loci above, supported by multiple lines of evidence

including GWAS, TWAS, colocalization and gene functions, should be prioritized in follow-up

studies. Another two novel loci were connected to pseudogene *RN7SKP16* and *PDXDC2P* of

which the function is less clear and may also be worth future investigation. However, Dashti et

al found a transcription factor site variant rs915416 to be associated with sleep duration, which

is approximately 930Kb away from the transcription start site of *RN7SKP1* and might have

potential long range regulatory effects which warrants further study. Among the genes located

in known loci, *PBX3* and *RP11-396F22.1* were highlighted by both TWAS and colocalization

results. *PBX3* is a member of the pre-B cell leukemia (PBX) family which have extensive roles in

early development and some adult processes (Morgan & Pandha, 2020), which could also

modulate its association with PA.

The novel phenotypes in this study provided important insights into the genetic architecture of

PA, which may have been overlooked by previous GWASs on a small number of phenotypes.

The accelerometry-based study by Doherty et al identified the genetic associations with overall

activity, sleep duration and sedentary time(Doherty et al., 2018); the study by Klimentidis et al

studied the average acceleration and the duration of active states(Klimentidis et al., 2018). Our

results found that there can be different genetic architecture for PA during different times of

the day, and there can be unique variants that only affect certain PA patterns, like ASTP, LIPA

and relative amplitude, but not others (**Table 2**). The heritability and genetic correlation can

also vary across different PA phenotypes (**Figures S4 and S7**).

TWAS and tissue-specific heritability enrichment analysis suggested that in addition to the CNS,

the blood and immune system could be also associated with PA. This finding was further

supported by colocalization, gene-set enrichment and genetic correlation analyses. A previous

study(Doherty et al., 2018), which explored enrichment of heritability for PA traits by tissue-

specific gene expression patterns, identified potential modulating role of the CNS,

adrenal/pancreatic and skeletal muscle tissues. Our study, which used a more extended set of

phenotypes and chromatin-state-based annotations, confirmed previous findings and further

highlighted the role of the blood and immune system. Though there is lack of studies on the

effect of immune functions on PA, previous medical literature has established the effect of PA

on immune functions. A study showed that higher PA is associated with elevation of T-

regulatory cells and lower risk for autoimmune diseases(Sharif et al., 2018). Multiple studies

showed that regular moderately intense PA boost immune functions in older adults and

protects against age-related inflammatory disorders(Dhalwani et al., 2016; Duggal, Niemiro,

Harridge, Simpson, & Lord, 2019; Vancampfort et al., 2017).Though the direction of causal

effect may not be the same as that suggested by genetic analyses, these studies supported

broad connections between PA and immune functions. Future studies are needed to better

understand the underlying mechanisms and causal directions.

In addition to the blood and immune system, TWAS and enrichment analysis also suggested

that the digestive system and endocrine system could be involved in modulating the genetic

effects on PA. The literature has also established broad connections between PA and digestive

and endocrine tissues. A previous study found that PA has complex effects on gastrointestinal

health(Peters, De Vries, Vanberge-Henegouwen, & Akkermans, 2001): acute strenuous activity

may provoke gastrointestinal symptoms while low-intensity activity could have benefits.

Interestingly, three TWAS loci that were significant in digestive tissues were associated with PA

phenotypes that are proxies for meal-time activity: *PDXDC2P* with TLA 6am-8am, *RERE* with TLA

6pm-8PM and *KANSL1* with TLA 4pm-6pm (**Table 3**). It was also known that multiple organs in

the endocrine system produce hormones that regulate physiological functions of the body,

which can have complex bidirectional relationships with PA(Ciloglu et al., 2005; Hawkins et al.,

2008; Ennour-Idrissi, Maunsell, & Diorio, 2015; Alessa et al., 2017; Hackney & Saeidi, 2019).

Among the endocrine tissues, thyroid appeared to be modulating the genetic effect of the

largest number of loci. Previous studies showed that TWAS lead genes *PBX3*, *Corf62* and

KANSL1 were highly expressed in thyroid (GTEx et al., 2017), and SNPs near *KANSL1* were found

to be associated with thyroid-stimulating hormone levels (Teumer et al., 2018). Our TWAS

analysis indicated that the genes associated with PA via the blood and immune system tended

to also be associated with the digestive and endocrine systems, but do not usually overlap with

the genes associated with the CNS. This suggests that the blood and immune, digestive and

endocrine systems may be involved in the same broad pathway that affects PA, which is

different from that of CNS.

It is noteworthy that the accelerometry-derived PA phenotypes in this study are not limited to

exercise, but include a variety of broad-sense activity patterns. In fact, we do find that several

phenotypes have strong genetic correlation with sleep, chronotype and other behavioral traits

(**Figure S7**). Therefore, when defining novel loci, we have further excluded those variants

previously reported to be associated with PA, sleep and circadian rhythm. However, due to the

nature of accelerometry, which captures the acceleration of human body, those phenotypes

are still essentially and broadly PA phenotypes, though they can indirectly reflect and be related

to other traits. Hence we still address all of them by PA phenotypes throughout the paper.

This study has a number of limitations. Though we derived a more extensive set of PA

phenotypes than previous studies, information was still lost when collapsing a 7-day continuous

times series of wrist accelerometry into 31 PA phenotypes. The ideal approach would be to

conduct a GWAS utilizing all the information across the 7 days of accelerometry measurements.

Results could outline genetic regulation of a continuous course of PA over time. The current

analysis of TLA during 12 non-overlapping two-hour time intervals during the day, indicated

that different genetic variants may affect PA during different times of the day (**Tables 2 and 3**).

Another limitation is that some of the phenotypes are not directly interpretable. For example,

the PCs of log acceleration are less interpretable than other phenotypes, such as TLA and ASTP.

However, they do reflect important features of physical activity and warrant further

investigations. A potential solution is to obtain proxy measurements that are interpretable and

highly correlated with PC scores.

In conclusion, we conducted association studies on a wide range of PA phenotypes and

identified 5 novel loci associated with PA. We found that in addition to the CNS, the blood and

immune system may also play an important role in the genetic mechanisms of PA, and the

digestive and endocrine systems could also be involved in the blood and immune pathway.

Data Availability

Data supporting the findings of this paper are available upon application to the UK Biobank

study. The summary statistics are publicly available via the GWAS Catalog

(https://www.ebi.ac.uk/gwas/) under accession numbers GCST90061408-GCST90061434.

Supplemental Data

Supplemental Data include seven figures and four tables.

Declaration of Interests

Dr. Ciprian Crainiceanu is consulting with Bayer and Johnson and Johnson on methods

development for wearable devices in clinical trials. The details of the contracts are disclosed

through the Johns Hopkins University eDisclose system and have no direct or apparent

relationship with the current paper. The other authors declare no conflict of interest.

Acknowledgements

The UK Biobank data was accessed via application ID 17712. Research of Drs. Guanghao Qi,

Diptavo Dutta and Nilanjan Chatterjee was supported by an R01 grant from the National

Human Genome Research Institute [1 R01 HG010480-01].

Web Resources

UK Biobank: https://www.ukbiobank.ac.uk/

fastGWA software: https://cnsgenomics.com/software/gcta/#fastGWA

FUSION TWAS software: http://gusevlab.org/projects/fusion/

COLOC R package: https://cran.r-project.org/web/packages/coloc/index.html

LD score regression software: https://github.com/bulik/ldsc

LD Hub: http://ldsc.broadinstitute.org/ldhub/

FUMA GWAS: https://fuma.ctglab.nl/

PLINK: https://www.cog-genomics.org/plink/2.0/

Molecular Signatures Database: http://www.broadinstitute.org/msigdb

Open Targets Genetics: https://genetics.opentargets.org/

References

Alessa, H. B., Chomistek, A. K., Hankinson, S. E., Barnett, J. B., Rood, J., Matthews, C.

E., . . . Tobias, D. K. (2017). Objective Measures of Physical Activity and Cardiometabolic

and Endocrine Biomarkers. *Med Sci Sports Exerc*, *49*(9), 1817-1825.

doi:10.1249/MSS.0000000000001287

Bulik-Sullivan, B., Finucane, H. K., Anttila, V., Gusev, A., Day, F. R., Loh, P. R., . . . Neale, B. M.

(2015a). An atlas of genetic correlations across human diseases and traits. *Nature

Genetics*, *47*(11), 1236-1241. doi:10.1038/ng.3406

Bulik-Sullivan, B. K., Loh, P. R., Finucane, H. K., Ripke, S., Yang, J., Schizophrenia Working Group

of the Psychiatric Genomics Consortium, . . . Neale, B. M. (2015b). LD Score regression

distinguishes confounding from polygenicity in genome-wide association studies. *Nature

Genetics*, *47*(3), 291-295. doi:10.1038/ng.3211

Buniello, A., MacArthur, J. A. L., Cerezo, M., Harris, L. W., Hayhurst, J., Malangone,

C., . . . Parkinson, H. (2019). The NHGRI-EBI GWAS Catalog of published genome-wide

association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Research*,

47(D1), D1005-D1012. doi:10.1093/nar/gky1120

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L. T., Sharp, K., . . . Marchini, J. (2018). The

UK Biobank resource with deep phenotyping and genomic data. *Nature*, *562*(7726), 203-

209. doi:10.1038/s41586-018-0579-z

Ciloglu, F., Peker, I., Pehlivan, A., Karacabey, K., Ilhan, N., Saygin, O., & Ozmerdivenli, R. (2005).

Exercise intensity and its effects on thyroid hormones. *Neuro Endocrinol Lett*, *26*(6), 830-

834. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/16380698

Cohn, R. D., Henry, M. D., Michele, D. E., Barresi, R., Saito, F., Moore, S. A., . . . Campbell, K. P.

(2002). Disruption of DAG1 in differentiated skeletal muscle reveals a role for dystroglycan

in muscle regeneration. *Cell*, *110*(5), 639-648. doi:10.1016/s0092-8674(02)00907-8

Dashti, H. S., Jones, S. E., Wood, A. R., Lane, J. M., van Hees, V. T., Wang, H., . . . Saxena, R.

(2019). Genome-wide association study identifies genetic loci for self-reported habitual

sleep duration supported by accelerometer-derived estimates. *Nature Communications*,

10(1), 1100. doi:10.1038/s41467-019-08917-4

De Moor, M. H., Liu, Y. J., Boomsma, D. I., Li, J., Hamilton, J. J., Hottenga, J. J., . . . Deng, H. W.

(2009). Genome-wide association study of exercise behavior in Dutch and American

adults. *Med Sci Sports Exerc*, *41*(10), 1887-1895. doi:10.1249/MSS.0b013e3181a2f646

Dhalwani, N. N., O'Donovan, G., Zaccardi, F., Hamer, M., Yates, T., Davies, M., & Khunti, K.

(2016). Long terms trends of multimorbidity and association with physical activity in older

English population. *Int J Behav Nutr Phys Act*, *13*, 8. doi:10.1186/s12966-016-0330-9

Doherty, A., Jackson, D., Hammerla, N., Plötz, T., Olivier, P., Granat, M. H., . . . Wareham, N. J.

(2017). Large Scale Population Assessment of Physical Activity Using Wrist Worn

Accelerometers: The UK Biobank Study. *PLoS One*, *12*(2), e0169649.

doi:10.1371/journal.pone.0169649

Doherty, A., Smith-Byrne, K., Ferreira, T., Holmes, M. V., Holmes, C., Pulit, S. L., & Lindgren, C.

M. (2018). GWAS identifies 14 loci for device-measured physical activity and sleep

duration. *Nature Communications*, *9*(1), 5257. doi:10.1038/s41467-018-07743-4

Duggal, N. A., Niemiro, G., Harridge, S. D. R., Simpson, R. J., & Lord, J. M. (2019). Can physical

activity ameliorate immunosenescence and thereby reduce age-related multi-morbidity.

Nat Rev Immunol, *19*(9), 563-572. doi:10.1038/s41577-019-0177-9

ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human

genome. *Nature*, *489*(7414), 57-74. doi:10.1038/nature11247

Ennour-Idrissi, K., Maunsell, E., & Diorio, C. (2015). Effect of physical activity on sex hormones in

women: a systematic review and meta-analysis of randomized controlled trials. *Breast*

Cancer Res, *17*(1), 139. doi:10.1186/s13058-015-0647-3

Finucane, H. K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P. R., . . . Price, A. L.

(2015). Partitioning heritability by functional annotation using genome-wide association

summary statistics. *Nature Genetics*, *47*(11), 1228-1235. doi:10.1038/ng.3404

Finucane, H. K., Reshef, Y. A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., . . . Price, A. L.

(2018). Heritability enrichment of specifically expressed genes identifies disease-relevant

tissues and cell types. *Nature Genetics*, *50*(4), 621-629. doi:10.1038/s41588-018-0081-4

Gamazon, E. R., Wheeler, H. E., Shah, K. P., Mozaffari, S. V., Aquino-Michaels, K., Carroll, R.

J., . . . Im, H. K. (2015). A gene-based association method for mapping traits using

reference transcriptome data. *Nature Genetics*, *47*(9), 1091-1098. doi:10.1038/ng.3367

Giambartolomei, C., Vukcevic, D., Schadt, E. E., Franke, L., Hingorani, A. D., Wallace, C., &

Plagnol, V. (2014). Bayesian test for colocalisation between pairs of genetic association

studies using summary statistics. *PLoS Genetics*, *10*(5), e1004383.

doi:10.1371/journal.pgen.1004383

GTEx Consortium (2017). Genetic effects on gene expression across human tissues. *Nature*,

550(7675), 204-213. doi:10.1038/nature24277

Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B. W., . . . Pasaniuc, B. (2016).

Integrative approaches for large-scale transcriptome-wide association studies. *Nature*

Genetics, *48*(3), 245-252. doi:10.1038/ng.3506

Hackney, A. C., & Saeidi, A. (2019). The thyroid axis, prolactin, and exercise in humans. *Curr*

Opin Endocr Metab Res, *9*, 45-50. doi:10.1016/j.coemr.2019.06.012

Hara, M., Hachiya, T., Sutoh, Y., Matsuo, K., Nishida, Y., Shimanoe, C., . . . Wakai, K. (2018).

Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults.

Med Sci Sports Exerc, *50*(12), 2433-2441. doi:10.1249/MSS.0000000000001712

Hawkins, V. N., Foster-Schubert, K., Chubak, J., Sorensen, B., Ulrich, C. M., Stancyzk, F.

Z., . . . McTiernan, A. (2008). Effect of exercise on serum sex hormones in men: a 12-

month randomized clinical trial. *Med Sci Sports Exerc*, *40*(2), 223-233.

doi:10.1249/mss.0b013e31815bbba9

Jiang, L., Zheng, Z., Qi, T., Kemper, K. E., Wray, N. R., Visscher, P. M., & Yang, J. (2019). A

resource-efficient tool for mixed model association analysis of large-scale data. *Nature*

Genetics, *51*(12), 1749-1755. doi:10.1038/s41588-019-0530-8

Jones, S. E., Lane, J. M., Wood, A. R., van Hees, V. T., Tyrrell, J., Beaumont, R. N., . . . Weedon,

M. N. (2019). Genome-wide association analyses of chronotype in 697,828 individuals

provides insights into circadian rhythms. *Nature Communications*, *10*(1), 343.

doi:10.1038/s41467-018-08259-7

Kim, J., Min, H., Oh, S., Kim, Y., Lee, A. H., & Park, T. (2014). Joint identification of genetic

variants for physical activity in Korean population. *Int J Mol Sci*, *15*(7), 12407-12421.

doi:10.3390/ijms150712407

Klimentidis, Y. C., Raichlen, D. A., Bea, J., Garcia, D. O., Wineinger, N. E., Mandarino, L.

J., . . . Going, S. B. (2018). Genome-wide association study of habitual physical activity in

over 377,000 UK Biobank participants identifies multiple variants including CADM2 and

APOE. *Int J Obes (Lond)*, *42*(6), 1161-1176. doi:10.1038/s41366-018-0120-3

Kyu, H. H., Bachman, V. F., Alexander, L. T., Mumford, J. E., Afshin, A., Estep, K., . . . Forouzanfar,

M. H. (2016). Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic

heart disease, and ischemic stroke events: systematic review and dose-response meta-

analysis for the Global Burden of Disease Study 2013. *BMJ*, *354*, i3857.

doi:10.1136/bmj.i3857

Leroux, A., Di, J., Smirnova, E., Mcguffey, E. J., Cao, Q., Bayatmokhtari, E., . . . Crainiceanu, C.

(2019). Organizing and analyzing the activity data in NHANES. *Stat Biosci*, *11*(2), 262-287.

doi:10.1007/s12561-018-09229-9

Leroux, A., Xu, S., Kundu, P., Muschelli, J., Smirnova, E., Chatterjee, N., & Crainiceanu, C. (2020).

Quantifying the Predictive Performance of Objectively Measured Physical Activity on

Mortality in the UK Biobank. *J Gerontol A Biol Sci Med Sci*. doi:10.1093/gerona/glaa250

Liberzon, A., Subramanian, A., Pinchback, R., Thorvaldsdóttir, H., Tamayo, P., & Mesirov, J. P.

(2011). Molecular signatures database (MSigDB) 3.0. *Bioinformatics*, *27*(12), 1739-1740.

doi:10.1093/bioinformatics/btr260

Lightfoot, J. T., DE Geus, E. J. C., Booth, F. W., Bray, M. S., DEN Hoed, M., Kaprio,

J., . . . Bouchard, C. (2018). Biological/Genetic Regulation of Physical Activity Level:

Consensus from GenBioPAC. *Med Sci Sports Exerc*, *50*(4), 863-873.

doi:10.1249/MSS.0000000000001499

Lin, A. E., Ebert, G., Ow, Y., Preston, S. P., Toe, J. G., Cooney, J. P., . . . Pellegrini, M. (2013).

ARIH2 is essential for embryogenesis, and its hematopoietic deficiency causes lethal

activation of the immune system. *Nature Immunology*, *14*(1), 27-33. doi:10.1038/ni.2478

Loh, P. R., Kichaev, G., Gazal, S., Schoech, A. P., & Price, A. L. (2018). Mixed-model association

for biobank-scale datasets. *Nature Genetics*, *50*(7), 906-908. doi:10.1038/s41588-018-

0144-6

Loh, P. R., Tucker, G., Bulik-Sullivan, B. K., Vilhjálmsson, B. J., Finucane, H. K., Salem, R.

M., . . . Price, A. L. (2015). Efficient Bayesian mixed-model analysis increases association

power in large cohorts. *Nature Genetics*, *47*(3), 284-290. doi:10.1038/ng.3190

McGregor, D. E., Palarea-Albaladejo, J., Dall, P. M., Stamatakis, E., & Chastin, S. F. M. (2019).

Differences in physical activity time-use composition associated with cardiometabolic

risks. *Prev Med Rep*, *13*, 23-29. doi:10.1016/j.pmedr.2018.11.006

Moore-Harrison, T., & Lightfoot, J. T. (2010). Driven to be inactive? The genetics of physical

activity. *Prog Mol Biol Transl Sci*, *94*, 271-290. doi:10.1016/B978-0-12-375003-7.00010-8

Morgan, R., & Pandha, H. S. (2020). PBX3 in Cancer. *Cancers (Basel)*, *12*(2).

doi:10.3390/cancers12020431

Peters, H. P., De Vries, W. R., Vanberge-Henegouwen, G. P., & Akkermans, L. M. (2001).

Potential benefits and hazards of physical activity and exercise on the gastrointestinal

tract. *Gut*, *48*(3), 435-439. doi:10.1136/gut.48.3.435

Piercy, K. L., Troiano, R. P., Ballard, R. M., Carlson, S. A., Fulton, J. E., Galuska, D. A., . . . Olson, R.

D. (2018). The Physical Activity Guidelines for Americans. *JAMA*, *320*(19), 2020-2028.

doi:10.1001/jama.2018.14854

Roadmap, E. C., Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen, A., . . . Kellis, M. (2015).

Integrative analysis of 111 reference human epigenomes. *Nature*, *518*(7539), 317-330.

doi:10.1038/nature14248

Rock, P., Goodwin, G., Harmer, C., & Wulff, K. (2014a). Daily rest-activity patterns in the bipolar

phenotype: A controlled actigraphy study. *Chronobiol Int*, *31*(2), 290-296.

doi:10.3109/07420528.2013.843542

Rock, P., Goodwin, G., Harmer, C., & Wulff, K. (2014b). Daily rest-activity patterns in the bipolar

phenotype: A controlled actigraphy study. *Chronobiol Int*, *31*(2), 290-296.

doi:10.3109/07420528.2013.843542

Rovio, S., Kåreholt, I., Helkala, E. L., Viitanen, M., Winblad, B., Tuomilehto, J., . . . Kivipelto, M.

(2005). Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's

disease. *Lancet Neurol*, *4*(11), 705-711. doi:10.1016/S1474-4422(05)70198-8

Schrack, J. A., Kuo, P. L., Wanigatunga, A. A., Di, J., Simonsick, E. M., Spira, A. P., . . . Zipunnikov,

V. (2019). Active-to-Sedentary Behavior Transitions, Fatigability, and Physical Functioning

in Older Adults. *J Gerontol A Biol Sci Med Sci*, *74*(4), 560-567. doi:10.1093/gerona/gly243

Sharif, K., Watad, A., Bragazzi, N. L., Lichtbroun, M., Amital, H., & Shoenfeld, Y. (2018). Physical

activity and autoimmune diseases: Get moving and manage the disease. *Autoimmun Rev*,

17(1), 53-72. doi:10.1016/j.autrev.2017.11.010

Smirnova, E., Leroux, A., Cao, Q., Tabacu, L., Zipunnikov, V., Crainiceanu, C., & Urbanek, J.

(2019). The predictive performance of objective measures of physical activity derived

from accelerometry data for 5-year all-cause mortality in older adults: NHANES 2003-

2006. *J Gerontol A Biol Sci Med Sci*. doi:10.1093/gerona/glz193

Teumer, A., Chaker, L., Groeneweg, S., Li, Y., Di Munno, C., Barbieri, C., . . . Medici, M. (2018).

Genome-wide analyses identify a role for SLC17A4 and AADAT in thyroid hormone

regulation. *Nature Communications*, *9*(1), 4455. doi:10.1038/s41467-018-06356-1

Vancampfort, D., Koyanagi, A., Ward, P. B., Rosenbaum, S., Schuch, F. B., Mugisha,

J., . . . Stubbs, B. (2017). Chronic physical conditions, multimorbidity and physical activity

across 46 low- and middle-income countries. *Int J Behav Nutr Phys Act*, *14*(1), 6.

doi:10.1186/s12966-017-0463-5

Watanabe, K., Taskesen, E., van Bochoven, A., & Posthuma, D. (2017). Functional mapping and

annotation of genetic associations with FUMA. *Nature Communications*, *8*(1), 1826.

doi:10.1038/s41467-017-01261-5

Watanabe, K., Umićević Mirkov, M., de Leeuw, C. A., van den Heuvel, M. P., & Posthuma, D.

(2019). Genetic mapping of cell type specificity for complex traits. *Nature*

Communications, *10*(1), 3222. doi:10.1038/s41467-019-11181-1

Young, D. R., & Haskell, W. L. (2018). Accumulation of Moderate-to-Vigorous Physical Activity

and All-Cause Mortality. *J Am Heart Assoc*, *7*(6). doi:10.1161/JAHA.118.008929

Zheng, J., Erzurumluoglu, A. M., Elsworth, B. L., Kemp, J. P., Howe, L., Haycock, P. C., . . . Neale,

B. M. (2017). LD Hub: a centralized database and web interface to perform LD score

regression that maximizes the potential of summary level GWAS data for SNP heritability

and genetic correlation analysis. *Bioinformatics*, *33*(2), 272-279.

doi:10.1093/bioinformatics/btw613

Table 1. Description of physical activity phenotypes.

Category	Phenotype abbreviation	Phenotype full name and description
Total volume of activity	TA*	Total acceleration.
	TLA	Total log acceleration. Sum of log(1+acceleration).
Activity during 2-hour windows of the day	TLA 12am-2am, TLA 2am-4am, TLA 4am-6pm, …, TLA 8pm-10pm, TLA 10pm-12am	Total log acceleration in the k-th two-hour window of the day, starting from midnight (12 intervals in total)
Duration of activity patterns	ST*	Duration of sedentary state. Time when the acceleration is less than 30 milli-gravity (mg) is defined as sedentary state.
	LIPA	Duration of light-intensity physical activity (LIPA). LIPA is defined as the time when the acceleration is >=30mg but <100mg.
	MVPA	Duration of moderate to vigorous physical activity (MVPA). MVPA is defined as the time when the acceleration if >=100mg.
Principal components	PC1*, PC2, …, PC6	First 6 principal component of the log-transformed minute-level activity profiles.
Active-sedentary transition patterns	SATP	Sedentary to active transition probability. SATP is the probability of transitioning to active state if the subject is currently sedentary. Active state is defined as acceleration >=30mg and sedentary state is defined as acceleration <30mg.
	ASTP	Active to sedentary transition probability. SATP is the probability of transitioning to sedentary state if the subject is currently active.
Circadian rhythm proxies	DARE	Dynamic activity ratio estimate. Total log-acceleration during 8am-8pm as proportion of TLA for the whole day.
	M10*	Average log acceleration during the ten most active hours of the day
	L5	Average log acceleration during the five least active hours of the day
	Timing of M10	Mid-point of the ten most active hours of the day
	Timing of L5	Mid-point of the five least active hours of the day
	Relative amplitude	Relative amplitude = (M10-L5)/(M10+L5). It measures the difference between daytime activity and activity during sleep.

All the phenotypes are first calculated at the day level and then averaged across 7 days.

* Excluded from genetic association analyses due to high correlation (>0.8) with TLA.

Table 2. Significant loci associated with physical activity in single-variant analysis.

Lead variant[a]	Chromosome region	Base pair	Minor allele	Major allele	Minor allele frequency	Nearest coding gene and distance[d]	Significantly associated traits[c]	Previous studies that discovered the locus
Novel loci[b]								
rs3836464	3p25.3	10454772	CA	C	0.274	SEC13 (92kb)	ASTP (p=2.2e-09, b=-0.032)	
rs9818758	3p21.31	49382925	A	G	0.171	USP4 (4.8kb)	Relative amplitude (p=2.1e-09, b=-0.036)	NA
3:131647162_TA_T	3q22	131647162	T	TA	0.466	CPNE4 (357kb)	TLA 2am-4am (p=2.2e-09, b=0.029)	
Known loci								
rs1144566	1q25.3	182569626	T	C	0.030	RGS16 (3.9kb)	Timing of L5 (p=5e-10, b=-0.086)*	4
rs301799	1p36.23	8489302	C	T	0.422	SLC45A1 (111kb)	TLA 6pm-8pm (p=1.7e-09, b=0.027)*	OTG[e]
rs113851554	2p14	66750564	T	G	0.050	MEIS1 (90kb)	TLA 12am-2am (p=6.7e-37, b=0.138)*, TLA 2am-4am (p=7.9e-39, b=0.142)*, L5 (p=1.3e-33, b=0.13)*, Relative amplitude (p=6.9e-15, b=-0.082)*, Timing of L5 (p=5.4e-22, b=0.105)*	1,4
rs2909950	5q33.1	151886147	A	G	0.418	NMUR2 (73kb)	TLA 6pm-8pm (p=9.4e-10, b=-0.028)*	4
rs12717867	5q33.1	152412845	G	A	0.453	GRIA1 (456kb)	LIPA (p=6.2e-10, b=-0.029)*	4
rs9369062	6p21.2	38437303	C	A	0.292	BTBD9 (171kb)	TLA 12am-2am (p=1.5e-12, b=-0.037)*, TLA 2am-4am (p=2.3e-10, b=-0.033)*	4
rs2006810	7q11.22	69902152	C	T	0.395	GALNT17 (695kb)	TLA 8pm-10pm (p=8.1e-11, b=-0.031)*	1,4
rs1268539	9q33.3	128195657	A	C	0.419	GAPVD1 (172kb)	TLA (p=1.1e-10, b=0.03), LIPA (p=3.2e-10, b=0.03)*	1
rs564819152	10p12.31	21820650	G	A	0.320	SKIDA1 (5kb)	TLA 8am-10am (p=1.4e-09, b=-0.031)*	1,3
rs2138543	12q12	39298423	A	T	0.477	CPNE8 (2.8kb)	TLA 6am-8am (p=3.9e-11, b=0.03)*, PC2 (p=1.5e-09, b=-0.029)*	4
rs12927162	16q12.2	52684916	G	A	0.277	TOX3 (103kb)	TLA 10pm-12am (p=1e-09, b=0.032)*	4
rs2532402	17q21.31	44304130	G	C	0.221	KANSL1 (1.4kb)	TLA (p=1.5e-12, b=0.04), MVPA (p=1.9e-10, b=0.035)	1,2,3,4
rs3837946	19p13.2	9955920	TTTTG	T	0.475	PIN1 (10kb)	TLA (p=1.3e-11, b=-0.032), LIPA (p=1.6e-09, b=-0.029)*	1,3

[a]Significant variants (single nucleotide polymorphism (SNP) or insertion/deletion) are defined as those with fastGWA p-value $< 2.63 \times 10^{-9}$ (Bonferroni corrected for 19 independent traits). LD clumping was performed at $r^2 < 0.1$ and lead variants of different loci were required to be >500kb apart.

[b]A locus is defined as novel if its lead variant is >500kb from the lead variant of any loci identified in one of the previous GWAS: 1) Doherty et al study on a smaller set of accelerometry-based physical activity; 2) Klimentidis et al study on self-reported and accelerometry based physical activity; 3) Dashti et al study on self-reported sleep duration; 4) Jones et al study on circadian rhythm. Loci that were discovered by the four above studies are marked with 1, 2, 3 and 4 in the last column, respectively. Novel phenotypes associated with known loci are marked with *.

[c]Traits are followed by p-values (p) and effect sizes (b). The effects size b is for the effect of minor allele vs major allele, e.g. a positive b indicates the minor allele is the trait-increasing allele. TLA: total log-acceleration. ASTP: active-to-sedentary transition probability. MVPA: moderate-to-vigorous physical activity. PC2: second principal component. LIPA: light-intensity physical activity.

[d]Nearest coding gene is defined as the protein-coding gene whose transcription start site (TSS) is closest to the variant.

[e]This locus was not reported by the four studies we catalogued but was associated with daytime nap, as reported by Open Targets Genetics (OTG). See Table S4 for details.

Table 3. Significant loci associated with physical activity identified in transcriptome-wide association studies (TWAS).

Most significant gene[a]	CHR	Gene start	Gene end	Locus	Minimum p-value across traits and tissues	Secondary genes in the locus with comparable associations[c]	Trait and tissue[d]	Previous studies that discovered the locus
Novel loci[b]								
RN7SKP16	1	33802167	33802465	1p35.1	2.72E-09	-	MVPA (CNS)	
PDXDC2P	16	70069541	70098679	16q22.1	1.81E-09	-	TLA 6am-8am (Blood/Immune, Digestive)	NA
Known loci								
RERE†	1	8412457	8877702	1p36.23	5.28E-09	ENO1-IT1, RP5-1115A15.1†	TLA 6pm-8pm (Digestive, Other, Blood/Immune)	GWAS
RP5-1160K1.6	1	110171118	110171939	1p13.3	7.68E-10	-	PC2 (CNS)	4
C3orf62†	3	49306219	49315263	3p21.31	3.28E-10	RP11-694I15.7, ARIH2, DAG1, SLC25A20	Relative amplitude (Digestive, Endocrine)	GWAS
CTC-467M3.3	5	87988462	87989789	5q14.3	1.39E-08	-	ASTP (CNS)	1,4
NMUR2	5	151771093	151812929	5q33.1	8.88E-10	-	TLA 6pm-8pm (CNS, Other)	4,GWAS
PBX3†	9	128509624	128729656	9q33.3	3.04E-10	MAPKAP1	LIPA (Blood/Immune, Digestive, Endocrine, Other); SATP (Blood/Immune, Other, Digestive, Endocrine); TLA (Blood/Immune, Other, Digestive, Endocrine)	1,GWAS
DNAJC1	10	22045466	22292698	10p12.31	3.20E-09	CASC10†	TLA 8am-10am (CNS)	1,3,GWAS
RP11-396F22.1†	12	39300253	39303394	12q12	9.89E-10	-	Timing of M10 (CNS), PC2 (CNS, Other), TLA 6am-8am (CNS, Other)	4,GWAS
FMNL1	17	43299590	43324633	17q21.31	2.82E-09	CTD-2020K17.1	TLA (CNS)	1,4
KANSL1	17	44107282	44302733	17q21.31	2.17E-13	KANSL1-AS1, RP11-798G7.8	MVPA (CNS, Blood/Immune, Other, Digestive, Musculoskeletal/connective); TLA 4pm-6pm (Blood/Immune, Other, Digestive); TLA (Adipose, CNS, Blood/Immune, Other, Digestive, Cardiovascular, Musculoskeletal/connective, Endocrine)	1,2,3,4,GWAS
ZNF846	19	9862669	9903856	19p13.2	2.22E-10	OLFM2, PIN1, CTD-2623N2.3	TLA (Other)	1,3,GWAS
JUND	19	18390563	18392432	19p13.11	3.18E-09	-	SATP (Other)	4
LINC00634	22	42348169	42354937	22q13.2	4.85E-11	-	TLA (CNS)	4

[a] For each gene, we apply Benjamini-Hochberg (BH) correction across all trait-tissue pairs. Significant genes are defined as those with BH corrected p-value $< 2.5 \times 10^{-6}$. Significant genes are clustered using an approach similar to LD clumping so that different loci, marked by the transcription start site (TSS) of the lead gene, are >1Mb apart. Gene symbols are italicized. † Genes of which the eQTL signal colocalizes with GWAS signal of the most significantly associated phenotype.

[b] A locus is defined as novel if its flanking region ($\pm$500kb from TSS) does not harbor a variant reported in Table 2 or any of the four previous studies considered in Table 2. Loci that were discovered by the above studies are marked with 1, 2, 3 and 4 in the last column, respectively. Loci that were discovered by our single-variant analysis (Table 2) are marked with "GWAS" in the first column.

[c] Other genes in the cluster with minimum p-value less that 10 times the minimum p-value of the lead gene are shown in column "Secondary genes in the locus with comparable associations". See Table S2 for all the significant tissue-trait pairs.

[d] CNS: central nervous system. TLA: total log-acceleration. ASTP: active-to-sedentary transition probability. SATP: sedentary-to-active transition probability. MVPA: moderate-to-vigorous physical activity. PC2: second principal component.

Figures

Figure 1. Manhattan plot for 18 traits that are significantly associated with at least one variant at $p < 2.63 \times 10^{-9}$ in single-variant analysis. The red dashed line is $p = 2.63 \times 10^{-9}$ accounting for the number of independent traits. The blue dashed line is the standard genome-wide significance threshold $p = 5 \times 10^{-8}$. Three novel loci that have not been discovered in previous GWAS of physical activity, sleep duration and circadian rhythm are circled out and annotated by the lead variant and nearest coding gene (see Table 2).

Figure 2: TWAS Manhattan plots for three tissues that harbor the TWAS novel loci. Significant genes with FDR corrected p-values$< 2.5 \times 10^{-6}$ are circled (see Methods for details). Only the PA traits that are significantly associated with at least one variant at $p < 2.63 \times 10^{-9}$ in single-variant analysis are shown.

Figure 3. Tissue-specific heritability enrichment p-values for traits with significant enrichment at FDR < 0.05 in blood and immune tissues. The analysis was conducted using tissue/cell type specific stratified LD score regression based on 6 chromatin-based annotations in 111 tissues and cell types described in Finucane et al, Nature Genetics 2018 (PMID: 29632380). Each dot corresponds to an annotation in a tissue or cell type. A complete list of tissue and cell types is provided in Supplementary Table 7 of the above paper. Black line corresponds to FDR < 0.05 (-log(p-value)=3.83) across all combinations of trait, tissue, and histone mark. Red line corresponds to p = 0.05. See Figure S6 for the enrichment p-values for the rest of the traits. CNS: central nervous system.