Title: Worries about COVID-19 infection and psychological distress at work and

while commuting

Masamichi Uehara[1], M.D., Tomohiro Ishimaru[1], M.D., M.P.H., Ph.D., Hajime Ando[2],

M.D., M.O.H., Ph.D., Seiichiro Tateishi[3], M.D., Ph.D., Hisashi Eguchi[4], M.D., M.B.A.,

Ph.D., Mayumi Tsuji[5], M.D., Ph.D., Koji Mori[6], M.D., Ph.D., Shinya Matsuda[7], M.D.,

Ph.D., and Yoshihisa Fujino[1], M.D., M.P.H., Ph.D., for the CORoNaWork Project

[1] Department of Environmental Epidemiology, Institute of Industrial Ecological

Sciences, University of Occupational and Environmental Health, Japan

[2] Department of Work Systems and Health, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

[3] Department of Occupational Medicine, School of Medicine, University of

Occupational and Environmental Health, Japan

[4] Department of Mental Health, Institute of Industrial Ecological Sciences, University of

Occupational and Environmental Health, Japan

[5] Department of Environmental Health, School of Medicine, University of Occupational

and Environmental Health, Japan

[6] Department of Occupational Health Practice and Management, Institute of Industrial

Ecological Sciences, University of Occupational and Environmental Health, Japan

[7] Department of Preventive Medicine and Community Health, School of Medicine,

University of Occupational and Environmental Health, Japan

Author contribution:

MU; Writing the manuscript, TI; creating the questionnaire, review of manuscripts, and

advice on interpretation, HA; review of manuscripts and advice on interpretation, ST,

HE, MT, KM and SM; Review of manuscripts, advice on interpretation, and funding for

research, YF; Overall survey planning, creating the questionnaire, analysis, and drafting

the manuscript

Correspondence to Yoshihisa Fujino[1], M.D., M.P.H., Ph.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

Email: zenq@med.uoeh-u.ac.jp

Funding Sources

This study was supported and partly funded by the University of Occupational and

Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan); The

Development of Educational Materials on Mental Health Measures for Managers at

Small-sized Enterprises; Health, Labour and Welfare Sciences Research Grants;

Comprehensive Research for Women's Healthcare (H30-josei-ippan-002); Research for

the Establishment of an Occupational Health System in Times of Disaster

(H30-roudou-ippan-007), scholarship donations from Chugai Pharmaceutical Co., Ltd.,

the Collabo-Health Study Group, and Hitachi Systems, Ltd.

Conflict of Interest:

Dr. Fujino received personal fees from Sompo Health Support Inc., outside the

submitted work.

Acknowledgements

The current members of the CORoNaWork Project, in alphabetical order, are as

follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira

Ogami, Dr. Arisa Harada, Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr.

Kazunori Ikegami, Dr. Kei Tokutsu, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke

Mafune, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Ms. Ning Liu,

Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr. Shinya Matsuda, Dr.

Tomohiro Ishimaru, and Dr. Tomohisa Nagata. All members are affiliated with the

University of Occupational and Environmental Health, Japan.

Ethical approval: This study was approved by the ethics committee of the University

of Occupational and Environmental Health, Japan（reference No. R2-079 and R3-006）.

Running title: worries about COVID-19 infection and psychological distress

Abstract

Objective: This study examined the relationship between worry about COVID-19

infection in the workplace and while commuting to work and psychological distress in

Japan.

Methods: An internet monitor study was conducted. Out of a total of 33,302

participants, 26,841 people were included. The subjects were asked single-item

questions about whether they were worried about COVID-19 infection in general, at

work and while commuting to work. Kessler 6 (K6) was used to assess psychological

distress.

Results: The OR was significantly higher in association with worry about infection in

the workplace at 1.71 (95%CI 1.53–1.92) and worry about infection while commuting

at 1.49 (95%CI 1.32–1.67).

Conclusions: This study suggests the need for psychological intervention to reduce

worry about infection in response to public mental health challenges associated with the

COVID-19 pandemic.

Keywords: Anxiety, COVID-19, Japan**,** Occupational Health, Psychological Distress

Introduction

Introduction

Mental health problems associated with the COVID-19 pandemic are an

emerging public health issue[1]. Outbreaks increase generalized fear, community anxiety,

and panic symptoms[2,3]. Previous studies have reported a higher prevalence of

psychological symptoms such as depression and anxiety among individuals who

experienced lockdown during the COVID-19 pandemic [1,4–7]

Containment measures against pandemics like COVID-19 have a strong impact

on individuals' daily lives and psychological well-being[8]. "A new normal" of avoiding

the "three Cs" (crowded places, close-contact settings, and confined and enclosed

spaces) has been recommended as an effective countermeasure against infection[8].

However, such measures also distance people from each other, which can lead to

decreased communication and socializing. In addition, COVID-19 has also brought

about economic recession and job insecurity. Together, these effects are thought to have

contributed to an increase in loneliness, worry, fear, anxiety, and stress among

individuals.

In situations like the current pandemic, worry of infection is a common feeling

among the majority of the population, given that various aspects of daily life pose

potential risk of infection. An increase in infections can trigger psychological symptoms

such as generalized fear and widespread community anxiety[2,3]. Psychological symptoms

such as persistent worrying, feeling overwhelmed by emotions, restlessness and

irritability emerge within the population in the form of anxiety, panic attacks,

depression and suicide[9,10]. Worry of infection is an early sign of psychological distress.

For workers, the workplace and commuting[11–14] are considered particularly risky

opportunities for infection. For this reason, most workplaces are taking various infection

control measures, including limiting the number of visitors, restricting business trips,

encouraging telecommuting, ensuring sufficient office ventilation, and installing

partitions in customer-facing spaces.[15,16] However, some workplaces are not taking

sufficient recommended measures for business reasons or due to cost.[17,18] Naturally,

infection control efforts in the workplace influence workers' worry of infection. When

infection control does not meet expectations, workers' anxiety and depression are

high.[19] In contrast, intensive workplace measures against COVID-19 are associated with

lower psychological distress among employees.[15,19] In addition, anxiety and depression

are higher among workers who do not telecommute and those with interpersonal tasks.[19]

Therefore, we hypothesized that worry of infection in the workplace and while

commuting affect the psychological distress of workers. In this study, we examined the

relationship between workers' worry about infection and psychological distress in the

midst of a rapid outbreak of COVID-19 in Japan.

Method

Study design and subjects

This cross-sectional, internet monitor study was conducted on December 22–26,

2020, when Japan experienced its third wave of COVID-19 infection. Details of the

protocol of this survey are reported elsewhere[20]. Briefly, data were collected from

workers who had employment contracts at the time of the survey and were selected

based on prefecture, job type, and sex.

Out of a total of 33,302 people who participated in the survey, 27,036 people

were included in the study after removing those found to have provided fraudulent

responses. Participants were excluded if they had exceptionally short response times (≤6

minutes), exceptionally low body weight (<30 kg), exceptionally short height (<140

cm), provided inconsistent responses to comparable questions (e.g., inconsistent

responses to questions on marital status and area of residence), and gave incorrect

responses to a question included to identify fraudulent responses (select the third largest

number from among the following five numbers). After further excluding 195

individuals who indicated in the survey that they had already been infected with

COVID-19, a total of 26,841 individuals (13,713 males and 13,128 females) were

included in the current analysis. This study was approved by the ethics committee of the

University of Occupational and Environmental Health, Japan (reference No. R2-079).

Informed consent was obtained in the form of the website.

Assessment of worry about COVID-19 infection

Three single-item questions were used to determine whether or not the subjects

were worried about infection. One question inquired about general worry of COVID-19

infection, while the other two were situation dependent, inquiring about worry of

infection at work and while commuting to work. The questions were, "Are you worried

about being infected with COVID-19?" "Are you worried about being infected while

working at your workplace?" and "Are you worried about being infected while

commuting to work?" The participants responded "yes" or "no" to the questions.

Assessment of psychological distress

Kessler 6 (K6) was used to assess psychological distress[21]. The validity of the

Japanese version of the K6 has been confirmed[22]. In the present study, a K6 score of 5

or higher was used as the cutoff for mild psychological distress, and a score of 13 or

higher as the cutoff for severe psychological distress.

Other covariates

We selected potential confounding factors which are possibly related to anxiety

and psychological distress. The following survey items were considered confounding

factors: age, sex, marital status (married, unmarried, bereaved/divorced), occupation

(mainly desk work, jobs mainly involving interpersonal communication, and mainly

labor), number of employees, educational background, equivalent income (household

income divided by the square root of household size), smoking status, alcohol

consumption, frequency of telecommuting, and use of public transportation when

commuting. The questionnaire also asked the following questions: "Have you been a

close contact of someone infected with COVID-19?" and "Do you know of anyone

close to you (friends or family) who has been infected with COVID-19?"

The questionnaire also asked the participants to rate their company's infection

control measures using the question "Do you think your company has taken adequate

infection control measures for its employees?" Participants responded on a four-point

scale: "yes," "somewhat," "not really," "no."

In addition, the cumulative incidence rate of COVID-19 infection one month

prior to conduct of the survey in the prefectures of residence was used as a

community-level variable. Information was collected from the websites of public

institutions.

Statistical analysis.

The odds ratios (ORs) of psychological distress associated with worry about

infection were estimated using a multilevel logistic model nested in the prefectures of

residence. Psychological distress was defined as a K6 score of 5 or higher and 13 or

higher.

The multivariate model was adjusted for sex, age, education, equivalent

household income, occupation, number of business establishments, smoking status,

alcohol consumption, frequency of telecommuting, use of public transportation when

commuting, perceived assessment of workplace infection control efforts, presence of

infection among acquaintances, and experience of being a close contact. The incidence

rate of COVID-19 by prefecture was also used as a prefecture-level variable.

We further estimated the multivariate ORs of psychological distress associated

with use of public transportation, with adjustment for all factors except worry about

infection during commuting, because adjusting for worry about infection would be an

over-adjustment.

A p value less than .05 was considered statistically significant. All analyses

were conducted using Stata (Stata Statistical Software: Release 16; StataCorp LLC, TX,

USA).

Results

Table 1 shows the characteristics of the subjects according to their worries about

infection. A total of 75% of the participants were worried about infection. While 50% of

participants were worried about being infected in the workplace, 32% were worried

about being infected while commuting to work. Women were more worried about

infection than men. There were no substantial differences in income, educational

background, or lifestyle by worry about COVID-19 infection in general, while at work,

or while commuting. Participants who indicated that they worried about infection in the

workplace and while commuting shared similar characteristics.

Table 2 shows the ORs of psychological distress. In the age-adjusted model for

moderate psychological distress defined by a K6 score of 5 or higher, general worry

about COVID-19 infection (OR=1.29, 95%CI 1.22–1.37), worry about infection at work

(OR=1.79, 95%CI 1.70–1.88), and worry about infection while commuting (OR=1.64,

95%CI 1.55–1.73) all showed significant associations with psychological distress.

Multivariate analysis showed that the OR of moderate psychological distress associated

with general worry about COVID-19 infection was 0.97 (95% CI 0.91–1.04). The ORs

were significantly higher in association with worry about infection in the workplace at

1.57 (95%CI 1.57–1.48) and worry about infection while commuting at 1.49 (95%CI

1.38–1.60).

The sex- and age-adjusted OR of severe psychological distress, defined by a

K6 score of 13 or higher, associated with general worry about COVID-19 infection was

1.05 (95%CI 0.95–1.15). ORs were significantly higher in association with worry about

infection in the workplace (OR=1.76, 95%CI 1.61–1.92) and worry about infection

while commuting (OR=1.65, 95%CI 1.51–1.80). Multivariate analysis of severe

psychological distress showed similar results.

We also examined situations we predicted could enhance worry. Being in close

contact with an infected person increased the ORs of severe psychological distress;

however, having acquaintances who had been infected was not significantly associated

with severe psychological distress. The OR of severe psychological distress associated

with using public transportation for commuting to work was 1.25 (95%CI 1.25–1.39).

Discussion

This study showed that while general worry about COVID-19 infection was not

associated with psychological distress, worry about infection at work and while

commuting were associated with moderate and severe psychological distress.

The spread of COVID-19 infection is causing people to worry about infection in

a number of everyday situations. In the present study, 75% of participants were worried

about infection, suggesting that COVID-19 infection is a major source of public stress.

However, in this situation, worrying about infection is not unhealthy, but rather, a

natural reaction to current events. Such worry about infection may lead to preventive

behaviors[23–25]. In this situation, we found that having general worry about COVID-19

infection did not lead to psychological distress.

However, our study indicates that certain everyday situations, such as working

in the workplace and commuting to work, are particularly worrisome in relation to

infection. We found that worry about infection at work was associated with moderate

and severe psychological distress. There are many opportunities in the workplace for

direct contact with others, such as while communicating with colleagues and providing

services to customers, and for indirect contact with people through objects. Previous

studies have also reported an increase in anxiety and depression in interpersonal work.[19]

Additionally, in places like offices, workers share a relatively small space. Thus, the

fact that individuals have limited control over maintaining the recommended three Cs

strategy for infection prevention at work can be a major cause of stress. A sense of

security and trust that workplaces can be made safe in the event of a pandemic may

counteract any negative feelings.[18]

The present study also showed that worry about infection during commuting was

associated with moderate and severe psychological distress. This result is consistent

with previous studies that found a higher prevalence of anxiety and depressive

symptoms among workers who have to commute to work.[18,19] Commuting to work is

considered one way in which risk of infection can increase[26], because individuals come

into contact with an unspecified number of people, are unable to maintain social

distancing, and are present in a space that feels relatively enclosed, such as a bus or

train. In fact, 47% of participants who were worried about infection while commuting

experienced moderate psychological distress, and 23% experienced severe

psychological distress. These results suggest that reducing worries about infection in the

workplace and during commuting is important for reducing public mental health issues

caused by COVID-19[27].

However, our findings also suggest the difficulty of reducing psychological

distress due to worry about infection. We found that worries about infection in the

workplace and while commuting to work were associated with psychological distress

even after adjusting for assessment of adequate infection control measures in the

workplace and use of public transportation. This result is consistent with a previous

study that showed that workers' anxiety about infection differs in different workplaces,

and cannot be explained by workplace-related COVID-19 risk management strategies.[18]

These results suggest that workers worry regardless of whether or not they use public

transportation to commute to work or have adequate infection prevention measures

installed in their workplace, and that these worries are related to psychological distress.

In fact, 38% of those who were worried about their commute did not use public

transportation. While we found that public transportation use was associated with

psychological distress, this association disappeared after adjusting for general worry

about COVID-19 infection. This finding supports the idea that worry, rather than facts,

is associated with psychological distress. Worry about infection is not only based on

rational judgment, but is also an intuitive feeling. Therefore, to reduce excessive worry

about infection and the resulting psychological distress, it is necessary to both promote

infection prevention measures and adjust individuals' perception of their worries.

Recent studies have provided useful suggestions for reducing anxiety about

COVID-19 infection. Establishing infection control policies and implementing infection

control measures in the workplace can be beneficial for the mental health of workers,

and a sense of trust and security that infection control measures in the workplace are

adequate will alleviate anxiety.[15,18,19] Use of personal protective equipment when

conducting interpersonal tasks and access to testing in case of suspected infectious

contact can also ease anxiety.[18,19] Additionally, public awareness, such as through

access to correct information and avoiding excessive exposure to the media, may be

useful for reducing anxiety.[28–30] Cognitive-behavioral therapy is also considered an

effective method for such a purpose[31].

Limitations of this study warrant mention. First, the generalizability of the

results is uncertain because this study was conducted through internet monitors.

However, we attempted to reduce as much bias in the target population as possible by

sampling according to region, job type, and prefecture based on the infection incidence

rate. Second, the study asked simple questions about individuals' worries about

infection. Dichotomous yes/no responses can prevent a nuanced understanding of the

real sensations experienced by individuals. However, the questions were not intended to

diagnose psychiatric symptoms or psychiatric illnesses. Participants who indicated that

they were worried may also exhibit psychiatric symptoms, such as anxiety, depression,

and panic; however, distinguishing these symptoms was outside the scope of this study.

Third, worry about infection in the workplace is thought to be greatly influenced by the

type of job and the environment in which one works. Health care workers are a typical

example of a population that is at high risk of infection and are more likely to

experience psychological distress[10]. In this study, we only adjusted for simple variables

such as job type (desk work and physical work) and the number of employees in the

workplace. Fourth, because this was a cross-sectional study, the temporal relationship

between worry about infection and psychological distress is unclear. Excessive anxiety

and depression may be symptoms of psychological distress. Previous studies have

shown that people with a history of psychiatric disorders are more anxious and show

greater psychological distress during the COVID-19 pandemic[32]. However, because it is

easier to confirm feelings of worry than to make a psychiatric diagnosis, we believe that

it is useful to identify the presence of worry among workers for determining health

status.

In conclusion, while general worry about infection was not associated with

psychological distress, worry about infection in specific situations such as in the

workplace and while commuting was associated with moderate and severe

psychological distress. These worries and the associated psychological and emotional

responses may not be mediated by actual infection control measures taken in the

workplace or use of public transportation. Thus, despite the continued spread of

COVID-19 infection and an increase in anxiety in the population, situational fear, but

not generic fear, is correlated with poor mental health. Such knowledge will be useful

for designing future policies to manage mental health. The results of this study suggest

the need for psychological approaches to reduce worries about infection in response to

the mental health challenges associated with the COVID-19 pandemic.

References:

1.   Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated
Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19)
Epidemic among the General Population in China. *Int J Environ Res Public*
Health. 2020;17(5). doi:10.3390/ijerph17051729

2.   Varatharaj A, Thomas N, Ellul MA, et al. Neurological and neuropsychiatric
complications of COVID-19 in 153 patients: a UK-wide surveillance study. *Lancet*
Psychiatry. 2020;7(10):875-882.

3.   Serafini G, Parmigiani B, Amerio A, Aguglia A, Sher L, Amore M. The
psychological impact of COVID-19 on the mental health in the general population.
QJM. Published online June 22, 2020. doi:10.1093/qjmed/hcaa201

4.   Tang W, Hu T, Hu B, et al. Prevalence and correlates of PTSD and depressive
symptoms one month after the outbreak of the COVID-19 epidemic in a sample of
home-quarantined Chinese university students. *J Affect Disord*. 2020;274:1-7.

5.   Losada-Baltar A, Jiménez-Gonzalo L, Gallego-Alberto L, Pedroso-Chaparro MDS,
Fernandes-Pires J, Márquez-González M. "We're staying at home". Association of
self-perceptions of aging, personal and family resources and loneliness with
psychological distress during the lock-down period of COVID-19. *J Gerontol B*
Psychol Sci Soc Sci. Published online April 13, 2020. doi:10.1093/geronb/gbaa048

6.   Mazza C, Ricci E, Biondi S, et al. A Nationwide Survey of Psychological Distress
among Italian People during the COVID-19 Pandemic: Immediate Psychological
Responses and Associated Factors. *Int J Environ Res Public Health*. 2020;17(9).
doi:10.3390/ijerph17093165

7.   Moccia L, Janiri D, Pepe M, et al. Affective temperament, attachment style, and
the psychological impact of the COVID-19 outbreak: an early report on the Italian
general population. *Brain Behav Immun*. 2020;87:75-79.

8.   World Health Organization. Coronavirus disease (COVID-19) advice for the
public. Accessed January 27, 2021.
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-pub
lic

9.   Greenberg N, Brooks SK, Wessely S, Tracy DK. How might the NHS protect the
mental health of health-care workers after the COVID-19 crisis? *Lancet*
Psychiatry. 2020;7(9):733-734.

10.   Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsi E, Katsaounou P.
Prevalence of depression, anxiety, and insomnia among healthcare workers during
the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav*
Immun. 2020;88:901-907.

11.   Tan L, Ma C. Choice behavior of commuters' rail transit mode during the
COVID-19 pandemic based on logistic model. *Journal of Traffic and*
Transportation Engineering (English Edition). 2021;8(2):186-195.

12.   Hu M, Lin H, Wang J, et al. Risk of Coronavirus Disease 2019 Transmission in
Train Passengers: an Epidemiological and Modeling Study. *Clin Infect Dis*.
2021;72(4):604-610.

13.   Ontario Agency for Health Protection and Promotion (Public Health Ontario).
Public Transport and COVID-19 – What We Know so Far. Queen's Printer for
Ontario; 2020.

14.   World Health Organization. Supporting healthy urban transport and mobility in the
context of COVID-19. Published November 2, 2020. Accessed June 3, 2021.
https://www.who.int/publications/i/item/9789240012554

15.   Sasaki N, Kuroda R, Tsuno K, Kawakami N. Workplace responses to COVID-19
associated with mental health and work performance of employees in Japan. *J*
Occup Health. 2020;62(1):e12134.

16.   Ministry of Health Labour and Welfare, Japan. Checklist for the prevention of 257
COVID-19   spreading   at   workplaces.   Accessed   June   3,   2021.
https://www.mhlw.go.jp/content/000694987.pdf

17.   Ishimaru T, Nagata M, Hino A, et al. Workplace measures against COVID-19
during the winter third wave in Japan: Company size-based differences. *J Occup*
Health. 2021;63(1):e12224.

18.   Nabe-Nielsen K, Nilsson CJ, Juul-Madsen M, Bredal C, Hansen LOP, Hansen
ÅM. COVID-19 risk management at the workplace, fear of infection and fear of

transmission of infection among frontline employees. *Occup Environ Med*.
2021;78(4):248-254.

19. Smith PM, Oudyk J, Potter G, Mustard C. Labour market attachment, workplace
infection control procedures and mental health: A cross-sectional survey of
Canadian non-healthcare workers during the COVID-19 pandemic. *Ann Work*
Expo Health. 2021;65(3):266-276.

20. Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based
health survey in workers during the COVID-19 pandemic in 2020. *medRxiv*.
Published online January 1, 2021:2021.02.02.21249309.

21. Kessler RC, Andrews G, Colpe LJ, et al. Short screening scales to monitor
population prevalences and trends in non-specific psychological distress. *Psychol*
Med. 2002;32(6):959-976.

22. Furukawa TA, Kawakami N, Saitoh M, et al. The performance of the Japanese
version of the K6 and K10 in the World Mental Health Survey Japan. *Int J*
Methods Psychiatr Res. 2008;17(3):152-158.

23. Kwok KO, Li KK, Chan HHH, et al. Community Responses during Early Phase of
COVID-19 Epidemic, Hong Kong. *Emerg Infect Dis*. 2020;26(7):1575-1579.

24. Bowman L, Kwok KO, Redd R, et al. Public perceptions and preventive
behaviours during the early phase of the COVID-19 pandemic: a comparative
study between Hong Kong and the United Kingdom. *bioRxiv*. Published online
August 7, 2020. doi:10.1101/2020.08.06.20169409

25. Magdy DM, Metwally A, Magdy O. Assessment of Community Psycho-behavioral
Responses during the outbreak of novel coronavirus (2019-nCoV): A cross
sectional study. *Research Square*. Published online April 29, 2020.
doi:10.21203/rs.3.rs-25146/v1

26. Zhang J. Transport policymaking that accounts for COVID-19 and future public
health threats: A PASS approach. *Transp Policy*. 2020;99:405-418.

27. Dam P, Mandal S, Mondal R, Sadat A, Chowdhury SR, Mandal AK. COVID-19:
Impact on transport and mental health. *J Transp Health*. 2020;19:100969.

28. Sasaki N, Kuroda R, Tsuno K, Kawakami N. Exposure to media and fear and
worry about COVID-19. *Psychiatry Clin Neurosci*. 2020;74(9):501-502.
doi:10.1111/pcn.13095

29. Torales J, O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of
COVID-19 coronavirus and its impact on global mental health. *Int J Soc*
Psychiatry. 2020;66(4):317-320.

30. Mowbray H. In Beijing, coronavirus 2019-nCoV has created a siege mentality.
BMJ. 2020;368:m516.

31. Yue J-L, Yan W, Sun Y-K, et al. Mental health services for infectious disease
outbreaks including COVID-19: a rapid systematic review. *Psychol Med*.
2020;50(15):2498-2513.

32. Yamamoto T, Uchiumi C, Suzuki N, Yoshimoto J, Murillo-Rodriguez E. The
Psychological Impact of "Mild Lockdown" in Japan during the COVID-19
Pandemic: A Nationwide Survey under a Declared State of Emergency. *Int J*
Environ Res Public Health. 2020;17(24). doi:10.3390/ijerph17249382

Table 1. The characteristics of the subjects according to their worries about infection.

	Worry about infection		Worry about infection at work		Worry about infection while commuting	
	No	Yes	No	Yes	No	Yes
Number of subjects	6967	19874	13323	13518	18091	8750
Age, mean (SD)	47.0 (10.3)	47.0 (10.6)	48.3 (10.2)	45.8 (10.7)	47.4 (10.4)	46.2 (10.8)
Sex, male	4099 (58.8%)	9614 (48.4%)	7819 (58.7%)	5894 (43.6%)	9551 (52.8%)	4162 (47.6%)
Marriage status, married	3702 (53.1%)	11207 (56.4%)	7519 (56.4%)	7390 (54.7%)	10196 (56.4%)	4713 (53.9%)
Job type						
Mainly desk work	3402 (48.8%)	9970 (50.2%)	7257 (54.5%)	6115 (45.2%)	8594 (47.5%)	4778 (54.6%)
Jobs mainly involving interpersonal communi-cation	1707 (24.5%)	5163 (26.0%)	2856 (21.4%)	4014 (29.7%)	4570 (25.3%)	2300 (26.3%)
Mainly labor	1858 (26.7%)	4741 (23.9%)	3210 (24.1%)	3389 (25.1%)	4927 (27.2%)	1672 (19.1%)
Equivalent income (million JPY)						
50-249	1612 (23.1%)	4061 (20.4%)	2920 (21.9%)	2753 (20.4%)	4013 (22.2%)	1660 (19.0%)
250-374	1897 (27.2%)	5607 (28.2%)	3578 (26.9%)	3926 (29.0%)	5128 (28.3%)	2376 (27.2%)
375-489	1565 (22.5%)	5013 (25.2%)	3170 (23.8%)	3408 (25.2%)	4426 (24.5%)	2152 (24.6%)
≥490	1893 (27.2%)	5193 (26.1%)	3655 (27.4%)	3431 (25.4%)	4524 (25.0%)	2562 (29.3%)
Educational background						
Junior high	115 (1.7%)	252 (1.3%)	205 (1.5%)	162 (1.2%)	273 (1.5%)	94 (1.1%)
High school	1892 (27.2%)	5018 (25.2%)	3599 (27.0%)	3311 (24.5%)	5113 (28.3%)	1797 (20.5%)
University, graduate school, vocational school, junior college	4960 (71.2%)	14604 (73.5%)	9519 (71.4%)	10045 (74.3%)	12705 (70.2%)	6859 (78.4%)
Current smoker	1818 (26.1%)	5136 (25.8%)	3560 (26.7%)	3394 (25.1%)	4807 (26.6%)	2147 (24.5%)
Alcohol consumption						
6 to 7 days per week	1523 (21.9%)	4092 (20.6%)	3089 (23.2%)	2526 (18.7%)	3932 (21.7%)	1683 (19.2%)
4 to 5 days per week	530 (7.6%)	1513 (7.6%)	1012 (7.6%)	1031 (7.6%)	1309 (7.2%)	734 (8.4%)
2 to 3 days per week	826 (11.9%)	2420 (12.2%)	1635 (12.3%)	1611 (11.9%)	2127 (11.8%)	1119 (12.8%)
Less than 1 day per week	1106 (15.9%)	3417 (17.2%)	2117 (15.9%)	2406 (17.8%)	2960 (16.4%)	1563 (17.9%)
Almost none	2982 (42.8%)	8432 (42.4%)	5470 (41.1%)	5944 (44.0%)	7763 (42.9%)	3651 (41.7%)
Number of employees in the workplace						
1-29	1736 (24.9%)	4386 (22.1%)	3927 (29.5%)	2195 (16.2%)	4615 (25.5%)	1507 (17.2%)
30-99	1768 (25.4%)	5126 (25.8%)	3245 (24.4%)	3649 (27.0%)	4893 (27.0%)	2001 (22.9%)
100-999	1725 (24.8%)	5375 (27.0%)	3095 (23.2%)	4005 (29.6%)	4640 (25.6%)	2460 (28.1%)
≤1000	1738 (24.9%)	4987 (25.1%)	3056 (22.9%)	3669 (27.1%)	3943 (21.8%)	2782 (31.8%)
Experience of being a close contact	56 (0.8%)	148 (0.7%)	79 (0.6%)	125 (0.9%)	122 (0.7%)	82 (0.9%)
Presence of infection among acquaintances	364 (5.2%)	1778 (8.9%)	855 (6.4%)	1287 (9.5%)	1248 (6.9%)	894 (10.2%)
Use of public transportation for commuting	1358 (19.5%)	5884 (29.6%)	2730 (20.5%)	4512 (33.4%)	1761 (9.7%)	5481 (62.6%)
General worry about infection	0 (0.0%)	19874 (100.0%)	7221 (54.2%)	12653 (93.6%)	11895 (65.8%)	7979 (91.2%)
Worry about infection at work	865 (12.4%)	12653 (63.7%)	0 (0.0%)	13518 (100.0%)	6568 (36.3%)	6950 (79.4%)
Worry about infection while commuting	771 (11.1%)	7979 (40.1%)	1800 (13.5%)	6950 (51.4%)	0 (0.0%)	8750 (100.0%)
Moderate psychological distress (K6≥5)	2447 (35.1%)	8240 (41.5%)	4264 (32.0%)	6423 (47.5%)	6516 (36.0%)	4171 (47.7%)
Severe psychological distress (K6≥13)	605 (8.7%)	1814 (9.1%)	860 (6.5%)	1559 (11.5%)	2926 (16.2%)	2047 (23.4%)

Table 2. Associations of worry about infection and psychological distress

	OR of moderate psychological distress (K6≥5)								OR of severe psychological distress (K6≥13)							
	Age-sex adjusted				Multivariate*				Age-sex adjusted				Multivariate*			
	OR	95% CI		p	OR	95% CI		p	OR	95% CI		p	OR	95% CI		p
General worry about infection	1.29	1.22	1.37	<0.001	0.97	0.91	1.04	0.412	1.05	0.95	1.15	0.365	1.10	1.00	1.22	0.060
Worry about infection at work	1.79	1.70	1.88	<0.001	1.57	1.48	1.68	<0.001	1.76	1.61	1.92	<0.001	1.71	1.53	1.92	<0.001
Worry about infection during commuting	1.64	1.55	1.73	<0.001	1.49	1.38	1.60	<0.001	1.65	1.51	1.80	<0.001	1.49	1.32	1.67	<0.001
Experience of being a close contact	1.42	1.08	1.89	0.014	1.47	1.10	1.97	0.010	1.45	0.96	2.19	0.074	1.57	1.02	2.41	0.039
Presence of infection among acquaintances	1.07	0.97	1.17	0.163	1.06	0.96	1.16	0.271	0.97	0.83	1.14	0.730	0.97	0.82	1.14	0.689
Use of public transportation for commuting	1.11	1.04	1.17	0.001	0.86	0.80	0.93	<0.001	1.18	1.07	1.31	0.001	0.94	0.83	1.06	0.336
					1.15**	1.08	1.23	<0.001					1.25**	1.12	1.39	<0.001
"Do you think your company has taken adequate infection control measures for its employees?"																
Yes	reference				reference				reference				reference			
Somewhat	1.23	1.14	1.31	<0.001	1.23	1.14	1.32	<0.001	0.90	0.79	1.01	0.080	0.86	0.76	0.98	0.022
Not really	1.99	1.83	2.16	<0.001	1.93	1.77	2.10	<0.001	1.48	1.29	1.71	<0.001	1.35	1.17	1.56	<0.001
No	2.70	2.44	2.98	<0.001	2.50	2.25	2.77	<0.001	3.21	2.78	3.70	<0.001	2.74	2.36	3.18	<0.001

* The multivariate model included sex, age, education, equivalent household income, occupation, number of business establishments, smoking status, alcohol consumption, frequency of telecommuting, and use of public transportation when commuting, perceived assessment of workplace infection control efforts, presence of infection among acquaintances, and experience of being a close contact. The incidence rate of COVID-19 by prefecture was also used as a prefecture-level variable

** ORs were derived from the multivariate model without worry about infection at work and while commuting.