Probiotic supplements reduce antipsychotic-induced metabolic disturbances in drug-naïve first-episode schizophrenia

Dongyu Kang, M.D.[a#], Fengyu Zhang, PhD. MS[b,c#], Ye Yang, M.D.[a], Chenchen Liu, M.D.[a], Jingmei Xiao, M.D.[a], Yujun Long, M.D.[a], Jing Huang, M.D. PhD.[a], Xingjie Peng, M.D.[a], Weiyan Wang, M.D.[a], Xiaoyi Wang, M.D.[a], John M. Davis, M.D.[d], Jingping Zhao M.D. PhD.[a], Renrong Wu M.D. PhD.[a,e*]

Affiliation/address:

a. National Clinical Research Center for Metabolic Diseases, and Department of Psychiatry, the Second Xiangya Hospital of Central South University, Changsha 410011, Hunan, China.
b. Beijing Huilongguan Hospital and Peking University Huilongguan Clinical Medical School, Beijing, China
c. Global Clinical and Translational Research Institute, Bethesda, MD, USA
d. Psychiatric Institute, University of Illinois, Chicago, IL, USA
e. Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China

[#] Authors contributed equally to this work.

*Corresponding author and E-mail address:

Renrong Wu, MD PhD
Mental Health Institute of the Second Xiangya Hospital
Central South University
Changsha, Hunan
China
E-mail: wurenrong@csu.edu.cn
Tel: +86 15874179855；Fax: +86 73185295214

Word count 4108, 4 Table, 2 Figure.

Keywords: schizophrenia, probiotic, antipsychotic, metabolic dysfunction

ABSTRACT

Probiotic supplements have demonstrated efficacy in improving metabolic abnormalities and may prevent antipsychotic-induced metabolic disturbance and weight gain. A few studies in rodents have found that antipsychotic-induced metabolic dysfunctions are associated with the altered composition of gut microbiota. Here, we conducted a randomized-controlled clinical trial to determine the effectiveness and safety of probiotic supplements on antipsychotic-induced metabolic disturbance and weight gain. Patients with drug-naïve first-episode schizophrenia were randomized to receive either olanzapine plus probiotics or olanzapine monotherapy and scheduled to evaluate with follow-ups for clinical and metabolic profiles. After a treatment of 12 weeks with addition of probiotics, the increase of mean fasting insulin was significantly lower than olanzapine monotherapy. Insulin resistance increased considerably in the olanzapine plus probiotics, but also significantly lower than olanzapine monotherapy. We noted a difference in the increase in body mass index and body weight between treatments at a nominal level of significance, but it became non-significant after adjusting for appetite increase. Probiotics concurrently used with olanzapine is effective and safe in attenuating antipsychotic-induced elevation of fasting insulin and insulin resistance, but not the weight gain in drug-naïve first-episode schizophrenia. Further study is warranted to assess the longer-term maintenance of efficacy and safety.

1. Introduction

The second-generation antipsychotics (SGAs) are widely used as first-line medications to treat and manage patients with schizophrenia. However, antipsychotics-induced weight gain and metabolic dysregulations such as hyperglycemia and hyperlipidemia are of significant concern (Newcomer, 2007). These side effects not only affect the patients' compliance with antipsychotic medications but also cause substantial medical morbidities such as diabetes, cardiovascular disease, stroke, and premature death (De Hert et al., 2011; Foley and Morley, 2011). Antipsychotic drugs have sharply increased in the prescription of children and adolescents and are shown to cause significant weight gain and risk for type 2 diabetes (Maayan and Correll, 2011). Of the SGAs, olanzapine is one of the commonly used and has better efficacy than other drugs in patients with chronic schizophrenia (Lieberman et al., 2005). In a large cohort study about the SGAs Treatment Indications, Effectiveness, and Tolerability in Youth Study (SATIETY), the proportion of individuals who gain more than 7% of weight over the baseline during the first three months is 84% in olanzapine and 64% in risperidone (Correll et al., 2009).

The exact mechanism of antipsychotic-induced weight gain and metabolic disturbance has not been elucidated (Nasrallah, 2008; Newcomer, 2005). Animal studies have indicated that the initial weight gain in individuals using olanzapine is primarily driven by the increases in appetite and food intakes (Davoodi et al., 2009), which is likely caused by the antagonism of histamine H_1 receptors and $5\text{-}HT_{2C}$ in the presence of dopamine D_2 receptors (Kirk et al., 2009); whereas others show that risperidone induced weight gain is partly attributable to the alteration of gut microbiome and

suppression of energy expenditure (Bahra et al., 2015). In addition, antipsychotics can

inhibit the insulin-signaling pathway in the target cells such as muscle cells,

hepatocytes and adipocytes from using glucose, and, therefore, cause insulin

resistance, which is independent of weight gain and food intake that has been

indicated in both patients and healthy controls using antipsychotics (Hahn et al., 2013;

van Winkel et al., 2008). Further, the induced obesity can result in high levels of free

fatty acids (FFA) and inflammation, which can cause insulin resistance, and

antipsychotics can cause direct damage to β-cells, leading to dysfunction and

apoptosis of β-cells that increase insulin resistance (Chen et al., 2017).

Antipsychotic-induced metabolic dysfunctions are likely linked to the altered

composition of the gut microbiota caused by using antipsychotics. The gut microbiota

contains about 500 to 1000 different microorganisms, and the *firmicutes,*

bacteroidetes, actinobacteria, and *proteobacteria* are four dominant bacterial phyla in

the human gut (Khanna and Tosh, 2014). In germ-free mice, the gut microbiota has

been demonstrated as an important environmental factor that regulates energy harvest

and storage in the host and plays a critical role in the healthy weight gain and fat

composition (Bäckhed et al., 2004; Backhed et al., 2007). Compared with

conventionally raised mice, germ-free mice tend to have less total fat and are resistant

to diet-induced obesity, which is likely through two complementary, but independent

mechanisms of 1) elevated levels of the fasting-induced adipose factor (Fiaf), which

induces a key regulator (Pgc-1a) of energy metabolism and 2) increased adenosine

monophosphate activated protein kinase (AMPK) activity that increases the

metabolism of fatty acids (Backhed et al., 2007). In a double-blind, randomized,

placebo-controlled parallel-group clinical trial, glycemic conditions are improved by

probiotics and particularly synbiotic supplements in pre-diabetic individuals

(Kassaian et al., 2018). In addition, in rats chronically treated with olanzapine, altered

fecal microbiota profiles were observed, and they might be associated with metabolic

effects induced by olanzapine (Davey et al., 2012). Supplementation of vitamin D

plus probiotics also showed clinical and metabolic improvements compared with the

placebo in patients with schizophrenia (Ghaderi et al., 2019) who had been treated

with chlorpromazine and anticholinergic agents but not concurrently treated with

probiotics in the past six months.

The investigations of agents such as antibiotics and probiotics in olanzapine-induced

metabolic dysfunctions (Davey et al., 2013), metabolic diseases, and obesity have

been encouraging in animal models (Cani et al., 2008; Murphy et al., 2013).

Although the number of existing studies about antipsychotics on the gut microbiota

is limited, a few studies in rodents have found that olanzapine- and

risperidone-induced weight gain and metabolic dysfunction were associated with the

changes in the gut microbiota (Bahra et al., 2015; Bretler et al., 2019). However, a

minimal number of studies have been conducted in humans (Kang et al., 2019), in

particular on antipsychotics-induced metabolic disturbances. A longitudinal study of

five male adolescents with psychiatric disorders treated with risperidone showed that

a gradual increase in the ratio of *Firmicutes* to *Bacteroidetes* was associated with an

increase in body mass index (BMI) compared with antipsychotic-naïve psychiatric

controls, and the gut microbiome of the participants was enriched with genes for

short-chain fatty acids (SCFA) and serotonin metabolism (Bahr et al., 2015).

However, no other outcome measures were evaluated. In this study, we assessed the

efficacy of probiotics supplements in preventing olanzapine-induced metabolic

disturbance and weight gain in drug-naive first-episode patients with schizophrenia.

2. Material and methods

2.1 Trial design

This study was a randomized-controlled clinical trial designed to evaluate the efficacy of probiotics for reducing olanzapine-induced metabolic disturbance and weight gain. Participants were randomly assigned (without any restriction or stratification) through a computer-generated random number to the treatment of either olanzapine plus probiotics (olanzapine +) or olanzapine monotherapy in blocks of 4 to ensure approximately equal numbers of participants within the two treatment groups. The treatment assignment (randomization) was determined after the completion of all assessments and acceptance into the study.

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human patients were approved by the ethics committee of the Second Xiangya Hospital of Central South University (2017-027). Written informed consent was obtained from all patients.

2.2 Participants

All participants were recruited (November 30, 2017 to February 01, 2019) by clinical investigators from the Department of Psychiatry of the Second Xiangya Hospital at Central South University, China. Patients were aged 18 to 50 years, the first psychotic episode of schizophrenia and met diagnostic criteria of the Diagnostic and Statistical Manual of Mental Disorders Fifth Edition (DSM-V). All patients were ensured not having used any antipsychotics or recreational drugs at least three

months before enrollment and without abnormal in physical examinations, laboratories, and electrocardiogram (ECG) tests.

Exclusions: 1) female patients who were pregnant or lactating at the time of enrollment, 2) patients with mental retardation, addictive disorders, and 3) patients with specific systemic diseases, metabolic disorders, or medical conditions such as diabetes mellitus, dyslipidemia, cardiovascular diseases, and hypertension.

2.3 Randomization

Participants were randomly assigned 1:1 to olanzapine+ or olanzapine through a computer-generated random number. A research nurse conducted the randomization. Treating clinicians were not blinded to trial groups, but independent assessors who were blinded to the information on the treatment group evaluated all primary and most of the secondary outcomes.

2.4 Intervention

The participants were randomly assigned by investigators to a 12-week treatment of olanzapine+, with olanzapine (15-20mg/day at 8:00 p.m.) plus probiotics (Bifico, triple live bacteria oral capsule, 1680 mg/day, 840mg bid), or olanzapine (15-20mg/day at 8:00 p.m.). Bifico (Shanghai Xinyi Pharmaceutical Inc., Shanghai, China), supplied by the manufacturer, is one kind of commonly used probiotics supplement in China and contains live bacteria of *Bifidobacterium, Lactobacillus* and *Enterococcus* at least 5.0×10^7 CFU/g. The participants' adherences to probiotics and olanzapine treatment for each visit was defined as taking more than 80% of the study drug dosage prescribed for that interval. If a participant was non-adherent, both the patient and caregiver were counseled on the importance of taking the

prescribed amount of study medication.

2.5 Outcome Measures

Primary outcomes were the change in fasting serum insulin, insulin resistance, body

mass index (BMI), and weight gain. BMI is calculated as weight in kilograms

divided by height in meter squared, and insulin resistance was calculated based on a

formula of homeostatic model assessment for insulin resistance (HOMA-IR), by

fasting insulin [mIU/L]x fasting glucose [mmol/L]/22.5.

Secondary outcomes were 1) the increase in the levels of lipids, including

high-density lipoprotein cholesterol (HDL), low-density lipoprotein cholesterol

(LDL), total cholesterol, and triglyceride, 2) Psychopathologic symptoms measured

by positive and negative symptom scales (PANSS), and 3) appetite increase. We

used a self-reported scale to evaluate appetite. The scale included four items: how

hungry you are, how full you are, how strong desire to eat, and how much food you

think you can eat. They were measured at a scale of 0-10. A composite score was

calculated to measure the overall appetite increase. Compared with baseline, the

composite score increased ≥1 was defined as appetite increase, decreased ≥1 was

defined as appetite decrease, and an increase or decrease less than 1 was defined as

no change in appetite.

2.6 Sample size

The sample size was calculated for the primary outcome of HOMA-IR based on the

Hotelling-Lawley F test with a linear exponent autoregressive correlation structure

(Xu et al., 2020), a base correlation of 0.8, and decay of 0.3 over the time. The total

sample size required for a power of 80%, a level of 0.05 with a standard deviation of

0.6 and 0.8 were 44 and 74, respectively; and our actual sample of 76 patients met

the calculation.

2.7 Data collection

All patients who received the treatment were scheduled to having a clinical

evaluation through the scheduled follow-up at week 4, 8, and 12. The baseline

assessments included demographics, a comprehensive medical history, physical

examination, anthropometric measurements (weight and height), and PANSS score.

The laboratory tests at baseline included fasting glucose, lipids and insulin, liver and

renal function, blood counts, and electrocardiogram. At each follow-up visit, all

baseline clinical evaluations, including physical examination, anthropometric

measurements, and laboratory tests, were repeated. The Treatment Emergent

Symptom Scale (TESS) (Guy) was used to record adverse events throughout the

clinical trial. PANSS were also evaluated at week 12, the end of the trial.

We assessed the appetite half an hour before lunch every day within the first two

weeks of the treatments and at each follow-up visit.

2.8 Statistical Analysis

Data were examined and processed before the analysis. Summary statistics including

mean, median, and standard deviation were calculated for all variables at baseline

and each follow-up. Through a comparison of mean and median, one can have a

quick assessment of any severe violation of the fair assumption for the statistical

methods used for analysis (Zhang and Hughes, 2019). General linear random-effect

model(José A. Apud et al., 2012; Wu et al., 2016; Xu et al., 2020) was employed to

estimate the fixed-effect of treatments on primary and secondary outcomes over time

after adjustment for the baseline characteristics of age, sex, duration of illness, which

provide a better analysis in presence of missing or dropouts than the last observation

carried forward. The covariance matrix chooses first-order autoregressive [AR (1)]

over the time of treatment. Post-hoc least square estimates of mean and standard

errors were obtained for graphic presentation; correction for multiple testing was

performed using Dunnett's test to compare the outcomes at each follow-up visit with

the baselines. The analysis was performed using SPSS and SAS/STAT 13.2; a

two-sided level of 0.05 was considered as statistically significant.

Ancillary analyses were also performed on the four primary outcomes while

adjusting for the status of appetite increase, which was coded as a binary outcome.

The main effect of appetite increase, plus the interaction between appetite increase

and treatment time was included in the regression model to assess the treatment

effect over time. The report of the study followed the CONSORT 2010 Statement

updated guidelines for reporting parallel group randomized trials and related

recommendations for reporting of clinical trials as well as calls for transparency in

the report for clinical and preclinical research (Landis et al., 2012; Zhang and

Hughes, 2019).

3.   Results

Participants were recruited from November 30, 2017, to February 01, 2019. Of 126

patients screened for eligibility (**Figure 1**), 17 patients did not meet the inclusion or

met exclusion criteria, 19 patients declined to participate; 90 patients were randomly

and equally assigned to the group of olanzapine+ or olanzapine monotherapy. During

the trial period, 14 patients did not receive intervention with medication as

randomized or withdrew the consent; so, 76 patients were evaluated with follow-up

to collect the outcome measurements, which included 39 patients in the olanzapine+

and 37 in the olanzapine, who were followed at least one time. Finally, 88.2% of

patients completed the 12-week treatment. We examined the nine individuals who

did not complete the follow-up at week 8 or 12, found no significant difference in

baseline characteristics of patients from those patients who completed all follow-ups

(**Table S1**).

Table 1 presents the demographics and characteristics of the patients by treatment

group. There was no significant difference in all the variables between groups,

which were expected due to the randomized assignments. About 65-72% of the

patients were males, and 35-36% were cigarette smokers. The mean age was 24

years, and the duration of illness was about 11 months; the total PANSS score was

about 80. The baseline characteristics and outcome measures were not different

($p>0.05$) between two treatments except for a slight difference in LDL ($p=0.0017$)

and total cholesterol ($p=0.037$), which were higher in the olanzapine+ than in the

olanzapine monotherapy. The higher level of LDL and total cholesterol at baseline

assured at least no over-estimation of the interventional effect of probiotic

supplements on the secondary outcomes when olanzapine+ are assumed to reduces

the metabolic disturbance of lipid profiles. The summary statistics of the key

outcomes at baseline and follow-up time points by treatment group were presented

in the supplementary (**Table S2**)

3.1 Primary outcomes

Figure 2 provides an intuitive display of the treatment effects over time. The

addition of probiotic supplements concurrently with olanzapine medication

significantly slowed the increase in fasting insulin ($p<0.0001$) and HOMA-IR

($p=0.0005$), but only had a slight effect on reducing the increase in BMI ($p=0.077$)

and weight gain ($p=0.051$).

Table 2 provides the detailed estimates of treatment effect. Specifically, patients

received treatment of olanzapine+ had an increase in the levels of fasting insulin by

2.45 uIU/mL, from 7.81 at baselines to 10.26 at week 12 ($p<0.0001$), which was much

lower than an increase by 6.03 uIU/mL (p<0.0001), from 7.69 at baselines to 13.72

at week 12 in the olanzapine monotherapy, and there was a significant difference in

the level of fasting insulin at the end of the trial between two groups (difference,

-3.46, 95% CI [-5.0,-1.91]; $p<0.0001$). Similarly, the HOMA-IR increased by 0.63

($p<0.0001$), from 1.54 at the baseline to 2.17 at week 12 in the olanzapine+ group,

which was lower than the increase by 1.27 ($p<0.0001$), from 1.51 at the baselines to

2.78 at week 12 in the olanzapine monotherapy; also, there was a significant

difference in the HOMA-IR at the end of the trial between two groups (difference,

-0.61, 95% CI [-0.94,-0.28]; $p=0.0004$). Both BMI and body weight significantly

increased at week 4 compared with baseline and continued to increase throughout

the study in both the olanzapine+ and olanzapine group, but overall, these increases

were not significantly different between the two treatment groups ($p>0.05$).

3.2 Secondary outcomes

Appetite increase

Olanzapine plus probiotic supplements slightly reduced appetite increase. In the

olanzapine+ group, 38.5% (15/39) of patients had appetite increase, which was

lower than 56.8% (21/37) in the olanzapine group (log-rank test, $p=0.0666$;

likelihood ratio test, $p=0.0255$) (**Figure S1**). The difference in the appetite increase

became no significant (p=0.1211) between two treatments after adjustment for sex,

age, duration of illness, and dosage of olanzapine taken in the Cox regression

analysis. However, on average, patients treated with olanzapine+ were about 40%

less likely (Hazard Ratio, HR=0.589) to have an increase in appetite than those who

received olanzapine monotherapy (**Table S3**).

Psychopathological symptoms

The psychopathological symptoms improved in both groups after treatment for 12

weeks but the improvement was not significantly different between groups. The

PANSS total score was reduced by 24.5 (p<0.0001) from 80.18 at the baseline to

55.67, after olanzapine+ for 12 weeks and by 22.7 (p<0.0001), from 79.5 at the

baseline to 56.8, after olanzapine treatment. However, the improvement of PANSS

total score was not significantly different between two groups (p=0.48). We also

observed a similar pattern in positive, negative, and the general psychopathology

(**Table S4**).

Lipid profiles

The lipid profiles that comprised triglyceride, total cholesterol, HDL, and LDL

increased very significantly (p<0.0001) in both the olanzapine+ and the olanzapine

group, but no significant difference in triglyceride (p=0.21), HDL (p=0.60) and LDL

(p=0.55), except total cholesterol (p=0.028) between two treatment groups (**Table

S5**).

3.3 Appetite modulated antipsychotics-induced metabolic change

While the addition of probiotic supplements slightly reduced the likelihood and

timing of the change in appetite, which had a significant effect on the metabolic

disturbance in both groups. Patients with appetite increase had a significant

($p<0.0001$) elevation of all primary outcomes of fasting insulin, HOMA-IR, BMI,

and weight than those patients without appetite increase (**Table 3**). The metabolic

disturbance increased gradually with the duration of treatment and became

significant at week 12 in BMI ($p=0.028$), fasting insulin ($p<0.0001$), and HOMA-IR

($p<0.0001$). However, we noted that the significant differences between two groups

in weight gain were primarily driven by the baseline measurements, in which the

weight was significantly different ($p=0.0065$) at baseline but no difference after

treatment at week 12 ($p=0.28$). Of note, appetite also modulated the

antipsychotic-induced changes in the secondary outcomes such as triglyceride

($p=0.019$), total cholesterol ($p=0.0002$), and LDL ($p<0.0001$), but not in HDL

($p=0.054$).

3.4 Ancillary analysis

We further investigated if the effect of probiotic supplements on the metabolic

disturbance was through the less increase in appetite (**Table S6**). After adjustment

for the status of appetite increase, probiotic supplements still significantly reduced

the increase in fasting insulin ($p<0.0001$) and HOMA-IR ($p=0.0012$); the effect also

started at week 8 after treatment, suggesting that the impact of probiotic supplements

on the increase in fasting insulin and HOMA-IR were mostly independent of appetite

increase. However, two treatments had no significant impact on the change in BMI

($p=0.143$) and body weight ($p=0.11$) during the trial after adjustment for appetite

increase.

3.5 Adverse events

Adverse events considered grade 3 or higher that were reported at least 5% of the

evaluable patients (**Table 4**). The most common adverse events in both groups were

hypoactivity (26-30%), somnolence (26-27%), and abnormal in liver function test

(15-16%), but they were not significantly different between the two treatment groups.

The only adverse event reported as possibly related to the probiotic supplements was

diarrhea, which was 10% in the olanzapine+ but only 3% in the olanzapine group. In

contrast, constipation was 3% in olanzapine+, but 14% in the olanzapine

monotherapy. None of these was suitable for pursuing a formal statistical test due to

small sample size.

4. Discussion

The main findings of this trial were that probiotic supplements with olanzapine

attenuated antipsychotics-induced elevation of fasting serum insulin and insulin

resistance as measured by HOMA-IR while retaining the treatment effect on

psychopathological symptoms. However, we did not find a significant benefit on

BMI and weight gain, nor the secondary outcomes of appetite increase as well as

lipid profiles. Our study provided the first evidence for probiotic supplements in

preventing antipsychotics-induced insulin resistance in patients with drug-naive

first-episode schizophrenia while concurrently using antipsychotics.

Our findings that probiotic supplements reduced the olanzapine-induced increase in

fasting insulin and insulin resistance were consistent with previous multiple studies

in individuals with diabetic conditions. 1) One randomized, double-blind,

placebo-controlled trial in Thailand finds that supplementation of probiotics

containing *Bifidobacterium* (*B*.) and *Lactobacillus* (*L*.) daily for four consecutive

weeks improves the plasma fasting insulin and HOMA-IR in women diagnosed with

gestational diabetes mellitus (GDM) between 24–28 weeks (Kijmanawat et al.,

2019), compared to the placebo group; 2) In another randomized clinical trial (RCT)

in Ukraine, supplementation of multi-probiotics, including concentrated biomass of

14 probiotic bacteria genera *Bifidobacterium, Lactobacillus, Lactococcus,*

Propionibacterium for eight weeks, was associated with the reduction of HOMA-IR

in individuals with type 2 diabetes (Kobyliak et al., 2018). Both trials used probiotics

that contain *Bifidobacterium* and *Lactobacillus,* two common bacteria for probiotics,

as that used in our study; 3) A meta-analysis of six RCTs involved 830 patients also

confirmed that supplementation of probiotics significantly reduced the serum fasting

insulin and HOMA-IR in women with GDM (Pan et al., 2019). In addition, in a

double blind, randomized, placebo-controlled parallel-group clinical trial for 24

weeks, glycemic conditions can be improved significantly by probiotics, particularly

synbiotic (prebiotic plus probiotic) supplements in pre-diabetic individuals (Kassaian

et al., 2018). Insulin resistance plays a crucial role in the olanzapine-induced

metabolic disturbance, including lipid profiles and weight gain (Wu et al., 2016; Wu

et al., 2008), which at least partly depends on appetite increase (Davoodi et al., 2009).

However, we found that the benefit of probiotic supplements on preventing the

olanzapine-induced insulin resistance was still significant regardless of appetite

increase.

Our study did not find a significant effect of probiotic supplements on the lipid

profiles, although reports have shown that probiotics have a significant effect on

lipid metabolism. In the most recent meta-analysis of ten RCTs with 1,139

participants, probiotics supplements were shown to have improved effect on glucose,

HOMA-IR, and lipid metabolism in pregnant women (Han et al., 2019). A

meta-analysis of 13 RCTs with 485 patients with high, borderline high, or healthy

cholesterol also shows that a diet rich in probiotics decreases the total cholesterol

and LDL cholesterol concentration in plasma (Guo et al., 2011). In addition, an RCT

study of vitamin D plus probiotic supplements ($8x10^9$ CFU), containing *B.bifidum,*

L.acidophilus, L.reuteri, and *L.fermentum* (each 2×10^9) for 12 weeks in patients

with chronic schizophrenia who had used but not concurrently with antipsychotics,

showed a significant effect on the improvement of metabolic profiles including

fasting glucose, fasting insulin and HOMA-IR, triglycerides and total cholesterol

levels (Ghaderi et al., 2019). The limited effect of probiotic supplements on lipid

profiles could be due to the short duration of treatment in this trial because our

previous trials of metformin on antipsychotic-induced dyslipidemia in schizophrenia

indicate that the treatment effect on lipids requires a prolonged duration of treatment

(24 weeks) than on insulin or insulin resistance (12 weeks) (Wu et al., 2016).

Increasing evidence shows that there is a strong metabolic risk intrinsic to

schizophrenia even in the absence of antipsychotic drug treatment (Freyberg et al.,

2017). As in this study with drug-naive patients, we noted elevated levels of lipids in

one of the groups. This would make a study of definitive mechanisms more

challenging for antipsychotic-induced metabolic disturbance. There have been

studies indicating that genetic variants at *TCF7L2* and *PSMD9* were independently

associated with schizophrenia and type 2 diabetes (Freyberg et al., 2017). In addition,

the central nervous system (e.g. hypothalamus) where the neurotransmitter and

neuropeptide system such as dopamine, serotonin, and histamine may play a role in

the intrinsic metabolic risk. This is because that almost all antipsychotics act on

dopamine receptors (D2 and D3) and/or serotonin receptors, and polymorphism in

DRD2, DRD3, 5HT2a, and *5HT2c* have been associated with both

antipsychotic-induced metabolic dysfunction and schizophrenia (Freyberg et al.,

2017). In fact, some polymorphisms in *TCF7L2* and *DRD2* have been found to be

shared by schizophrenia, bipolar, and major depressive disorders (Xia et al., 2019).

The effect of probiotic supplements on the levels of fast insulin and insulin

resistance may support that the gut-brain axis plays a role in modulating

antipsychotic-induced metabolic dysfunction and might have implications for

reducing the risk of schizophrenia at least for a partial population of the patients with

schizophrenia.

To our knowledge, this was the first study to assess probiotic supplements on

preventing olanzapine-induced metabolic disturbance and weight gain in drug naïve

first-episode patients with schizophrenia. Supplementation of probiotics was in favor

of preventing olanzapine-induced elevation of fasting serum insulin and insulin

resistance, and was shown to be well tolerated and safe in the participants.

4.1 Limitation

This study had some limitations. First, we did not use a placebo in this trial, although

the treatment information was blinded to the clinical evaluators. Second, this was a

12-week trial, and about 20% of patients were lost to follow-up due to the inability

to take the medication as a randomized assignment. In such a limited duration of

treatment, we did not know whether the improved insulin sensitivity could sustain

after patients stopped taking probiotic supplements. Third, this was a single-center

clinical trial, which had the advantage of homogenous samples but may not allow us

to address the variability across regions. Therefore, it may limit the generalizability

of the study results to the study population in a south-central province of China. In

addition, some related measurements such as adiposity and leptin levels were not

measured; we did not examine the microbiome that would strengthen the study and

help reveal the underlying mechanism of treatment response. These may require

additional effort such as funding support and development of study protocols.

4.2 Generalizability

Despite some limitations, our study provided the first evidence that probiotic

supplements can be used for preventing antipsychotic-induced metabolic disturbance

in particular elevation of fasting insulin and insulin resistance. Given that probiotic

supplements were generally safe for use, as shown in this trial, the study results may

have implications for clinical practice, the supplementations of probiotics can be

recommended for concurrent use with antipsychotic treatment in patients with

schizophrenia in the study population in the area where the study was conducted. In

addition, further research is needed to assess the effect of the probiotic supplements

on antipsychotic-induced weight gain and metabolic disturbance in a longer-term

and potentially synergistic effect on antipsychotic response.

Role of the funding source

Funding. The research was supported by the Key R&D Program Projects, National

Science Foundation of China (Grant No.2016YFC1306900), and the National Natural

Science Foundation of China (Grant No.81622018), Beijing Haiju Scholarship

(BHTO201511097).

Declarations of Interest: None.

Contributors: RW, JZ applied the funding for the trial. RW, JD conceived the trial. DK,

YY, CL, JX, YL, JH, XP, WW, and XW were involved in patient recruitment. RW,

FZ, and JZ were involved in study quality control. Statistical analyses were performed

by RW and FZ. DK and FZ wrote the manuscript. All authors provided critical

comments. All authors approved the final version of the manuscript.

Acknowledgement: We would like to thank all the participants and researchers who

have been involved in the research.

REFERENCES

Bäckhed, F., Ding, H., Wang, T., Hooper, L.V., Koh, G.Y., Nagy, A., Semenkovich, C.F., Gordon, J.I., 2004. The gut microbiota as an environmental factor that regulates fat storage. Proceedings of the National Academy of Sciences of the United States of America 101(44), 15718-15723.

Backhed, F., Manchester, J.K., Semenkovich, C.F., Gordon, J.I., 2007. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. Proceedings of the National Academy of Sciences of the United States of America 104(3), 979-984.

Bahr, S.M., Tyler, B.C., Wooldridge, N., Butcher, B.D., Burns, T.L., Teesch, L.M., Oltman, C.L., Azcarate-Peril, M.A., Kirby, J.R., Calarge, C.A., 2015. Use of the second-generation antipsychotic, risperidone, and secondary weight gain are associated with an altered gut microbiota in children. Transl Psychiatry 5, e652.

Bahra, S.M., Weidemann, B.J., Castro, A.N., Walsh, J.W., deLeon, O., Burnett, C.M., Pearson, N.A., Murry, D.J., Grobe, J.L., Kirby, J.R., 2015. Risperidone-induced weight gain is mediated through shifts in the gut microbiome and suppression of energy expenditure. EBioMedicine 2(11), 1725-1734.

Bretler, T., Weisberg, H., Koren, O., Neuman, H., 2019. The effects of antipsychotic medications on microbiome and weight gain in children and adolescents. BMC medicine 17(1), 112.

Cani, P.D., Bibiloni, R., Knauf, C., Waget, A., Neyrinck, A.M., Delzenne, N.M., Burcelin, R., 2008. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. Diabetes 57(6), 1470-1481.

Chen, J., Huang, X.F., Shao, R., Chen, C., Deng, C., 2017. Molecular Mechanisms of Antipsychotic Drug-Induced Diabetes. Frontiers in neuroscience 11, 643.

Correll, C.U., Manu, P., Olshanskiy, V., Napolitano, B., Kane, J.M., Malhotra, A.K., 2009. Cardiometabolic risk of second-generation antipsychotic medications during first-time use in children and adolescents. Jama 302(16), 1765-1773.

Davey, K.J., Cotter, P.D., O'Sullivan, O., Crispie, F., Dinan, T.G., Cryan, J.F., O'Mahony, S.M., 2013. Antipsychotics and the gut microbiome: olanzapine-induced metabolic dysfunction is attenuated by antibiotic administration in the rat. Translational psychiatry 3, e309.

Davey, K.J., O'Mahony, S.M., Schellekens, H., O'Sullivan, O., Bienenstock, J., Cotter, P.D., Dinan, T.G., Cryan, J.F., 2012. Gender-dependent consequences of chronic olanzapine in the rat: effects on body weight, inflammatory, metabolic and microbiota parameters. Psychopharmacology 221(1), 155-169.

Davoodi, N., Kalinichev, M., Korneev, S.A., Clifton, P.G., 2009. Hyperphagia and increased meal size are responsible for weight gain in rats treated sub-chronically with olanzapine. Psychopharmacology 203(4), 693-702.

De Hert, M., Detraux, J., van Winkel, R., Yu, W., Correll, C.U., 2011. Metabolic and cardiovascular adverse effects associated with antipsychotic drugs. Nature reviews. Endocrinology 8(2), 114-126.

Foley, D.L., Morley, K.I., 2011. Systematic review of early cardiometabolic outcomes of the first treated episode of psychosis. Arch Gen Psychiatry 68(6), 609-616.

Freyberg, Z., Aslanoglou, D., Shah, R., Ballon, J.S., 2017. Intrinsic and Antipsychotic Drug-Induced Metabolic Dysfunction in Schizophrenia. Frontiers in neuroscience 11(432).

Ghaderi, A., Banafshe, H.R., Mirhosseini, N., Moradi, M., Karimi, M.A., Mehrzad, F., Bahmani, F., Asemi, Z., 2019. Clinical and metabolic response to vitamin D plus probiotic in schizophrenia patients. BMC psychiatry 19(1), 77.

Guo, Z., Liu, X.M., Zhang, Q.X., Shen, Z., Tian, F.W., Zhang, H., Sun, Z.H., Zhang, H.P., Chen, W., 2011. Influence of consumption of probiotics on the plasma lipid profile: a meta-analysis of randomised

controlled trials. Nutrition, metabolism, and cardiovascular diseases : NMCD 21(11), 844-850.

Guy, W., ECDEU Assessment Manual for Psychopharmacology, Revised. US Department of Health,
Education, and Welfare: Public Health Service, Alcohol, Drug Abuse and Mental Health
Administration., Rockville, MD.

Hahn, M.K., Wolever, T.M., Arenovich, T., Teo, C., Giacca, A., Powell, V., Clarke, L., Fletcher, P., Cohn, T.,
McIntyre, R.S., Gomes, S., Chintoh, A., Remington, G.J., 2013. Acute effects of single-dose olanzapine
on metabolic, endocrine, and inflammatory markers in healthy controls. Journal of clinical
psychopharmacology 33(6), 740-746.

Han, M.M., Sun, J.F., Su, X.H., Peng, Y.F., Goyal, H., Wu, C.H., Zhu, X.Y., Li, L., 2019. Probiotics improve
glucose and lipid metabolism in pregnant women: a meta-analysis. Annals of translational medicine
7(5), 99.

José A. Apud, M.D., Ph.D. ,, Fengyu Zhang, Ph.D. ,, Heather Decot, B.S. ,, Kristin L. Bigos, Ph.D. , and,
Daniel R. Weinberger, M.D., 2012. Genetic Variation in KCNH2 Associated With Expression in the
Brain of a Unique hERG Isoform Modulates Treatment Response in Patients With Schizophrenia.
American Journal of Psychiatry 169(7), 725-734.

Kang, D.-Y., Li, S.-J., Liu, C.-C., Wu, R.-R., 2019. Gut microbiota and antipsychotic induced metabolic
alteration. Glob Clin Transl Res 1(4), 131-143.

Kassaian, N., Feizi, A., Aminorroaya, A., Jafari, P., Ebrahimi, M.T., Amini, M., 2018. The effects of
probiotics and synbiotic supplementation on glucose and insulin metabolism in adults with
prediabetes: a double-blind randomized clinical trial. Acta diabetologica 55(10), 1019-1028.

Khanna, S., Tosh, P.K., 2014. A clinician's primer on the role of the microbiome in human health and
disease. Mayo Clinic proceedings 89(1), 107-114.

Kijmanawat, A., Panburana, P., Reutrakul, S., Tangshewinsirikul, C., 2019. Effects of probiotic
supplements on insulin resistance in gestational diabetes mellitus: A double-blind randomized
controlled trial. Journal of diabetes investigation 10(1), 163-170.

Kirk, S.L., Glazebrook, J., Grayson, B., Neill, J.C., Reynolds, G.P., 2009. Olanzapine-induced weight gain
in the rat: role of 5-HT2C and histamine H1 receptors. Psychopharmacology 207(1), 119-125.

Kobyliak, N., Falalyeyeva, T., Mykhalchyshyn, G., Kyriienko, D., Komissarenko, I., 2018. Effect of alive
probiotic on insulin resistance in type 2 diabetes patients: Randomized clinical trial. Diabetes &
metabolic syndrome 12(5), 617-624.

Landis, S.C., Amara, S.G., Asadullah, K., Austin, C.P., Blumenstein, R., Bradley, E.W., Crystal, R.G.,
Darnell, R.B., Ferrante, R.J., Fillit, H., Finkelstein, R., Fisher, M., Gendelman, H.E., Golub, R.M.,
Goudreau, J.L., Gross, R.A., Gubitz, A.K., Hesterlee, S.E., Howells, D.W., Huguenard, J., Kelner, K.,
Koroshetz, W., Krainc, D., Lazic, S.E., Levine, M.S., Macleod, M.R., McCall, J.M., Moxley, R.T., 3rd,
Narasimhan, K., Noble, L.J., Perrin, S., Porter, J.D., Steward, O., Unger, E., Utz, U., Silberberg, S.D.,
2012. A call for transparent reporting to optimize the predictive value of preclinical research. Nature
490(7419), 187-191.

Lieberman, J.A., Stroup, T.S., McEvoy, J.P., Swartz, M.S., Rosenheck, R.A., Perkins, D.O., Keefe, R.S.,
Davis, S.M., Davis, C.E., Lebowitz, B.D., Severe, J., Hsiao, J.K., Clinical Antipsychotic Trials of
Intervention Effectiveness, I., 2005. Effectiveness of antipsychotic drugs in patients with chronic
schizophrenia. The New England journal of medicine 353(12), 1209-1223.

Maayan, L., Correll, C.U., 2011. Weight gain and metabolic risks associated with antipsychotic
medications in children and adolescents. Journal of child and adolescent psychopharmacology 21(6),
517-535.

Murphy, E.F., Cotter, P.D., Hogan, A., O'Sullivan, O., Joyce, A., Fouhy, F., Clarke, S.F., Marques, T.M.,
O'Toole, P.W., Stanton, C., Quigley, E.M., Daly, C., Ross, P.R., O'Doherty, R.M., Shanahan, F., 2013.
Divergent metabolic outcomes arising from targeted manipulation of the gut microbiota in
diet-induced obesity. Gut 62(2), 220-226.

Nasrallah, H.A., 2008. Atypical antipsychotic-induced metabolic side effects: insights from
receptor-binding profiles. Mol Psychiatry 13(1), 27-35.

Newcomer, J.W., 2005. Second-generation (atypical) antipsychotics and metabolic effects: a
comprehensive literature review. CNS drugs 19 Suppl 1, 1-93.

Newcomer, J.W., 2007. Metabolic considerations in the use of antipsychotic medications: a review of
recent evidence. The Journal of clinical psychiatry 68 Suppl 1, 20-27.

Pan, J., Pan, Q., Chen, Y., Zhang, H., Zheng, X., 2019. Efficacy of probiotic supplement for gestational
diabetes mellitus: a systematic review and meta-analysis. The journal of maternal-fetal & neonatal
medicine : the official journal of the European Association of Perinatal Medicine, the Federation of
Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet 32(2), 317-323.

van Winkel, R., De Hert, M., Wampers, M., Van Eyck, D., Hanssens, L., Scheen, A., Peuskens, J., 2008.
Major changes in glucose metabolism, including new-onset diabetes, within 3 months after initiation
of or switch to atypical antipsychotic medication in patients with schizophrenia and schizoaffective
disorder. The Journal of clinical psychiatry 69(3), 472-479.

Wu, R.R., Zhang, F.Y., Gao, K.M., Ou, J.J., Shao, P., Jin, H., Guo, W.B., Chan, P.K., Zhao, J.P., 2016.
Metformin treatment of antipsychotic-induced dyslipidemia: an analysis of two randomized,
placebo-controlled trials. Mol Psychiatry 21(11), 1537-1544.

Wu, R.R., Zhao, J.P., Jin, H., Shao, P., Fang, M.S., Guo, X.F., He, Y.Q., Liu, Y.J., Chen, J.D., Li, L.H., 2008.
Lifestyle intervention and metformin for treatment of antipsychotic-induced weight gain: a
randomized controlled trial. Jama 299(2), 185-193.

Xia, L., Xia, K., Weinberger, D.R., Zhang, F., 2019. Common genetic variants shared among five major
psychiatric disorders: a large-scale genome-wide combined analysis. Glob Clin Transl Res. 1(1), 21-30.

Xu, Z., Xie, C., Xia, L., Yuan, Y., Zhu, H., Huang, X., Li, C., Tao, Y., Qu, X., Zhang, F., Zhang, Z., 2020.
Targeted exome sequencing identifies five novel loci at genome-wide significance for modulating
antidepressant response in patients with major depressive disorder. Translational psychiatry 10(1),
30.

Zhang, F., Hughes, C., 2019. Reporting standards for clinical and translational research. Glob Clin
Transl Res 1(2), 69-73.

Table 1. Baseline Characteristics of the Patients by Treatment Group

	Olanzapine plus probiotics (N=39)							Olanzapine monotherapy (N=37)							Normal range
	Mean	Med	SD	Min	5%	95%	Max	Mean	Med	SD	Min	5%	95%	Max	
Male	72%							65%							
Smoking	36%							35%							
Age, year	24.82	22.00	5.64	18.00	18.00	36.00	38.00	23.43	22.00	4.89	18.00	18.00	35.00	36.00	
Duration of illness, month	11.49	11.00	3.51	6.00	6.00	18.00	18.00	11.08	10.00	3.64	5.00	6.00	18.00	18.00	
Dose of olanzapine, mg	19.36	20.00	1.59	15.00	15.00	20.00	20.00	18.85	20.00	1.92	15.00	15.00	20.00	20.00	
Weight, kg	55.63	53.00	7.54	47.00	48.00	72.00	74.70	56.75	56.00	8.35	45.00	45.00	70.00	72.00	
Body mass index, kg/m²	21.26	20.70	2.59	18.22	18.29	29.26	29.92	21.25	21.33	1.69	18.03	18.73	23.94	23.94	18.5-23.9 kg/m²
Fasting glucose, mmol/L	4.41	4.34	0.39	3.39	3.96	5.30	5.30	4.40	4.30	0.37	3.90	3.90	5.30	5.30	3.9-6.1 mmol/L
Fasting insulin, uIU/mL	8.07	7.60	2.88	3.50	3.60	14.50	15.60	7.85	7.50	1.72	4.90	4.90	11.20	12.30	6.4-15.0 uIU/mL
HOMA-Insulin Resistance	1.59	1.41	0.59	0.70	0.81	2.84	2.96	1.54	1.53	0.38	0.85	0.91	2.34	2.35	<2.67
Triglyceride, mmol/L	0.86	0.76	0.41	0.28	0.32	1.78	1.84	0.72	0.68	0.26	0.32	0.38	1.22	1.23	<1.71 mmol/L
Total cholesterol, mmol/L[a]	3.75	3.60	0.58	2.80	2.89	4.70	4.80	3.70	3.57	0.48	3.10	3.11	4.56	4.57	2.9-5.2 mmol/L
HDL, mmol/L	1.21	1.17	0.18	0.92	0.93	1.60	1.67	1.23	1.20	0.14	1.04	1.05	1.56	1.60	>1.04 mmol/L
LDL, mmol/L[a]	2.39	2.40	0.44	1.60	1.65	3.10	3.23	2.10	2.04	0.36	1.45	1.65	2.89	3.10	<3.12 mmol/L
Score of appetite scale	5.7	5.8	0.60	4.5	5	7	7.5	6.0	6.0	0.5	5	5	6.8	7	
Total score of PANSS	79.77	78.00	5.37	70.00	70.00	88.00	89.00	79.65	80.00	7.37	68.00	70.00	93.00	93.00	

Abbreviations: Med, median; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

[a], $p<0.05$, compared between the two treatment groups.

Table 2. Estimates of Treatment Effect on Primary Outcomes by Group

		Olanzapine+				Olanzapine				Group difference			
	N	LS Mean	SE	*P_adj*	N	LS mean	SE	*P_adj*	Beta	SE	95%CI	*P*	*P* [a]
Fasting Insulin, uIU/mL													
Baseline	35	7.81	0.564	ref	36	7.69	0.553	ref	0.119	0.772	-1.394, 1.632	0.8775	<.0001
4 Weeks	35	7.61	0.564	0.7352	34	8.22	0.562	0.6315	-0.612	0.78	-2.141, 0.917	0.434	
8 Weeks	33	8.15	0.575	0.5475	34	10.21	0.562	0.0012	-2.061	0.788	-3.605, -0.517	0.0098	
12 Weeks	33	10.26	0.575	<.0001	34	13.72	0.562	<.0001	-3.455	0.788	-4.999, -1.911	<.0001	
HOMA-IR													
Baseline	35	1.54	0.121	ref	36	1.51	0.118	ref	0.037	0.165	-0.286, 0.360	0.8254	0.0005
4 Weeks	35	1.47	0.121	0.5875	34	1.63	0.12	0.577	-0.151	0.167	-0.478, 0.176	0.3665	
8 Weeks	33	1.6	0.123	0.8024	34	2.06	0.12	0.0007	-0.459	0.169	-0.790, -0.128	0.0073	
12 Weeks	33	2.17	0.123	<.0001	34	2.78	0.12	<.0001	-0.611	0.169	-0.942, -0.280	0.0004	
Body mass index, kg/m^2													
Baseline	39	21.54	0.355	ref	37	21.47	0.359	ref	0.068	0.493	-0.898, 1.034	0.8905	0.077
4 Weeks	39	21.95	0.355	0.0036	37	22.41	0.359	<.0001	-0.465	0.493	-1.431, 0.501	0.3488	
8 Weeks	37	23.32	0.357	<.0001	34	23.92	0.362	<.0001	-0.6	0.497	-1.574, 0.374	0.2308	
12 Weeks	33	24.33	0.361	<.0001	34	24.55	0.362	<.0001	-0.217	0.5	-1.197, 0.763	0.6649	
Weight, kg													
Baseline	39	57.98	0.897		37	58.79	0.907		-0.817	1.247	-3.261, 1.627	0.514	0.051
4 Weeks	39	59.02	0.897	0.0035	37	61.29	0.907	0.0035	-2.273	1.247	-4.717, 0.171	0.0718	
8 Weeks	37	62.55	0.903	<.0001	34	65.16	0.915	<.0001	-2.608	1.257	-5.072, -0.144	0.0409	
12 Weeks	33	65.14	0.913	<.0001	34	66.78	0.915	<.0001	-1.64	1.264	-4.117, 0.837	0.1976	

Abbreviations: Olanzapine+ , olanzapine plus probiotics; HOMA-IR, HOMA insulin resistance.

LS mean, post-hoc least square estimate of mean.

P [a] , p value for testing the difference in insulin and body mass index over time between treatments.

P_adj, p value for testing the metabolic change versus the baseline within the treatment group.

Table 3. Appetite Modulated Metabolic Disturbance

	No appetite increase		Appetite increase		Group difference				
	LS mean	SE	LS mean	SE	Beta	SE	95%CI	P	P [a]
Fasting insulin, uIU/mL									
Baseline	7.69	0.52	7.88	0.57	-0.194	0.762	-1.688,1.300	0.7996	<.0001
4 Weeks	7.81	0.52	8.08	0.57	-0.267	0.768	-1.772,1.238	0.7283	
8 Weeks	8.66	0.53	9.81	0.57	-1.146	0.772	-2.659,0.367	0.1402	
12 Weeks	10.27	0.53	13.93	0.57	-3.664	0.772	-5.177,-2.151	<.0001	
HOMA-IR									
Baseline	1.52	0.11	1.54	0.12	-0.012	0.162	-0.330,0.306	0.943	<.0001
4 Weeks	1.55	0.11	1.57	0.12	-0.02	0.163	-0.339, 0.299	0.904	
8 Weeks	1.68	0.11	1.99	0.12	-0.312	0.164	-0.633, 0.009	0.06	
12 Weeks	2.1	0.11	2.91	0.12	-0.815	0.164	-1.136, -0.494	<.0001	
Body mass index, kg/m^2									
Baseline	21.93	0.34	21.05	0.36	0.874	0.495	-0.096, 1.844	0.0812	<.0001
4 Weeks	22.21	0.34	22.16	0.36	0.052	0.495	-0.918, 1.022	0.9159	
8 Weeks	23.22	0.34	24.1	0.37	-0.889	0.497	-1.863, 0.085	0.0774	
12 Weeks	23.91	0.34	25.03	0.37	-1.119	0.499	-2.097, -0.141	0.0276	
Body weight, kg									
Baseline	60.05	0.87	56.54	0.93	3.515	1.257	1.051, 5.979	0.0065	<.0001
4 Weeks	60.81	0.87	59.41	0.93	1.394	1.257	-1.070, 3.858	0.2707	
8 Weeks	63.46	0.87	64.34	0.93	-0.874	1.262	-3.348, 1.600	0.4904	
12 Weeks	65.31	0.87	66.69	0.94	-1.381	1.267	-3.864, 1.102	0.2788	
Triglycerides, mmol/L									
Baseline	0.84	0.066	0.7	0.071	0.145	0.096	-0.043, 0.333	0.1356	0.019
4 weeks	0.89	0.067	0.84	0.072	0.045	0.097	-0.145, 0.235	0.6447	
8 weeks	1.09	0.067	1.06	0.072	0.033	0.097	-0.157, 0.223	0.732	
12 Weeks	1.23	0.068	1.3	0.073	-0.067	0.098	-0.259, 0.125	0.4983	
Total Cholesterol, mmol/L									
Baseline	3.58	0.091	3.802	0.098	-0.225	0.133	-0.486, 0.036	0.0929	0.0002
4 weeks	3.56	0.092	3.78	0.1	-0.221	0.134	-0.484, 0.042	0.1021	
8 weeks	3.67	0.092	4.07	0.099	-0.407	0.134	-0.670, -0.144	0.003	
12 Weeks	3.72	0.093	4.36	0.1	-0.643	0.136	-0.910, -0.376	<.0001	
HDL, mmol/L									
Baseline	1.19	0.031	1.26	0.033	-0.062	0.045	-0.150, 0.026	0.1654	0.054
4 weeks	1.22	0.031	1.26	0.034	-0.034	0.046	-0.124, 0.056	0.4543	
8 weeks	1.19	0.031	1.2	0.034	-0.006	0.046	-0.096, 0.084	0.8894	
12 Weeks	1.13	0.032	1.07	0.034	0.058	0.046	-0.032, 0.148	0.2085	
LDL, mmol/L									
Baseline	2.32	0.079	2.14	0.085	0.179	0.115	-0.046, 0.404	0.1222	<.0001
4 weeks	2.32	0.08	2.23	0.087	0.096	0.117	-0.133, 0.325	0.4134	
8 weeks	2.49	0.079	2.57	0.086	-0.082	0.116	-0.309, 0.145	0.4803	
12 Weeks	2.51	0.082	2.83	0.088	-0.315	0.119	-0.548, -0.082	0.0089	

Abbreviations: HOMA-IR, HOMA insulin resistance; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

P [a] , p value for testing the difference in outcomes over time between treatments.

5   **Tables 4** Common Adverse Events (>5%) in the two Treatment Groups [a]

	Olanzapine plus probiotic (N=39)		Olanzapine monotherapy (N=37)	
Adverse events	N	%	N	%
Hypoactivity	10	26	11	30
Somnolence	10	26	10	27
Abnormal in liver function test	6	15	6	16
Akathisia	2	5	3	8
Constipation	1	3	5	14
Diarrhea	4	10	1	3
Dizzy	2	5	3	8
Weight gain (7% from baseline)	24	62	27	73

6   [a], Assessment was based on 76 patients who took at least one dose of olanzapine or olanzapine

7   plus probiotics and were evaluable for adverse event assessment.

Figure 1. Flowchart of Patients in Randomized Trial of Probiotic Supplements for Olanzapine-induced Metabolic Disturbances and Weight Gain

Figure 2. The Estimated Effect of Probiotic supplements on Olanzapine-induced Elevation of Fasting Insulin, Insulin Resistance, Body Mass Index, and Weight

Plots were presented with LS mean, and SEM;

Body mass index, which is calculated as weight in kilograms divided by height in meters squared;

HOMA-IR, Homeostatic Model Assessment of Insulin Resistance, which is calculated as insulin level (mIU/L)×fasting glucose(mmol/L)/22.5;

[a] P value for testing the difference in outcomes over time between treatments.