1 **SARS-CoV-2 Seroprevalence in a University Community: A Longitudinal Study of the Impact of**
2 **Student Return to Campus on Infection Risk Among Community Members**
3

4 *Arnold, Callum R.K. (MSc)[1,2]; *Srinivasan, Sreenidhi (PhD)[2,3]; Herzog, Catherine M.
5 (PhD[2,3]; Gontu, Abhinay (MVSc)[4]; Bharti, Nita (PhD)[1,2]; Small, Meg (PhD)[5,8]; Rogers,
6 Connie J. (PhD)[5]; Schade, Margeaux M. (PhD)[5]; Kuchipudi, Suresh V[2,4] (PhD); Kapur,
7 Vivek (PhD)[2,3,7]; Read, Andrew (D.Phil)[1,2,3]; Ferrari, Matthew J. (PhD)[1,2]
8

Author affiliations:
[1]Department of Biology, Pennsylvania State University, University Park, PA, USA 16802
[2]Center for Infectious Disease Dynamics, Pennsylvania State University, University Park,
PA, USA 16802
[3]Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA,
USA 16802
[4]Department of Veterinary and Biomedical Sciences, Pennsylvania State University,
University Park, PA, USA 16802
[5]College of Health and Human Development, Pennsylvania State University, University Park,
PA, USA 16802
[6]Department of Nutritional Sciences, Pennsylvania State University, University Park, PA,
USA 16802
[7]Department of Animal Science, Pennsylvania State University, University Park, PA, USA
16802
[8]Social Science Research Institute, Pennsylvania State University, University Park, PA, USA
16802
25

26 * *Contributed equally*
27

Corresponding author: Ferrari, Matthew J. mjf283@psu.edu
Word count: 3056

ABSTRACT (250 WORDS)

Background
Returning university students represent large-scale, transient demographic shifts and a
potential source of transmission to adjacent communities during the COVID-19 pandemic.
35

Methods
In this prospective longitudinal cohort study, we tested for IgG antibodies against SARS-
CoV-2 in a non-random cohort of residents living in Centre County prior to the Fall 2020
term at the Pennsylvania State University and following the conclusion of the Fall 2020 term.
We also report the seroprevalence in a non-random cohort of students collected at the end of
the Fall 2020 term.
42

Findings
Of 345 community participants, 19 (5·5%) were positive for SARS-CoV-2 IgG antibodies at
their first visit between 7 August and 2 October. Of 625 student participants who returned to
campus for fall instruction, 195 (31·2%) were positive for SARS-CoV-2 antibodies between
26 October and 23 November. Twenty-eight (8·1%) community participants returned a
positive IgG antibody result by 9 December. Only contact with known SARS-CoV-2-positive

individuals and attendance at small gatherings (20-50 individuals) were significant predictors
of detecting IgG antibodies among returning students (aOR, 95% CI: 3·24, 2·14-4·91; 1·62,
1·08-2·44; respectively).

Interpretation

53 Despite high seroprevalence observed within the student population, seroprevalence in a
54 longitudinal cohort of community residents was low and stable from before student arrival for
55 the Fall 2020 term to after student departure. The study implies that heterogeneity in SARS-
56 CoV-2 transmission can occur in geographically coincident populations.

Funding

The Pennsylvania State University Office of the Provost, Social Science Research Institute,
Huck Institute for the Life Sciences, and Clinical and Translational Science Institute;
National Institutes of Health.

RESEARCH IN CONTEXT

Evidence before this study
This study was initiated in Spring of 2020, during the pandemic but prior to the return of
students to US universities. There was no prior research on the consequences of the return to
in-person university instruction on SARS-CoV-2 transmission. While the explicit case of
transmission of SARS-CoV-2 between student and resident communities was poorly
understood at the start of this study, there is a well-established literature that has documented
the role that large population movements, consistent with the return of students to campuses,
play in the introduction and maintenance of pathogen transmission or diseases. Several
studies documented the role of travelers in the introduction of SARS-CoV-2 to populations.
The return of students to colleges and universities creates the potential for SARS-CoV-2
introduction and transmission with a sustained increase in population size, density, and
mixing. No prior study has considered the long-term transmission consequences of such
demographic change on the pre-influx population. We searched PubMed, medRxiv, bioRxiv,
and Google Scholar between May 2020 and September 2020. We used search terms
"seroprevalence", "anti-SARS-CoV-2", "IgG", "COVID antibodies" and "university",
"college". Several modeling studies had been conducted to project likely prevalence of anti-
SARS-CoV-2 antibodies in student and resident populations resulting from the return of
students to in-person instruction in the Fall 2020 term in the US and UK. Several studies
conducted after September 2020 described the county-level incidence of SARS-CoV-2 in
counties that contain colleges and universities. None of these studies specifically
disaggregated the incidence of COVID in resident and student populations.
84
Added value of this study
There is clear evidence that in-person university instruction is correlated with high incidence
in students and higher incidence in counties with colleges and universities in counties those
without such institutions. However, this is the first study to specifically quantify the change
in seroprevalence in non-student residents associated with the return of students to in-person
instruction in a large university setting. Despite high COVID-19 incidence in the student
population, seroprevalence in the community resident cohort was low and stable from before
student arrival for the Fall 2020 term until after student departure. The results of this study
highlight the potential for significant heterogeneity in incidence in geographically coincident
sub-populations.
95
Implications of all the available evidence
The influx of students into college and university towns reflects a significant increase in
population size and density that could result in increases in COVID-19 risk in surrounding
communities. Despite the potential risk, it is possible that SARS-CoV-2 transmission may lag
behind in surrounding communities This implies the potential to minimize the transmission
of SARS-CoV-2 from high prevalence sub-populations with targeted interventions. Future
work should investigate the relative efficacy of specific measures to reduce risk of
transmission between subpopulations and guide communities with high levels of episodic
migration and similar high-density subpopulations.
105
106

BACKGROUND

Demographic shifts, high population densities, and population mobility are known to impact the spread of infectious diseases.[1–5] While this has been well characterized at large scales[6–8], it has proved more challenging to demonstrate at smaller geographic scales.[9–11] In the context of the COVID-19 pandemic, the return of college and university students to in-person and hybrid (in-person and online) instruction in the Fall 2020 term represented a massive demographic shift in many communities in the United States (US); specifically, increased total population and proportion living in high density living facilities, with a concomitant increase in person-to-person interactions.[12] This shift had the potential to increase SARS-CoV-2 transmission within returning student populations and to surrounding community resident populations, particularly for non-urban campuses where incidence lagged behind larger population centers.[13] Modeling analyses conducted prior to student return raised concerns that university re-opening would result in significant SARS-CoV-2 transmission in both the returning student and community resident populations.[14,15]

During the Fall 2020 term, many universities in the US experienced high rates of COVID-19 cases among students,[16] with a 56% increase in incidence among counties home to large colleges or universities relative to matched counties without such institutions.[12] While there is strong evidence of high incidence rates associated with a return to campus at US colleges and universities,[12] the increase in risk in surrounding communities, and the rate of transmission from campuses to communities, have been less well characterized. The observed increases in COVID-19 cases in these communities cannot be explicitly attributed to campus origin, absent detailed contact tracing.

This investigation reports the results of a longitudinal serosurvey of community residents in Centre County, Pennsylvania, USA, which is home to The Pennsylvania State University

(PSU), University Park (UP) campus. The return of approximately 35,000 students to the UP

campus in August 2020 represented a nearly 20% increase in the county population size.[17]

During the Fall 2020 term, more than 4,500 cases of SARS-CoV-2 infections were detected

among the student population.[18] Between 7 August and 2 October 2020 (before and just after

student return), we enrolled a cohort of community residents and tested serum for the

presence of anti-Spike Receptor Binding Domain (S/RBD) IgG, which would indicate prior

SARS-CoV-2 exposure.[19] This was repeated in the same cohort during December 2020 (post-

departure of students), and we present seroprevalence for both sampling waves. Additionally,

returning students were enrolled in a longitudinal cohort, and IgG seroprevalence results are

presented from the first wave of sampling (between October and November 2020, prior to the

end of the term). The hypothesis tested was that the rapid influx of students during the Fall

2020 term would be correlated with increased prevalence of SARS-CoV-2 within the

geographically co-incident community.

METHODS

Design, Setting, and Participants
This human subjects research was conducted with PSU Institutional Review Board approval

and in accordance with the Declaration of Helsinki. The study uses a longitudinal cohort

design, with two separate cohorts: community residents and returning students. We report on

measures from the first two clinic visits for the community resident cohort and the first clinic

visit for the returning student cohort.

To assist with recruitment into studies under the Data4Action (D4A) Centre County COVID

Cohort Study umbrella, a REDCap survey was distributed to residents of Centre County,

where respondents could indicate interest in participating in future studies and provide

demographic data. Returning students received a similar survey to express interest in study

participation and were also recruited through cold-emails and word-of-mouth.

Individuals were eligible for participation in the community resident cohort if they were: ≥18

years old, residing in Centre County at the time of recruitment (June through September

2020); expecting to reside in Centre County until June 2021; fluent in English; and capable of

providing their own consent. PSU students who remained in Centre County through spring

and summer university closure were eligible for inclusion in the community resident cohort

as this group experienced similar geographic COVID-19 risks as community residents

(versus returning students who experienced variable risk during spring and summer

depending on their residence).

Participants were eligible for participation in the returning student cohort if they were: ≥18

years old; fluent in English; capable of providing their own consent; residing in Centre

County at the time of recruitment (October 2020); officially enrolled as PSU UP students for

the Fall 2020 term; and intended to be living in Centre County through April 2021.

In both cohorts, individuals were invited to participate in the survey-only portion of the study

if they were: lactating, pregnant, or intended to become pregnant in the next 12 months;

unable to wear a mask for the clinic visit; demonstrated acute COVID-19 symptoms within

the previous 14 days; or reported a health condition that made them uncomfortable with

participating in the clinic visit.

Upon enrollment, returning students were supplied with a REDCap survey to examine socio-

behavioral phenomena, such as attendance at gatherings and adherence to non-

pharmaceutical interventions, in addition to information pertaining to their travel history and

contact with individuals who were known or suspected of being positive for SARS-CoV-2.

Community residents received similar surveys at both time points with questions relating to

potential SARS-CoV-2 household exposures. All eligible participants were scheduled for a

clinical visit at each time interval where blood samples were collected for ELISA testing.

Outcomes

The primary outcome was the presence of S/RBD IgG antibodies, measured using an indirect isotype-specific (IgG) screening ELISA developed at PSU.[20] Further details in the Supplement. The presence of anti-SARS-CoV-2 antibodies has been documented in prior seroprevalence studies as a method of quantifying cumulative exposure.[21–23]

Statistical Methods

Community resident and returning student cohorts' seroprevalence are presented with binomial 95% confidence intervals. We estimated each subgroup's true prevalence, accounting for imperfect sensitivity and specificity of the IgG assay, using the *truePrev()* function in the *prevalence* package in R. We calculated a 95% binomial confidence interval for test sensitivity of the IgG assay for detecting prior self-reported positive tests in the returning student cohort (students had high access to testing from a common University provider) and assumed a uniform prior distribution between these limits. Estimates of the true prevalence were then calculated across all possible values of specificity between 0·85 and 0·99. We did not estimate prevalence corrected for demographics as participants were not enrolled using a probability-based sample. We assessed demographic characteristics of the tested participants relative to all study participants to illustrate potential selection biases (Table 1).

For contingency tables and raw odds ratios, we used two-sided Fisher's exact test to determine significance at alpha = 0·05 and present 95% confidence intervals. Pairwise Fisher's exact test of proportions was used with the Bonferroni-Holm correction in the *r* x *c* case, and Welch's two-sample t-test to compare distributions.

Missing values were deemed "Missing At Random" and imputed, as described in the Supplement. We estimated the adjusted odds ratios (aOR) of IgG positivity in the returning student subgroup using multivariable logistic regression implemented with the *mice* and

finalfit packages. We considered the following risk factors: close proximity (6 feet or less) to

an individual who tested positive for SARS-CoV-2; close proximity to an individual showing

key COVID-19 symptoms (fever, cough, shortness of breath); attendance at a small gathering

(20-50 people) in the past 3 months; attendance at a medium gathering (51-200 people) in the

past 3 months; live in University housing; ate in a restaurant in the past 7 days; ate in a dining

hall in the past 7 days; only ate in their room/apartment in the past 7 days; travelled in the 3

months prior to returning to campus; and travelled since returning to campus for the Fall

term. Given the airborne transmission of SARS-CoV-2, all the above variables are potential

exposure risks due to the increased contact with individuals outside of a participants'

household.[24–27]

All statistical analyses were conducted using R – version 4.03 (2020-10-10), with a pipeline

created using the *targets* package.

Role of the Funding Source

This research was supported by funding from the Office of the Provost and the Clinical and

Translational Science Institute, Huck Life Sciences Institute, and Social Science Research

Institutes at the Pennsylvania State University. The project described was supported by the

National Center for Advancing Translational Sciences, National Institutes of Health, through

Grant UL1 TR002014. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the NIH. The funding sources had no role in the

collection, analysis, interpretation, or writing of the report.

RESULTS

A total of 10,369 community residents were identified through an initial REDCap survey that

collected eligibility, demographic, and contact information. Of those who identified

willingness to participate in a later study, 1,540 were contacted and enrolled. 1,432

completed a first clinic visit between 7 August and 2 October 2020, and 345 of those

completed a second clinic visit between 30 November and 9 December 2020 and for whom

both visit 1 and visit 2 samples were analysed using the in-house ELISA assay; the remaining

participants' samples will be analysed in the future. 1,349 returning students were recruited

using volunteer sampling and 707 enrolled; of these, 625 completed clinic visits for serum

collection between 26 October and 23 November. 642 students indicated willingness to

participate but did not complete intake procedures or did not attend their scheduled clinic

visit.

Among participants with serum samples: the median age community residents was 47 years

(IQR: 35·0-60·0), with 81·7% between the ages 18-65 years, and for the returning students

the median age was 20·9 years (IQR: 19·7-21·6), with 100% between the ages 18-65 years;

62·0% of the community residents identified as female and 33·6% as male; 64·6% of the

returning students identified as female and 34·4% as male; 90·1% of the community residents

identified as white, as did 81·3% of the students. Similar proportions were seen in those

enrolled without samples, and among the initial REDCap survey respondents (Table 1; Table

2).

Although all county residents were eligible for participation, 77·0% of community resident

participants were from the 5 townships (College, Ferguson, Harris, Half Moon, Patton) and 1

borough (State College) that form the "Centre Region" and account for ~59% of Centre

County's population[17] (Figure 1). The median household income group in the community

residents providing samples was $100,000-149,999 USD (IQR: $50,000-74,999; $100,000-

149,999). The median household income in the county is $60,403.[17] 47.4% of the county is

female, 87·9% white, and 70·3% are between the ages of 18-65 years old.[17] The study cohort

is moderately older and more affluent (in part because of the exclusion of returning students),

and disproportionately female compared to the general Centre County population.

Of the returning student participants, 665 (94·1%) had at least one test prior to enrollment in

the study; of these, 105 (15·8%) self-reported a positive result (Table 3). Of these, 98 (93·3%)

indicated that this test result occurred after their return to campus (median: 26 September;

IQR: 13 September, 9 October). Of the 625 returning students with an ELISA result, 84 of

the 90 (93·3%) with a self-reported prior positive test result were positive for SARS-CoV-2

IgG antibodies. Of the 535 returning students with ELISA results who did not report a

positive SARS-CoV-2 test, 111 (20·7%) were positive for SARS-CoV-2 IgG antibodies. Of

the total 625 returning students with ELISA results, 195 (31·2%) were positive for SARS-

CoV-2 IgG antibodies (Figure 2).

Among the community resident participants, 19 of 345 (5·5%) were positive for SARS-CoV-

2 antibodies at their first visit (Figure 2). Between their first and second visit, 9 participants

converted from negative to positive and 9 converted from positive to negative; 28 (8·2%)

were positive for SARS-CoV-2 IgG antibodies at either visit (Figure 2).

To estimate true prevalence of SARS-CoV-2 exposure in our cohorts, the seroprevalence was

corrected for the sensitivity and specificity of the assay. 93·3% (95% CI: 86-97%) of

returning students with a self-reported prior positive SARS-CoV-2 test had positive IgG

antibodies; this was used as an estimate of sensitivity of the IgG assay for detecting

previously detectable infection (see Supplement for an alternative calculation of sensitivity

that includes community resident responses). For all assumed levels of specificity, the 95%

credible intervals for the true prevalence in the community residents overlapped for the pre-

and post-term time points, and neither overlapped the 95% credible interval for the true

prevalence estimate in the returning student subgroup (Figure 3).

Among the returning student subgroup, only close proximity to a known SARS-CoV-2-

positive individual and attending small gatherings in the past 3 months (20-50 individuals)

were significantly associated with a positive ELISA classification at alpha = 0·05 (aOR: 3·24,

2·14-4·91; aOR: 1·62, 1·08-2·44; respectively) in the multivariable model (Table 4). Taken

individually, attending small gatherings (20-50 people) (OR: 2·38, 1·66-3·43), attending

medium gatherings (51-200 people) (OR:1·88, 1·19-2·94), and close proximity to an

individual showing key COVID-19 symptoms (OR: 1·81, 1·24-2·64) were all associated with

the IgG positivity in crude calculations of association. Given the low prevalence observed

within the community subgroup, and lack of significant effects observed during crude

analysis, models were not fit to the community resident data.

Both the returning student and community resident participants self-reported high compliance

with masking; 86·4% and 76·9%, respectively, reported always wearing mask or cloth face

covering when in public (Table S3, Table S4). By contrast, less than one third of both groups

(28·8% and 31·0%, respectively) self-reported always maintaining 6-feet of distance from

others in public. Less than half (42·9%) of returning students indicated that they always

avoided groups of 25 or greater, in contrast with 65·7% of community residents.

DISCUSSION
The return of students to in-person instruction on the PSU UP campus was associated with a

large increase in COVID-19 incidence in the county, as evidenced by over 4,500 student

cases at PSU.[18] In a sample of 625 returning students, 31% were positive for SARS-CoV-2

antibodies. Out of approximately 35,000 students who returned to campus, this implies that

the detected cases may account for ~40% of all infections among PSU UP students. Despite

this high incidence of SARS-CoV-2 infection in the county during the Fall 2020 term, a

cohort of community residents, who disproportionately identified as female and lived in close

proximity to campus, saw only a modest increase in the prevalence of SARS-CoV-2 IgG

antibodies (5·5 to 8·1%) between September and December 2020, aligning with a nation-wide

estimate of seroprevalence for the summer of 2020.[23] The true prevalence of prior SARS-

CoV-2 infection in the cohorts depends on the assumed sensitivity and specificity. However,

for all realistic values of sensitivity and specificity, there was no evidence of a significant

increase in true prevalence in the community resident sample. While in-person student

instruction has been associated with an increase in per-capita COVID-19 incidence at the

aggregate level,[12] these results suggest that outbreaks in the returning student and the

community resident cohort we studied were asynchronous, implying limited between-cohort

transmission. A recent analysis of age-specific movement and transmission patterns in the US

suggested that individuals between the ages of 20-34 disproportionately contributed to spread

of SARS-CoV-2;[28] however, despite close geographic proximity to a college-aged

population, transmission in our community resident sample appears distinctly lagged;

suggestive of the potential for health behaviours to prevent infection.

Within the student group, presence of SARS-CoV-2 antibodies was significantly associated

with close proximity to known SARS-CoV-2-positive individuals and attendance of small

events, however, no other risk factors were correlated with an increase in SARS-CoV-2 IgG

test positivity, aligning with other research.[23] However, it is not possible to discern how much

the likelihood of contact with a SARS-CoV-2 positive individual is due to the high campus

prevalence versus individual behaviours. When considered independently, eating in dining

halls within the past 7 days was weakly associated with testing positive for SARS-CoV-2

antibodies and participation in medium-sized events (51-200 individuals) was significantly

associated with testing positive for SARS-CoV-2 antibodies, which is consistent with

patterns observed elsewhere.[26,27] Both community residents and returning students reported

high adoption of masking (>75%) and low adoption of distancing in public (31·0% and

28·8%, respectively). Returning students reported lower adherence to avoiding gatherings of

greater than 25 individuals than community residents.

Neither the resident nor the student participants were selected using a probability-based

sample. Thus, these participants may not be representative of any particular population.

Those who chose to participate in this study may have been more cognizant and compliant

with public health mitigation measures than non-participants. Specifically, the resident

participants disproportionately lived in the townships immediately surrounding the UP

campus, where extensive messaging[29] and preventative campaigns were enacted, and they

have a higher median income than the residents of Centre County overall.

Though the participants reflect a convenience sample, the large differences in SARS-CoV-2

seroprevalence suggest that the cohorts did not experience a synchronous, well-mixed

epidemic despite their close geographic proximity. College campuses have been observed to

have high COVID-19 attack rates, and counties containing colleges and universities have

been observed to have significantly higher COVID-19 incidence than demographically

matched counties without such institutions.[12] Thus, while college and university operations

may present a significant exposure risk (because of large numbers of returning students, high

density housing, and frequent in-person socialization), this analysis suggests the possibility

that local-scale heterogeneity in mixing may allow for asynchronous transmission dynamics

despite close geographic proximity. Thus, the disproportionately high incidence in the student

population, which comprises less than one quarter of the county population, may bias

assessment of risk in the non-student resident population. Risk assessment in spatial units

(e.g., counties) that have strong population sub-structuring should consider these

heterogeneities and their consequences for policy.

SARS-CoV-2 transmission between the student and community resident populations is likely

to have occurred (perhaps multiple times) and these results do not preclude that possibility.

However, the large difference in seroprevalence between the student and resident participants

after the Fall term are consistent with either rare transmission events between students and

residents, non-persistent transmission with the community residents, or both. This suggests

that it is possible to minimize risks brought about by sub-populations with high SARS-CoV-2

incidence using behavioral interventions. This observation may have implications for

outbreak management in other high risk, highly mobile populations (e.g., displaced

populations, seasonal workers, military deployment). However, we note that this was

achieved in the context of disproportionate investment in prevention education, testing,

contact tracing, and infrastructure for isolation and quarantine by PSU in the high-prevalence

sub-population (students).

With respect to the health behaviors measured, the community resident and returning student

groups differed only in their small social group contact, and thus a next step is to identify

factors that may explain this difference (e.g., differences in leisure time activity norms and/or

perceptions of age-related risk; business closures; university policies and sanctions).

Minimizing risk, however, may come at significant social, psychological, educational,

economic, and societal costs.[30] Thus, operational planning for both institutions of higher

education and the communities in which they are embedded should consider both the risk of

SARS-CoV-2 transmission and the costs of mitigation efforts.

ACKNOWLEDGEMENTS
371

Author Contributions
Conceptualization: MJF, NB, MS, AR, VK
Data curation: MJF, CA
Formal analysis: CA, MJF, CMH
Funding acquisition: MJF, AR
Investigation: SS, AG, MMS, CJR
Methodology: CA, MJF
Project administration: MJF, MMS
Software: CA, MJF, CMH
Supervision: MJF, VK, SK
Validation: CA, MJF, CMH, SS
Visualization: CA, MJF
Writing - original draft: CA, SS, CMH, VK, MJF
Writing - review and editing: all authors.

Conflicts of Interest and Financial Disclosures
The authors declare no conflicts of interest.

Financial and Material Support
This research was supported by funding from the Office of the Provost and the Clinical and
Translational Science Institute, Huck Life Sciences Institute, and Social Science Research
Institutes at the Pennsylvania State University. The project described was supported by the
National Center for Advancing Translational Sciences, National Institutes of Health, through
Grant UL1 TR002014. The content is solely the responsibility of the authors and does not
necessarily represent the official views of the NIH.
395

Data Access, Responsibility, and Analysis
Callum Arnold and Dr. Matthew J. Ferrari had full access to all the data in the study and take
responsibility for the integrity of the data and the accuracy of the data analysis. Callum
Arnold, Dr. Matthew J. Ferrari (Department of Biology, Pennsylvania State University), and
Dr. Catherine M. Herzog (Huck Institutes of the Life Sciences, Pennsylvania State
University) conducted the data analysis.

Collaborators
1. Florian Krammer, Mount Sinai, USA for generously providing the transfection plasmid
pCAGGS-RBD
2. Scott E. Lindner, Allen M. Minns, Randall Rossi produced and purified RBD
3. The D4A Research Group: Dee Bagshaw, Clinical & Translational Science Institute,
Cyndi Flanagan, Clinical Research Center and the Clinical & Translational Science
Institute, Thomas Gates, Social Science Research Institute, Margeaux Gray, Dept. of
Biobehavioral Health, Stephanie Lanza, Dept. of Biobehavioral Health and Prevention
Research Center, James Marden, Dept. of Biology and Huck Institutes of the Life
Sciences, Susan McHale, Dept. of Human Development and Family Studies and the
Social Science Research Institute, Glenda Palmer, Social Science Research Institute,
Rachel Smith, Dept. of Communication Arts and Sciences and Huck Institutes of the Life

Sciences, and Charima Young, Penn State Office of Government and Community
Relations.
4. The authors thank the following for their assistance in the lab: Liz D. Cambron,
Elizabeth M. Schwartz, Devin F. Morrison, Julia Fecko, Brian Dawson, Sean Gullette,
Sara Neering, Mark Signs, Nigel Deighton, Janhayi Damani, Mario Novelo, Diego
Hernandez, Ester Oh, Chauncy Hinshaw, B. Joanne Power, James McGee, Riëtte van
Biljon, Andrew Stephenson, Alexis Pino, Nick Heller, Rose Ni, Eleanor Jenkins, Julia
Yu, Mackenzie Doyle, Alana Stracuzzi, Brielle Bellow, Abriana Cain, Jaime Farrell,
Megan Kostek, Amelia Zazzera, Sara Ann Malinchak, Alex Small, Sam DeMatte,
Elizabeth Morrow, Ty Somberger, Haylea Debolt, Kyle Albert, Corey Price, Nazmiye
Celik
425

REFERENCES

Bengtsson L, Gaudart J, Lu X, *et al.* Using Mobile Phone Data to Predict the Spatial Spread of Cholera. *Scientific Reports* 2015; **5**: 8923.

Wilson ME. Travel and the emergence of infectious diseases. *Emerg Infect Dis* 1995; **1**: 39–46.

Chinazzi M, Davis JT, Ajelli M, *et al.* The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science* 2020; **368**: 395–400.

Viboud C, Bjørnstad ON, Smith DL, Simonsen L, Miller MA, Grenfell BT. Synchrony, Waves, and Spatial Hierarchies in the Spread of Influenza. *Science* 2006; **312**: 447–51.

Bharti N, Tatem AJ, Ferrari MJ, Grais RF, Djibo A, Grenfell BT. Explaining seasonal fluctuations of measles in Niger using nighttime lights imagery. *Science* 2011; **334**: 1424–7.

Wells CR, Sah P, Moghadas SM, *et al.* Impact of international travel and border control measures on the global spread of the novel 2019 coronavirus outbreak. *PNAS* 2020; **117**: 7504–9.

Bogoch II, Creatore MI, Cetron MS, *et al.* Assessment of the potential for international dissemination of Ebola virus via commercial air travel during the 2014 west African outbreak. *The Lancet* 2015; **385**: 29–35.

Jia JS, Lu X, Yuan Y, Xu G, Jia J, Christakis NA. Population flow drives spatio-temporal distribution of COVID-19 in China. *Nature* 2020; **582**: 389–94.

Funk S, Ciglenecki I, Tiffany A, *et al.* The impact of control strategies and behavioural changes on the elimination of Ebola from Lofa County, Liberia. *Philos Trans R Soc Lond B Biol Sci* 2017; **372**. DOI:10.1098/rstb.2016.0302.

Sniadack DH, Moscoso B, Aguilar R, Heath J, Bellini W, Chiu MC. Measles epidemiology and outbreak response immunization in a rural community in Peru. *Bull World Health Organ* 1999; **77**: 545–52.

Stoddard ST, Forshey BM, Morrison AC, *et al.* House-to-house human movement drives dengue virus transmission. *PNAS* 2012; published online Dec 27. DOI:10.1073/pnas.1213349110.

Leidner AJ, Barry V, Bowen VB, *et al.* Opening of Large Institutions of Higher Education and County-Level COVID-19 Incidence — United States, July 6–September 17, 2020 Vaughn Barry, PhD1; Virginia B. Bowen, PhD1; Rachel Silver, MPH1; Trieste Musial, MS2; Gloria J. Kang, PhD1; Matthew D. Ritchey, DPT3; Kelly Fletcher, MPH2; Lisa Barrios, DrPH1; Eric Pevzner, PhD. *MMWR Morb Mortal Wkly Rep* 2021; **70**. DOI:10.15585/mmwr.mm7001a4.

Goetz S, Tian Z, Schmidt C, Meadowcroft D, NERCRD, Pennsylvania State University. Rural COVID-19 Cases Lag Urban Areas but Are Growing Much More Rapidly. Pennsylvania State University Department of Agricultural Economics, Sociology, and

Education, 2020 https://aese.psu.edu/nercrd/publications/covid-19-issues-briefs/rural-
covid-19-cases-lag-urban-areas-but-are-growing-much-more-rapidly (accessed Feb 6,
2021).

14 Christensen H, Turner K, Trickey A, *et al.* COVID-19 transmission in a university setting:
a rapid review of modelling studies. *medRxiv* 2020; : 2020.09.07.20189688.

15 Andersen MS, Bento AI, Basu A, Marsicano C, Simon K. College Openings, Mobility,
and the Incidence of COVID-19 Cases. *medRxiv* 2020; : 2020.09.22.20196048.

16 The New York Times. Tracking the Coronavirus at U.S. Colleges and Universities - The
New York Times. Tracking the Coronavirus at U.S. Colleges and Universities. 2020;
published online Dec 11. https://www.nytimes.com/interactive/2020/us/covid-college-
cases-tracker.html (accessed Jan 30, 2021).

17 United States Census Bureau. U.S. Census Bureau QuickFacts: Centre County,
Pennsylvania. QuickFacts: Centre County, Pennsylvania. 2019.
https://www.census.gov/quickfacts/centrecountypennsylvania (accessed Jan 30, 2021).

18 Pennsylvania State University. COVID-19 Dashboard. Virus Info. 2021; published online
Jan 29.
https://app.powerbi.com/view?r=eyJrIjoiNDY3NjhiMDItOWY0Mi00NzBmLWExNTAtZ
GIzNjdkMGI0OTM0IiwidCI6IjdjZjQ4ZDQ1LTNkZGItNDM4OS1hOWMxLWMxMTU
1MjZlYjUyZSIsImMiOjF9 (accessed Jan 30, 2021).

19 Long Q-X, Liu B-Z, Deng H-J, *et al.* Antibody responses to SARS-CoV-2 in patients with
COVID-19. *Nat Med* 2020; **26**: 845–8.

20 Gontu A, Srinivasan S, Nair MS, *et al.* Quantitative Estimation of IgM and IgG Antibodies
Against SARS-CoV-2. *protocols.io* 2020; published online Aug 20.
DOI:dx.doi.org/10.17504/protocols.io.bivgke3w.

21 Uyoga S, Adetifa IMO, Karanja HK, *et al.* Seroprevalence of anti–SARS-CoV-2 IgG
antibodies in Kenyan blood donors. *Science* 2021; **371**: 79–82.

22 Stringhini S, Zaballa M-E, Perez-Saez J, *et al.* Seroprevalence of anti-SARS-CoV-2
antibodies after the second pandemic peak. *The Lancet Infectious Diseases* 2021; **0**.
DOI:10.1016/S1473-3099(21)00054-2.

23 Kalish H, Klumpp-Thomas C, Hunsberger S, *et al.* Mapping a Pandemic: SARS-CoV-2
Seropositivity in the United States. *medRxiv* 2021; : 2021.01.27.21250570.

24 Huang Y-T, Tu Y-K, Lai P-C. Estimation of the secondary attack rate of COVID-19 using
proportional meta-analysis of nationwide contact tracing data in Taiwan. *Journal of*
Microbiology, Immunology and Infection 2020; published online June 11.
DOI:10.1016/j.jmii.2020.06.003.

25 Cheng H-Y, Jian S-W, Liu D-P, *et al.* Contact Tracing Assessment of COVID-19
Transmission Dynamics in Taiwan and Risk at Different Exposure Periods Before and
After Symptom Onset. *JAMA Internal Medicine* 2020; **180**: 1156–63.

26 Leclerc QJ, Fuller NM, Knight LE, Funk S, Knight GM. What settings have been linked to
SARS-CoV-2 transmission clusters? *Wellcome Open Res* 2020; **5**.
DOI:10.12688/wellcomeopenres.15889.2.

27 Brooks-Pollock E, Read JM, House T, Medley GF, Keeling MJ, Danon L. The Population
Attributable Fraction (PAF) of cases due to gatherings and groups with relevance to
COVID-19 mitigation strategies. *medRxiv* 2020; : 2020.03.20.20039537.

28 Monod M, Blenkinsop A, Xi X, *et al.* Age groups that sustain resurging COVID-19
epidemics in the United States. *Science* 2021; : eabe8372.

29 Pennsylvania State University. Mask Up or Pack Up. Virus Info. 2021.
https://virusinfo.psu.edu/mask-up-or-pack-up/ (accessed Feb 6, 2021).

30 Brooks SK, Webster RK, Smith LE, *et al.* The psychological impact of quarantine and
how to reduce it: rapid review of the evidence. *Lancet* 2020; **395**: 912–20.

515
516
517

 TABLES

Table 1
Demographic characteristics of study participants. Non-D4A participants are all participants
in the initial anonymous survey from which Data4Action participants were drawn. D4A
participants are divided into subsets for which antibody assays were conducted (N=345) and
those for which assays were not conducted (N=1195).

Variable	D4A Participant (N=1540)		Not D4A Participant (N=8826)
	Assay Subset (N=345)	**Non-Assay Subset (N=1195)**	
Age			
Median [IQR]	47·0 [35·0, 60·0]	46·0 [36·0, 58·0]	46·0 [33·0, 59·0]
Missing	1 (0·3%)	11 (0·9%)	335 (3·8%)
Race			
White	311 (90·1%)	1057 (88·5%)	6075 (68·8%)
Aggregated Category*	18 (5·2%)	84 (7·0%)	1289 (14·6%)
Listed more than one race or ethnicity	15 (4·3%)	41 (3·4%)	159 (1·8%)
Missing	1 (0·3%)	13 (1·1%)	1303 (14·8%)
Gender			
Female	214 (62·0%)	784 (65·6%)	0 (0%)
Male	116 (33·6%)	337 (28·2%)	0 (0%)
Non-binary / Transgender / Self-described	6 (1·7%)	8 (0·7%)	0 (0%)
Prefer not to answer	1 (0·3%)	1 (0·1%)	0 (0%)
Missing	8 (2·3%)	65 (5·4%)	8826 (100%)
Household Income			
$200,000 and over	28 (8·1%)	134 (11·2%)	677 (7·7%)
$150,000 to $199,999	49 (14·2%)	158 (13·2%)	767 (8·7%)
$100,000 to $149,999	91 (26·4%)	311 (26·0%)	1502 (17·0%)
$75,000 to $99,999	41 (11·9%)	171 (14·3%)	1091 (12·4%)
$50,000 to $74,999	47 (13·6%)	149 (12·5%)	963 (10·9%)
$25,000 to $49,999	46 (13·3%)	123 (10·3%)	1462 (16·6%)
Under $25,000	9 (2·6%)	44 (3·7%)	259 (2·9%)
Prefer not to answer	33 (9·6%)	93 (7·8%)	801 (9·1%)
Missing	1 (0·3%)	12 (1·0%)	1304 (14·8%)

* Asian; Hispanic, Lantino/a, or Spanish; Black or African American; Middle Eastern or
North African; Native American or Alaska Native; other race or ethnicity. This category is
aggregated to protect participant identities because no single group comprised >4% of
participants.

528 **Table 2**
529 Demographic characteristics of the returning student participants.
530

Variable	PSU Subset (N=625)
Age	
Median [IQR]	20·8 [19·7, 21·6]
Missing	10 (1·6%)
Race	
White	508 (81·3%)
Aggregated Category*	81 (13·0%)
Listed more than one race	30 (4·8%)
Missing	6 (1·0%)
Gender	
Female	404 (64·6%)
Male	215 (34·4%)
Genderqueer/nonconforming/ transgender/different identity	5 (0·8%)
Missing	1 (0·2%)
University Housing	
Not Uni housing	445 (71·2%)
Uni housing	178 (28·5%)
Missing	2 (0·3%)

531 * Asian; Black or African American; American Indian or Alaska Native; Native Hawaiian or
532 other Pacific Islander; some other race. This category is aggregated to protect participant
533 identities because no single group comprised >4% of participants.
534
535
536

Table 3

IgG ELISA results as a function of self-reported prior SARS-CoV-2 diagnostic test outcome among returning student cohort participants.

ELISA Result	Prior Test (N=665)		No Prior Test (N=42)
	Prior Positive (N=105)	No Prior Positive (N=560)	
Positive	84	106	5
Negative	6	390	34
NA	15	64	3

Table 4
Crude and adjusted odds ratios (aOR) of risk factors among returning PSU UP student cohort

Variable/Response	No/total (%) with SARS-CoV-2 antibodies	Crude OR (95% CI)	MICE Multivariable OR (95% CI)
Close proximity to known COVID-19 Positive Individual			
No	57/316 (18·04)	1 [Reference]	1 [Reference]
Yes	137/307 (44·63)	3·65 (2·51-5·37)***	3·24 (2·14-4·91)***
Attended a gathering of 20-50 people since arrival for the Fall Semester			
No	75/332 (22·59)	1 [Reference]	1 [Reference]
Yes	119/290 (41·03)	2·38 (1·66-3·43)***	1·62 (1·08-2·44)*
Ate in a dining hall in the past 7 days			
No	150/500 (30·00)	1 [Reference]	1 [Reference]
Yes	43/121 (35·54)	1·29 (0·82-1·99)	1·40 (0·79-2·48)
Attended a gathering of 51-200 people since arrival for the Fall Semester			
No	146/510 (28·63)	1 [Reference]	1 [Reference]
Yes	46/107 (42·99)	1·88 (1·19-2·94) **	1·26 (0·77-2·06)
Travelled in the 3 months prior to campus arrival			
No	74/263 (28·13)	1 [Reference]	1 [Reference]
Yes	114/339 (33·63)	1·29 (0·90-1·87)	1·15 (0·78-1·70)
Ate in a restaurant in the past 7 days			
No	90/317(28·39)	1 [Reference]	1 [Reference]
Yes	102/305 (33·44)	1·27 (0·89-1·81)	1·10 (0·75-1·62)
Only ate in their room in the past 7 days			
No	70/214 (32·71)	1 [Reference]	1 [Reference]
Yes	123/409 (30·07)	0·88 (0·61-1·28)	0·97 (0·64-1·46)
Close proximity to individual showing COVID-19 symptoms			
No	121/442 (27·38)	1 [Reference]	1 [Reference]

Yes	73/180 (40·56)	1·81 (1·24-2·64)**	0·89 (0·59-1·36)
Travelled since campus arrival			
No	80/255 (31·37)	1 [Reference]	1 [Reference]
Yes	114/368 (30·98)	0·98 (0·69-1·41)	0·87 (0·59-1·27)
Lives in University housing			
No	140/445 (31·46)	1 [Reference]	1 [Reference]
Yes	54/178 (30·34)	0·95 (0·64-1·40)	0·82 (0·50-1·36)

543
544
545 * p < 0·05 (2-tailed)
546 ** p < 0·01 (2-tailed)
547 *** p < 0·001 (2-tailed)

FIGURES

Figure 1
Map of Centre County, Pennsylvania, USA. Blue indicates the 5 townships and 1 borough
that comprise the Centre Region. Red indicates the location of The Pennsylvania State
University (PSU), University Park (UP) Campus. Inset illustrates the proportion of the county
population in each region; PSU indicates the estimated student population that returned to
campus for the Fall 2020 term.

Dot Indicates PSU Campus
Each square represents 1000 individuals

568
569
570

571   **Figure 2**
572   Raw seroprevalence (circles) with 95% binomial confidence intervals for the community
573   residents at the first visit at the start of the Fall 2020 term (light blue), returning students at
574   the end of the fall 2020 term (red), and community residents at either the first or the second
575   visit after student departure (dark blue).
576

Figure 3
Estimated true prevalence (circles, with 95% confidence intervals) among participants at each
sampling interval corrected for estimated assay sensitivity as a function of the assumed assay
specificity (x-axis). Light blue indicates community residents at the first visit at the start of
the Fall 2020 term, red indicates returning students at the end of the Fall 2020 term, and dark
blue indicates community residents at the second visit after student departure.
584