

Factors influencing self-harm thoughts and self-harm behaviours over the first 45 weeks of the COVID-19 pandemic in the UK: a longitudinal analysis of 48,446 adults

Elise Paul, PhD and Daisy Fancourt, PhD

**Keywords** Self-harm behaviours, longitudinal studies, COVID-19, self-harm thoughts, adversity

## Background

There is concern that the COVID-19 pandemic and its aftermath will result in excess suicides by increasing known risk factors such as self-harm. However, evidence on how pandemic-related risk factors contribute to changes in these outcomes is lacking.

## Aims

We sought to examine how different adverse experiences and worries about adversity brought about by the COVID-19 pandemic contribute to changes in self-harm thoughts and behaviours over the first 45 weeks of the pandemic.

## Method

Data from 48,446 UK adults in the UCL COVID-19 Social Study were analysed across the first 45 weeks (1 April to 7 February 2021) of the pandemic. Fixed effects regressions explored associations between weekly within-person variation in five categories of adversity experiences and adversity worries with changes in self-harm thoughts and behaviours across age groups (18-29, 30-44, 45-59, and 60+ years).

## Results

More adverse experiences were more strongly related to outcomes than worries. Specific categories of worries and adversities had stronger associations with outcomes than others. The largest contributing factor to increases in both self-harm thoughts (odds ratio (OR) range = 3.38 to 4.00) and behaviours (OR range = 3.72 to 6.82) was physical or psychological abuse amongst all ages. Having had COVID-19 and financial worries were associated with increased likelihood of both outcomes, the latter being more consistently associated with outcomes than actual financial adversity.

## Conclusions

These findings suggest that a significant portion of UK adults may be at increased risk for self-harm thoughts and behaviours during the pandemic, which is concerning given the link between self-harm and suicide. Policies currently place to protect and reassure citizens may be insufficient in assuaging fears and anxieties. Given the likelihood that as time goes on, the economic and social consequences of the COVID-19 pandemic will accumulate, policy makers can begin adapting a variety of existing evidence-based suicide prevention strategies and other social policies to help mitigate its consequences.

## Introduction

Numerous studies have shown that the COVID-19 pandemic is having a detrimental impact on population mental health, and although not inevitable, there are concerns that suicide rates will subsequently increase (1,2). Although other high-income countries have reported either no meaningful change or a decrease in suicide rates in the first months of the pandemic (3–5), temporarily lowered suicide rates have been observed in the early phases of other crises such as natural disasters and epidemics that were then followed by increases (6,7). This pattern has already been observed during the COVID-19 pandemic in Japan, where the first five months were marked by a 14% reduction in suicides, followed by a 16% increase in overall suicides, with a 49% increase in children and adolescents and a 37% increase in females (8). There are several reasons why suicides may increase once the immediate crisis has passed. First, the COVID-19 pandemic has involved the exacerbation of known risk factors for suicide such as unemployment, mental health problems, intimate partner violence, and insufficient access to mental health care that may not immediately resolve as the pandemic abates (1,2,9). Second, the cumulative effects of lockdowns, job losses, and uncertainty during the pandemic itself may start to take a toll over time (1,9). Third, the International Monetary Fund predicts that the global recession resulting from the COVID-19 pandemic will be the worst since the Great Depression (10), and research has consistently identified links between economic recessions, large scale unemployment, and increases in suicide rates (11,12). All of these stressful circumstances and life events have the potential to increase risk for suicide through increasing mental health difficulties such as depression, defeat, anxiety, and a sense of entrapment (13,14).

One reason for concern about a potential future increase in suicide deaths as a result of the pandemic is that there is already evidence that risk factors for death by suicide have been increasing. Thinking about self-harming, suicide or death and intentionally damaging or injuring oneself have been widely observed to be risk factors for death by suicide (15–17). A number of studies have suggested that prevalence rates for thinking about self-harm or suicide or engaging in self-harming have been higher during the pandemic than previously (18–20). Although clinical presentations for self-harm have been significantly lower in the early months of the COVID-19 pandemic compared to prior trends (22,23), this could have been due to fears of contracting COVID-19 in hospitals and not wanting to be a burden on the healthcare system (9,24). Even pre-pandemic, the majority of individuals who self-harm or consider suicide do not seek help from clinical services (25).

In considering why these risk factors for suicide may have increased in the first months of the current pandemic, several studies have identified potential predictors. Experiencing physical/psychological abuse and receiving a COVID-19 diagnosis (18), legal problems, ongoing arguments with a partner, and worries about a life-threatening illness or injury in a family member or close friend have been associated with thinking about and carrying out self-harm (26), as have new and exacerbated mental health problems and insufficient access to mental health care (27). However, studies exploring predictors of self-harm thoughts and behaviours have been limited so far in which predictors they have considered and considering predictors at a single moment in time. As the social and economic circumstances of the pandemic are changing so fast, predictors identified early on in the pandemic may no longer be relevant. So, it is important to have updated information on what is causing people to think about harming themselves and to actually do so as the pandemic continues. Finally, it is important to identify which factors are associated over time not just with an increased overall risk in self-harm thoughts and behaviours but dynamic changes (both increases and decreases) so that modifiable targets to reduce self-harm can become the subject of future interventions.

Therefore, the aim of this study is to establish which factors are associated with changes over time in thoughts of death or self-harm (hereafter referred to as “self-harm thoughts”) and self-harm behaviours in a large sample of UK adults across the first 45 weeks of the COVID-19 pandemic. Specifically, we

explore the time-varying longitudinal relationships between (i) adverse experiences and ii) worries about adverse experiences with changes in self-harm thoughts and behaviours, and how these associations vary by age. Identifying specific concerns and adversities that are risk factors for self-harm thoughts and behaviours will provide an opportunity for policy makers to address those issues by designing policies to mitigate the impact of the COVID-19 pandemic and the anticipated upcoming economic recession.

## Methods

### Study design and participants

Data were drawn from the COVID-19 Social Study; a large panel study of the psychological and social experiences of over 75,000 adults (aged 18+) in the UK during the COVID-19 pandemic. The study commenced on 21 March 2020 and involves online weekly data collection for the first 22 weeks of the COVID-19 lockdown in the UK (until 15 August 2020) then monthly collection thereafter. Sampling was not random and therefore is not representative of the UK population, but the sample is heterogeneous. More information on sampling methods can be found in the Supplementary Materials.

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human subjects were approved by the UCL Ethics Committee. Written informed consent was obtained from all participants.

As questions asked about adverse experiences and worries about adversity in the prior week, we focused on data collected from 1 April 2020 (one week after lockdown commenced in the UK) to 7 February 2021 ( $n = 65,218$ , observations = 811,163). We then limited our analysis to participants who participated on three or more occasions during this period ( $n = 50,770$  observations = 791,249). We further excluded participants who had missing data on any study variable for at least three interviews ( $n = 2,324$ ). This resulted in the final sample of 48,446 participants totalling 746,123 observations.

## Outcomes

### Self-harm thoughts and behaviours

Self-harm thoughts were measured with an item from the Patient Health Questionnaire (PHQ-9) (28) and self-harm behaviours were measured with a similar study-developed item (See Supplementary Materials). Responses to both items were collapsed into the presence (one or two days, more than half the days, or nearly every day) vs absence (not at all) at each time point.

## Exposures

### Adversity experiences

Five categories of adversities measured weekly for the first 22 weeks of the study (1 April to 21 August 2020) and then monthly to 7 February 2021 were considered: financial adversity, COVID-19 illness, family/friend illness or bereavement, experiencing physical or psychological abuse, and not being able to access essential items. Each category of adversity was treated as binary (absent vs. present). More detailed description of these measures can be found in the Supplementary Materials.

### Worries about adversity

Worries about adverse experiences were measured at the same time as the adversity measures and selected to correspond with these variables. Each category of worry was operationalised as binary (absent vs present): financial worries, COVID-19 illness, social and relationship worries, concerns about safety and security, and worries about accessing essentials. See the Supplementary Materials for further description of these measures.

## Statistical Analysis

First, we describe weekly patterns in our outcome, adversity, and worries about adversity variables from 1 April 2020 through 7 February 2021. We then use fixed-effects regression to analyse the time-varying associations between changes in both experiences of and worries about adversity with changes in self-harm thoughts and behaviours across these 45 weeks. In this approach, individuals serve as their own reference point, which accounts for any confounding associations between time-invariant (stable) covariates such as socio-economic status, genetics, personality, and history of mental illness between predictors and outcomes (29). Our analyses consisted of regressing each outcome measure on i) the total number of adversity experiences and adversity worries jointly and ii) individual categories of adversity and worry about adversity for the total sample then stratified by age. All regression models adjusted for day of week (categorical) and days since lockdown commenced (continuous). Resulting regression coefficients were exponentiated and presented as odds ratios along with 95% confidence intervals. See Supplementary Materials for more detail, including model equation.

Sensitivity analyses included: models which included continuous measures of i) weekly depression symptoms, ii) weekly anxiety symptoms, and iii) the physical/psychological abuse variable separated into

physical abuse and psychological abuse. To increase representativeness of the UK general population, data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics (30). Analyses were conducted using Stata version 16 (31).

## Results

### Sample characteristics

The average proportion of the sample reporting self-harm thoughts, self-harm behaviours, adversities, and worries about adversities over the course of the first 45 weeks of the COVID-19 pandemic are presented in Figure 1. Self-harm thoughts and self-harm behaviours were relatively stable over the first 21 weeks, but fluctuations were seen around the end of August to the beginning of November 2020 in self-harm thoughts and the end of August to the first week of October 2020 for self-harm behaviours (Supplemental Figure S1). The total number of worries about adversity were consistently about three times higher than actual adversity experiences across the 45 weeks, and fluctuations in both corresponded to the pattern of variation in self-harm thoughts and behaviours.

Nearly a quarter (24.6%) of respondents in the total sample reported having self-harm thoughts over the first 45 weeks of the pandemic (Supplemental Table S1), and nearly one in ten (7%) had self-harmed at least once (Supplemental Table S2). There was within-individual variation in self-harm thoughts and behaviours outcome measures in 10,456 and 3,360 individuals (Supplemental Table S3), respectively, suggesting fixed effects was a valid approach. Socio-demographic characteristics of these two groups are presented in Supplemental Table S4.

### Associations between total number of adversities and worries with self-harm thoughts and behaviours

Each additional adverse event experienced was associated with 1.60 (95% CI = 1.55 to 1.64) times higher odds of self-harm thoughts (Figure S2 & Table 1), whilst each additional adversity was associated with a nearly two-fold (OR = 1.84; 95% CI = 1.76 to 1.93) increased likelihood of self-harm behaviours (Figure S3 & Table 2). Increased likelihood of both outcomes was smaller in magnitude for the total number of worries about adversity (self-harm thoughts: OR = 1.26; 95% CI = 1.26 to 1.31; self-harm behaviours: OR = 1.15; 95% CI = 1.11 to 1.18) than actual adversity experiences. This also applied to age-stratified analyses, where the total number of adversity experiences (self-harm thoughts OR range = 1.50 to 1.65; self-harm behaviours OR range = 1.67 to 2.11) were more strongly associated with both outcomes than adversity worries in each of the four age groups (self-harm thoughts OR range = 1.19 to 1.36; self-harm behaviours OR range = 1.08 to 1.23).

### Associations between individual categories of adversities and worries with self-harm thoughts and behaviours

When examining individual categories of adversities and adversity worries, four of the five categories were associated with increased likelihood of outcomes in the total sample (Figures S4 and S5). The exception was COVID-19 illness; increases in concerns about becoming seriously ill with COVID-19 were not associated with changes in either outcome (Tables 3 and 4). However, having already had COVID-19 associated with increased odds of each outcome (self-harm thoughts: OR = 1.21; 95% CI = 1.12 to 1.31; self-harm behaviours: OR = 1.25; 95% CI = 1.09 to 1.43). The largest increases in the likelihood of both outcomes were for experiences of physical or psychological abuse (self-harm thoughts: odds ratio (OR) = 3.61; 95% confidence interval (CI) = 3.41 to 3.83; self-harm behaviours: OR = 5.12; 95% CI = 4.70 to 5.59). These associations were larger than those for worries about personal safety with outcomes (self-harm thoughts: OR = 1.43; 95% CI = 1.36 to 1.50; self-harm behaviours: OR = 1.25; 95% CI = 1.15 to 1.35). Similarly, odds of increase in both outcomes were higher for worries about financial

adversity, compared to actually experiencing financial adversity, the latter of which was not associated with change in self-harm behaviours (OR = 1.02; 95% CI = 0.93 to 1.12). Concerns about social relationships also impacted increases in both outcomes, more so for self-harm thoughts (OR = 1.62; 95% CI = 1.55 to 1.69) than for self-harm behaviours (OR = 1.23; 95% CI = 1.15 to 1.35).

There was heterogeneity in the associations of adversities and worries with outcomes when models were stratified by age. The largest associations with both outcomes were found for experiencing physical or psychological abuse (self-harm thoughts: OR ranges = 3.38 to 4.00; self-harm behaviours: OR ranges = 3.72 to 6.82) in all age groups. In the two younger age groups (18-29 and 30-44), social and relationship concerns had the second strongest associations with self-harm thoughts (OR range = 1.64 to 1.87), after experiencing physical/psychological abuse. In the oldest two age groups, however, the second strongest associations with self-harm thoughts after physical/psychological abuse were for not being able to access essential items (OR range = 1.50 to 1.67).

In the older age categories, the second strongest associations with self-harm behaviours were seen for threats to personal safety in the 30-44 age group (OR = 1.72; 95% CI = 1.44 to 2.05), having had COVID-19 in adults ages 45-59 (OR = 1.72; 95% CI = 1.37 to 2.15), and not having been able to access essential items in the oldest age group (OR = 1.68; 95% CI = 1.31 to 2.14). As in the total sample, concerns about financial adversity were more strongly related to both outcomes than actual financial adversity experiences across age groups, and for self-harm behaviours, there was no evidence for a relationship of the latter with self-harm behaviours.

### Sensitivity analyses

When accounting for anxiety and depression symptoms within models, results were largely similar (Supplemental Tables S5-S8). Analyses examining physical abuse and psychological abuse as individual adversity experiences showed different patterns of association for each abuse type with outcomes (Supplemental Tables S9 and S10).

## Discussion

Both experiencing adversities and worrying about adversities were associated with an increased likelihood of self-harm thoughts and actually engaging in self-harm behaviours during the first 45 weeks of the COVID-19 pandemic. These results were found across age groups, with strongest associations for the total number of adversity experiences compared to the number of worries.

The largest predictor by far of both thinking about and engaging in self-harm was experiencing physical or psychological abuse, and this finding was consistent across all four age groups examined. Sensitivity analyses suggested that physical abuse is making larger contributions than psychological abuse to self-harm behaviours, whilst the size of the associations of both abuse types with self-harm thoughts were similar. A range of literature outside of pandemic circumstances shows that different forms of abuse including domestic violence are predictors of self-harm behaviours and suicide (32–34). That increases in domestic abuse would occur during stay-at-home orders was anticipated early in the pandemic (35,36), and has been demonstrated in countries internationally. For example, the number of calls to emergency domestic abuse hotlines in the European Union had increased 60% by the end of April 2020 (37), whilst intimate partner violence against women increased 23% over the first three months of the first lockdown in Spain (38).

Worsening economic circumstances have been identified as one of the causes of this increase in domestic abuse (38), and financial stress was another predictor of self-harm thoughts and behaviours identified by our study. Notably, it was worrying about financial adversity such as losing one's job rather than actually

experiencing such an adversity that was consistently associated with self-harm thoughts and behaviours across age groups. This suggests that, thus far, economic uncertainty rather than actual adversities are negatively impacting people's mental health, and it is possible that these associations may change in magnitude as the economic consequences of the pandemic and recession unfold. Considerable evidence indicates that economic recessions are associated with increases in rates of self-harm and suicide, particularly in the working-age population (13). Risk for both attempted suicide and suicide death is higher for those unemployed over the long-term compared to shorter term unemployment (39,40). However, an increase in self-harm and suicides during economic recessions is not inevitable (1).

Lessons learnt from prior economic recessions suggest multiple opportunities for how governments can respond with policies to mitigate the mental health impact of the upcoming recession (9). The increases in suicides that correspond to unemployment rates are not uniform across all countries but are instead modified by differential investment in social programmes to mitigate these effects (41). For example, following the 2008 financial crisis, an increase of 1% per capita in government spending designed to mitigate the effect of financial hardship was associated with a 0.2% decrease in suicide in Japan (42). In the three decades leading up to the 2008 recession, every \$10 USD invested per person on programmes aimed to increase chances of gainful employment resulted in a decrease of the effect of unemployment on suicides decreased by 0.04% in EU countries (41). Thus, our findings highlight the potential danger of the economic impacts of COVID-19 on self-harm thoughts and behaviours and, as self-harm is an important risk factor for suicide, potentially for suicide too, and suggest the importance of addressing economic concerns amongst individuals urgently.

Whilst many of our findings were consistent across age groups, there were some discrepancies. For example, worrying about catching COVID-19 was associated with reduced likelihood of having self-harm thoughts and self-harm behaviours in adults aged 30 to 44, but this pattern was reversed in the oldest age group (60 years plus), where worries about falling ill were associated with increased likelihood of thoughts of harming themselves or that they would be better off dead (self-harm thoughts). This could have been influenced by public health messaging highlighting that older adults are at higher risk than younger adults for dying of the illness; a death that news coverage often portrayed as occurring alone and without the ability to say goodbye to loved ones. In contrast, having already had the illness was related to increased likelihood of both outcomes in the total sample, and this remained true in young (18-29) and middle aged (45-59) adults for self-harm thoughts and for self-harm behaviours. It is therefore possible that the disease itself is causing people to increase their self-harm thoughts and behaviours, while those who are particularly worried about falling ill with COVID-19 are more protective of their health and less likely to want to harm themselves. Evidence for the former, that COVID-19 illness leads to increased risk for mental health problems such as depression, self-harm thoughts and self-harm behaviours has been documented across a range of studies (18,43–45).

This study has a number of strengths including the use of a large, well-stratified sample on socio-demographic groups which were weighted on the basis of population estimates of core demographics, and its longitudinal follow-up with repeated assessments of adversities, worries and self-harm thoughts and behaviours. We also used robust statistical methods to account for unobserved stable participant characteristics. However, sampling was not random and the data are therefore not representative of the general UK population. It is possible that individuals who were experiencing greater adversity and were more likely to have self-harm thoughts and behaviours were more likely to participate in the study. Nevertheless, this study did not aim to report prevalence of such experiences, but rather to identify the time-varying relationship between exposures and self-harm thoughts and behaviours. Crucially, the sample was heterogeneous and maintained its heterogeneity over time. There may have been other relevant forms of adversity and worry not captured in the current study, which may have resulted in an overestimation of the adversities and worries we did include. Finally, fixed effects regression does not address direction of causality. Arellano Bond models can be used to follow up fixed effects models to

account for this lack of directionality (46), but because linear estimation is used, we could not utilise this approach as our outcomes were binary. However, whilst the relationship between worries and self-harm thoughts and behaviours may have involved some bidirectionality, there is little evidence to suggest that self-harm thoughts or behaviours increase the likelihood of individuals experiencing adversities.

Across the COVID-19 pandemic, there are concerns about potential future increases in suicide levels. Self-harm thoughts and behaviours are important and strong predictors of future suicide risk, so identifying modifiable risk factors for self-harm that can be addressed through public health interventions during the COVID-19 pandemic and beyond is vital (1,9). Our findings suggest that increases in self-harm thoughts and behaviours across the first 45 weeks of the pandemic were related to financial uncertainty, physical or psychological abuse, concern for others, not being able to access essential items, and worries about one's personal safety. Suggestions have already been made for how to adapt evidence-based suicide prevention strategies to current the pandemic (Hawton et al., 2021; Moutier, 2020; Niederkrotenthaler et al., 2020). For example, it has been recommended that universal interventions to mitigate the impact of poverty and unemployment on suicide risk should be implemented (9). Our data suggest the importance of following such strategies to try and reduce self-harm thoughts and behaviours that have the potential to drive increasing suicide rates of the coming months. The findings here also suggest the need for ongoing surveillance of how these well-established risk factors for suicide and self-harm may be exacerbated by the upcoming recession and as public health measures such as social distancing continue.

Author details: Elise Paul and Daisy Fancourt are with the Department of Behavioural Science and Health, University College London, 1-19 Torrington Place, London, WC1E 7HB, United Kingdom. The corresponding author is Elise Paul ([e.paul@ucl.ac.uk](mailto:e.paul@ucl.ac.uk)).

Declaration of interests: none.

Funding: This work was supported by the Nuffield Foundation [DF, WEL/FR-000022583], the MARCH Mental Health Network funded by the Cross-Disciplinary Mental Health Network Plus initiative supported by UK Research and Innovation [DF, ES/S002588/1], and the Wellcome Trust [DF, 221400/Z/20/Z and DF, 205407/Z/16/Z].

Acknowledgements: The researchers are grateful for the support of a number of organisations with their recruitment efforts including: the UKRI Mental Health Networks, Find Out Now, UCL BioResource, SEO Works, FieldworkHub, and Optimal Workshop.

Author contribution: DF and EP conceptualised and designed the study. DF acquired funding, led the investigation, provided oversight on the methodology, provided software, and supervised the project. Data were curated, validated, and formally analysed by EP. EP created visualisations, wrote the original manuscript draft with input from all authors, who then reviewed and edited the manuscript. All authors approved the final version of the manuscript and had full access to and verified the data.

Data availability: The COVID-19 Social Study documentation and codebook are available for download at <https://www.covidsocialstudy.org/>. Statistical code is available upon request from Elise Paul ([e.paul@ucl.ac.uk](mailto:e.paul@ucl.ac.uk)).

## References

1. Gunnell D, Appleby L, Arensman E, Hawton K, John A, Kapur N, et al. Suicide risk and prevention during the COVID-19 pandemic. *Lancet Psychiatry*. 2020;7(6):468–71.
2. Moutier C. Suicide Prevention in the COVID-19 Era: Transforming Threat Into Opportunity. *JAMA Psychiatry*. 2020;
3. John A, Pirkis J, Gunnell D, Appleby L, Morrissey J. Trends in suicide during the covid-19 pandemic. *BMJ*. 2020 Nov 12;m4352.
4. Leske S, Kølves K, Crompton D, Arensman E, De Leo D. Real-time suicide mortality data from police reports in Queensland, Australia, during the COVID-19 pandemic: an interrupted time-series analysis. *Lancet Psychiatry*. 2020;8(1):58–63.
5. Qin P, Mehlum L. National observation of death by suicide in the first 3 months under COVID-19 pandemic. *Acta Psychiatr Scand*. 2021;143(1):92–3.
6. Kølves K, Kølves KE, De Leo D. Natural disasters and suicidal behaviours: a systematic literature review. *J Affect Disord*. 2013;146(1):1–14.
7. Lester D. Suicide during war and genocide. In: Wasserman D, Wasserman C, editors. *Oxford textbook of suicidology and suicide prevention*. Oxford: Oxford University Press; 2009. p. 215–8.
8. Tanaka T, Okamoto S. Increase in suicide following an initial decline during the COVID-19 pandemic in Japan | *Nature Human Behaviour*. *Nat Hum Behav* [Internet]. 2021 [cited 2021 Feb 4]; Available from: <https://www.nature.com/articles/s41562-020-01042-z>
9. Wasserman D, Iosue M, Wuestefeld A, Carli V. Adaptation of evidence-based suicide prevention strategies during and after the COVID-19 pandemic. *World Psychiatry*. 2020;19(3):294–306.
10. International Monetary Fund. *World Economic Outlook, April 2020: The Great Lockdown* [Internet]. 2020 [cited 2021 Jan 12]. Available from: <https://www.imf.org/en/Publications/WEO/Issues/2020/04/14/weo-april-2020>
11. Oyesanya M, Lopez-Morinigo J, Dutta R. Systematic review of suicide in economic recession. *World J Psychiatry*. 2015;5(2):243.
12. Parmar D, Stavropoulou C, Ioannidis JP. Health outcomes during the 2008 financial crisis in Europe: systematic literature review. *Br Med J*. 2016;354.
13. Haw C, Hawton K, Gunnell D, Platt S. Economic recession and suicidal behaviour: Possible mechanisms and ameliorating factors. *Int J Soc Psychiatry*. 2015;61(1):73–81.
14. Turecki G, Brent DA, Gunnell D, O'Connor RC, Oquendo MA, Pirkis J, et al. Suicide and suicide risk. *Nat Rev Dis Primer*. 2019 Oct 24;5(1):1–22.
15. Geulayov G, Casey D, Bale L, Brand F, Clements C, Farooq B, et al. Suicide following presentation to hospital for non-fatal self-harm in the Multicentre Study of Self-harm: a long-term follow-up study. *Lancet Psychiatry*. 2019 Dec 1;6(12):1021–30.

16. Hawton K, Bergen H, Kapur N, Cooper J, Steeg S, Ness J, et al. Repetition of self-harm and suicide following self-harm in children and adolescents: Findings from the Multicentre Study of Self-harm in England. *J Child Psychol Psychiatry*. 2012;53(12):1212–9.
17. Owens D, Horrocks J, House A. Fatal and non-fatal repetition of self-harm: Systematic review. *Br J Psychiatry*. 2002 Sep;181(3):193–9.
18. Iob E, Steptoe A, Fancourt D. Abuse, self-harm and suicidal ideation in the UK during the COVID-19 pandemic. *Br J Psychiatry*. 2020;217(4):543–6.
19. O'Connor RC, Wetherall K, Cleare S, McClelland H, Melson AJ, Niedzwiedz CL, et al. Mental health and wellbeing during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 Mental Health & Wellbeing study. *Br J Psychiatry*. 2020;1–17.
20. Centers for Disease Control and Prevention (CDC). Mental Health, Substance Use, and Suicidal Ideation During the COVID-19 Pandemic — United States, June 24–30, 2020 [Internet]. National Center for Injury Prevention and Control; 2020 [cited 2021 Jan 7]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7440121/>
21. Killgore WD, Cloonan SA, Taylor EC, Allbright MC, Dailey NS. Trends in suicidal ideation over the first three months of COVID-19 lockdowns. *Psychiatry Res*. 2020;293:113390.
22. Carr MJ, Steeg S, Webb RT, Kapur N, Chew-Graham C, Abel K, et al. Primary Care Contact for Mental Illness and Self-Harm Before, During and After the Peak of the COVID-19 Pandemic in the UK: Cohort Study of 13 Million Individuals. *Peak COVID-19 Pandemic UK Cohort Study Of*. 2020;13.
23. Pignon B, Gourevitch R, Tebeka S, Dubertret C, Cardot H, Dauriac-Le Masson V, et al. Dramatic reduction of psychiatric emergency consultations during lockdown linked to COVID-19 in Paris and suburbs. *medRxiv*. 2020;
24. Zortea TC, Brenna CTA, Joyce M, McClelland H, Tippett M, Tran MM, et al. The Impact of Infectious Disease-Related Public Health Emergencies on Suicide, Suicidal Behavior, and Suicidal Thoughts. *Crisis*. 2020 Oct 16;1–14.
25. Geulayov G, Casey D, McDonald KC, Foster P, Pritchard K, Wells C, et al. Incidence of suicide, hospital-presenting non-fatal self-harm, and community-occurring non-fatal self-harm in adolescents in England (the iceberg model of self-harm): a retrospective study. *Lancet Psychiatry*. 2018 Feb 1;5(2):167–74.
26. Bryan C, Bryan AO, Baker JC. Associations among state-level physical distancing measures and suicidal thoughts and behaviors among US adults during the early COVID-19 pandemic. 2020;
27. Hawton K, Lascelles K, Brand F, Casey D, Bale E, Ness J, et al. Self-harm and the COVID-19 pandemic: a study of factors contributing to self-harm during lockdown restrictions. *medRxiv*. 2020;
28. Löwe B, Kroenke K, Herzog W, Gräfe K. Measuring depression outcome with a brief self-report instrument: sensitivity to change of the Patient Health Questionnaire (PHQ-9). *J Affect Disord*. 2004;81(1):61–6.

29. Allison PD. Fixed effects regression models. SAGE publications; 2009.
30. Office for National Statistics. Population estimates for the UK, England and Wales, Scotland and Northern Ireland [Internet]. 2020 May [cited 2020 Sep 30]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2018>
31. StataCorp. Stata Statistical Software: Release 16. College Station, TX: StataCorp LP; 2019.
32. Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. 2013;10(5):e1001439.
33. Zatti C, Rosa V, Barros A, Valdivia L, Calegario VC, Freitas LH, et al. Childhood trauma and suicide attempt: A meta-analysis of longitudinal studies from the last decade. *Psychiatry Res*. 2017;256:353–8.
34. Rodway C, Tham S-G, Ibrahim S, Turnbull P, Windfuhr K, Shaw J, et al. Suicide in children and young people in England: a consecutive case series. *Lancet Psychiatry*. 2016 Aug 1;3(8):751–9.
35. Bradbury-Jones C, Isham L. The pandemic paradox: The consequences of COVID-19 on domestic violence. *J Clin Nurs*. 2020;
36. Usher K, Bhullar N, Durkin J, Gyamfi N, Jackson D. Family violence and COVID-19: Increased vulnerability and reduced options for support. *Int J Ment Health Nurs*. 2020;29:549–52.
37. Mahase E. Covid-19: EU states report 60% rise in emergency calls about domestic violence - ProQuest. *Br Med J Online [Internet]*. 2020 [cited 2021 Feb 4];369. Available from: <https://search.proquest.com/openview/11255fd2e5ba902a8cb651a20eeb035f/1?pq-origsite=gscholar&cbl=2043523>
38. Arenas-Arroyo E, Fernandez-Kranz D, Nollenberger N. Intimate partner violence under forced cohabitation and economic stress: Evidence from the COVID-19 pandemic. *J Public Econ*. 2021 Feb 1;194:104350.
39. Mäki N, Martikainen P. A register-based study on excess suicide mortality among unemployed men and women during different levels of unemployment in Finland. *J Epidemiol Community Health*. 2012;66(4):302–7.
40. Platt S, Hawton K. Suicidal Behaviour and the Labour Market. In: Hawton K, van Heeringen K, editors. *The International Handbook of Suicide and Attempted Suicide*. Wiley Online Library; 2000. p. 309–84.
41. Stuckler D, Basu S, Suhrcke M, Coutts A, McKee M. The public health effect of economic crises and alternative policy responses in Europe: an empirical analysis. *The Lancet*. 2009 Jul 25;374(9686):315–23.
42. Matsubayashi T, Sekijima K, Ueda M. Government spending, recession, and suicide: evidence from Japan. *BMC Public Health*. 2020;20(1):1–8.

43. Hossain MM, Tasnim S, Sultana A, Faizah F, Mazumder H, Zou L, et al. Epidemiology of mental health problems in COVID-19: a review. F1000Research [Internet]. 2020 Jun 23 [cited 2021 Feb 14];9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7549174/>
44. Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: Systematic review of the current evidence. *Brain Behav Immun*. 2020 Oct 1;89:531–42.
45. Wright L, Steptoe A, Fancourt D. How are adversities during COVID-19 affecting mental health? Differential associations for worries and experiences and implications for policy. medRxiv. 2020;
46. Arellano M, Bond S. Some tests of specification for panel data: Monte Carlo evidence and an application to employment equations. *Rev Econ Stud*. 1991;58(2):277–97.
47. Hawton K, Marzano L, Fraser L, Hawley M, Harris-Skillman E, Lainez YX. Reporting on suicidal behaviour and COVID-19—need for caution. *Lancet Psychiatry*. 2021 Jan 1;8(1):15–7.
48. Niederkrotenthaler T, Gunnell D, Arensman E, Pirkis J, Appleby L, Hawton K, et al. Suicide research, prevention, and COVID-19. Hogrefe Publishing; 2020.




Figure 1. Average number of adversities and worries and proportion with self-harm thoughts and self-harm behaviours over time for the duration of the study period (1 April 2020 to 7 February 2021). Increased variability in data starting in August may be due to the change in data collection frequency from weekly to monthly that occurred at this time.

Table 1. Fixed-effects logistic regression models predicting within-individual change in self-harm thoughts from the total number of adversity experiences and adversity worries

Variable	Self-harm thoughts														
	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Total number of adversity experiences	1.60	1.55	1.64	1.65	1.54	1.78	1.50	1.42	1.59	1.62	1.55	1.71	1.62	1.53	1.71
Total number of worries	1.26	1.26	1.31	1.36	1.30	1.42	1.19	1.15	1.24	1.27	1.23	1.31	1.33	1.29	1.38
Number of observations	169,954			14,286			44,659			61,051			49,958		
Number of individuals	10,456			1,081			3,055			3,664			2,656		

Note. Adversity experiences (0-5) and worries (0-5) variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table 2. Fixed-effects logistic regression models predicting within-individual change in self-harm behaviours from the total number of adversity experiences and adversity worries

Self-harm behaviours															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Total number of adversity experiences	1.84	1.76	1.93	1.67	1.52	1.84	1.70	1.54	1.88	1.88	1.74	2.04	2.11	1.91	2.32
Total number of worries	1.15	1.11	1.18	1.23	1.16	1.31	1.08	1.01	1.15	1.17	1.10	1.24	1.10	1.03	1.18
Number of observations	52,074			6,463			15,277			19,294			11,040		
Number of individuals	3,360			475			1,085			1,184			616		

Note. Adversity experiences (0-5) and worries (0-5) variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table 3. Fixed-effects logistic regression models predicting within-individual change in self-harm thoughts from individual categories of adversity experiences and worries

Self-harm thoughts															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Adversity experiences															
Financial	1.18	1.12	1.25	1.24	1.09	1.42	1.27	1.13	1.42	1.09	0.99	1.20	1.20	1.06	1.35
Accessing essentials	1.35	1.25	1.47	1.17	0.94	1.45	0.99	0.85	1.16	1.67	1.46	1.93	1.50	1.30	1.73
COVID-19 illness	1.21	1.12	1.31	1.32	1.09	1.61	1.08	0.93	1.26	1.35	1.18	1.54	1.11	0.96	1.30
Illness of others/bereavement	1.15	1.08	1.23	1.42	1.20	1.69	1.10	0.96	1.27	1.20	1.07	1.35	1.03	0.92	1.17
Physical/psychological abuse	3.61	3.41	3.83	4.00	3.38	4.75	3.92	3.45	4.46	3.56	3.22	3.95	3.38	3.05	3.75
Worries															
Financial	1.43	1.36	1.49	1.34	1.21	1.49	1.35	1.23	1.48	1.61	1.47	1.75	1.37	1.25	1.50
Accessing essentials	1.19	1.14	1.25	1.47	1.30	1.65	1.12	1.03	1.23	1.00	0.92	1.09	1.32	1.22	1.44
COVID-19 illness	0.95	0.91	1.00	0.91	0.82	1.01	0.79	0.72	0.86	0.94	0.86	1.02	1.18	1.08	1.29
Social/ relationship concerns	1.62	1.55	1.69	1.87	1.67	2.09	1.64	1.49	1.79	1.66	1.53	1.81	1.42	1.30	1.55
Threats to safety	1.43	1.36	1.50	1.68	1.49	1.89	1.40	1.27	1.54	1.39	1.27	1.52	1.38	1.27	1.51
Number of observations	169,954			14,286			44,659			61,051			49,958		
Number of individuals	10,456			1,081			3,055			3,664			2,656		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table 4. Fixed-effects logistic regression models predicting within-individual change in self-harm behaviours from individual categories of adversity experiences and worries

Variable	Self-harm behaviours														
	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Adversity experiences															
Financial	1.02	0.93	1.12	1.04	0.85	1.26	0.81	0.65	0.99	1.14	0.96	1.34	1.09	0.87	1.37
Accessing essentials	1.30	1.16	1.45	1.24	0.99	1.56	0.92	0.72	1.19	1.46	1.19	1.79	1.68	1.31	2.14
COVID-19 illness	1.25	1.09	1.43	1.53	1.08	2.17	0.93	0.71	1.22	1.72	1.37	2.15	0.91	0.67	1.24
Illness of others/ bereavement	1.24	1.11	1.38	1.44	1.15	1.79	1.20	0.92	1.56	1.15	0.95	1.39	1.17	0.93	1.48
Physical/ psychological abuse	5.12	4.70	5.59	3.72	3.04	4.55	6.82	5.63	8.27	4.37	3.77	5.07	6.35	5.35	7.53
Worries															
Financial	1.25	1.15	1.36	1.20	1.02	1.41	1.23	1.03	1.46	1.53	1.31	1.78	1.04	0.86	1.25
Accessing essentials	1.18	1.09	1.27	1.36	1.16	1.59	1.13	0.95	1.34	1.14	0.99	1.31	1.10	0.93	1.30
COVID-19 illness	0.93	0.86	1.01	1.01	0.85	1.20	0.76	0.64	0.91	0.95	0.82	1.10	1.08	0.91	1.30
Social/relationship concerns	1.23	1.13	1.34	1.60	1.34	1.90	1.08	0.91	1.29	1.20	1.03	1.40	1.10	0.91	1.33
Threats to safety	1.25	1.15	1.35	1.07	0.90	1.27	1.72	1.44	2.05	1.19	1.03	1.37	1.17	0.99	1.39
Number of observations	52,074			6,463			15,277			19,294			11,040		
Number of individuals	3,360			475			1,085			1,184			616		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

## Supplementary Materials

### Sampling methods

Three primary recruitment approaches were used. First, convenience sampling was used, including promoting the study through existing networks and mailing lists (including large databases of adults who had previously consented to be involved in health research across the UK), print and digital media coverage, and social media. Second, more targeted recruitment was undertaken focusing on groups who were anticipated to be less likely to take part in the research via our first strategy, including (i) individuals from a low-income background, (ii) individuals with no or few educational qualifications, and (iii) individuals who were unemployed. Third, the study was promoted via partnerships with third sector organisations to vulnerable groups, including adults with pre-existing mental health conditions, older adults, carers, and people experiencing domestic violence or abuse. Recruitment was refreshed in August when participants who were lost-to-follow-up were recontacted. The study was approved by the UCL Research Ethics Committee [12467/005] and all participants gave informed consent. Participants were not compensated for participation.

### Outcomes

#### Self-harm thoughts and behaviours

Two outcome variables were assessed weekly for the first 22 weeks of the study (1 April to 21 August 2020) and then monthly to 7 February 2021. Thoughts of death or self-harm (“self-harm thoughts”) were measured with an item from the Patient Health Questionnaire (PHQ-9) (1), an instrument often used as a diagnostic tool for depression in primary care practice: “Over the last week, how often have you been bothered by: Thoughts that you would be better off dead or hurting yourself in some way?”. Second, self-harm behaviours were measured with a similar study-developed item: “Over the last week, how often have you been bothered by: Self-harming or deliberately hurting yourself?”. Responses to both items were rated on a four-point scale from “not at all” to “nearly every day”. Analyses focused on binary variables indicating the presence of at least some self-harm thoughts and self-harm behaviours at each time point.

### Exposure variables

#### Adversities

Financial adversity consisted of: (i) loss of job/been unable to do paid work; (ii) spouse/partner lost their job/was unable to do paid work; (iii) major cut in household income (e.g., due to you or your partner being furloughed/put on leave/ not receiving sufficient work); (iv) unable to pay bills/rent/mortgage; and (v) evicted/lost accommodation. Illness with COVID-19 was measured with an item asking if participants had either suspected or been diagnosed with the illness. Family/friend illness or bereavement was assessed as: (i) the participant having someone close to them ill in hospital (due to COVID-19 or another cause), and (ii) lost somebody close (due to COVID-19 or another cause). Participants indicated whether they had been “physically harmed or hurt by someone else” or “bullied, controlled, intimidated, or psychologically hurt by someone else” during the last week. Responses were rated on a four-point scale ranging from “not at all” to “nearly every day”. A response to either item indicating physical or psychological abuse was categorised as present. Finally, inability to access essential items was measured by asking if participants had been unable to access sufficient food and required medication. Total weekly adversities scores were calculated by summing each of the five binary variables (range 0-5).

#### Worries about adversity

Two questions asked participants to select which of a list of items had caused them (i) stress (however minor) in the past week, and (ii) significant stress (something being constantly on their mind or keeping them awake at night in the past week). Financial worries was measured with three items: (i)

losing your job/unemployment, (ii) finances, and (iii) work, COVID-19 illness with two: (i) catching COVID-19 and (ii) becoming seriously ill from COVID-19, social and relationship worries with five: (i) marriage or other romantic relationship, (ii) friends or family living in your household, (iii) friends or family living outside your household, (iv) neighbours, and (v) your pet), safety and security concerns with one: your own safety/security', and worries about access essentials with two (i) food and (ii) medication. Total weekly worries scores were computed by summing these five binary variables (range 0-5).

## Statistical Analysis

The basic model can be expressed as follows:

$$\text{Outcome}_{it} = \beta_0 + \beta_1 E_{ikt} + \beta_2 W_{ikt} + \beta_3 D_t + \beta_4 N_t + \alpha_i + \varepsilon_{it}$$

where  $\text{Outcome}_{it}$  is a measure of individual  $i$ 's self-harm thoughts or self-harm behaviours at time  $t$ ,  $E$  is whether an individual  $i$  was experiencing adversity  $k$  at time  $t$ ,  $W$  is whether an individual  $i$  was worrying about adversity  $k$  at time  $t$ ,  $D_t$  is a vector of indicator variables for day,  $N_t$  is a continuous variable for days since lockdown, is unobserved time invariant confounding factors, and  $\varepsilon$  is error.

Three sets of sensitivity analyses were conducted to ensure the robustness of our results. First, models with individual categories of adversities and worries were re-run which included continuous measures of weekly depression symptoms assessed with the PHQ-9 (1), without the inclusion of the self-harm thoughts item. Second, these models were estimated again, but this time with continuous measures of weekly anxiety symptoms using the Generalized Anxiety Disorder (GAD-7) scale (2). Third, we repeated our fixed effects models using individual categories of worry and adversity with the physical/psychological abuse variable separated into two separate variables: physical abuse and psychological abuse.

To account for the non-random nature of the sample and increase representativeness of the UK general population, all data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics (3). Weights were constructed using a multivariate reweighting method using the Stata user written command 'ebalance' (4). Analyses were conducted using Stata version 16 (5) and fixed effects model results plotted with the user-written command 'coefplot' (6).

## Results

### Sensitivity analyses

When accounting for anxiety and depression symptoms within models, results were largely similar. But financial adversities were no longer associated with increased risk for either outcome in any age group and were related to decreased likelihood of outcomes in some age groups (Supplemental Tables S8-S11). Worries about financial adversity, however, remained associated with increased likelihood of both outcomes in the total sample and across some age groups. Analyses examining physical abuse and psychological abuse as individual adversity experiences showed different patterns of association for each abuse type with outcomes (Supplemental Tables S12 and S13). Whilst physical abuse (OR range = 6.04 to 18.23) had much larger associations than psychological abuse (OR range = 1.83 to 2.83) with self-harm behaviours in all age groups, the associations for these two abuse types were more similar in magnitude across age groups for self-harm thoughts.

## References

1. Löwe B, Kroenke K, Herzog W, Gräfe K. Measuring depression outcome with a brief self-report instrument: sensitivity to change of the Patient Health Questionnaire (PHQ-9). *J Affect Disord.* 2004;81(1):61–6.
2. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166(10):1092–7.
3. Office for National Statistics. Population estimates for the UK, England and Wales, Scotland and Northern Ireland [Internet]. 2020 May [cited 2020 Sep 30]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2018>
4. Hainmueller J, Xu Y. Ebalance: A Stata package for entropy balancing. *J Stat Softw.* 2013;54(7).
5. StataCorp. Stata Statistical Software: Release 16. College Station, TX: StataCorp LP; 2019.
6. Jann B. Plotting Regression Coefficients and other Estimates - Ben Jann, 2014. Stata J [Internet]. 2014 [cited 2021 Feb 3]; Available from: <https://journals.sagepub.com/doi/10.1177/1536867X1401400402>




Figure S1. Average proportion of the sample with self-harm thoughts and self-harm behaviours over time for the duration of the study period (1 April 2020 to 7 February 2021). Increased variability in data starting in August may be due to the change in data collection frequency from weekly to monthly that occurred at this time.




Figure S2. Associations between the total number adversity experiences and worries about adversity and change over time in self-harm thoughts derived from fixed effects models. Experiences and worries were entered simultaneously into the same model. Analyses were further adjusted for day of the week and time since lockdown began.




Figure S3. Associations between the total number adversity experiences and worries about adversity and change over time in self-harm behaviours derived from fixed effects models stratified by age. Experiences and worries were entered simultaneously into the same model. Analyses were further adjusted for day of the week and time since lockdown began.




Figure S4. Associations between individual adversity experiences and worries about adversity and change over time in self-harm thoughts derived from fixed effects models. Experiences and worries were entered simultaneously into the same model. Analyses were further adjusted for day of the week and time since lockdown began.




Figure S5. Associations between individual adversity experiences and worries about adversity and change over time in self-harm behaviours derived from fixed effects models. Experiences and worries were entered simultaneously into the same model. Analyses were further adjusted for day of the week and time since lockdown began.

Table S1. Descriptive statistics by number of times self-harm thoughts were reported in total sample (N = 48,446), weighted

		Self-harm thoughts					
		Never n = 36,519 (75.4%)		Once or twice n = 5,342 (11.0%)		Thrice or more n = 6,585 (13.6%)	
Variable		M	SD	M	SD	M	SD
Gender (ref male)	Female	0.49	0.50	0.56	0.50	0.53	0.50
Age	18-29	0.07	0.25	0.13	0.34	0.16	0.37
	30-44	0.14	0.35	0.23	0.42	0.22	0.41
	45-59	0.27	0.44	0.30	0.46	0.32	0.46
	60+	0.52	0.50	0.34	0.47	0.30	0.46
Ethnicity (ref white)	Ethnic minority	0.07	0.25	0.11	0.31	0.10	0.30
Education	Up to GCSE	0.33	0.47	0.30	0.46	0.31	0.46
	A-levels or vocational	0.31	0.46	0.33	0.47	0.35	0.48
	Undergraduate	0.22	0.41	0.22	0.41	0.20	0.40
	Postgraduate	0.14	0.35	0.15	0.36	0.14	0.35
Income	£90,000+	0.08	0.27	0.07	0.26	0.05	0.21
	£60,000-90,000	0.12	0.32	0.12	0.32	0.08	0.27
	£30,000-60,000	0.34	0.47	0.29	0.45	0.29	0.45
	£16,000-30,000	0.30	0.46	0.30	0.46	0.29	0.45
	< £16,000	0.17	0.38	0.22	0.42	0.30	0.46
Living arrangement	Alone	0.20	0.40	0.22	0.42	0.27	0.44
	With others, not children	0.62	0.49	0.55	0.50	0.55	0.50
	With others, including children	0.18	0.39	0.23	0.42	0.18	0.38
Keyworker (ref not)	Keyworker	0.18	0.38	0.20	0.40	0.18	0.39
Long-term physical health condition	Present	0.45	0.50	0.47	0.50	0.53	0.50
Long-term mental health condition	Present	0.09	0.29	0.25	0.43	0.46	0.50

Note. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Ethnic minority refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education.

Table S2. Descriptive statistics by number of times self-harm behaviours were reported in total sample (N = 48,446), weighted

		Self-harm behaviours					
		Never n = 44,852 (92.6%)		Once or twice n = 2,246 (4.6%)		Thrice or more n = 1,348 (2.8%)	
Variable		M	SD	M	SD	M	SD
Gender (ref male)	Female	0.50	0.50	0.55	0.50	0.55	0.50
Age	18-29	0.08	0.27	0.18	0.39	0.25	0.43
	30-44	0.16	0.37	0.24	0.43	0.21	0.41
	45-59	0.28	0.45	0.31	0.46	0.33	0.47
	60+	0.49	0.50	0.26	0.44	0.21	0.41
	Ethnicity (ref white)	Ethnic minority	0.07	0.26	0.11	0.31	0.13
Education	Up to GCSE	0.32	0.47	0.33	0.47	0.34	0.47
	A-levels or vocational	0.32	0.47	0.35	0.48	0.36	0.48
	Undergraduate	0.22	0.41	0.18	0.39	0.18	0.39
	Postgraduate	0.14	0.35	0.13	0.34	0.12	0.32
Income	£90,000+	0.07	0.26	0.06	0.24	0.03	0.16
	£60,000-90,000	0.12	0.32	0.09	0.28	0.06	0.23
	£30,000-60,000	0.33	0.47	0.27	0.44	0.22	0.41
	£16,000-30,000	0.30	0.46	0.28	0.45	0.25	0.43
	< £16,000	0.18	0.39	0.30	0.46	0.46	0.50
Living arrangement	Alone	0.21	0.41	0.26	0.44	0.27	0.44
	With others, not including children	0.61	0.49	0.53	0.50	0.55	0.50
	With others, including children	0.18	0.39	0.21	0.41	0.18	0.39
Keyworker (ref not)	Keyworker	0.18	0.39	0.19	0.39	0.15	0.36
Long-term physical health condition	Present	0.46	0.50	0.54	0.50	0.58	0.49
Long-term mental health condition	Present	0.13	0.34	0.46	0.50	0.65	0.48

Note. Data were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Ethnic minority refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education.

Table S3. Descriptive statistics for predictors and outcome variables amongst individuals with variation in each outcome variable

		Sample with variation in self-harm thoughts				Sample with variation in self-harm behaviours			
Variable		Overall Mean	Overall SD	Between SD	Within SD	Overall Mean	Overall SD	Between SD	Within SD
Outcomes	Self-harm thoughts	0.29	0.45	0.26	0.36	0.48	0.50	0.37	0.34
	Self-harm behaviours	0.04	0.20	0.14	0.16	0.21	0.41	0.23	0.34
Adversity experiences									
	Total adversities (0-5)	0.50	0.72	0.59	0.46	0.62	0.81	0.68	0.51
	Financial	0.16	0.36	0.30	0.23	0.17	0.38	0.31	0.25
	Accessing essentials	0.04	0.21	0.15	0.16	0.07	0.26	0.20	0.20
	COVID-19 illness	0.13	0.33	0.29	0.18	0.13	0.34	0.30	0.19
	Illness of others/bereavement	0.06	0.23	0.13	0.20	0.07	0.25	0.15	0.22
	Physical/psychological abuse	0.11	0.32	0.24	0.22	0.18	0.38	0.28	0.26
Worries									
	Total worries (0-5)	2.22	1.22	0.97	0.77	2.39	1.29	1.04	0.80
	Financial	0.60	0.49	0.38	0.31	0.64	0.48	0.36	0.31
	Accessing essentials	0.19	0.40	0.27	0.31	0.26	0.44	0.31	0.33
	COVID-19 illness	0.50	0.50	0.39	0.31	0.50	0.50	0.39	0.31
	Social/relationship concerns	0.77	0.42	0.27	0.32	0.79	0.40	0.27	0.30
	Threats to safety	0.15	0.36	0.24	0.27	0.20	0.40	0.28	0.30
Number of observations		16,9954				52,074			
Number of individuals		10,456				3,360			

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Outcome variables and individual adversity experiences and worries variables are binary.

Table S4. Socio-demographic characteristics for entire sample and for those with variation in self-harm thoughts and self-harm behaviours, unweighted and weighted

Variable	Whole sample N = 48,446				Sample with variation in self-harm thoughts n = 10,456				Sample with variation in self-harm behaviours n = 3,360			
	Unweighted		Weighted		Unweighted		Weighted		Unweighted		Weighted	
	M	SD	M	SD	M	SD	M	SD	M	SD	M	SD
Female (ref male)	0.75	0.43	0.51	0.50	0.78	0.41	0.55	0.50	0.78	0.41	0.54	0.50
Age												
18-29	0.05	0.22	0.09	0.28	0.08	0.28	0.15	0.35	0.12	0.33	0.21	0.41
30-44	0.21	0.40	0.16	0.37	0.26	0.44	0.22	0.42	0.29	0.46	0.22	0.42
45-59	0.34	0.47	0.28	0.45	0.36	0.48	0.31	0.46	0.37	0.48	0.32	0.47
60+	0.40	0.49	0.47	0.50	0.29	0.46	0.33	0.47	0.21	0.41	0.25	0.43
Ethnic minority (ref white)	0.04	0.19	0.08	0.27	0.05	0.21	0.11	0.31	0.06	0.23	0.12	0.33
Education												
Up to GCSE	0.14	0.34	0.32	0.47	0.13	0.33	0.30	0.46	0.15	0.36	0.33	0.47
A-levels or vocational	0.17	0.38	0.32	0.47	0.17	0.38	0.33	0.47	0.19	0.39	0.36	0.48
Undergraduate	0.42	0.49	0.22	0.41	0.41	0.49	0.22	0.41	0.38	0.49	0.18	0.39
Postgraduate	0.27	0.45	0.14	0.35	0.29	0.45	0.15	0.36	0.27	0.45	0.13	0.34
Income												
£90,000+	0.10	0.30	0.07	0.26	0.08	0.27	0.06	0.24	0.06	0.25	0.05	0.21
£60,000-90,000	0.15	0.35	0.11	0.32	0.14	0.34	0.10	0.31	0.12	0.33	0.08	0.27
£30,000-60,000	0.35	0.48	0.32	0.47	0.34	0.47	0.30	0.46	0.31	0.46	0.25	0.43
£16,000-30,000	0.25	0.44	0.30	0.46	0.26	0.44	0.29	0.46	0.24	0.43	0.27	0.44
< £16,000	0.15	0.35	0.20	0.40	0.19	0.39	0.25	0.43	0.27	0.44	0.36	0.48
Living arrangement												
Alone	0.21	0.41	0.21	0.41	0.24	0.43	0.24	0.43	0.25	0.44	0.26	0.44
With others, not including children	0.57	0.49	0.60	0.49	0.53	0.50	0.55	0.50	0.53	0.50	0.54	0.50
With others, including children	0.22	0.41	0.19	0.39	0.22	0.42	0.21	0.41	0.22	0.41	0.20	0.40
Keyworker (ref not)	0.21	0.40	0.18	0.39	0.22	0.41	0.19	0.39	0.22	0.41	0.18	0.38
Long-term physical health condition	0.42	0.49	0.47	0.50	0.46	0.50	0.50	0.50	0.51	0.50	0.55	0.50

Long-term mental health condition	0.16	0.37	0.16	0.37	0.33	0.47	0.33	0.47	0.51	0.50	0.53	0.50
-----------------------------------	------	------	------	------	------	------	------	------	------	------	------	------

Note. Data in the weighted samples were weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Ethnic minority refers to Black, Asian and minority ethnicity. GCSE refers to General Certificate of Secondary Education.

Table S5. Fixed-effects logistic regression models predicting within-individual change in self-harm thoughts from individual categories of adversity experiences and worries accounting for anxiety symptoms

Self-harm thoughts															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Anxiety symptoms	1.34	1.33	1.34	1.32	1.30	1.34	1.32	1.30	1.33	1.33	1.32	1.35	1.36	1.35	1.38
Adversity experiences															
Financial	1.01	0.95	1.07	1.04	0.90	1.19	1.12	0.99	1.26	0.94	0.85	1.04	0.98	0.86	1.11
Accessing essentials	1.24	1.14	1.35	1.01	0.79	1.28	0.88	0.74	1.04	1.55	1.33	1.80	1.41	1.21	1.64
COVID-19 illness	1.15	1.06	1.25	1.27	1.02	1.57	1.05	0.89	1.24	1.30	1.12	1.50	1.04	0.88	1.23
Illness of others/ bereavement	0.97	0.90	1.04	1.13	0.94	1.36	0.99	0.85	1.16	1.01	0.89	1.14	0.85	0.75	0.97
Physical/psychological abuse	2.78	2.61	2.96	3.66	3.05	4.40	3.11	2.71	3.57	2.68	2.40	2.99	2.43	2.18	2.72
Worries															
Financial	1.14	1.09	1.20	1.12	1.00	1.26	1.06	0.96	1.17	1.21	1.10	1.33	1.18	1.07	1.31
Accessing essentials	1.03	0.98	1.08	1.20	1.05	1.36	0.99	0.90	1.10	0.87	0.79	0.95	1.14	1.04	1.25
COVID-19 illness	0.84	0.80	0.88	0.82	0.73	0.92	0.71	0.65	0.79	0.82	0.75	0.90	0.99	0.90	1.08
Social/relationship concerns	1.31	1.25	1.38	1.51	1.33	1.71	1.31	1.19	1.45	1.33	1.22	1.46	1.19	1.09	1.31
Threats to safety	1.12	1.06	1.18	1.30	1.14	1.49	1.06	0.95	1.17	1.12	1.02	1.23	1.10	1.00	1.21
Number of observations	169,939			14,285			44,658			61,045			49,951		
Number of individuals	10,454			1,081			3,055			3,663			2,655		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table S6. Fixed-effects logistic regression models predicting within-individual change in self-harm behaviours from individual categories of adversity experiences and worries accounting for anxiety symptoms

		Self-harm behaviours														
		Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
Variable		OR	95% CI	95% CI	OR	95% CI	95% CI	OR	95% CI	95% CI	OR	95% CI	95% CI	OR	95% CI	95% CI
			lower	upper		lower	upper		lower	upper		lower	upper		lower	upper
	Anxiety symptoms	1.18	1.17	1.19	1.15	1.13	1.17	1.18	1.15	1.20	1.18	1.16	1.20	1.23	1.20	1.25
Adversity experiences	Financial	0.97	0.87	1.07	1.08	0.88	1.32	0.78	0.63	0.97	1.08	0.91	1.29	0.83	0.65	1.06
	Accessing essentials	1.31	1.16	1.48	1.17	0.92	1.48	1.01	0.77	1.33	1.40	1.13	1.74	1.87	1.44	2.43
	COVID-19 illness	1.21	1.04	1.40	1.45	1.00	2.10	0.85	0.64	1.14	1.55	1.21	1.99	1.02	0.72	1.44
	Illness of others/bereavement	1.12	0.99	1.26	1.23	0.97	1.56	1.08	0.82	1.43	1.10	0.90	1.36	1.00	0.77	1.30
Worries	Physical/ psychological abuse	4.24	3.86	4.65	2.84	2.29	3.52	6.05	4.93	7.42	3.87	3.30	4.53	4.76	3.95	5.73
	Financial	1.11	1.02	1.22	1.08	0.91	1.28	1.08	0.89	1.31	1.34	1.14	1.58	0.93	0.76	1.14
	Accessing essentials	1.08	1.00	1.18	1.26	1.07	1.49	1.07	0.90	1.28	1.07	0.92	1.25	0.95	0.79	1.13
	COVID-19 illness	0.87	0.79	0.95	0.95	0.79	1.14	0.76	0.63	0.91	0.91	0.77	1.07	0.90	0.74	1.10
	Social/relationship concerns	1.04	0.95	1.14	1.31	1.09	1.58	0.97	0.80	1.17	1.01	0.85	1.19	0.93	0.76	1.14
	Threats to safety	1.08	0.99	1.18	0.92	0.77	1.11	1.44	1.19	1.73	1.01	0.87	1.17	1.09	0.91	1.31
	Number of observations	52,071			6,463			15,277			19,293			11,038		
	Number of individuals	3,360			475			1,085			1,184			616		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table S7. Fixed-effects logistic regression models predicting within-individual change in self-harm thoughts from individual categories of adversity experiences and worries accounting for depressive symptoms

Variable	Self-harm thoughts														
	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Depressive symptoms	1.55	1.54	1.56	1.52	1.49	1.54	1.52	1.50	1.54	1.56	1.54	1.58	1.60	1.58	1.62
Adversity experiences															
Financial	0.90	0.84	0.97	0.95	0.81	1.11	0.94	0.82	1.08	0.86	0.77	0.96	0.92	0.79	1.07
Accessing essentials	1.15	1.04	1.26	0.85	0.66	1.11	0.78	0.64	0.95	1.53	1.30	1.82	1.30	1.10	1.53
COVID-19 illness	1.03	0.94	1.13	1.43	1.12	1.83	0.90	0.75	1.09	1.14	0.97	1.35	0.86	0.71	1.03
Illness of others/ bereavement	0.92	0.85	1.00	1.02	0.84	1.25	0.90	0.76	1.07	0.95	0.82	1.09	0.90	0.78	1.04
Physical/ psychological abuse	2.39	2.23	2.57	2.96	2.42	3.62	2.41	2.07	2.80	2.32	2.05	2.62	2.29	2.02	2.60
Worries															
Financial	1.11	1.05	1.17	1.17	1.03	1.33	1.02	0.91	1.14	1.10	0.99	1.22	1.16	1.04	1.29
Accessing essentials	0.94	0.89	0.99	1.17	1.01	1.34	0.89	0.80	1.00	0.78	0.70	0.86	1.02	0.92	1.13
COVID-19 illness	0.88	0.83	0.93	0.88	0.78	1.01	0.77	0.69	0.86	0.87	0.78	0.96	0.99	0.89	1.09
Social/ relationship concerns	1.22	1.15	1.29	1.49	1.30	1.70	1.18	1.05	1.32	1.22	1.10	1.35	1.13	1.02	1.26
Threats to safety	1.09	1.03	1.15	1.36	1.17	1.57	0.99	0.88	1.11	1.07	0.97	1.19	1.09	0.98	1.21
Number of observations	169,939			14,285			44,658			61,045			49,951		
Number of individuals	10,454			1,081			3,055			3,663			2,655		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table S8. Fixed-effects logistic regression models predicting within-individual change in self-harm behaviours from individual categories of adversity experiences and worries accounting for depressive symptoms

Self-harm behaviours															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Depressive symptoms	1.21	1.20	1.22	1.18	1.16	1.20	1.22	1.20	1.25	1.22	1.20	1.24	1.23	1.21	1.25
Adversity experiences															
Financial	0.91	0.83	1.01	0.98	0.80	1.20	0.75	0.60	0.94	1.01	0.84	1.20	0.88	0.69	1.12
Accessing essentials	1.17	1.04	1.32	1.04	0.82	1.32	0.90	0.69	1.17	1.29	1.05	1.60	1.55	1.20	2.00
COVID-19 illness	1.18	1.02	1.36	1.80	1.24	2.62	0.75	0.56	0.99	1.65	1.31	2.09	0.84	0.61	1.15
Illness of others/bereavement	1.10	0.98	1.24	1.35	1.07	1.69	1.04	0.79	1.37	1.02	0.84	1.25	1.05	0.82	1.34
Physical/psychological abuse	3.83	3.49	4.19	3.06	2.49	3.76	5.50	4.48	6.75	3.20	2.74	3.73	4.33	3.62	5.19
Worries															
Financial	1.09	1.00	1.19	1.06	0.90	1.26	1.06	0.88	1.28	1.33	1.13	1.56	0.90	0.74	1.09
Accessing essentials	1.08	0.99	1.17	1.28	1.09	1.51	1.00	0.84	1.20	1.05	0.91	1.22	0.98	0.82	1.16
COVID-19 illness	0.94	0.86	1.03	1.03	0.86	1.23	0.80	0.67	0.96	0.95	0.81	1.11	1.07	0.88	1.30
Social/ relationship concerns	1.06	0.97	1.16	1.40	1.17	1.68	0.89	0.74	1.07	1.05	0.89	1.24	0.93	0.76	1.14
Threats to safety	1.06	0.97	1.15	0.96	0.80	1.15	1.49	1.24	1.79	1.00	0.86	1.16	0.94	0.79	1.12
Number of observations	52,071			6,463			15,277			19,184			11,038		
Number of individuals	3,360			475			1,085			1,184			616		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table S9. Fixed-effects logistic regression models predicting within-individual change in self-harm thoughts from individual categories of adversity experiences and worries with physical abuse and psychological abuse as individual adversity experiences

Self-harm thoughts															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
Adversity experiences															
Financial	1.18	1.12	1.25	1.24	1.09	1.41	1.26	1.13	1.41	1.09	0.99	1.20	1.20	1.06	1.35
Accessing essentials	1.36	1.26	1.47	1.19	0.95	1.48	0.99	0.84	1.16	1.68	1.46	1.93	1.50	1.30	1.74
COVID-19 illness	1.20	1.11	1.30	1.32	1.09	1.61	1.08	0.92	1.26	1.33	1.16	1.52	1.12	0.96	1.31
Illness of others/bereavement	1.16	1.09	1.24	1.43	1.21	1.69	1.11	0.96	1.28	1.21	1.07	1.36	1.04	0.92	1.17
Physical abuse	2.93	2.62	3.28	6.46	4.16	10.02	4.61	3.52	6.03	2.85	2.32	3.50	2.23	1.88	2.65
Psychological abuse	3.11	2.92	3.31	3.46	2.90	4.11	3.06	2.66	3.52	3.09	2.78	3.44	3.03	2.71	3.39
Worries															
Financial	1.42	1.36	1.49	1.32	1.19	1.47	1.36	1.24	1.5	1.61	1.48	1.76	1.37	1.25	1.50
Accessing essentials	1.19	1.14	1.25	1.46	1.29	1.64	1.13	1.03	1.23	1.00	0.91	1.09	1.32	1.21	1.44
COVID-19 illness	0.96	0.92	1.00	0.92	0.83	1.02	0.79	0.72	0.87	0.93	0.86	1.01	1.19	1.09	1.30
Social/relationship concerns	1.62	1.55	1.70	1.87	1.67	2.10	1.65	1.51	1.81	1.66	1.53	1.81	1.42	1.31	1.55
Threats to safety	1.41	1.34	1.48	1.68	1.49	1.90	1.40	1.27	1.55	1.37	1.25	1.50	1.35	1.24	1.48
Number of observations	169,954			14,286			44,659			61,051			49,958		
Number of individuals	10,456			1,081			3,055			3,664			2,656		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.

Table S10. Fixed-effects logistic regression models predicting within-individual change in self-harm behaviours from individual categories of adversity experiences and worries with physical abuse and psychological abuse as individual adversity experiences

Self-harm behaviours															
Variable	Total sample			Ages 18-29			Ages 30-44			Ages 45-59			Ages 60+		
	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper	OR	95% CI lower	95% CI upper
<b>Adversity experiences</b>															
Financial	1.01	0.92	1.12	0.99	0.81	1.21	0.78	0.63	0.97	1.15	0.97	1.36	1.08	0.86	1.37
Accessing essentials	1.32	1.17	1.48	1.25	0.98	1.58	0.95	0.73	1.24	1.44	1.17	1.77	1.81	1.40	2.34
COVID-19 illness	1.22	1.06	1.40	1.52	1.06	2.17	0.88	0.66	1.17	1.63	1.30	2.04	0.94	0.69	1.29
Illness of others/bereavement	1.25	1.11	1.39	1.51	1.20	1.90	1.21	0.93	1.59	1.13	0.93	1.37	1.18	0.93	1.49
Physical abuse	8.73	7.69	9.92	15.96	10.95	23.25	18.23	13.04	25.48	6.04	4.90	7.45	7.09	5.67	8.85
Psychological abuse	2.42	2.20	2.67	1.84	1.49	2.28	2.29	1.83	2.86	2.62	2.22	3.09	2.83	2.33	3.44
<b>Worries</b>															
Financial	1.26	1.16	1.38	1.21	1.03	1.43	1.30	1.08	1.56	1.54	1.31	1.80	1.00	0.83	1.21
Accessing essentials	1.19	1.09	1.28	1.49	1.26	1.74	1.14	0.96	1.36	1.12	0.97	1.30	1.07	0.90	1.27
COVID-19 illness	0.95	0.87	1.04	1.04	0.88	1.24	0.79	0.66	0.94	0.93	0.80	1.09	1.15	0.95	1.38
Social/relationship concerns	1.21	1.11	1.33	1.40	1.17	1.68	1.14	0.95	1.37	1.19	1.02	1.40	1.11	0.92	1.35
Threats to safety	1.21	1.11	1.31	1.07	0.89	1.27	1.75	1.46	2.10	1.13	0.98	1.31	1.08	0.91	1.29
Number of observations	52,074			6,463			15,277			19,294			11,040		
Number of individuals	3,360			475			1,085			1,184			616		

Note. Adversity experiences and worries variables are weighted to the proportions of gender, age, ethnicity, country, and education obtained from the Office for National Statistics. Individual adversity experiences and worries variables are binary. Analyses were further adjusted for day of the week and time since lockdown began. OR = odds ratio. CI = confidence interval.