TITLE:

Microbial signatures in the lower airways of mechanically ventilated COVID19 patients associated with poor clinical outcome

Imran Sulaiman[1,2], Matthew Chung[3], Luis Angel[1,2], Jun-Chieh J. Tsay[1,2,4], Benjamin G. Wu[1,2,4], Stephen T. Yeung[5], Kelsey Krolikowski[1,2], Yonghua Li[1,2], Ralf Duerr[5], Rosemary Schluger[1,2], Sara A. Thannickal[5], Akiko Koide[2,6], Samaan Rafeq[1,2], Clea Barnett[1,2], Radu Postelnicu[1,2], Chang Wang[7], Stephanie Banakis[3], Lizzette Perez-Perez[8], George Jour[9], Guomiao Shen[9], Peter Meyn[10], Joseph Carpenito[1,2], Xiuxiu Liu[1,10], Kun Ji[1,11], Destiny Collazo[1,2], Anthony Labarbiera[1,2], Nancy Amoroso[1,2], Shari Brosnahan[1,2], Vikramjit Mukherjee[1,2], David Kaufman[1,2], Jan Bakker[1,2], Anthony Lubinsky[1,2], Deepak Pradhan[1,2], Daniel H. Sterman[1,2], Michael Weiden[1,2], Adriana Hegu[9,12], Laura Evans[13], Timothy M. Uyeki[14], Jose C. Clemente[15], Emmie De wit[8], Ann Marie Schmidt[16], Bo Shopsin[17], Ludovic Desvignes [5], Chan Wang[18], Huilin Li[18], Bin Zhang[19], Christian V. Forst[19], Shohei Koide[6,20], Kenneth A. Stapleford[5], Kamal M. Khanna[5,6], Elodie Ghedin[3,7], and Leopoldo N. Segal[1,2,6*]*

[1]Division of Pulmonary and Critical Care Medicine, New York University Grossman School of Medicine, NYU Langone Health, New York, NY
[2] Department of Medicine, New York University Grossman School of Medicine, NYU Langone Health, New York, NY
[3]Systems Genomics Section, Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD
[4]Division of Pulmonary and Critical Care Medicine, VA New York Harbor Healthcare System, NY
[5] Department of Microbiology, New York University Grossman School of Medicine, NYU Langone Health, New York, NY.
[6] Laura and Isaac Perlmutter Cancer Center, New York University School of Medicine, NYU Langone Health, New York, NY.
[7] Center for Genomics & Systems Biology, Department of Biology, New York University New York, NY
[8] Molecular Pathology Unit, Laboratory of Virology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rocky Mountain Laboratories, Hamilton, MT
[9] Department of Pathology, New York University Grossman School of Medicine, NYU Langone Health, New York, NY.
[10] Division of Pediatrics, Longhua Hospital Affiliated to Shanghai University of Chinese Medicine, CN
[11] Dongfang Hospital affiliated to Beijing University of Traditional Chinese Medicine, Beijing, China
[12] NYU Langone Genome Technology Center, New York University Grossman School of Medicine, NYU Langone Health, New York, NY
[13] Pulmonary, Critical Care and Sleep Medicine, University of Washington, Seattle, USA
[14] Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA
[15] Department of Genetics and Genomic Sciences and Immunology Institute, Icahn School of Medicine at Mount Sinai, NY.

[16] Diabetes Research Program, Division of Endocrinology, Diabetes and Metabolism,
Department of Medicine, New York University Grossman School of Medicine, NYU
Langone Health, New York, NY
[17] Division of Infectious Diseases, Department of Medicine, New York University School
of Medicine, NYU Langone Health, New York, NY
[18] Department of Population Health, New York University School of Medicine, NYU
Langone Health, New York, NY
[19] Department of Genetics and Genomic Sciences, Icahn Institute of Genomics and
Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, NY
[20] Department of Biochemistry and Molecular Pharmacology, New York University
Grossman School of Medicine, NYU Langone Health, New York, NY
and Perlmutter Cancer Center, New York, NY 10016
59
60
Imran Sulaiman, MD, PhD   SheikMohammadImran.Sulaiman@nyumc.org
Matthew Chung, PhD   matt.chung@nih.gov
Luis Angel, MD   Luis.Angel@nyulangone.org
Jun-Chieh J. Tsay, MD   jun-chieh.tsay@nyumc.org
Benjamin G. Wu, MD   Benjamin.Wu@nyumc.org
Stephen Yeung, PhD   Stephen.Yeung@nyulangone.org
Kelsey Krolikowski   Kelsey.Krolikowski@nyulangone.org
Yonghua Li, MD, PhD   Yonghua.Li@nyumc.org
Ralf Duerr, MD, PhD   Ralf.Duerr@nyulangone.org
Rosemary Schluger, RN   rosemary.schluger@nyumc.org
Sara A. Thannickal, BS   Sara.thannickal@nyulangone.org
Akiko Koide, Phd,   Akiko.Koide@nyulangone.org
Samaan Rafeq, MD   Samaan.Rafeq@nyulangone.org
Clea Barnett, MD   Clea.Barnett@nyulangone.org
Radu Postelnicu, MD   Radu.Postelnicu@nyulangone.org
Chang Wang, PhD   cw2505@nyu.edu
Stephanie Banakis   stephanie.banakis@nih.gov
Lizzette Perez-Perez   lizzette.perez-perez@nih.gov
George Jour, PhD   George.Jour@nyulangone.org
Guomiao Shen, Phd   Guomiao.Shen@nyulangone.org
Peter Meyn   peter.meyn@nyumc.org
Joseph Carpenito   Joseph.carpenito@nyumc.org
Xiuxiu Liu, MS   liuxiuxiu_longhua@outlook.com
Kun Ji, MD, PhD   Kun.Ji@nyulangone.org
Destiny Collazo   Destiny.Collazo@nyulangone.org
Anthony Labarbiera   Anthony.Labarbiera@nyulangone.org
Nancy Amoroso, MD   Nancy.Amoroso@nyulangone.org
Shari Brosnahan, MD   Shari.Brosnahan@nyulangone.org
Vikramjit Mukherjee, MD   Vikramjit.Mukherjee@nyulangone.org
David Kaufman, MD   David.A.Kaufman@nyulangone.org
Jan Bakker, MD, PhD   Jan.Bakker@nyulangone.org
Anthony Lubinsky, MD   Anthony.Lubinsky@nyulangone.org
Deepak Pradhan, MD   Deepak.Pradhan@nyulangone.org
Daniel H. Sterman, MD   Daniel.sterman@nyumc.org

Michael Weiden, MD   Michael.Weiden@nyulangone.org
Adriana Heguy, PhD   Adriana.heguy@nyumc.org
Laura Evans, MD   leevans@uw.edu
Timothy M. Uyeki, MD, MPH   tmu0@cdc.gov
Jose C. Clemente, PhD   jose.clemente@mssm.edu
Emmie De wit   emmie.dewit@nih.gov
Ann Marie Schmidt, MD   AnnMarie.Schmidt@nyulangone.org
Bo Shopsin, MD, PhD   Bo.Shopsin@nyulangone.org
Ludovic Desvignes, PhD   Ludovic.Desvignes@nyulangone.org
Chan Wang, PhD   Chan.Wang@nyulangone.org
Huilin Li, PhD   Huilin.Li@nyulangone.org
Bin Zhang, PhD   bin.zhang@mssm.edu
Christian V. Forst, PhD   christian.forst@mssm.edu
Shohei Koide, PhD   Shohei.Koide@nyulangone.org
Kenneth A. Stapleford, PhD   Kenneth.Stapleford@nyulangone.org
Kamal Khanna, PhD   Kamal.Khanna@nyulangone.org
Elodie Ghedin, PhD   elodie.ghedin@nih.gov
Leopoldo N. Segal, MD   Leopoldo.Segal@nyumc.org
113
114 *Corresponding Authors: Elodie Ghedin (elodie.ghedin@nih.gov) and Leopoldo N.
115 Segal, (Leopoldo.Segal@nyumc.org)

116
117
Authors Contributions:
Study conception and design: IS, LA, LD, SK, KS, KK, EG, LNS
Data was obtained by: IS, LA, SY, KK, YL, RD, RS, SAT, AK, SR, CB, RP, GJ, GS, PM,
JC, XL, KJ, DC, AL, NA, SB, VM, DK, JB, AL, DP, LPP, EDW, DHS, AH, LNS
Data were analyzed by: IS, MC, JCJT, BGW, SY, KK, RD, CW, JCC, CW, HL, CVF, BZ,
SK, KS, EG, LNS
The first draft of the manuscript was written by: IS, MC, LA, EG, LNS
All authors read, critically revised and approved the final manuscript.
126

Research support funding:
R37 CA244775 (LNS, NCI/NIH); R01 HL125816 (LNS, SBK, NHLBI/NIH); Colton Pilot
Project Grant (LNS, LA, SBL, KK); UWSC12030 (LNS, RP, LE, CDC Foundation);
PACT grant (LNS, FNIH); R21 AI158997 (SK); R01 AI143861 (KMK, NIAID/NIH); R01
AI143861-02S1 (KMK, NIAID/NIH); R01 DK110014 (HL and CW, NIDDK/NIH); P20
CA252728 (CW and HL, NCI/NIH) American Association for Cancer Research Grant
(HP/LNS); The Genome Technology Center is partially supported by the Cancer Center
Support Grant P30CA016087 at the Laura and Isaac Perlmutter Cancer Center (AH,
AT); FAMRI Young Clinical Scientist Award (BGW), Stony Wold-Herbert Fund Grant-in-
Aid/Fellowship (IS, CB). This work was supported in part by the Division of Intramural
Research (DIR) of the NIAID/NIH (EG, EDW).
138
139 **Acknowledgement:**
140 We would like to thank the Genome Technology Center (GTC) for expert library
141 preparation and sequencing, and the Applied Bioinformatics Laboratories (ABL) for
142 providing bioinformatics support and helping with the analysis and interpretation of the

data. Experimental Pathology Research Laboratory for histopathology services and imaging. GTC and ABL are shared resources partially supported by the Cancer Center Support Grant P30CA016087 at the Laura and Isaac Perlmutter Cancer Center. This work has used computing resources at the NYU School of Medicine High Performance Computing Facility (HPCF). Financial support for the PACT project is possible through funding support provided to the FNIH by: AbbVie Inc., Amgen Inc., Boehringer-Ingelheim Pharma GmbH & Co. KG, Bristol-Myers Squibb, Celgene Corporation, Genentech Inc., Gilead, GlaxoSmithKline plc, Janssen Pharmaceutical Companies of Johnson & Johnson, Novartis Institutes for Biomedical Research, Pfizer Inc., and Sanofi. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Running Head: COVID-19

Word Count: 4869
Body of the manuscript:

Financial Disclosure: None

Key words: microbiome, virome, severe acute respiratory syndrome

Abstract:

Mortality among patients with COVID-19 and respiratory failure is high and there are no known

lower airway biomarkers that predict clinical outcome. We investigated whether bacterial

respiratory infections and viral load were associated with poor clinical outcome and host

immune tone. We obtained bacterial and fungal culture data from 589 critically ill subjects with

COVID-19 requiring mechanical ventilation. On a subset of the subjects that underwent

bronchoscopy, we also quantified SARS-CoV-2 viral load, analyzed the microbiome of the lower

airways by metagenome and metatranscriptome analyses and profiled the host immune

response. We found that isolation of a hospital-acquired respiratory pathogen was not

associated with fatal outcome. However, poor clinical outcome was associated with enrichment

of the lower airway microbiota with an oral commensal (*Mycoplasma salivarium*), while high

SARS-CoV-2 viral burden, poor anti-SARS-CoV-2 antibody response, together with a unique

host transcriptome profile of the lower airways were most predictive of mortality. Collectively,

these data support the hypothesis that 1) the extent of viral infectivity drives mortality in severe

COVID-19, and therefore 2) clinical management strategies targeting viral replication and host

responses to SARS-CoV-2 should be prioritized.

Introduction

The earliest known case of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) infection causing coronavirus virus disease (COVID-19) is thought to have occurred on November 17, 2019[1]. As of February 20, 2021, 110.3 million confirmed cases of COVID-19 and 2.4 million deaths have been reported worldwide[2]. As the global scientific community rallied in a concerted effort to understand SARS-CoV-2 infections, our background knowledge is rooted in previous experience with the related zoonotic betacoronaviruses Middle East Respiratory Syndrome coronavirus (MERS-CoV) and SARS-CoV-1 that have caused severe pneumonia with 34.4% and 9% case fatality, respectively[3]. As observed for these related coronaviruses, SARS-CoV-2 infection can result in an uncontrolled inflammatory response[4] leading to acute respiratory distress syndrome (ARDS) and multi-organ failure, both associated with increased mortality. While a large proportion of the SARS-CoV-2 infected population is asymptomatic or experiences mild illness, a substantial number of individuals will develop severe disease and require hospitalization, with some progressing to respiratory failure. Mortality among hospitalized COVID-19 patients is estimated to be approximately 20%, which can go up to 70% among those requiring invasive mechanical ventilation [5-12].

Mortality in other viral pandemics, such as the 1918 H1N1 and 2009 H1N1 influenza pandemics, has been attributed in part to bacterial co-infection or super-infection [13,14]. To determine if this is also the case for COVID-19, we can use next generation sequencing (NGS) to probe the complexity of the microbial environment (including RNA and DNA viruses, bacteria and fungi) and how the host (human) responds to infection.

Recent studies have used this approach to uncover microbial signatures in patients with

ARDS.[15,16] Increased bacterial burden and the presence of gut-associated bacteria in

the lung were shown to worsen outcomes in these critically ill patients [15,17], highlighting

the potential role of the lung microbiome in predicting outcomes in ARDS. In a recent

study using whole genome sequencing to profile the gut microbiome of 69 patients from

Hong Kong, investigators identified an increased abundance of opportunistic fungal

pathogens among patients with confirmed COVID-19[18]. While there is emerging interest

in understanding the microbial environment in patients with SARS-CoV-2 infections, few

studies have attempted to characterize this at the primary site of the disease activity:

the lower airways[19,20]. Furthermore, no study has yet determined whether microbial

differences in the airways of COVID-19 patients could be contributing to the different

outcomes in patients receiving mechanical ventilation.

In this investigation, we accessed a large prospective cohort of critically ill patients with

SARS-CoV-2 infection who required invasive mechanical ventilation, and from whom

bronchoalveolar lavage (BAL) samples were collected. We characterized the lung

microbiome of these patients in parallel with analyses of lower airway markers of host

immunity. While we did not find that isolation of a secondary respiratory pathogen was

associated with prolonged mechanical ventilation (>28 days) or fatal outcome, we did

identify critical microbial signatures—characterized by enrichment of oral commensals,

high SARS-CoV-2 load, and decreased anti-SARS-CoV-2 IgG response—associated

with fatal outcome, suggesting a need for more targeted antiviral therapeutic

approaches for the care of critically ill COVID19 patients.

Results

Cohort

From March 3rd to June 18th 2020, a total of 589 patients with laboratory-confirmed SARS-CoV-2 infection were admitted to the intensive care units of two academic medical centers of NYU Langone Health in New York (Long Island and Manhattan) and required invasive mechanical ventilation (MV). This included a subset of 142 patients from the Manhattan campus who underwent bronchoscopy for airway clearance and/or tracheostomy from which we collected and processed lower airway samples for this investigation (**Supplementary Fig. 1**). **Table 1** shows demographics and clinical characteristics of the 142 patients who underwent bronchoscopy divided into three clinical outcomes: survivors with ≤28 Days on MV; survivors with >28 Days on MV; and deceased. The median post admission follow-up time was 232 days (CI=226-237 days). **Supplementary Tables 1** and **2** compare similar data across all 589 subjects, divided per site and sub-cohorts. Patients at the Manhattan campus who underwent bronchoscopy were younger, had lower body mass index (BMI), and a lower prevalence of chronic obstructive pulmonary disease (COPD; **Supplementary Table 1**). Among the cohort that provided lower airway samples through bronchoscopy, 37% of the subjects were successfully weaned within 28 days of initiation of MV and survived hospitalization, 39% required prolonged MV but survived hospitalization, and 23% died. Patients within the bronchoscopy cohort had a higher overall survival than the rest of the NYU COVID-19 cohort since most critically ill patients were not eligible for bronchoscopy or tracheostomy. Mortality among those in the no-bronchoscopy cohort was 77%. In the overall NYU cohort, higher age and BMI were associated with increased mortality (**Supplementary Table 2**). There was a similar, albeit non-

significant, trend for the bronchoscopy cohort. Among the clinical characteristics of this

cohort, patients within the deceased group more commonly had a past medical history

of chronic kidney disease and cerebrovascular accident.

Study patients were admitted during the first wave of the pandemic in the US, prior to

current standardized management of COVID-19. Within the bronchoscopy cohort, more

than 90% of the subjects received hydroxychloroquine and anticoagulation (therapeutic

dose), 69% received corticosteroids, 41% received tocilizumab (anti-Interleukin (IL)-6

receptor monoclonal antibody), 21% required dialysis, and 18.9% were started on

extracorporeal membrane oxygenation (ECMO) (**Table 1**). Antimicrobial therapy

included use of antivirals (lopinavir/ritonavir in 16% and remdesivir in 10%), antifungals

(fluconazole in 40% and micafungin in 57%), and antibiotics (any, in 90% of the

subjects). Among the factors associated with clinical outcome within the bronchoscopy

cohort, patients who survived were more commonly placed on ECMO whereas patients

who died had frequently required dialysis (**Table 1**); these trends were also observed

across the whole NYU cohort. Neither hydroxychloroquine or azithromycin were

significantly associated with clinical outcome; however, patients who survived were

more frequently treated with the combination antibiotic piperacillin/tazobactam.

Within the first 48hrs from admission, respiratory bacterial cultures were rarely obtained

(n=70/589, 12%) with very few positive results (n=12, 17%). Blood cultures were more

commonly obtained (n=353/589, 60%) but the rate of bacterial culture positivity was

much lower (n=5, 1.4%). These data support that community acquired bacterial co-

infection was not a common presentation among critically ill COVID-19 patients.

During their hospitalization, most patients had respiratory and/or blood specimens collected for bacterial cultures (**Table 1** and **Supplementary Table 1**). The proportions of positive bacterial respiratory cultures and blood cultures were 73% and 43%, respectively. We evaluated whether respiratory or blood culture results obtained as per clinical standard of care were associated with clinical outcome. Risk analyses for the culture results during hospitalization for the whole cohort (n=589) demonstrated that bacterial culture positivity was not associated with increased odds of dying but was associated with prolonged mechanical ventilation in the surviving patients (**Figure 1**). Since length of stay could potentially affect these results (patients who died could have a shorter hospitalization, and therefore may have had fewer specimens collected for cultures), we repeated the analysis using culture data obtained during the first two weeks of hospitalization. This analysis showed that bacterial pathogen culture positivity (both respiratory and blood) during the early period of hospitalization was not associated with worse outcome (**Figure 1** and **Supplementary Table 3**). Interestingly, identification of oral bacteria in respiratory culture, commonly regarded as procedural contaminants, was associated with higher odds of prolonged mechanical ventilation (>28 days) among survivors. Similar trends were noted when analysis was performed on subjects from NYU LI and NYU Manhattan separately, or for the bronchoscopy cohort (**Supplementary Table 2**). Among the bronchoscopy cohort, there was no statistically significant association between culture results and clinical outcome, but there was a trend towards an increased rate of positive respiratory cultures for *Staphylococcus aureus* (including MRSA), *S. epidermidis*, and *Klebsiella pneumoniae* in the survival groups (**Table 1**). These data suggest that in critically ill patients with COVID-19 requiring MV, hospital isolation of a secondary respiratory bacterial pathogen

is not associated with worse clinical outcome.

SARS-CoV-2 load in the lower airways is associated with poor clinical outcome

Using supraglottic and BAL samples from patients undergoing bronchoscopy (n=142),

we evaluated the viral load by rRT-PCR for the SARS-CoV-2 N gene, adjusted by levels

of human ribosomal protein (RP). Of note, the majority of samples were largely obtained

in the second week of hospitalization (**Table 1**, median[IQR] = 10[6-14], 13[8-16], and

13[8-16] for the ≤28-days MV, >28-days MV, and deceased groups, respectively, p=ns).

Paired analysis of upper and lower airway samples revealed that, while there was a

positive association between SARS-CoV-2 viral load of the paired samples (rho = 0.60,

p<0.0001), there was a subset of subjects (21%) for which the viral load was greater in

the BAL than in the supraglottic area, indicating topographical differences in SARS-

CoV-2 replication (**Figure 2a**). Importantly, while the SARS-CoV-2 viral load in the

upper airway samples was not associated with clinical outcome (**Supplementary Fig.**

2), patients who died had higher viral load in their lower airways than patients who

survived (**Figure 2b**). We then evaluated virus viability in BAL samples by measuring

levels of subgenomic RNA (sgRNA) targeting the E gene of SARS-CoV-2. This mRNA

is only transcribed inside infected mammalian cells and is not packed into virions, thus,

its presence is indicative of viable infecting viral particles in a sample[21]. In BAL, levels of

sgRNA correlated with viral load as estimated by rRT-PCR for the SARS-CoV-2 N gene

(**Figure 2c**) and the highest percentage of measurable sgRNA was in the deceased

group followed by the ≤28-days MV group, and the >28-days MV group (17,7%, 11.5%,

and 3.7%, respectively, chi-square p=0.028 for the comparison deceased vs. >28-days

MV group). Thus, while in most cases levels of sgRNA were not measurable in BAL

suggesting that there were no viable viral particles in the lower airways of COVID-19

patients at the time of bronchoscopy (overall median[IQR] = 12[7-16] days from

hospitalization), the lower airway viral burden, as estimated by rRT-PCR, is associated

with mortality in critically ill COVID-19 patients.

Microbial community structure of the lower airways is distinct from the upper airways in

critically ill patients.

Considering the bacterial species and the viral loads identified in the lower and upper

airways of this cohort and their association with outcomes, we profiled in detail their viral

and microbial composition. Microbial communities were evaluated using parallel

datasets of RNA and DNA sequencing from 118 COVID-19 patients with lower airway

samples that passed appropriate quality control and a subset of paired 64 upper airway

samples, along with background bronchoscope controls.

RNA sequencing (RNAseq) of the metatranscriptome provided insight into the RNA

virome as well as the transcriptomes of DNA viruses, bacteria, and fungi. Given the low

biomass of lower airway samples we first identified taxa as probable contaminants by

comparing the relative abundance between background bronchoscope and BAL

samples (**Supplementary Fig. 3a** and **Supplementary Table 4**). However, we did not

remove any taxa identified as probable contaminants from subsequent analyses. A

comparison of the microbial community complexity captured in these data, determined

using the Shannon diversity Index, showed there was significantly lower α diversity in

the lower airway samples than in the upper airways and background controls

(**Supplementary Fig. 4a**). Similarly, β diversity analysis based on the Bray Curtis

Dissimilarity index indicated that the microbial composition of the lower airways was

distinct from the upper airways and background controls (**Supplementary Fig. 4b**,

PERMANOVA p<0.01). Sequence reads indicated a much higher relative abundance of

SARS-CoV-2 in the lower than in the upper airways for this cohort (**Supplementary Fig.**

4c). Comparisons of the most dominant bacterial and fungal taxa that were functionally

active showed that *S. epidermidis*, *Mycoplasma salivarium, S. aureus*, *Prevotella oris*,

and *Candida albicans,* many often-considered oral commensals*,* were present in both

upper and lower airway samples (**Supplementary Fig. 4c**). Interestingly, the lytic phage

Proteus virus Isfahan, known to be active against biofilms of *Proteus mirabilis* [22], was

found to be highly transcriptionally active in the BAL.

DNA sequencing data provided insight into the DNA virome, as well as the bacterial and

fungal metagenomes. As for the metatranscriptome data, we first identified taxa as

probable contaminants but these were not removed for subsequent analyses

(**Supplementary Fig. 3b**). Both α and β diversity analyses of the metagenome support

distinct microbial community features in the lower airways as compared with the upper

airways and background controls (**Supplementary Fig. 5a, 5b**). Among the top 10 taxa

across lower and upper airway samples were *S. aureus*, *Salmonella enterica*,

Burkholderia dolosa, and *Klebsiella variicola*. *Candida albicans* only ranked #77 in the

BAL while it was ranked 5[th] in the metatranscriptome data indicating that while present

at low relative abundance, it was highly active (**Supplementary Table 4**). *K. variicola*,

while prevalent at a high relative abundance (#4 in BAL, and #5 in the upper airways) in

patients of this cohort, its ranking in the RNAseq data was not among the top 50,

indicating that it was not as active functionally as other bacteria. Conversely, while *S.*

epidermis ranked as the most highly functional taxon in both lower and upper airways,

based on RNAseq reads (**Supplemental Fig. 3c**), it was 33rd in relative abundance in

the BAL DNAseq data but was present at very high relative abundance in the upper

airways (ranked #3). These data suggest that microbes that colonize the upper airways

and the skin were common in the lower airways in this cohort of COVID-19 patients

requiring invasive mechanical ventilation.

Distinct microbial signatures are associated with different clinical outcomes.

To determine the potential impact of vertebrate viruses on outcome, we compared virus

enrichment differences in BAL samples across the three clinical outcome groups (≤28-

days MV, >28-days MV, and deceased). As it pertains to the vertebrate RNA virome

subfraction, there were significant differences (β diversity) between the three clinical

outcome groups (**Supplementary Fig. 6**, PERMANOVA p<0.01). There were no

significant differences for the vertebrate DNA virome or DNA virus transcriptome

subfractions of the sequence reads (data not shown). Consistent with the SARS-CoV-2

viral load assessed by RT-PCR, differential expression analysis (DESeq) of the RNA

virome identified SARS-CoV-2 as being enriched in the deceased group, as compared

with both ≤28-days and >28-days MV groups (fold change >5, **Figure 2d**). Cox

proportional hazards modeling supports that enrichment with SARS-CoV-2 was

associated with increased risk for death (HR 1.33, 95% CI= 1.07-1.67, pvalue=0.011,

FDR adjusted pvalue=0.06; **Supplementary Table 5**).

Analysis of differential DNA virus abundance using DEseq did not show statistically

significant differences. Because the virome includes viruses of bacteria and archaea,

we also analyzed the phage data (including viruses of archaea). Phages impact the

bacterial population—including bacterial pathogens—and so could be clinically relevant.

At a compositional level, the virome of DNA phages did not display statistically

significant differences or significant virus enrichment based on clinical outcome groups

(data not shown). However, while the phage metatranscriptome α and β diversity was

similar across the clinical outcome groups, there were various taxonomic differences at

the RNA level with enrichment of *Staphylococcus* phages CNPx in the deceased and

>28-day MV groups when compared with the ≤28-day MV group (**Figure 2e**).

Differential expression from two other *Staphylococcus* phages was also observed in the

>28-days MV group as compared with the ≤28-days MV group (**Figure 2e**). None of the

described taxa were identified as possible contaminants (**Supplementary Table 4**).

Enrichment of the lower airway microbiota with oral commensals is associated with poor

outcome

We evaluated the overall bacterial load by quantitative PCR, targeting the 16S rRNA

gene. As expected, the bacterial load in the lower airways was several folds lower than

in the upper airways but clearly higher than the background bronchoscope control

(**Supplementary Fig. 7**). Patients who died had higher total bacterial load in their lower

airways than patients who survived (**Figure 3a**).

While no statistically significant differences were noted in α or β diversity across clinical

outcome groups (**Figure 3b-c**), several differences were noted when differential

enrichment was evaluated using DESeq. For the comparisons made across the clinical

outcome groups we focused on consistent signatures identified in the lower airway

metagenome and metatranscriptome. Coherence of differentially enriched taxa was determined by gene set enrichment analysis (GSEA) (**Figure 3d**) and directionality of enrichment between the two datasets was evaluated (**Figure 3e)**. Among the most abundant taxa, the oral commensal *M. salivarum* was enriched in the deceased and >28-days MV groups as compared with the ≤28-days MV group. In contrast, a different oral commensal, *Prevotella oris*, was enriched in the ≤28-days MV group as compared with the deceased and >28-days MV groups. These data support that oral commensals are frequently found in the lower airways of critically ill COVID-19 patients and that differences between groups could be due to differential microbial pressures related to host factors. Interestingly, most of the statistically significant taxa were identified in the metatranscriptome rather than in the metagenome data, with only *P. oris* identified in both datasets. None of the described taxa were identified as possible contaminants (**Supplementary Table 4**). Overall, most of the microbial signatures identified as enriched in the deceased or in subjects on prolonged MV are regular colonizers of healthy skin and mucosal surfaces rather than frequent respiratory pathogens.

For the fungal data, there were no statistically significant differences in α or β diversity identified between clinical outcome groups in the metagenome or the metatranscriptome data (**Supplementary Fig. 8a** and **8c**). However, in the metagenome data, we identified *Candida glabrata* enriched in the deceased group as compared with the ≤28-days MV and the >28-days MV groups but this was not consistent in the metatranscriptome data (**Supplementary Fig. 8b** and **8d**).

Poor clinical outcomes are associated with enrichment of antimicrobial resistance genes

and glycosphingolipid biosynthesis

We used the gene annotation of the DNAseq and RNAseq data to profile the microbial

functional potential of the lower airway samples. For the comparisons made across the

clinical outcome groups, we focused on consistent functional signatures identified in the

lower airway metagenome and metatranscriptome. Coherence of differentially enriched

functions was determined using GSEA (**Figure 4a**) and directionality of enrichment was

also evaluated (**Figure 4b)**. Overall, there was coherence of directionality between the

metranscriptomics and metagenomics datasets for the comparisons between deceased

vs ≤28-days MV, and >28-days MV vs ≤28-days MV groups. Interestingly, statistically

significant differences were only noted in the metatranscriptome data and not in the

metagenome data. Among the top differentially expressed pathways in the poor

outcome groups were glycosylases, oxidoreductase activity, transporters, and two-

component system, among other genes. The two-component system is used by

bacteria and fungi for signaling. A specific analysis of antibiotic resistance genes shows

that there was significant gene enrichment and expression of biocide resistance in the

deceased group as compared to the two other MV groups (**Supplementary Fig. 9).**

There was also significant expression of genes resistant to trimethoprim and phenolic

compound, as well as multi drug resistance in the deceased group as compared to the

≤28-days MV group. Presence of the resistance gene against Trimethropim was not

significantly associated with prior exposure with Trimethoprim. However, only 7 patients

received this drug before sample collection.

 Lower airway host immune phenotype shows failure of adaptive and innate immune

response to SARS-CoV-2 among deceased subjects

To evaluate the host immune response to SARS-CoV-2 infection, we first measured

levels of anti-Spike and anti-RBD (receptor binding domain) antibodies in BAL samples.

For both anti-Spike and anti-RBD immunoglobulins, levels of IgG, IgA and IgM were

several logs higher than levels found in BAL samples from non-SARS-CoV-2 infected

patients. Importantly, IgG levels of anti-Spike and anti-RBD were significantly lower in

the deceased group as compared to the levels found in patients who survived (**Figure**

5a and **Supplementary Fig. 10a-c**, $p<0.05$). A neutralization assay performed using

BAL fluid showed varying levels of neutralization across all samples (as estimated by

EC50) but no statistically significant differences between the clinical outcome groups

(**Supplementary Fig. 10d**). We then evaluated whether levels of antibodies correlated

with viral load in BAL samples. While viral load levels of SARS-CoV-2 measured with

rRT-PCR did not correlate with BAL measurements of SARS-CoV-2 specific antibodies,

sgRNA viral load levels negatively correlated with BAL levels of Anti-Spike (IgG and

IgA), Anti-RBD (IgG and IgA) and the Neutralization assay (**Supplementary Table 6**).

These data suggest that the IgG subfraction is an important marker of the adaptive

immune response in the lung of critically ill COVID-19 patients and that both sub-

fractions of IgG and IgA anti-SARS-CoV-2 may contribute to the viral replication control

in the lower airways.

Host transcriptome analyses of BAL samples showed significant differences across

clinical outcome groups based on β diversity composition (**Supplementary Fig. 11)**. We

identified multiple differentially expressed genes across the clinical outcome groups

(**Supplementary Fig. 11b-d**). First, we noted that the lower airway transcriptomes

showed downregulation of heavy constant of IgG (IGHG3), and heavy constant of IgA

(IGHA1) genes in those with worse clinical outcome (**Supplementary Table 7**). We

then used IPA (Ingenuity Pathway Analysis) to summarize differentially expressed

genes across the three clinical outcome groups (**Figure 5b**). The sirtuin Signaling

Pathway (a pathway known to be involved in aging, gluconeogenesis/lipogenesis, and

host defense against viruses)[23] and the ferroptosis pathway (an iron-dependent form of

regulated cell death present in bronchial epithelium)[24,25] were both upregulated in those

with worse outcome. While this may reflect the host response to viral infection, other

differences in the transcriptomic data showed downregulation of mitochondrial oxidative

phosphorylation, HIF1α, STAT3, and Phospholipase C Signaling. Additional canonical

signaling pathways, including insulin secretion, multiple Inositol related pathways,

noradrenaline/adrenaline degradation signaling, and xenobiotic related metabolism

were significantly downregulated when comparing the >28-days MV vs. ≤28-days MV

groups. Upstream pathway prediction analyses of the host airway transcriptome support

previously reported mitochondria dysfunction[26] (inhibition in mitochondrial related

regulators NSUN3, MRPL14, MRPL12, LONP1, DAP3), and metabolic/gluconeogenesis

dysregulation[27,28] (SIRT3) in critically ill COVID-19 subjects with poor outcome

(Supplementary Table 8). We also observed decreased activation in the inflammatory

response in critically ill COVID-19 subjects with poor outcome (phagocytes, neutrophils,

and granulocytes, and leukocytes**; Supplementary Table 9**). A comparison of clinical

outcome between the >28-days MV vs. ≤28-days MV groups showed upstream

predicted inhibition in insulin, estrogen, beta-estradiol, EGF, EGFR, IL-5, and IL-10RA

in the >28-days MV group **(Supplementary Table 9)**. These differences suggest that, at

the stage that we sampled the lower airways of patients with critically COVID-19, an

overt inflammatory tone was not predictive of worst outcome.

To determine if the abundance of immune cells varies between different clinical

outcome groups, we estimated cell type abundance from the host transcriptome with

computational cell type quantification methods, including a deconvolution approach

implemented in CIBERSORTx [29] and a cell type signature enrichment approach

implemented in xCell [30]. As reported recently in other studies[31], among the cell types

detected in the BAL samples we observed a consistent enrichment of mast cells and

neutrophils in the >28-days MV and deceased groups compared with the ≤28-days MV

group (**Figure 5c** and **Supplementary Table 10**). We also identified significantly higher

inflammatory macrophages (M1), innate T-cells and memory T-cells (CCR7$^+$) among

subjects with worse clinical outcome.

Cross-kingdom network analyses identify bacteria, fungi, and host pathways functionally

impacted by SARS-CoV-2

To identify potential microbe-microbe and microbe-host interactions that could have an

effect on outcome, we used a multi-scale network analysis approach (Multiscale

Embedded Gene co-Expression Network Analysis, MEGENA)[32]. We first used the

relative abundance from the RNAseq data to capture co-expressing taxa in the

metatranscriptome network neighborhood of SARS-CoV-2 (SARS2-NWN). We

examined five such network neighborhoods (constructed by including nodes with

increasing distance 1 to 5 from SARS-CoV-2, i.e. neighborhood 1 to neighborhood 5)

that were significantly enriched for taxa functionally active in the deceased group when

compared with the ≤28-day MV group. Only the largest cluster, with 504 taxa, had

significantly enriched taxa in both the deceased and in the ≤28-day MV outcome groups

(**Supplementary Fig. 12a**) (FET P-value = 4.6e-45, 4.0 FE). Many of these taxa are

among the top 50 most abundant microbes we had previously identified in the

metatranscriptome dataset. Taxa present that are influenced by SARS-CoV-2 and

significantly differentially enriched in the deceased group include bacteria such as *M.*

salivarium, Bifidobacterium breve, and *Lactobacillus rhamnosus* (a gut commensal),

that we had previously identified by differential expression analysis (**Figure 3**e), but also

taxa such as *S. epidermis*, *Mycoplasma hominis* (urogenital bacteria)*,* and the phage

VB_PmiS-Isfahan (also referred to as Proteus virus Isfahan) that we had previously

only picked up as being highly abundant but not necessarily differentially enriched in the

deceased group. Most of the fungi, such as *C. albicans*, *C. glabrata* and *C. orthopsilosis*

were enriched in the ≤28-day MV group. Interestingly, our earlier analysis of the

metagenome (**Supplementary Fig. 8b**) had identified *C. glabrata* as being enriched in

the deceased group with no enrichment in the metatranscriptome. This analysis

indicates that some of these abundant taxa could be responding to SARS-CoV-2

disruption in a similar manner, or indirectly interacting functionally.

We further investigated the association of the network neighborhood with host network

modules using the host transcriptome data to identify groups of host genes that are co-

expressed in response to SARS-CoV-2 disruption. The 3 host modules with the most

significant correlations to SARS2-NWN are M175, M277 and M718. M277 is the parent

module of M718, and both are enriched with genes related to respiratory electron

transport, while M175 is enriched for IFN-γ signaling (**Supplementary Fig. 12b**).

Module M175 is positively correlated with the SARS2-NWN (ρ = 0.32, P-value = 2.1e-3).

While there was no collective enrichment of the module by differentially expressed

genes (DEGs) in the deceased vs ≤28-days MV, there was for >28-days vs ≤28-days

MV (FET P-value = 0.030, 4.5 FE). This module includes well-known antiviral IFN

stimulated genes (ISGs), such as *IRF7* and *OASL*. Investigating module response on an

individual gene level, Interleukin 4 induced 1 (IL4I1) appears as one of the most up-

regulated genes in this module when comparing the deceased group with the ≤28-day

MV group. The transporter 1, ATP binding cassette subfamily B member (TAP1) is also

upregulated and a key regulator (hub gene). Together with *TAP2*, *TAP1* plays a central

role in MHC I antigen presentation[33]. Transcriptional regulators *SP110* and *SP140*, both

ISGs and also identified as hub genes, were down-regulated. Module 718 was also

positively correlated with the SARS2-NWN (ρ = 0.31, P-value = 1.3e-3; enrichment FET

P-value = 0.029, 3.7 FE of M178 by differentially expressed genes in deceased vs ≤28-

days MV). The majority of genes in this module are down-regulated in the deceased

group compared with the ≤28-day group. Some of the genes encode subunits of the

mitochondrial ATP synthase, such as *ATP6* and *ATP8*, the cytochrome C oxidase, with

COX2 and *COX3* as well as the NADH dehydrogenase complex, such as *ND1-ND6*.

ND4L, ATP6, COX2, ND1, ND3, ND4L and *ND6* are key regulators, potentially

modulating the expression of the other genes in the module. These findings further

support mitochondria dysfunction[26], potentially disrupting processes indicated by the

module. Other down-regulated genes are humanin1 (*MTRNR2L1*) and R-spondin 1

(*RSPO1*). Humanin is known to protect against oxidative stress and mitochondrial

dysfunction[34]. *RSPO1* protects against cell stress by activating the Wnt/β-catenin

signaling pathway[35]. Non-coding RNAs, such as *MALAT1* and *RHOQ-AS*1 were found

to be up-regulated. *MALAT1* is known to suppress IRF3-initiated antiviral innate

immunity[36] while the function of *RHOQ-AS1* is unknown.

Metatranscriptome and Transcriptome signatures are predictive of mortality

We evaluated the strength of the metatranscriptomic, metagenomic and host

transcriptomic profiles to predict mortality in this cohort of critically ill COVID-19 patients.

To this end, we identified features in each of these datasets and constructed risk scores

that best predicted mortality. **Figure 6a** shows that the metatranscriptome data, alone or

combined with the other two datasets, was most predictive of mortality. Importantly, the

predictive power (as estimated by the area under the curve) of the metatranscriptome

data was improved by excluding probable contaminants and worsened when SARS-

CoV-2 was removed from the modeling. The selected features we used to construct the

metatranscriptome, metagenome and host transcriptome risk scores are reported in

Supplementary Table 11).   Using the means of the scores, we classified all subjects

into high risk and low risk groups for mortality. **Figure 6b** shows Kaplan-Meier survival

curve comparisons evaluating the predictive power of risk score stratification based on

metatranscriptome, metagenome and host transcriptome data. Combining risk scores

from different datasets showed an optimal identification of mortality when

metatranscriptome and host transcriptome were considered (**Figure 6c**). We then used

the gene signature found as being the most predictive of mortality to conduct IPA

analyses. Among the upstream regulators, mortality was associated with predicted

activation of interferon alpha while chemotaxis and infection by RNA virus were

predicted as activated in diseases and functions. These data highlight the importance of

SARS-CoV-2 abundance in the lower airways as a predictor for mortality, and the

significant contribution of the host cell transcriptome, which reflects the lower airway cell

response to infection.

Discussion

A limited number of studies to date have evaluated the lower airway microenvironment in patients with SARS-CoV-2 infection because of the increased risk of virus transmission to healthcare providers during sampling[19,20,37-42]. This has limited molecular investigations into the primary site of the disease. Having built a substantial biorepository of lower airway samples among COVID-19 patients on mechanical ventilation recruited during the first wave of SARS-CoV-2 infections in New York City, we used a metagenomic approach to characterize the microbiome in the lower airways and assessed its impact on clinically meaningful outcomes. In this analysis of 142 critically ill hospitalized patients with confirmed SARS-CoV-2 infection and lower airway biorepository samples available, we determined that higher SARS-CoV-2 viral load, higher relative abundance of *Mycoplasma salivarium*, and limited anti-SARS-CoV-2 Spike protein IgG response in the lower airways were associated with increased mortality. This signature was supported by the metatranscriptome data of the lower airway samples where SARS-CoV-2 sequence reads were significantly enriched in those patients who died compared to those who survived after developing respiratory failure requiring mechanical ventilation. Importantly, although we observed changes in other microbial components of the lower airway microbiome in our analysis of lower airway samples from 118 patients and by clinical laboratory culture results obtained from 589 patients, we did not find evidence to support the hypothesis that co-infection with common (bacterial, viral, fungi) respiratory pathogens was associated with poor outcome—although most patients received empiric treatment with broad spectrum antibiotics and anti-fungals.

Several studies have explored the relationship between SARS-CoV-2 viral load and

mortality[43-48]. Severe influenza requiring hospitalization has also been associated with

higher viral loads[49,50]. It has been argued that high viral load might merely be a

reflection of an individual's immune response[43]. In fact, in SARS-CoV-1, clinical

progression was not associated with increased viral load or uncontrolled viral replication

in the nasopharynx but rather with an upregulated immune profile in these patients[51]. In

a large cohort of 1145 patients with confirmed SARS-CoV-2, viral load measured in

nasopharyngeal swab samples was found to be significantly associated with mortality,

even after adjusting for age, sex, race and several co-morbidities [48]. Similar results were

found in a cohort of patients in New York City with or without cancer, where in-hospital

mortality was significantly associated with a high SARS-CoV-2 viral load in the upper

airways [47]. The data presented here through the use of direct quantitative methods

(RT-PCR) and a semiquantitative untargeted approach (metatranscriptome sequencing)

support the hypothesis that the SARS-CoV-2 viral load in the lower airways plays a

critical role in the clinical progression of critically ill COVID-19 patients. It is important to

note that current guidelines for treatment of COVID-19 do not recommend treatment

with remdesivir for patients receiving invasive mechanical ventilation[52]. The results of

this investigation suggest that antivirals might still have a role in the treatment of

critically ill COVID-19 patients.

We investigated the possibility that mortality with SARS-CoV-2 infection was related to

co-infection with other pathogens. To this point several investigations have shown

evidence of SARS-CoV-2 co-infection with other viruses, bacteria and fungi identified by

culture-based techniques [18,53-60]. In a cohort of 116 specimens positive for SARS-CoV-

2, 21% were positive for one or more additional respiratory pathogens including

rhinovirus/enterovirus and respiratory syncytial virus[53]. In a meta-analysis of 3,338

patients with COVID-19, only 3.5% of patients had an identified bacterial co-infection at

admission, while 14.3% were found later to have a secondary bacterial infection[55]. The

most common pathogens identified included species in the genera *Mycoplasma*,

Hemophilus, and *Pseudomonas*. In another study, the most commonly identified co-

infections were with *Streptococcus pneumoniae*, *Klebsiella pneumoniae*, and

Haemophilus influenzae[57]. Using detailed clinical laboratory culture data available for

589 subjects hospitalized with respiratory failure due to COVID-19, we showed that

higher rates of respiratory infection with other organisms, especially early in their

hospitalization, did not occur among subjects with poor clinical outcome. Further, we did

not observe an association between positive cultures for any pathogen tested and

increased odds of dying in critically ill COVID-19 patients.

In the subset of COVID-19 patients with BAL samples, we used NGS to identify all

potential pathogens and commensals in the lower airways beyond microbial cultures

routinely obtained as per clinical care. The RNA virome data showed that SARS-CoV-2

dominates the lower airways and was significantly associated with death. A small

number of samples had a few sequences that mapped to influenza A or B viruses,

suggesting that co-infection with influenza did not occur frequently during this first wave

of SARS-CoV-2 infections. Within the DNA virome, there was no significant difference in

viruses between the three outcome groups despite the frequent finding of HSV-1.

Similarly, when evaluating the metatranscriptome of DNA viruses, there were few

differences between the three outcome groups. Although analysis of the phage

metagenome data showed no differential enrichment between the three cohorts, we did

identify in the metatranscriptome data differentially active phages when comparing the

three cohorts, suggesting that changes in the bacterial microbiome may be occurring in

critically ill patients with COVID-19. Certain *Staphylococcus* phages were differentially

active in those who were ventilated for more than 28 days and in those who died.

Interestingly, the bacterial signatures also identified *Mycoplasma salivarium*, a known

oral commensal that has previously been associated with ventilator-acquired

pneumonia[61], as differentially active in those who died and those who were ventilated

for more than 28 days when compared to those ventilated less than 28 days. From

previous data published by us, enrichment of the lower airway microbiota with oral

commensals was seen to be associated with a pro-inflammatory state in several

diseases including lung cancer[62,63] and non-tuberculosis mycobacterium related

bronchiectasis[64].

With the use of metagenomic and metatranscriptomic analyses it is also possible to

examine how functionally active microbes impact the host[65]. In this cohort of patients,

we evaluated the functional profile of the microbiome within the lower airways and its

effect on mortality, something that, to our knowledge, had not yet been assessed in

COVID-19 patients. The only significant gene function enrichment was found with the

metatranscriptome data suggesting that functional activation of microbes can provide

further insights into the lower airway microbial environment of patients with worst

outcome. Among the pathways that were differentially expressed in those patients with

poor outcome, we identified genes associated with degradation, transport, and

antimicrobial resistance genes, as well as with signaling. These differences may

indicate important functional differences leading to a different metabolic environment in

the lower airways that could impact host immune responses. It could also be

representative of differences in microbial pressure in patients with higher viral loads and

different inflammatory environments.

In the current investigation, we also characterized the immune response within the

lower airways by measuring anti-SARS-CoV-2 Spike antibodies and profiling the host

RNA transcriptome. We observed that low levels of anti-Spike and anti-RBD IgG in the

lung were associated with poor outcome. Although we did not find a statistically

significant association between SARS-CoV-2 neutralizing capacity and poor outcome,

levels of SARS-CoV-2 neutralizing antibodies, anti-Spike and anti-RBD antibodies (both

IgG and IgA) were negatively correlated with SARS-CoV-2 viability. Prior investigations

have suggested that IgA levels are a key driver of neutralization in the mucosa[66-68]. The

differences noted in the current investigation in the IgG pools are intriguing and future

work investigating the antibodies generated during SARS-CoV-2 infections will be

essential.

When examining host transcriptomic differences across the different clinical outcome

groups, Sirtuin and Ferroptosis signaling pathways were found to be upregulated in the

most critically ill COVID-19 patients. Upregulation in the Sirtuin pathway demonstrates

an increased host inflammatory response to viral infection[23]. In addition, ferroptosis, a

recently identified form of non-apoptotic regulated cell death through iron-dependent

accumulation of lipid peroxides, has been shown to cause direct lung injury[69] or

pulmonary ischemia-reperfusion injury[70,71]. Interestingly, there is evidence to support

that STAT3[69] and ACSL4[70] alleviated ferroptosis-mediated acute lung injury

dysregulation, which are both down-regulated in COVID-19 patients with worse clinical

outcome. Further analysis showed that there appeared to be an inactivation of

phagocytes, neutrophils, granulocytes, and leukocytes, including downregulation of IgG

expression levels, with additional mitochondria dysfunction, and down-regulation of

Inositol related pathways and noradrenaline/adrenaline degradation. There is evidence

that in the neonatal lung, inositol related components exert an anti-inflammatory effect

and can prevent acute lung injury[72,73].

Collectively, these data suggest that an imbalance rather than an elevated inflammatory

state in the lung is an important marker that predicts poor outcomes in critically ill

COVID-19 patients. Indeed, the inferred cell composition analysis from the bulk

transcriptome data overall points to a tepid immune response. Memory T cells have

been implicated with a robust immune response in SARS-CoV-2.[74] The deficiency of

these memory T cells that we found in the lungs of COVID-19 patients with worse

outcome further supports the presence of an ineffective immune response or presence

of immune exhaustion. IL4I1, found in the network analysis to be up-regulated in the

deceased group in association with SARS-CoV-2, is an immunosuppression enzyme

that plays a role in infection and the control of immunopathology[75]. IL4I1 induction has

been reported in viral infections with influenza virus[76]. The ISGs and transcriptional

regulators *SP110* and *SP140*, both downregulated in the deceased group. play

important roles in resisting intracellular pathogens[77].

Strikingly, interrogation of the host transcriptomic analysis identified survival-associated

differences in interferon-related responses. Our host transcriptomic risk stratified model

seems to point to a predictive activation of type I interferon as a prediction for mortality.

This might be inconsistent with the current suggestion that, based on systemic levels,

early interferon responses are associated with poor outcome in COVID19.[78,79] Others

have suggested that a robust interferon response may lead to a hyperinflammatory

state that could be detrimental in the disease process, justifying the use of Janus kinase

inhibitor inhibitors in patients with COVID-19.[80] Studies comparing transcriptomic

signatures in BAL of patients with severe COVID-19 and controls have shown activation

of type 1 interferons.[81] While further longitudinal data will be needed to clarify the role of

interferon signaling on the disease, the data presented here suggest that combining

microbial and host signatures could help understand the increase risk for mortality in

critically ill COVID-19 patients.

By collecting BAL samples rather than endotracheal aspirate specimens we were able

to ensure extensive sampling of the lower respiratory tract in intubated patients.

However,

we were limited to samples from intubated patients in whom a clinically indicated

bronchoscopy was done to place a percutaneous tracheostomy or for airway clearance.

Although this included a large number of patients with various clinical outcomes, those

sampled may not be representative of the extremes in the spectrum of disease severity

who were most likely not eligible for bronchoscopy. For example, patients that

presented with very rapid clinical deterioration and died within the first few days of

hospitalization or those who were quickly weaned from mechanical ventilation did not

receive bronchoscopy. However, extensive and detailed clinical data were also obtained

from intubated COVID-19 patients without bronchoscopy performed within the

Manhattan Campus (no bronchoscopy cohort) and from the Long Island cohort for

whom bronchoscopies were done without collecting research samples. In both of these

cohorts, clinical laboratory culture data did not identify untreated secondary pathogen

infections associated with poor outcome.

The samples used in this investigation were obtained during the first surge of cases of

COVID-19 in New York City, and management reflected clinical practices at that time.

Among the differences with current therapeutic approaches in COVID-19 patients,

corticosteroids and remdesivir, two medications that likely affect the lower airway

microbial landscape, were rarely used during the first surge. Other medications, such as

antibiotics and anti-inflammatory drugs could affect our findings and we therefore

considered them as potential confounders. However, the use of these medications was

not found to be associated with clinical outcome. The cross-sectional study design

precluded evaluation of the temporal dynamics of the microbial community or the host

immune response in this cohort, which could provide important insights into the

pathogenesis of this disease. Performing repeated bronchoscopies without a clinical

indication would be challenging in these patients and other less invasive methods might

need to be considered to study the lower airways at earlier timepoints and serially over

time in patients with respiratory failure. It is important to note that there were no

statistically significant differences in the timing of sample collection across the three

outcome groups.

In summary, we present here the first evaluation of the lower airway microbiome using a

metagenomic and metatranscriptomic approach, along with host immune profiling in

critically ill patients with COVID-19 requiring invasive mechanical ventilation. The RNA

metatranscriptome analysis showed an association between the abundance of SARS-

CoV-2 and mortality, consistent with the signal found when viral load was assessed by

targeted rRT-PCR. These viral signatures correlated with lower anti-SARS-CoV-2 Spike

IgG and host transcriptomic signatures in the lower airways associated with poor

outcome. Importantly, both through culture and NGS data, we did not find evidence for

an association between untreated infections with secondary respiratory pathogens and

mortality. Together, these data suggest that active lower airway SARS-CoV-2

replication and poor SARS-CoV-2-specific antibody responses are the main drivers of

increased mortality in COVID-19 patients requiring mechanical ventilation. The potential

role of oral commensals such as *Mycoplasma salivarium* need to be explored further. It

is possible that *M. salivarium* can impact key immune cells and has recently been

reported at a high prevalence in patients with ventilator-acquired pneumonia[61]. Critically,

our finding that SARS-CoV-2 evades and/or derails effective innate/adaptive immune

responses indicates that therapies aiming to control viral replication or induce a targeted

antiviral immune response may be the most promising approach for hospitalized

patients with SARS-CoV-2 infection requiring invasive mechanical ventilation.

Methods

Subjects

Enrolled subjects were 18 years or older, admitted to the intensive care units (ICUs) at NYU Langone Health from March 10[th] to May 10[th], 2020 with a nasal swab confirmed diagnosis of SARS-CoV-2 infection by reverse transcriptase polymerase chain reaction (RT-PCR) assay and respiratory failure requiring invasive mechanical ventilation. Samples were obtained during clinically indicated bronchoscopy performed for airway clearance or for percutaneous tracheostomy placement. Surviving subjects signed informed consent to participate in this study. Samples and metadata from subjects who died or were incapacitated were de-identified and included in this study. Comprehensive demographic and clinical data were collected. We also collected longitudinal data on clinical laboratory culture results and treatment. **Supplementary figure 1** shows the distribution of subjects and sampling strategy used for this study. The study protocol was approved by the Institutional Review Board of New York University.

Lower airway bronchoscopic sampling procedure

Both background and supraglottic (buccal) samples were obtained prior to the procedure, as previously described[62]. The background samples were obtained by passing sterile saline through the suctioning channel of the bronchoscope prior to the procedure. Bronchoalveolar lavage (BAL) samples were obtained from one lung segment as per discretion of the treating physician as clinically indicated. Samples were then transferred to a BSL3 laboratory for processing. Once there, 2 mL of whole BAL was stored in a tube prefilled with 2 mL of Zymo Research's DNA/RNA Shield™ (R1100-250, https://www.zymoresearch.com/pages/covid-19-efforts) for RNA/DNA

preservation and virus inactivation. In addition, background control samples (saline

passed through the bronchoscope prior to bronchoscopy) and supraglottic aspirates

were stored in the same RNA/DNA shield. A subset of samples underwent BAL cell

separation by centrifugation and cells were cryopreserved in DMSO while acellular BAL

fluid was aliquoted for cytokine measurements. A paired blood sample was also

obtained in EDTA tubes (Becton Dickinson, ref# 366450) and PAXgene Blood RNA

tubes (PreAnalytiX) ref# 762165).

Viral load detection targeting the N gene

SARS-CoV-2 viral load was measured by quantitative real-time reverse transcription

polymerase chain reaction (rRT-PCR) targeting the SARS-CoV-2 nucleocapsid (N)

gene and an additional primer/probe set to detect the human RNase P gene (RP).

Assays were performed using Thermo Fisher Scientific (Waltham, MA) TaqPath 1-Step

RT-qPCR Master Mix, CG (catalog number A15299) on the Applied Biosystems (Foster

City, CA) 7500 Fast Dx RealTime PCR Instrument. Using the positive controls provided

by the CDC, which are normalized to 1000 copies/mL, we converted the different Ct

positive to copies/mL. This was done using the DDCT method, applying the formula:

Power [2, (CT (sample, N1 gene) - CT (PC, N1 gene)] – [CT (sample, RP gene) - CT

(PC, RP gene)]*1000.

SARS-CoV-2 viral viability through measurement of subgenomic transcripts

Viral subgenomic mRNA (sgRNA) is transcribed in infected cells and is not packaged

into virions. Thus, presence of sgRNA is indicative of active infection of a mammalian

cell in samples. We therefore measure sgRNA in all BAL samples obtained targeting the

E gene as previously described.[21] Briefly, five µl RNA was used in a one-step real-time

RT-PCR assay to sgRNA (forward primer 5'- CGATCTCTTGTAGATCTGTTCTC-3';

reverse   primer   5'-   ATATTGCAGCAGTACGCACACA-3';   probe   5'-FAM-

ACACTAGCCATCCTTACTGCGCTTCG-ZEN-IBHQ-3') and using the Quantifast Probe

RT-PCR kit (Qiagen) according to instructions of the manufacturer. In each run,

standard dilutions of counted RNA standards were run in parallel to calculate copy

numbers in the samples.

DNA/RNA isolation, library preparation and sequencing

DNA and RNA were isolated in parallel using zymoBIOMICS™ DNA/RNA Miniprep Kit

(Cat: R2002) as per manufacturer's instructions. DNA was then used for whole genome

shotgun (WGS) sequencing using it as input into the NexteraXT library preparation kit

following the manufacturer's protocol. Libraries were purified using the Agencourt

AMPure XP beads (Beckman Coulter, Inc.) to remove fragments below 200 bp. The

purified libraries were quantified using the Qubit dsDNA High Sensitivity Assay kit

(Invitrogen) and the average fragment length for each library was determined using a

High Sensitivity D1000 ScreenTape Assay (Agilent). Samples were added in an

equimolar manner to form two sequencing pools. The sequencing pools were quantified

using the KAPA Library Quantification Kit for Illumina platforms. The pools were then

sequenced on the Illumina Novaseq 6000 in one single run. For RNA sequencing, RNA

quantity and integrity were tested with a BioAnalyzer 2100 (Agilent). Among

bronchoscope control (BKG) samples, only 5 yielded RNA with sufficient quality and

quantity to undergo library preparation and sequencing. The automated Nugen Ovation

Trio Low Input RNA method was used for library prep with 3ng total RNA input of each

sample. After 6 amplification cycles, samples were sequenced using 2x Novaseq 6000

S4 200 cycle Flowcells using PE100 sequencing.

Microbial community characterization using whole genome shotgun sequencing (WGS)

and RNA metatranscriptome

For all metagenomic and metatranscriptomic reads, Trimmomatic v0.36[82], with leading

and trailing values set to 3 and minimum length set to 36, was used to remove adaptor

sequences. All rRNA reads were then removed from the metatranscriptomic reads using

SortMeRNA v4.2.0[83] with default settings. Metagenomic and filtered metatranscriptomic

reads were mapped to the human genome using Bowtie2 v2.3.4.1[84] with default settings

and all mapping reads were excluded from subsequent microbiome, mycobiome, and

virome metagenomic and metatranscriptomic analysis. Technical replicates for each

biological sample were pooled together for subsequent analyses. Taxonomic profiles for

all metagenomic and metatranscriptomic samples were generated using Kraken

v2.0.7[85] and Bracken v2.5 [https://doi.org/10.7717/peerj-cs.104] run with default

settings. The database used for quantifying taxonomic profiles was generated using a

combined database containing human, bacterial, fungal, archaeal, and viral genomes

downloaded from NCBI RefSeq on January 8, 2021. Additionally, genomes for Candida

auris (Genbank: GCA_003013715.2, GCA_008275145.1) and Pneumocystic jirovecii

(Genbank: GCA_001477535.1) were manually added to the database. Differentially

abundant bacterial and viral taxa were identified for the BAL and UA samples groups

individually using DESeq2 v1.28.1[86] with the three group clinical outcome meta-data

readouts set as the sample groupings. Significantly differentially abundant taxa

contained at a minimum an aggregate of 5 reads across samples and had an FDR

<0.2[87,88].

For functional microbial profiling, processed sequencing reads were further depleted of

human-mapping reads by removing all reads classified as human by Kraken v2.0.7[85]

using KrakenTools v0.1-alpha (https://github.com/jenniferlu717/KrakenTools). FMAP

v0.15[89] was run on both the metagenomic and metatranscriptomic reads to profile the

metabolic pathways present in each sample. FMAP_mapping.pl paired with diamond

v0.9.24[90] and FMAP_quantification.pl were used with default settings to identify and

quantify proteins in the Uniref90 database. Using DESeq2 v1.28.1[86], differentially

expressed genes were identified for the BAL samples individually using the three group

clinical outcome-metadata readouts for all genes that had an aggregate 5 reads across

all samples.

Antibiotic resistance genes were quantified in all metagenome and metatranscriptome

samples using Salmon v1.3.0[91] run with --keepDuplicates for indexing and --libtype A --

allowDovetail --meta for quantification. Genes were filtered such that only genes that

actively conferred antibiotic resistance were kept. To assess differentially expressed

classes of antibiotic resistance genes, gene counts for individual antibiotic resistance

genes were collapsed by their conferred antibiotic resistance.

Supplementary Figure 1 shows a summary of depth achieved with the parallel WGS

and metatranscriptome approach across sample types and the number of reads

assigned to different microbial subfractions (bacteria, fungi, DNA viruses, RNA viruses

and phages). Further analysis was also done to identify possible contaminants in the

metatranscriptome and metagenome datasets. To this end, we compared the relative

abundance of taxa between background bronchoscope control and BAL samples. Taxa

with median relative abundance greater in background than in BAL were identified as

probably contaminant and listed in **Supplementary Table 4).** None of the taxa identified

as possible contaminants were removed from the analyzed data but are shown for

comparison with signatures identified in the rest of the analyses.

Anti-Spike SARS-CoV-2 antibody profiling in BAL

BAL samples were heat-treated at 56°C for one hour, and centrifuged at 14000g for 5

min. The supernatant was collected and diluted 50-fold in PBST containing 1% skim

milk. The diluted samples were incubated at room temperature (R.T.) for 30 min with

QBeads DevScreen: SAv (Streptavidin) (Sartorius 90792) that had been loaded with

biotinylated Spike, biotinylated RBD or biotin (negative control) in wells of a 96 well HTS

filter plate (MSHVN4550). As positive controls, we used CR3022 antibody, that

recognizes SARS-CoV-2 Spike and RBD, in human IgG, IgA and IgM formats (Absolute

Antibody). After washing the beads, bound antibodies were labeled with anti IgG-

DyLight488, anti IgA-PE and anti IgM-PECy7, and the fluorescence intensities were

measured in Intellicyt IQue3 (Sartorius). The acquired data [median fluorescence

intensity (MFI)] were normalized using the MFI values of the CR3022 antibodies to

compensate for variations across plates. **Supplementary Figure 10** shows that the

levels of these antibodies were higher in BAL samples of patients with SARS-CoV-2

than in BAL samples from 10 uninfected healthy smokers recruited for research

bronchoscopy. Details of method development and validation will be described

elsewhere (Koide et al. in preparation).

SARS-CoV-2 preparation and neutralization assay

icSARS-CoV-2-mNG (isolate USA/WA/1/2020, obtained from the UTMB World

Reference Center for Emerging Viruses and Arboviruses) was amplified once in Vero E6

cells (P1 from the original stock). Briefly, 90-95% confluent T175 flask (Thomas

Scientific) of Vero E6 ($1x10^7$ cells) was inoculated with 50 µL of icSARS-CoV-2-mNG in

5 mL of infection media (DMEM, 2% FBS, 1% NEAA, and 10 mM HEPES) for 1 hour.

After 1 hour, 20 mL of infection media was added to the inoculum and cells were

incubated 72 hours at 37 °C and 5% CO_2. After 72 hours, the supernatant was collected

and the monolayer was frozen and thawed once. Both supernatant and cellular fractions

were combined, centrifuged for 5 min at 1200 rpm, and filtered using a 0.22 µm Steriflip

(Millipore). Viral titers were determined by plaque assay in Vero E6 cells. In brief,

220,000 Vero E6 cells/well were seeded in a 24 well plate, 24 hours before inoculation.

Ten-fold dilutions of the virus in DMEM (Corning) were added to the Vero E6

monolayers for 1 hour at 37 °C. Following incubation, cells were overlaid with 0.8%

agarose in DMEM containing 2% FBS (Atlanta biologicals) and incubated at 37 °C for

72 h. The cells were fixed with 10% formalin, the agarose plug removed, and plaques

visualized by crystal violet staining. All procedures including icSARS-CoV-2-mNG virus

were performed using Biosafety Level 3 laboratory conditions.

1007

1008 For SARS-CoV-2 neutralization assays, Vero E6 cells (30,000 cells/well) were seeded

1009 in a 96 well plate 24 h before infection. Two-fold serial dilutions of BAL lysates were

mixed with mixed 1:1 (vol/vol) with SARS-CoV-2 mNG virus (multiplicity of infection,

MOI 0.5), and incubated for 1 h at 37 °C. After incubation, 100 µL of the mixtures of the

antibody and SARS-CoV-2 mNG were added to the Vero E6 monolayers, and cells

were incubated at 37°C. After 20 h, cells were fixed with 4 % formaldehyde (Electron

Microscopy Sciences) at room temperature for 1 h. After fixation, cells were washed

twice with PBS and permeabilized with 0.25% triton-100, stained with DAPI (Thermo),

and quantified on a CellInsight CX7 High-content microscope (Thermo) using a cut-off

for three standard deviations from negative to be scored as an infected cell.

Transcriptome of BAL cells

RNA-Seq was performed on bronchial epithelial cells obtained by airway brushing, as

described[92-94], using the Hi-seq/Illumina platform at the NYU Langone Genomic

Technology Center (data available at Sequence Read Archive: # PRJNA592149).

KEGG[95,96] annotation was summarized at levels 1 to 3. Genes with an FDR-corrected

adjusted p-value <0.25 were considered significantly differentiated, unless otherwise

specified. Pathway analysis using differentially regulated genes (FDR<0.25) was done

using Ingenuity Pathway Analysis, RRID:SCR_0- at least 1 count per million in at least

two samples were retained. For digital cytometry with CIBERSORTx, a signature matrix

derived from single-cell transcriptome of BAL cells collected from patients with COVID-

19[31] was first generated with the "Create Signature Matrix" module in the CIBERSORTx

online tool. A maximum of 10 cells per cell type per patient were initially sampled from

the original data and 20 cells per cell type were then used to build the single-cell

reference with the default parameters. Then the "Impute Cell Fractions" module was

used to estimate the absolute cell fraction score of different cell types in bulk

transcriptomes using the single-cell signatures with "S-mode" batch correction and 100

permutations in the absolute mode. Bulk transcriptomes with a significant deconvolution

p-value (≤0.05) were retained. For xCell cell type signature enrichment analysis, the

enrichment scores were inferred with built-in signature of cell types detected in the BAL

samples as reported previously [31]. The two-tailed Wilcoxon rank sum test with

Benjamini-Hochberg correction were computed between groups of samples for

comparison.

1041

Microbial and Host predictive modeling

Cox proportional hazards model was used for investigating the association between the

time to death and the relative abundance of each taxon quantified using

metatranscriptomic and metagenomic data separately. We first performed the univariate

screening test to identify significant features associated with the time to death using the

Cox proportion hazards regression model for the relative abundance of taxa from the

RNA and DNA data, and log-transformed count of host transcriptome data, respectively.

Within each type of data, given the p-value cutoff, the features with a p-value less than

the cutoff were selected and integrated as a sub-community. For the RNA and DNA

data, the alpha diversity (Shannon index) was calculated for each sample on the

selected sub-community and the negative of the value was defined as the microbial risk

score, because high alpha diversity indicates low risk of death.   For the host

transcriptome data, the log-transformed total count of all selected candidate

transcriptome for each sample was defined as the risk score, since most selected

candidate transcriptomes increased the risk of death. The leave-one-out cross-

validation (LOOCV) was used for the predictions. The p value cutoff was set at the

value which produces the largest AUC (area under the receiver operating characteristic

curve) in predicting the death/survival status using the risk score we constructed over

these features. The additive model was used to integrate when more than one scores

are used for the prediction.

1062

Multiscale and co-expression network analyses

Raw counts from the human transcriptome were normalized and converted to log2-

counts per million using limma[97]/voom[98] (v3.44.1 with R v4.0.0) with standard

parameters. Microbiome abundance information was converted to relative abundance.

Low abundance taxa were removed based on average abundance across all samples to

yield a minimum of 1000 taxa for each metatranscriptome dataset. All datasets were

batch adjusted. Differentially expressed genes (DEGs) and differentially abundant taxa

were called using the DESeq2 package[86] (v1.28.1), based on the negative binomial (i.e.

Gamma-Poisson) distribution. According to the recommendation by the authors, we

used non-normalized data (i.e. raw gene counts and abundance data), as DESeq2

internally corrects data and performs normalization steps. For this purpose, raw

microbiome abundance data were converted to DESeq2 dds objects using the phyloseq

R library (V1.32.0). Contrasts are based on outcome groups (≤ 28 days MV, > 28 days

MV or death). Differentially expressed genes and differentially abundant tax with FDR of

0.2 or less are considered significant.

Multiscale Embedded Gene Co-Expression Network Analysis (MEGENA) [32] was

performed to identify host modules of highly co-expressed genes in SARS-CoV-2

infection. The MEGENA workflow comprises four major steps: 1) Fast Planar Filtered

Network construction (FPFNC), 2) Multiscale Clustering Analysis (MCA), 3) Multiscale

Hub Analysis (MHA), 4) and Cluster-Trait Association Analysis (CTA). The total

relevance of each module to SARS-CoV-2 infection was calculated by using the Product

of Rank method with the combined enrichment of the differentially expressed gene

(DEG) signatures as implemented: $G_j = \prod_i g_{ji}$, where, g_{ji} is the relevance of a

consensus **j** to a signature **i;** and g_{ji} is defined as $(max_j(r_{ji}) + 1 - r_{ji})/\sum_j r_{ji}$, where r_{ji}

is the ranking order of the significance level of the overlap between the module **j** and the

signature.

1089

To functionally annotate gene signatures and gene modules derived from the host

transcriptome data, we performed an enrichment analysis of the established pathways

and signatures—including the gene ontology (GO) categories and MSigDB. The hub

genes in each subnetwork were identified using the adopted Fisher's inverse Chi-

square approach in MEGENA; Bonferroni-corrected p-values smaller than 0.05 were set

as the threshold to identify significant hubs. The correlation between modules, modules

and clinical traits as well as modules and individual taxa were performed using

Spearman correlation. Other correlation measures, such as Pearson correlation or the

Maximal Information Coefficient (MIC)[99] proved to be inferior for this task. Categorical

trait data was converted to numerical values as suitable.

1100

1101   *Data availability*

1102   Sequencing data are available in NCBI's Sequence Read Archive under project

1103   numbers PRJNA688510 and PRJNA687506 (RNA and DNA sequencing, respectively).

1104   Codes used for the analyses presented in the current manuscript are available at

1105   *https://github.com/segalmicrobiomelab/SARS_CoV2.*

Figure 1. Associations between culture positivity and clinical outcome. Odds ratios and corresponding 95% confidence intervals for rates of culture positivity for the whole cohort (n=589) during the length of their hospitalization (left) and during the first 2 weeks of hospitalization (right).

Figure 2. SARS-CoV-2 viral load and virus metatranscriptome analyses. Copies of the SARS-CoV-2 N gene per ml, normalized by the Human RNase P gene, comparing paired upper and lower airway samples **(a)** and levels in BAL comparing clinical outcome groups **(b**, *= Mann–Whitney U $p<0.05$, **= Mann–Whitney U $p<0.01$**). **(c)** PCoA analysis based on Bray Curtis Dissimilarity index of BAL Metatranscriptome data comparing clinical outcome (PERMANOVA p-value). Bubble plot showing DESeq results of RNA viruses **(d)** and expressed DNA phages **(e)** enriched in each clinical outcome comparisons (bubble size based on median relative abundance for those found statistically significant).

Figure 3. Bacteria load and taxonomic compositional analyses. (a) Bacterial load measured by ddPCR targeting 16S rRNA gene (**= Mann–Whitney U $p<0.01$). PCoA analysis based on Bray Curtis Dissimilarity index of BAL Metagenome **(b)** and Metatranscriptome **(c)** data comparing clinical outcome (PERMANOVA p-value). **(d)** Gene Set Enrichment Analysis (GSEA) was used to compare the taxonomic signatures identified in BAL metagenome (diamonds) and metatranscriptome (circles) as distinctly enriched for comparisons between clinical outcome groups (differential enrichment

performed based on DESeq2 analysis). **(e)** Bubble plot showing DESeq results of bacteria found concordantly differentially enriched between clinical outcome groups (bubble size based on median relative abundance for those found statistically significant).

Figure 4. Functional microbial compositional analyses. KOs were summarized to associated pathways and differential expression was calculated based on DESeq2 analysis. **(a)** Gene Set Enrichment Analysis (GSEA) was used to compare the functional signatures identified in BAL metagenome and metatranscriptome as distinctly enriched for comparisons between clinical outcome groups. **(b)** Bubble plot showing DESeq results of microbial functions found concordantly differentially enriched between clinical outcome groups (bubble size based on median relative abundance for those found statistically significant).

Figure 5. Lower airway host immune profiling in severely ill COVID-19. (a) Levels of anti-SARS-CoV-2 Spike antibodies in BAL (*= Mann–Whitney U p<0.05)**. (b)** Heat-map of canonical pathway analysis based on Ingenuity Pathway Analysis (IPA, RRID:SCR_008653) using the lower airway host transcriptome comparing clinical outcome groups. Orange shows up-regulation of pathway, blue shows down-regulation of pathway**. (c)** Cell type abundance quantification plots. Comparison of abundance of mast cells and neutrophils among outcome groups in the BAL fluids of critically ill patients with COVID-19. Cell type abundance was estimated from the host transcriptome with CIBERSORTx. Each dot denotes the quantification score of a sample and boxes depict median and inter-quartile range (*= Mann–Whitney U p<0.05).

1155

Figure 6. Mortality predictive power of metatranscriptome, metagenome and host

transcriptome. (a) Area under the curved median and confidence interval for receiver

operating characteristic curve analyses calculated from each sequencing datasets as

predictor and mortality as outcome. **(b)** Kaplan-meier survival analyses based on a

cutoff value estimated from features selected from each sequencing dataset. The "High

risk" and "Low risk" groups is the mean of predicted risk scores in all samples. **(c)**

Scatterplot among risk scores from metatranscriptome, metagenome, and host

transcriptome. Dotted line denotes the mean of the risk scores across all subjects,

which is also the threshold for dividing the samples into "High risk" and "Low risk"

groups. **(d)** IPA analyses of host transcriptomic signatures identified as most predictive

of mortality.

1167

1168

1169

 Supplementary Figure Legends:

 Supplementary Figure 1. Description of patient cohort, samples obtained,

 analyses performed and sequencing depth.

 Supplementary Figure 2. SARS-CoV-2 viral load in upper airway samples. Copies

 of the SARS-CoV-2 N gene per ml, normalized by the Human RNase P gene, in upper

 airways comparing clinical outcome groups **(**Mann–Whitney U p-value**)**.

 Supplementary Figure 3. Identification of top taxa found in background samples

 as compared with BAL and upper airway samples. Boxplots showing the relative

 abundance values in log10 relative abundance of taxa ranked ordered based on

 dominance of Background bronchoscope control samples and compared to abundances

 in BAL and Upper Airway samples within metatranscriptome **(a)** and metagenome **(b)**

 data. Red labels indicate taxa where relative abundance is higher in background

 samples than in BAL and therefore considered possible contaminant.

 Supplementary Figure 4. Topographical analyses of Metatranscriptome data.

 Comparison of alpha diversity (Shannon Index, **a**) and beta diversity (Bray Curtis

 Dissimilarity index, **b**) across background negative controls (bronchoscope),

 bronchoalveolar lavage (BAL) and upper airway (UA) samples (Kruskal-Wallis and

 PERMANOVA p-values, respectively). **(c)** Boxplots showing the relative abundance

 values in log10 across all metatranscriptome samples for the BAL and Upper Airway

 samples. The 50 taxa with the highest relative abundance values in the BAL

 metatranscriptome data are displayed; the top 10 in the BAL are highlighted in bold.

Each column consists of four plots displaying in order from top to bottom, the most abundant RNA vertebrate viruses, DNA phages, bacteria, and fungi identified (from top to bottom). Numbers in parentheses next to the taxa labels display the ranking in relative abundance for either the BAL or UA metatranscriptome samples, respectively.

Supplementary Figure 5. Topographical analyses of Metagenome Data. Comparison of alpha diversity (Shannon Index, **a**) and beta diversity (Bray Curtis Dissimilarity index, **b**) across background negative controls (bronchoscope), bronchoalveolar lavage (BAL) and upper airway (UA) samples (Kruskal-Wallis and PERMANOVA p-values, respectively)**.** (c) Boxplots showing the relative abundance values in log10 across all metagenome samples for the BAL and Upper Airways. The 50 taxa with the highest relative abundance values in the BAL metagenome are displayed; the top 10 in the BAL are highlighted in bold. Each column consists of two plots displaying the most abundant bacteria and fungi identified. Numbers in parentheses next to the taxa labels displays its ranking in relative abundance for either the BAL or UA metagenome samples, respectively.

Supplementary Figure 6. Evaluation of associations between the lower airway RNA virome and clinical outcome. Comparisons between the three clinical outcome groups was performed for α diversity (Shannon Index, Kruskal-Wallis p-value, left panel), β diversity (based on Bray Curtis Dissimilarity Index, PERMANOVA p-value, right panel).

1218 **Supplementary Figure 7. Topographical analyses of the bacterial load.** Bacterial

1219 load measured by ddPCR targeting 16S rRNA gene in background bronchoscope

1220 controls (BKG), lower airway (BAL) and upper airway (UA) samples.

1221

Supplementary Figure 8. Evaluation of associations between the lower airway

mycobiome and clinical outcome. Fungal taxonomic data was subtracted from

metagenome and metatranscriptome data from lower airway samples. **(a)** Comparisons

between the three clinical outcome groups was performed for α diversity (Shannon

Index, Kruskal-Wallis p-value, left panel), β diversity (based on Bray Curtis Dissimilarity

Index, PERMANOVA p-value, right panel) on metagenome data. **(b)** Bubble plot

showing DESeq results of fungi enriched in each clinical outcome comparisons based

on metagenome data (bubble size based on median relative abundance for those found

statistically significant). **(c)** Comparisons between the three clinical outcome groups was

performed for α diversity (Shannon Index, Kruskal-Wallis p-value, left panel), β diversity

(based on Bray Curtis Dissimilarity Index, PERMANOVA p-value, right panel) on

metatranscriptome data. **(d)** Bubble plot showing DESeq results of fungi enriched in

each clinical outcome comparisons based on metatranscriptome data (bubble size

based on median relative abundance for those found statistically significant).

1236

Supplementary Figure 9. Evaluation of associations between the lower airway

antibiotic resistance genes and clinical outcome. Bubble plot showing DESeq

results of summarized categories of antibiotic resistant microbial genes taken from

MEGARes for the metagenome (top) and metatranscriptome (bottom) data sets for

each clinical outcome comparison (bubble size based on median relative abundance for

those found to be statistically significant). Colored bubbles indicate significantly

enriched antibiotic resistance groups.

1244

Supplementary Figure 10. Measurement of anti-SARS-CoV-2 Immunoglobulin

levels and neutralization activity. Levels of anti-SARS-CoV-2 Spike **(a) and** anti-

SARS-CoV-2 receptor binding domain (RBD, **b**) antibodies in BAL from non SARS-

CoV-2 infected smoker controls and severely ill COVID-19 intubated patients. Note that

the signals for different isotypes cannot be compared because they are detected with

different reagents. **(c)** Comparisons of levels of anti-SARS-CoV-2 RBD antibodies in

BAL across subjects in different clinical outcome groups (*= Mann–Whitney U $p<0.05$)**.**

(d) Neutralizing activity in BAL samples across subjects in different clinical outcome

groups.

1254

1255 **Supplementary Figure 11. Evaluation for associations between the lower airway**

1256 **host tanscriptome and clinical outcome. (a)** PCoA (based on Bray Curtis

1257 Dissimilarity Index, PERMANOVA p-value) comparing the three clinical outcome

1258 groups. (b, c, d) Volcano plot comparing lower airway host transcriptome between the

1259 three clinical outcome groups.

1260

Supplementary Figure 12. Multi-scale cross-kingdom and co-expression

networks. (a) The neighborhood 5 cross-kingdom metatranscriptome network centered

around SARS-CoV-2 is shown. Nodes refer to taxa, edges denote co-abundance after

MEGENA. The size of the nodes indicates abundance. Taxa with large nodes are highly

abundant. Node-shapes are according to the legend and refer to different microbial

kingdoms. The differential abundance of taxa in log2(fold change) between the

deceased group and the ≤28-day MV groups is shown by node color - red nodes are

taxa abundant in the deceased group compared to the ≤28-day MV group, blue colored

nodes denote the opposite. **(b)** Modules M175 and M718 of the host transcriptome are

shown. The node size refers to the absolute gene expression value. Nodes with wide

node border refer to key regulators/hub genes (see Methods). The differential gene

expression of taxa in log2(fold change) between the deceased group and the ≤28-day

MV groups is shown by node color - red nodes are up-regulated in the deceased group

compared to the ≤28-day MV group, blue colored nodes denote the opposite.

1275

Reference:

1. The, L. Emerging understandings of 2019-nCoV. *Lancet* **395**, 311 (2020).
2. WHO coronavirus disease (COVID-19) dashboard. *Geneva: World Health Organization, Available online: https://covid19.who.int/* (2020).
3. Rabaan, A.A., *et al.* SARS-CoV-2, SARS-CoV, and MERS-COV: A comparative overview. *Infez Med* **28**, 174-184 (2020).
4. Cao, X. COVID-19: immunopathology and its implications for therapy. *Nat Rev Immunol* **20**, 269-270 (2020).
5. Arentz, M., *et al.* Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA* **323**, 1612-1614 (2020).
6. Bhatraju, P.K., *et al.* Covid-19 in Critically Ill Patients in the Seattle Region - Case Series. *N Engl J Med* **382**, 2012-2022 (2020).
7. Cummings, M.J., *et al.* Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *The Lancet* **395**, 1763-1770 (2020).
8. European Society of Anesthesiology. Analysis of COVID-19 data on numbers in intensive care from Italy: European Society of Anaesthesiology (ESA). Vol. 2020 (2020).
9. Grasselli, G., *et al.* Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA* **323**, 1574-1581 (2020).
10. Intensive Care National Audit and Research Center UK. ICNARC report on COVID-19 in Critical Care., Vol. 2020 (2020).
11. Richardson, S., *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* **323**, 2052-2059 (2020).
12. Yang, X., *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* **8**, 475-481 (2020).
13. Morens, D.M. & Fauci, A.S. The 1918 influenza pandemic: insights for the 21st century. *J Infect Dis* **195**, 1018-1028 (2007).
14. Shieh, W.J., *et al.* 2009 pandemic influenza A (H1N1): pathology and pathogenesis of 100 fatal cases in the United States. *Am J Pathol* **177**, 166-175 (2010).
15. Dickson, R.P., *et al.* Enrichment of the lung microbiome with gut bacteria in sepsis and the acute respiratory distress syndrome. *Nat Microbiol* **1**, 16113 (2016).
16. Kitsios, G.D., *et al.* Respiratory Tract Dysbiosis Is Associated with Worse Outcomes in Mechanically Ventilated Patients. *Am J Respir Crit Care Med* **202**, 1666-1677 (2020).
17. Dickson, R.P., *et al.* Lung Microbiota Predict Clinical Outcomes in Critically Ill Patients. *Am J Respir Crit Care Med* **201**, 555-563 (2020).
18. Zuo, T., *et al.* Alterations in Fecal Fungal Microbiome of Patients With COVID-19 During Time of Hospitalization until Discharge. *Gastroenterology* **159**, 1302-1310 e1305 (2020).
19. Chen, L., *et al.* RNA based mNGS approach identifies a novel human coronavirus from two individual pneumonia cases in 2019 Wuhan outbreak. *Emerg Microbes Infect* **9**, 313-319 (2020).
20. Shen, Z., *et al.* Genomic Diversity of Severe Acute Respiratory Syndrome-Coronavirus 2 in Patients With Coronavirus Disease 2019. *Clin Infect Dis* **71**, 713-720 (2020).
21. Wolfel, R., *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**, 465-469 (2020).
22. Yazdi, M., Bouzari, M. & Ghaemi, E.A. Genomic analyses of a novel bacteriophage (VB_PmiS-Isfahan) within Siphoviridae family infecting Proteus mirabilis. *Genomics* **111**, 1283-1291 (2019).
23. Budayeva, H.G., Rowland, E.A. & Cristea, I.M. Intricate Roles of Mammalian Sirtuins in Defense against Viral Pathogens. *J Virol* **90**, 5-8 (2016).

1323   24.   Dar, H.H., *et al.* Pseudomonas aeruginosa utilizes host polyunsaturated
1324   phosphatidylethanolamines to trigger theft-ferroptosis in bronchial epithelium. *J Clin Invest* **128**,
1325   4639-4653 (2018).
1326   25.   Stoyanovsky, D.A., *et al.* Iron catalysis of lipid peroxidation in ferroptosis: Regulated enzymatic
1327   or random free radical reaction? *Free Radic Biol Med* **133**, 153-161 (2019).
1328   26.   Burtscher, J., Cappellano, G., Omori, A., Koshiba, T. & Millet, G.P. Mitochondria: In the Cross Fire
1329   of SARS-CoV-2 and Immunity. *iScience* **23**, 101631 (2020).
1330   27.   Smati, S., *et al.* Relationship between obesity and severe COVID-19 outcomes in patients with
1331   type 2 diabetes: Results from the CORONADO study. *Diabetes Obes Metab* (2020).
1332   28.   Wargny, M., *et al.* Type 1 Diabetes in People Hospitalized for COVID-19: New Insights From the
1333   CORONADO Study. *Diabetes Care* **43**, e174-e177 (2020).
1334   29.   Newman, A.M., *et al.* Determining cell type abundance and expression from bulk tissues with
1335   digital cytometry. *Nature Biotechnology* **37**, 773-782 (2019).
1336   30.   Aran, D., Hu, Z. & Butte, A.J. xCell: digitally portraying the tissue cellular heterogeneity
1337   landscape. *Genome Biology* **18**, 220 (2017).
1338   31.   Liao, M., *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19.
1339   *Nature Medicine* **26**, 842-844 (2020).
1340   32.   Song, W.M. & Zhang, B. Multiscale Embedded Gene Co-expression Network Analysis. *PLoS*
1341   *computational biology* **11**, e1004574 (2015).
1342   33.   Praest, P., *et al.* The influence of TAP1 and TAP2 gene polymorphisms on TAP function and its
1343   inhibition by viral immune evasion proteins. *Molecular immunology* **101**, 55-64 (2018).
1344   34.   Sreekumar, P.G., *et al.* The mitochondrial-derived peptide humanin protects RPE cells from
1345   oxidative stress, senescence, and mitochondrial dysfunction. *Investigative ophthalmology &*
1346   *visual science* **57**, 1238-1253 (2016).
1347   35.   Zhou, J.y., *et al.* Wnt/β-catenin-mediated heat exposure inhibits intestinal epithelial cell
1348   proliferation and stem cell expansion through endoplasmic reticulum stress. *Journal of cellular*
1349   *physiology* **235**, 5613-5627 (2020).
1350   36.   Liu, W., *et al.* LncRNA Malat1 inhibition of TDP43 cleavage suppresses IRF3-initiated antiviral
1351   innate immunity. *Proceedings of the National Academy of Sciences* **117**, 23695-23706 (2020).
1352   37.   American Association of Otolaryngology-Head and Neck Surgery. Tracheotomy
1353   Recommendations During the Covid-19 Pandemic. Vol. 2020 (2020).
1354   38.   Sommer, D.D., *et al.* Recommendations from the CSO-HNS taskforce on performance of
1355   tracheotomy during the COVID-19 pandemic. *J Otolaryngol Head Neck Surg* **49**, 23 (2020).
1356   39.   Shiba, T., Ghazizadeh, S., Chhetri, D., St John, M. & Long, J. Tracheostomy Considerations during
1357   the COVID-19 Pandemic. *OTO Open* **4**, 2473974X20922528 (2020).
1358   40.   Ear Nose and Throat Surgery United Kingdom. Tracheostomy guidance during the COVID-19
1359   Pandemic. Vol. 2020 (2020).
1360   41.   Michetti, C.P., Burlew, C.C., Bulger, E.M., Davis, K.A. & Spain, D.A. Performing tracheostomy
1361   during the Covid-19 pandemic: guidance and recommendations from the Critical Care and Acute
1362   Care Surgery Committees of the American Association for the Surgery of Trauma. *Trauma*
1363   *Surgery & Acute Care Open* **5**, e000482 (2020).
1364   42.   McGrath, B.A., *et al.* Tracheostomy in the COVID-19 era: global and multidisciplinary guidance.
1365   *Lancet Respir Med*, S2213-2600(2220)30230-30237 (2020).
1366   43.   Kelleni, M.T. SARS CoV-2 viral load might not be the right predictor of COVID-19 mortality. *J*
1367   *Infect* (2020).
1368   44.   Fajnzylber, J., *et al.* SARS-CoV-2 viral load is associated with increased disease severity and
1369   mortality. *Nat Commun* **11**, 5493 (2020).
1370   45.   Bitker, L., *et al.* Protracted viral shedding and viral load are associated with ICU mortality in
1371   Covid-19 patients with acute respiratory failure. *Ann Intensive Care* **10**, 167 (2020).

46. Magleby, R., *et al.* Impact of SARS-CoV-2 Viral Load on Risk of Intubation and Mortality Among
Hospitalized Patients with Coronavirus Disease 2019. *Clin Infect Dis* (2020).
47. Westblade, L.F., *et al.* SARS-CoV-2 Viral Load Predicts Mortality in Patients with and without
Cancer Who Are Hospitalized with COVID-19. *Cancer Cell* **38**, 661-671 e662 (2020).
48. Pujadas, E., *et al.* SARS-CoV-2 viral load predicts COVID-19 mortality. *Lancet Respir Med* **8**, e70
(2020).
49. Lee, N., *et al.* Viral loads and duration of viral shedding in adult patients hospitalized with
influenza. *J Infect Dis* **200**, 492-500 (2009).
50. Li, C.C., *et al.* Correlation of pandemic (H1N1) 2009 viral load with disease severity and
prolonged viral shedding in children. *Emerging infectious diseases* **16**, 1265-1272 (2010).
51. Peiris, J.S., *et al.* Clinical progression and viral load in a community outbreak of coronavirus-
associated SARS pneumonia: a prospective study. *Lancet* **361**, 1767-1772 (2003).
52. https://www.covid19treatmentguidelines.nih.gov/therapeutic-management/.
53. Kim, D., Quinn, J., Pinsky, B., Shah, N.H. & Brown, I. Rates of Co-infection Between SARS-CoV-2
and Other Respiratory Pathogens. *JAMA* **323**, 2085-2086 (2020).
54. Kreitmann, L., Monard, C., Dauwalder, O., Simon, M. & Argaud, L. Early bacterial co-infection in
ARDS related to COVID-19. *Intensive Care Med* **46**, 1787-1789 (2020).
55. Langford, B.J., *et al.* Bacterial co-infection and secondary infection in patients with COVID-19: a
living rapid review and meta-analysis. *Clin Microbiol Infect* **26**, 1622-1629 (2020).
56. Sepulveda, J., *et al.* Bacteremia and Blood Culture Utilization during COVID-19 Surge in New York
City. *J Clin Microbiol* **58**(2020).
57. Zhu, X., *et al.* Co-infection with respiratory pathogens among COVID-2019 cases. *Virus Res* **285**,
198005 (2020).
58. Zhang, H., *et al.* Risks and features of secondary infections in severe and critical ill COVID-19
patients. *Emerg Microbes Infect* **9**, 1958-1964 (2020).
59. Lansbury, L., Lim, B., Baskaran, V. & Lim, W.S. Co-infections in people with COVID-19: a
systematic review and meta-analysis. *J Infect* **81**, 266-275 (2020).
60. Koehler, P., *et al.* COVID-19 associated pulmonary aspergillosis. *Mycoses* **63**, 528-534 (2020).
61. Nolan, T.J., *et al.* Low-pathogenicity Mycoplasma spp. alter human monocyte and macrophage
function and are highly prevalent among patients with ventilator-acquired pneumonia. *Thorax*
71, 594-600 (2016).
62. Tsay, J.J., *et al.* Airway Microbiota Is Associated with Upregulation of the PI3K Pathway in Lung
Cancer. *Am J Respir Crit Care Med* **198**, 1188-1198 (2018).
63. Tsay, J.J., *et al.* Lower airway dysbiosis affects lung cancer progression. *Cancer Discov* (2020).
64. Sulaiman, I., *et al.* Evaluation of the airway microbiome in nontuberculous mycobacteria
disease. *Eur Respir J* **52**(2018).
65. Zhang, X., Li, L., Butcher, J., Stintzi, A. & Figeys, D. Advancing functional and translational
microbiome research using meta-omics approaches. *Microbiome* **7**, 154 (2019).
66. Sterlin, D., *et al.* IgA dominates the early neutralizing antibody response to SARS-CoV-2. *Science*
translational medicine **13**(2021).
67. Wang, Z., *et al.* Enhanced SARS-CoV-2 neutralization by dimeric IgA. *Science translational*
medicine **13**(2021).
68. Klingler, J., *et al.* Role of IgM and IgA Antibodies in the Neutralization of SARS-CoV-2. *J Infect Dis*
(2020).
69. Qiang, Z., *et al.* Nrf2 and STAT3 Alleviates Ferroptosis-Mediated IIR-ALI by Regulating SLC7A11.
Oxid Med Cell Longev **2020**, 5146982 (2020).
70. Xu, Y., Li, X., Cheng, Y., Yang, M. & Wang, R. Inhibition of ACSL4 attenuates ferroptotic damage
after pulmonary ischemia-reperfusion. *FASEB J* **34**, 16262-16275 (2020).

71. Dong, H., *et al.* Nrf2 inhibits ferroptosis and protects against acute lung injury due to intestinal
ischemia reperfusion via regulating SLC7A11 and HO-1. *Aging (Albany NY)* **12**, 12943-12959
(2020).
72. Hallman, M., Bry, K., Hoppu, K., Lappi, M. & Pohjavuori, M. Inositol supplementation in
premature infants with respiratory distress syndrome. *N Engl J Med* **326**, 1233-1239 (1992).
73. Preuss, S., *et al.* Inositol-trisphosphate reduces alveolar apoptosis and pulmonary edema in
neonatal lung injury. *Am J Respir Cell Mol Biol* **47**, 158-169 (2012).
74. Sekine, T., *et al.* Robust T Cell Immunity in Convalescent Individuals with Asymptomatic or Mild
COVID-19. *Cell* **183**, 158-168 e114 (2020).
75. Molinier-Frenkel, V., Prévost-Blondel, A. & Castellano, F. The IL4I1 Enzyme: A New Player in the
Immunosuppressive Tumor Microenvironment. *Cells* **8**(2019).
76. Baas, T., *et al.* Integrated molecular signature of disease: analysis of influenza virus-infected
macaques through functional genomics and proteomics. *Journal of virology* **80**, 10813-10828
(2006).
77. Regad, T. & Chelbi-Alix, M.K. Role and fate of PML nuclear bodies in response to interferon and
viral infections. *Oncogene* **20**, 7274-7286 (2001).
78. Bastard, P., *et al.* Autoantibodies against type I IFNs in patients with life-threatening COVID-19.
Science **370**(2020).
79. Zhang, Q., *et al.* Inborn errors of type I IFN immunity in patients with life-threatening COVID-19.
Science **370**(2020).
80. Kalil, A.C., *et al.* Baricitinib plus Remdesivir for Hospitalized Adults with Covid-19. *N Engl J Med*
(2020).
81. Zhou, Z., *et al.* Heightened Innate Immune Responses in the Respiratory Tract of COVID-19
Patients. *Cell host & microbe* **27**, 883-890 e882 (2020).
82. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114-2120 (2014).
83. Kopylova, E., Noe, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in
metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).
84. Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-
359 (2012).
85. Wood, D.E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2. *Genome*
biology **20**, 257 (2019).
86. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome biology* **15**, 550 (2014).
87. Pavel, A.B., *et al.* Alterations in Bronchial Airway miRNA Expression for Lung Cancer Detection.
Cancer Prev Res (Phila) **10**, 651-659 (2017).
88. Seumois, G., *et al.* Transcriptional Profiling of Th2 Cells Identifies Pathogenic Features Associated
with Asthma. *J Immunol* **197**, 655-664 (2016).
89. Kim, J., Kim, M.S., Koh, A.Y., Xie, Y. & Zhan, X. FMAP: Functional Mapping and Analysis Pipeline
for metagenomics and metatranscriptomics studies. *BMC Bioinformatics* **17**, 420 (2016).
90. Buchfink, B., Xie, C. & Huson, D.H. Fast and sensitive protein alignment using DIAMOND. *Nat*
Methods **12**, 59-60 (2015).
91. Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).
92. Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying
mammalian transcriptomes by RNA-Seq. *Nat Methods* **5**, 621-628 (2008).
93. Wilhelm, B.T., *et al.* Dynamic repertoire of a eukaryotic transcriptome surveyed at single-
nucleotide resolution. *Nature* **453**, 1239-1243 (2008).

94. Sultan, M., *et al.* A global view of gene activity and alternative splicing by deep sequencing of
the human transcriptome. *Science* **321**, 956-960 (2008).
95. Tanabe, M. & Kanehisa, M. Using the KEGG database resource. *Curr Protoc Bioinformatics*
Chapter 1, Unit1 12 (2012).
96. Kanehisa, M., Goto, S., Sato, Y., Furumichi, M. & Tanabe, M. KEGG for integration and
interpretation of large-scale molecular data sets. *Nucleic Acids Res* **40**, D109-114 (2012).
97. Law, C.W., *et al.* RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Res*
5(2016).
98. Law, C.W., Chen, Y., Shi, W. & Smyth, G.K. voom: Precision weights unlock linear model analysis
tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).
99. Reshef, D.N., *et al.* Detecting novel associations in large data sets. *Science* **334**, 1518-1524
(2011).

1480

Figure 1

Culture Positivity and risk for poor outcome (n=589)

Figure 1. Associations between culture positivity and clinical outcome. Odds ratios and corresponding 95% confidence intervals for rates of culture positivity for the whole cohort (n=589) during the length of their hospitalization (left) and during the first 2 weeks of hospitalization (right).

Figure 2

Figure 2. SARS-CoV-2 viral load and virus metatranscriptome analyses. Copies of the SARS-CoV-2 N gene per ml, normalized by the Human RNase P gene, comparing paired upper and lower airway samples (a) and levels in BAL comparing clinical outcome groups (b, *= Mann–Whitney U p<0.05, **= Mann–Whitney U p<0.01). (c) PCoA analysis based on Bray Curtis Dissimilarity index of BAL Metatranscriptome data comparing clinical outcome (PERMANOVA p-value). Bubble plot showing DESeq results of RNA viruses (d) and expressed DNA phages (e) enriched in each clinical outcome comparisons (bubble size based on median relative abundance for those found statistically significant).

Figure 3

Figure 3. Bacteria load and taxonomic compositional analyses. (a) Bacterial load measured by ddPCR targeting 16S rRNA gene (**= Mann–Whitney U p<0.01). PCoA analysis based on Bray Curtis Dissimilarity index of BAL Metagenome (b) and Metatranscriptome (c) data comparing clinical outcome (PERMANOVA p-value). (d) Gene Set Enrichment Analysis (GSEA) was used to compare the taxonomic signatures identified in BAL metagenome (diamonds) and metatranscriptome (circles) as distinctly enriched for comparisons between clinical outcome groups (differential enrichment performed based on DESeq2 analysis). (e) Bubble plot showing DESeq results of bacteria found concordantly differentially enriched between clinical outcome groups (bubble size based on median relative abundance for those found statistically significant).

Figure 4

Figure 4. Functional microbial compositional analyses. KOs were summarized to associated pathways and differential expression was calculated based on DESeq2 analysis. (a) Gene Set Enrichment Analysis (GSEA) was used to compare the functional signatures identified in BAL metagenome and metatranscriptome as distinctly enriched for comparisons between clinical outcome groups. (b) Bubble plot showing DESeq results of microbial functions found concordantly differentially enriched between clinical outcome groups (bubble size based on median relative abundance for those found statistically significant).

(a)

(b)

(c)

Figure 5. Lower airway host immune profiling in severely ill COVID-19. (a) Levels of anti-SARS-CoV-2 Spike antibodies in BAL (*= Mann–Whitney U p<0.05). (b) Heat-map of canonical pathway analysis based on Ingenuity Pathway Analysis (IPA, RRID:SCR_008653) using the lower airway host transcriptome comparing clinical outcome groups. Orange shows up-regulation of pathway, blue shows down-regulation of pathway. (c) Cell type abundance quantification plots. Comparison of abundance of mast cells and neutrophils among outcome groups in the BAL fluids of critically ill patients with COVID-19. Cell type abundance was estimated from the host transcriptome with CIBERSORTx. Each dot denotes the quantification score of a sample and boxes depict median and inter-quartile range (*= Mann–Whitney U p<0.05).

Figure 6

Figure 6. Mortality predictive power of metatranscriptome, metagenome and host transcriptome. (a) Area under the curved median and confidence interval for receiver operating characteristic curve analyses calculated from each sequencing datasets as predictor and mortality as outcome. (b) Kaplan-meier survival analyses based on a cutoff value estimated from features selected from each sequencing dataset. The "High risk" and "Low risk" groups is the mean of predicted risk scores in all samples. (c) Scatterplot among risk scores from metatranscriptome, metagenome, and host transcriptome. Dotted line denotes the mean of the risk scores across all subjects, which is also the threshold for dividing the samples into "High risk" and "Low risk" groups. (d) IPA analyses of host transcriptomic signatures identified as most predictive of mortality.