

1 **COVID-19 Vaccine hesitancy in Addis Ababa, Ethiopia: A mixed-methods study**

2 Nebiyu Dereje<sup>a\*</sup>, Abigel Tesfaye<sup>b</sup>, Beamlak Tamene<sup>b</sup>, Dina Alemeshet<sup>b</sup>, Haymanot Abe<sup>b</sup>,  
3 Nathnael Tesfa<sup>b</sup>, Saron Gedion<sup>b</sup>, Tigist Biruk<sup>b</sup>, Yabets Lakew<sup>b</sup>

4 <sup>a</sup>Department of Public Health, Myungsung Medical College, Addis Ababa, Ethiopia

5 <sup>b</sup>Department of Medicine, Myungsung Medical College, Addis Ababa, Ethiopia

6

7 \*Corresponding author contact information

8 Nebiyu Dereje

9 Email: [neba.jahovy@gmail.com](mailto:neba.jahovy@gmail.com)

10 P.O.Box 14972

11 Addis Ababa, Ethiopia

12 **List of Abbreviations**

13 aOR: Adjusted Odds Ratio

14 CI: Confidence Interval

15 COR: Crude Odds Ratio

16 SARS: Severe Acute Respiratory Syndrome

17 UK: United Kingdom

18 US: United States

## 19 **Abstract**

20 **Background:** COVID-19 infection is a global pandemic threatening the public health. Due to the  
21 development and initiation of vaccination, currently significant difference upon vaccine  
22 acceptance is seen between developed and developing countries. However, there are no data on  
23 the level of COVID-19 vaccine hesitancy and its associated factors in Addis Ababa, Ethiopia.

24 **Methods:** An embedded mixed method study [QUAN(quali)] was conducted among residents of  
25 Akaki Kality sub-city in Addis Ababa, Ethiopia. For the quantitative part, a multi-stage sampling  
26 technique was used to recruit the study participants (n = 422). Twenty four adults were included  
27 purposively for the qualitative in-depth interview. Data was collected by face-to-face interview  
28 by using a semi-structured questionnaire. Factors associated with COVID-19 vaccine hesitancy  
29 were identified by multivariable binary logistic regression model.

30 **Result:** The mean age of the participants was 34.1 years ( $\pm 12.9$ ). Nearly half (46.7%) of the  
31 participants exhibited poor level of knowledge and 51.8% had negative attitude towards COVID-  
32 19 and its preventive measures. One out five (19.1%) participants were not willing to get  
33 vaccinated when it becomes available. In the multivariable analysis, vaccine hesitancy was  
34 significantly associated with being female (aOR=1.97; 95% CI: 1.10 - 3.89, p=0.03), negative  
35 attitude towards COVID-19 and its preventive measures (aOR=1.75; 95% CI: 1.08 - 3.02,  
36 p=0.04), and information source being social media (internet) (aOR=3.59; 95% CI: 1.75 - 7.37, P  
37 <0.0001).

38 **Conclusion:** A considerable proportion of the people in Addis Ababa have concerns on COVID-  
39 19 vaccine and unwilling to accept once it is available. Several conspiracy theories were put

40 forth to justify their stance and this was mainly due to the misconceptions distributed from the  
41 use of social media as source of information. Overall, providing the community with health  
42 education and consistent government efforts in uphold the prevention measures are of paramount  
43 importance to tackle this pandemic.

44 **Key Words:** COVID-19, knowledge, attitude, Vaccine, Ethiopia

45

46

## 47 **Introduction**

48 Corona virus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome corona  
49 virus 2 (SARS-CoV-2) also known as Novel coronavirus (nCoV) [1]. The first case of COVID-  
50 19 was discovered in Wuhan city, Hubei province of China with unexplained pneumonia on  
51 December 12, 2019 [2]. The virus is transmitted through large droplets generated during  
52 coughing or sneezing of symptomatic and asymptomatic patients [3]. Therefore, frequent hand-  
53 washing with soap and water, using alcohol based hand rub or sanitizer, avoidance of hand  
54 shaking/public gathering and use of face mask are crucial to halt the spread of COVID-19 [4].  
55 COVID-19 was declared a pandemic by the World health organization on March 11, 2020 [5].  
56 Since its emergence, this pandemic has shown its capability to spread rapidly in the world  
57 causing the most dramatic global health crisis of our time resulting in devastating social,  
58 economic and political crises [6].

59 Globally, more than 210 countries/territories have been affected by the virus, with more than one  
60 hundred ten million people being infected and 2.5 million deaths reported as of February 24,  
61 2021. Ethiopia ranks 71<sup>st</sup> regarding COVID-19 with more than 154, 000 infected and 2,305 dead  
62 (February 24, 2021) [7]. Unfortunately, Ethiopia was found to be one of five African countries  
63 with the highest case burden of COVID-19 [7]. However, the government of Ethiopia has been  
64 striving to spread information on COVID-19 preventive measures via television, radio or social  
65 media outlets and declared a state of emergency [8].

66 Currently, COVID-19 vaccine has been made available but it is highly controversial. More than  
67 seven billion doses have been pre-purchased by countries and organizations of the world, of  
68 which more than half was sold-out to high income countries [9]. This figure is threatening to the  
69 global health as may be an indication of the disparities on the health delivery globally.

70 Myths and conspiracy theories on vaccinations have been spreading and can easily be accepted  
71 by the developing world. This may cause people to be reluctant and maleficent towards  
72 vaccination, which has been demonstrated by a study in Nigeria by a low vaccine acceptability  
73 rate [10]. WHO defined vaccine hesitancy as it is a difficulty in accepting or an outright refusal  
74 of vaccines, despite their availability. In 2019, before the COVID-19 pandemic, the World Health  
75 Organization listed vaccine hesitancy as one of the ten global threats to public health [11].  
76 Currently, Ethiopia has not started administering the vaccines.

77 Hence it is very crucial to understand the varying vaccine attitudes among the community to  
78 design a strategy to overcome the vaccine hesitancy. Furthermore, unraveling the specific fears  
79 and doubts of the community with regards to receiving the vaccine can help government and  
80 other concerned officials to adequately address the misconceptions and various conspiracy  
81 theories in their campaigns. Therefore, this study was aimed to assess the level of COVID-19  
82 vaccine acceptability among the population in Addis Ababa, the capital city of Ethiopia by  
83 employing a community based mixed methods study.

84

85

## 86 **Methods and Materials**

### 87 **Study design and participants**

88 A concurrent embedded mixed methods design [QUAN(qual)] was employed from January 20 –  
89 31, 2021 among adult population ( $\geq 18$  years) currently residing in Akaki Kality sub city of Addis  
90 Ababa, Ethiopia. The qualitative part of the study was embedded in the bigger quantitative cross-  
91 sectional study. The qualitative part was mainly intended to explain the reasons for COVID-19  
92 vaccine hesitancy, as a supplementary of the quantitative study.

93 A sample size for the quantitative part of the study ( $n = 422$ ) was determined by using a single  
94 population proportion formula, by taking 95% confidence interval, 5% margin of error, 50%  
95 proportion of vaccine hesitancy and adding up 10% non-response rate. For the qualitative part,  
96 24 participants were included into the study based on the information saturation.

97 Multi-stage sampling technique was employed to recruit the participants for the quantitative part.  
98 There were 13 districts in the sub-city; of which three of them were selected randomly (lottery  
99 method). The total sample was allocated proportionally to the districts. Then, the households  
100 from each district were selected by employing a systematic random sampling (sampling interval  
101 = every 4<sup>th</sup> house). From the specific households selected, only one randomly selected eligible  
102 individual was interviewed. For the qualitative part of the study, purposive sampling method was  
103 used to recruit participants who have reach information.

### 104 **Data collection tools and procedures**

105 Data was collected by using a semi-structured questionnaire which was adapted from reviewed  
106 literatures [10, 12, 13]. The questionnaire has 5 components: socio-demographic, knowledge  
107 towards COVID-19, attitude towards COVID-19, practice of COVID-19 prevention measures,

108 and COVID-19 vaccine acceptance. The questionnaire was in English and translated into  
109 Amharic for the interview. The questionnaire was administered face-to-face by the medical  
110 interns. For the qualitative part of the study, in-depth interviews were made by the investigators.

### 111 **Data management and analysis**

112 Data was coded and entered into SPSS-for windows version 25 for analysis. Frequency and  
113 proportions were used to summarize categorical variables, whereas mean and standard deviation  
114 were used to summarize continuous variables.

115 The primary outcome variable of the study was COVID-19 vaccine hesitancy which was  
116 assessed by asking a question “Will you get vaccinated if you get COVID-19 vaccine?” then the  
117 response was dichotomized as “Yes” or “No”.

118 Multivariable binary logistic regression analysis was carried out to identify factors associated  
119 with vaccine hesitancy, as expressed by adjusted odds ratio (aOR) along with its respective 95%  
120 confidence interval (CI). Variables with  $<0.25$  in bivariate analysis were considered for  
121 multivariable analysis. Variables having  $P$  value  $<0.05$  were considered statistically significant.  
122 Multicollinearity was assessed by the collenarity diagnostics (Variance Inflation Factor and the  
123 tolerance test). Goodness of the model was checked by the Hosmer Lemshow goodness of fit  
124 test. The qualitative data analysis was initiated by transcription and translating of the interviews,  
125 then coded and analyzed by thematic analysis. The findings of the qualitative study were used to  
126 supplement the findings of quantitative data.

### 127 **Ethical consideration**

128 Ethical approval of this study was obtained from the Institutional Review Board (IRB) of  
129 Myungsung Medical College. The participants of the study was informed about the purpose of

130 the study and provided their written consent. At the end of the interview, the data collectors have  
131 provided information with regard to the COVID-19 vaccine.

## 132 **Result**

### 133 **Socio-demographic characteristics**

134 A total of 409 respondents completed the questionnaire, with a response rate of 96.9%. Majority  
135 of the participants 294 (71.9%) were females and married (62.3%) (Table 1). The mean age of  
136 the participants was 34.1 years ( $\pm 12.9$ ), ranging from 18 - 85 years.

137 **Table 1: Socio-demographic characteristics of the study participants**

		<b>Frequency (N)</b>	<b>Percent (%)</b>
<b>Sex</b>	Male	115	28.1%
	Female	294	71.9%
<b>Age</b>	18-29	174	42.5%
	30-40	147	35.9%
	41-50	40	9.8%
	>50	48	11.7%
<b>Marital status</b>	Not married	123	30.1%
	Married	255	62.3%
	Widowed	20	4.9%
	Divorced	11	2.7%
<b>Religion</b>	Christian	349	85.3%
	Muslim	60	14.7%

Educational status	No formal education	39	9.5%
	Primary school	105	25.7%
	Secondary and above	265	64.8%
Occupation	Unemployed/housewife	190	46.5%
	Employed	219	53.5%
Monthly income*	≤3200 ETB (≤100 USD)	175	42.8%
	>3200 ETB (>100 USD)	228	57.2%

138 ETB: Ethiopian Birr USD: United States Dollar \*6 participants' data missing

### 139 **Knowledge towards COVID-19**

140 Almost all of the participants heard about COVID-19 from Television or Radio. However, the  
141 average knowledge score was  $56.7 \pm 3.7$ , with 46.7% (n=191) exhibited poor level of knowledge  
142 (Supplementary Table 1).

143 The qualitative part of the study found that some of the study participants stated that they do not  
144 possess enough knowledge and would like to know more about things like the prospective  
145 vaccines, the new strain and how to prevent getting infected from asymptomatic patients.

146 For example, one 39 year old male participant said that;

147 *“I want to know where the virus really came from initially; is it natural or*  
148 *manmade?”[39 year old, male]*

### 149 **Attitude towards COVID-19**

150 The mean attitude score was found to be  $20.3 \pm 1.2$ , with 51.8% of the participants have negative  
151 attitude regarding COVID-19 and its preventive measures (Supplementary Table 2).

152 This result is corroborated by the findings on the qualitative part of the study where the majority  
153 of the respondents stated that they were initially very concerned but now they were less so. In  
154 fact, the majority of the participants stated that they were very afraid and anxious initially  
155 believing that they were definitely getting infected and that they would certainly die if they  
156 caught it. Currently, most of the respondents reported feeling less anxious. While some stated  
157 that while they were still aware that it was a serious illness they were less concerned since they  
158 believed it was preventable. While others stated the reason for their lack or decreasing concern  
159 was that they believed they would recover if they got infected.

160 In a spectrum of similar answers, five of the participants stood out stating that they did not  
161 believe the disease exists anymore since they have not personally encountered an infected person.  
162 In contrast to these responses, we also found two participants who stated that while they initially  
163 were not concerned they are starting to get worried now since the number of infected individuals  
164 and the resulting death toll in this country is increasing.

165 One female participant stated the following to show that she is not concerned about COVID-19:

166 *“I am not scared because I expected this to happen; we brought this on ourselves and we*  
167 *are paying for our sins. It has been long time coming.” [Female, 50 year old]*

168 Another 45 year old female participant stated:


169 *“I have been through an outbreak before...I got sick and I had to be isolated from my*  
170 *family but I recovered easily and I don’t believe this would be any different.” [Female,*  
171 *47 year old]*

172 More than two-thirds (68.7%) of the participants believed Ethiopia would win the battle against  
173 the pandemic. This finding was also supported by the qualitative study.

174 *“I was afraid that everyone in Ethiopia would die because even white people could not*  
175 *handle it. I think the only reason we have survived is because Ethiopia is God’s country.”*  
176 *[Female, 70 year old]*

### 177 **Vaccine hesitancy**

178 More than 90% of the participants heard about the prospective COVID-19 vaccine mainly from  
179 TV/Radio. However, 78 (19.1%) were not willing to get vaccinated when it becomes available  
180 (Figure 1).




181  
182 *Figure 1: Prospective COVID-19 vaccine*  
183 Out of them, 43.6% don't take the vaccine due to fear of side effects and 41.0% of them believe  
184 that the vaccine may be biological weapon (Figure 2).

185 In the qualitative in-depth interview, some stated they did not have enough information about the  
186 vaccine and wanted to see other people take it first. Majority of the respondents feared it would  
187 not be effective or have too many side effects as there was not enough time to study it and as a

188 result stated that they preferred other methods to prevent COVID-19. A few of the participants  
189 thought that the vaccine that will be distributed in Africa would be of lower quality. Others  
190 thought it would be used as a biological weapon by the developed nations to cause infertility and  
191 control the population of poor countries. Moreover, it was also mentioned that the vaccines  
192 might be used as a weapon to insert microchips made by the illuminati, into the body as the mark  
193 of the beast “(666)” that would cause them to forsake their faith. A few others did not think they  
194 needed the vaccine because they had God’s protection.

195 *“I don’t think the vaccine will come to this country and even if it does I don’t need it;*  
196 *God will be my vaccine.” [Female, 45 year old]*

197 Close to 20% of the participants thought that children should not get vaccinated. Some of the  
198 participants did not recommend the vaccine to children even though they would take it  
199 themselves. These participants further expressed in the in depth interview that they thought the  
200 virus did not affect children or it would be too dangerous for them.



213 *Figure 2: Reasons of participant for refusing COVID-19 vaccination*

214 **Factors associated with vaccine hesitancy**

215 In the multi-variable analysis (Table 2), COVID-19 vaccine hesitancy was associated with sex,  
216 attitude and source of information about the vaccine.

217 It was found that the odds of vaccine hesitancy was 1.97 times (aOR=1.97; 95% CI: 1.10 - 3.89,  
218 p=0.03) higher among female participants as compared to male participants. The odds of vaccine  
219 hesitancy was 1.75 (aOR=1.75; 95% CI:1.08 - 3.02, p=0.04) times higher in those participants  
220 who were found to have a negative attitude towards COVID-19 and its preventive measures as  
221 compared to those who had a positive attitude. Similarly, the odds of vaccine hesitancy was 3.6  
222 times (aOR=3.59; 95% CI: 1.75 - 7.37, P <0.0001) higher among those participants that received  
223 their information from social media (internet) as compared to those who received information  
224 only from TV/radio.

225 *Table 2: Factors associated with COVID-19 vaccine hesitancy in Addis Ababa, Ethiopia, 2021*

Variables	Vaccine hesitancy		COR (95% CI)	aOR(95% CI)	P value
	Yes (%)	No (%)			
<b>Sex</b>					
Male	17 (21.8%)	98 (29.6%)	1.00	1.00	0.03
Female	61 (78.2%)	233 (70.4%)	1.49 (0.83-2.69)	1.97 (1.10-3.89)	
<b>Age</b>					
18-29	34 (43.6%)	140 (42.3%)	1.00	1.00	0.934
30-40	24 (30.8%)	122 (36.9%)	0.81 (0.46-1.44)	1.03 (0.55-1.92)	
41-50	12 (15.4%)	29 (8.8%)	1.77 (0.82-3.82)	2.22 (0.94-5.21)	

>50	8 (10.2%)	40 (12.0%)	0.82 (0.35-1.92)	1.08 (0.39-2.97)	0.892
<b>Religion</b>					
Christian	8 (10.3%)	52 (15.7%)	1.00	1.00	0.621
Muslim	70 (89.7%)	278 (84.3%)	1.64 (0.74-3.60)	1.23 (0.54-2.83)	
<b>Educational status</b>					
No formal education	9 (11.5%)	30 (9.1%)	1.11 (0.50-2.48)	1.11 (0.39-3.16)	0.840
Primary education	13 (16.7%)	93 (28.1%)	0.53 (0.27-1.01)	0.81 (0.40-1.63)	0.560
Secondary and above	56 (71.8%)	208 (62.8%)	1.00	1.00	
<b>Attitude</b>					
Positive attitude	28 (35.9%)	169 (51.1%)	1.00	1.00	0.04
Negative attitude	50 (64.1%)	162 (48.9%)	1.87 (1.12-3.12)	1.75 (1.08-3.02)	
<b>Primary source of information</b>					
TV/Radio	38 (48.7%)	255 (77.0%)	1.00	1.00	
Social media (internet)	40 (51.3%)	76 (23.0%)	3.53 (1.67-6.98)	3.59 (1.75-7.37)	0.0001

226

227

## 228 **Discussion**

229 For the COVID-19 battle, the population adherence to preventive measures is crucial; however,  
230 it is mainly affected by their KAP toward the disease [1]. The findings of this study showed that  
231 nearly half of the study participants demonstrated inadequate knowledge of COVID-19,  
232 indicating a great knowledge gap. This finding is higher than studies conducted in other parts of  
233 Ethiopia such as Arbaminch (23.5%) and Gedeo (39.5%), and other low income countries such  
234 as Ghana (34.9%), and Malaysia (22.7%) [14-17]. The discrepancies might be due to differences  
235 in the community awareness creation through mass media and social media. Further, in our  
236 study, more than half the participants had negative attitude towards COVID-19 and its preventive  
237 measures, which is higher than the finding of a study conducted in Southern Ethiopia [15, 18]  
238 and lower than study done among Dessie and Kombolcha town residents in Ethiopia [19]. The  
239 discrepancy in the findings may be due to differences in the study period. The later studies were  
240 conducted earlier in the pandemic when the declaration and enforcement of state of emergency  
241 and other measures were still in place. Our findings show a significant decrease in the  
242 community's attitude towards COVID-19 and its prevention measures which can lead people to  
243 become discouraged to consistently adhere to the measures set forth by the government and the  
244 world health organization. These findings of the study has an implication on the public health  
245 and underscore the need for urgent concerted efforts to promote the knowledge of the general  
246 public in Ethiopia towards COVID-19 preventive measures. If the current trend evidenced by  
247 this study continues in Ethiopia, COVID-19 will pose a devastating outcome on the medical,  
248 financial and social aspect of citizens besides the potential for new strains of disease developing.

249 As COVID-19 continues to ravage the world, vaccination offers the most reliable hope for a  
250 permanent solution to controlling the pandemic. However, a vaccine must be accepted and used

251 by a large majority of the population to create herd immunity [20]. The findings of this study  
252 showed that about one out of five participants are not willing to receive COVID-19 vaccine  
253 when it is available, which is higher than the findings reported from developed countries such as  
254 UK (3%) [9, 21, 22]. The discrepancies might be due to insufficient knowledge about the vaccine  
255 and difference in the perception of the seriousness of the pandemic. This implies that if the  
256 doubts and fears of the majority regarding the vaccine are not addressed properly, we may not be  
257 able to attain herd immunity. Surprisingly, the finding of this study was lower than a study  
258 conducted in the US (31%) and Nigeria (80%) [13, 20]. This might be due to difference in access  
259 to wide variety of conspiracy theories and doubts via internet.

260 Consistent to the study conducted in China [23], vaccine hesitancy was more likely among  
261 females as compared to males in our study. This could be due to higher exposure of males for  
262 different media as compared to females in Ethiopia. In the present study, increased likelihood of  
263 vaccine hesitancy was also indicated among those with negative attitude towards COVID-19 and  
264 its preventive measures. The qualitative aspects of this study also found that those participants  
265 who would not take the vaccine stated one of their reason to be their lack of implicit trust in the  
266 government and in health professionals. Thus, this lack of confidence in the government  
267 exhibited by 41.8% of our participants may be a potential hurdle we might face after the arrival  
268 of the vaccine to Ethiopia.

269 In our study, those participants who received their information from social media (internet) were  
270 more likely to have vaccine hesitancy as compared to those who got their information only from  
271 TV/radio. This finding of the study is in line with a study conducted to assess health protective  
272 behaviors and conspiracy theories during the pandemic found that there was significant  
273 association between holding a conspiracy belief and checking social media for news of COVID-

274 19 [24]. As a result, this finding is justified by our findings on both the quantitative and  
275 qualitative aspects of our study which found that the majority of the reasons given for hesitancy  
276 towards the vaccine were the belief in the conspiracy theories. Thus, the spread of these  
277 conspiracy theories is a potential issue that can cause problems when vaccine distribution starts  
278 in Ethiopia. Particularly, if these conspiracy theories start getting a wider audience thus there  
279 may be a need to act in haste and find a solution before this issue worsens.

280 This study is the first community based study to assess the Ethiopian community's perception  
281 towards COVID-19 vaccine and its level of acceptance. We employed a mixed methods design  
282 which enables us to make the deep understanding of the issue. However, the study might be  
283 limited due to the recall bias and social desirability bias during the data collection. In addition to  
284 this, our sample over represents female population because the majority of the study participants  
285 that were found at home during data collection time were housewives. Therefore, generalization  
286 of the study results need to be cautious.

## 287 **Conclusion**

288 A considerable proportion of the people have concerns of the COVID-19 vaccine and unwilling  
289 to accept once it is available. Several conspiracy theories were put forth to justify their stance  
290 and this was mainly due to the misconceptions distributed from the use of social media as source  
291 of information about the vaccine. Overall, providing the community with health education and  
292 consistent government efforts in uphold the prevention measures are of paramount importance to  
293 tackle this pandemic.

## 294 **Availability of data and material**

295 Data are available upon reasonable request from the corresponding author.

296 **Conflict of interests**

297 The authors declare that they have no conflict of interests.

298 **Funding**

299 This study was funded by Myungsung Medical College. However, the funder had no role in the  
300 design, conduct, analysis and interpretation of this study.

301 **Authors Contributions**

302 All the authors have substantially involved in the design of the study, collection of data,  
303 visualization of data, analysis of the data, and interpretation of the results, and drafted the initial  
304 manuscript and approved the final manuscript. All authors have read and approved the  
305 manuscript.

306 **Acknowledgement**

307 The authors would like to thank Myungsung Medical College for providing fund to conduct this  
308 study. The authors are grateful to the study participants for their contributions.

## 309 Reference

- 310 1. World Health Organization. Coronavirus disease 2019 (COVID-19): situation report, 82.  
311 2020.
- 312 2. Imai N, Dorigatti I, Cori A, Donnelly C, Riley S, Ferguson N. Report 2: Estimating the  
313 potential total number of novel Coronavirus cases in Wuhan City, China. 2020.
- 314 3. Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, et al.  
315 Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *New England*  
316 *journal of medicine*. 2020;382(10):970-1.
- 317 4. Ranney ML, Griffeth V, Jha AK. Critical supply shortages—the need for ventilators and  
318 personal protective equipment during the Covid-19 pandemic. *New England Journal of*  
319 *Medicine*. 2020;382(18):e41.
- 320 5. Assessment RR. Coronavirus disease 2019 (COVID-19) in the EU/EEA and the UK—  
321 ninth update. European Centre for Disease Prevention and Control: Stockholm. 2020.
- 322 6. COVID C. Global Cases by the Center for Systems Science and Engineering (CSSE) at  
323 Johns Hopkins University (JHU). JHU COVID-19 Resource Center. Johns Hopkins Coronavirus  
324 Resource Center. 2020.
- 325 7. Worldometer. Coronavirus Update (Live): Cases and Deaths from COVID-19 Virus  
326 Pandemic 2021 [cited 2021 Feb 24]. Available from:  
327 <https://www.worldometers.info/coronavirus/>.
- 328 8. Zikargae MH. COVID-19 in Ethiopia: Assessment of How the Ethiopian Government  
329 has Executed Administrative Actions and Managed Risk Communications and Community  
330 Engagement. *Risk management and healthcare policy*. 2020;13:2803.

- 331 9. Bell S, Clarke R, Mounier-Jack S, Walker JL, Paterson P. Parents' and guardians' views  
332 on the acceptability of a future COVID-19 vaccine: A multi-methods study in England. *Vaccine*.  
333 2020;38(49):7789-98.
- 334 10. Reuben RC, Danladi MM, Saleh DA, Ejembi PE. Knowledge, attitudes and practices  
335 towards COVID-19: an epidemiological survey in North-Central Nigeria. *Journal of community*  
336 *health*. 2020:1-14.
- 337 11. Graffigna G, Palamenghi L, Boccia S, Barello S. Relationship between citizens' health  
338 engagement and intention to take the covid-19 vaccine in italy: A mediation analysis. *Vaccines*.  
339 2020;8(4):576.
- 340 12. Ma L, Liu H, Tao Z, Jiang N, Wang S, Jiang X. Knowledge, Beliefs/Attitudes, and  
341 practices of rural residents in the prevention and control of COVID-19: an online questionnaire  
342 survey. *The American journal of tropical medicine and hygiene*. 2020;103(6):2357-67.
- 343 13. Enitan S, Oyekale A, Akele R, Olawuyi K, Olabisi E, Nwankiti A. Assessment of  
344 Knowledge, Perception and Readiness of Nigerians to participate in the COVID-19 Vaccine  
345 Trial. *International Journal of Vaccines and Immunization*. 2020;4:1-13.
- 346 14. Aweke Z, Jemal B, Mola S, Hussen R. Knowledge of COVID-19 and its prevention  
347 among residents of the Gedeo zone, South Ethiopia. Sources of information as a factor. *Current*  
348 *Medical Research and Opinion*. 2020;36(12):1955-60.
- 349 15. Nigussie TF, Azmach NN. Knowledge, attitude and practice towards covid-19 among  
350 Arba Minch town, southern Ethiopia. *GSJ*. 2020;8(6).
- 351 16. Nkansah C, Serwaa D, Adarkwah LA, Osei-Boakye F, Mensah K, Tetteh P, et al. Novel  
352 coronavirus disease 2019: knowledge, practice and preparedness: a survey of healthcare workers  
353 in the Offinso-North District, Ghana. *The Pan African Medical Journal*. 2020;35(79).

- 354 17. Mohamad EM, Azlan AA, Hamzah MR, Tham JS, Ayub SH. Public knowledge, attitudes  
355 and practices towards COVID-19: A cross-sectional study in Malaysia. medRxiv. 2020.
- 356 18. Mola S, Aweke Z, Jemal B, Hussein R, Hailu S, Neme D, et al. Magnitude and associated  
357 factors for attitude and practice of Southern Ethiopian residents toward COVID-19 and its  
358 preventions: A community based cross sectional study. 2020.
- 359 19. Kassa AM, Mekonen AM, Yesuf KA, Tadesse AW, Bogale GG. Knowledge level and  
360 factors influencing prevention of COVID-19 pandemic among residents of Dessie and  
361 Kombolcha City administrations, North-East Ethiopia: a population-based cross-sectional study.  
362 BMJ open. 2020;10(11):e044202.
- 363 20. Pogue K, Jensen JL, Stancil CK, Ferguson DG, Hughes SJ, Mello EJ, et al. Influences on  
364 attitudes regarding potential COVID-19 vaccination in the United States. Vaccines.  
365 2020;8(4):582.
- 366 21. Harapan H, Wagner AL, Yufika A, Winardi W, Anwar S, Gan AK, et al. Acceptance of a  
367 COVID-19 vaccine in southeast Asia: A cross-sectional study in Indonesia. Frontiers in public  
368 health. 2020;8.
- 369 22. Salali GD, Uysal MS. COVID-19 vaccine hesitancy is associated with beliefs on the  
370 origin of the novel coronavirus in the UK and Turkey. Psychological medicine. 2020:1-3.
- 371 23. Wang J, Jing R, Lai X, Zhang H, Lyu Y, Knoll MD, et al. Acceptance of COVID-19  
372 Vaccination during the COVID-19 Pandemic in China. Vaccines. 2020;8(3):482.
- 373 24. Allington D, Duffy B, Wessely S, Dhavan N, Rubin J. Health-protective behaviour,  
374 social media usage and conspiracy belief during the COVID-19 public health emergency.  
375 Psychological medicine. 2020:1-7.

376