Title

Reduced Frontal Cortical Tracking of Conflict between Selfish versus Prosocial Motives in Narcissistic

Personality Disorder

Authors

Dipl.-Psych. David S. Stolz (1,2)[+], Dr. Aline Vater (3), Dr. Björn H. Schott (4,5), Dr. Stefan Roepke(3)*, Dr.

Frieder M. Paulus (1,2)*, Dr. Sören Krach (1,2)*

[+] Corresponding author

*These authors contributed equally

Affiliations

1 Social Neuroscience Lab, Department of Psychiatry and Psychotherapy, University of Lübeck, Lübeck,

Germany

2 Translational Psychiatry Unit (TPU), Department of Psychiatry and Psychotherapy, University of Lübeck,

Germany

3 Charité – Universitätsmedizin Berlin, Campus Benjamin Franklin, Berlin, Germany

4 Leibniz Institute for Neurobiology Magdeburg, Magdeburg, Germany

5 Department of Psychiatry, University Medicine Göttingen, Göttingen, Germany

Corresponding author

David S. Stolz

Social Neuroscience Lab

Center of Brain, Behavior, and Metabolism (CBBM)

Universität zu Lübeck

Ratzeburger Allee 160

23562 Lübeck

Phone: +49 451 3101 7528

Email: david.stolz[at]uni-luebeck.de

Short title

Prosociality in Narcissistic Personality Disorder

Abstract

Narcissistic Personality Disorder (NPD) entails severe impairments in interpersonal functioning that are

likely driven by selfish and exploitative behavior. Here, we investigate the underlying motivational and

neural mechanisms of prosocial decision-making by experimentally manipulating motivational conflict

between selfish and prosocial incentives. One group of patients diagnosed with NPD and a group of healthy

controls (CTL) were scanned using functional magnetic resonance imaging while performing a prosocial

decision-making task. In this task, we systematically varied the level of conflict between selfish and

prosocial options on each trial. We analyzed choice behavior, response times, and neural activity in regions

associated with conflict monitoring to test how motivational conflict drives prosocial choice behavior.

Participants in the NPD group behaved less prosocially than the CTL group overall. Varying degrees of

motivational conflict between selfish and prosocial options induced response variability in both groups,

but more so in the CTL group. The NPD group responded faster than the CTL group, unless choosing

prosocially, which slowed response times to a level comparable to the CTL group. Additionally, neural

activity tracking motivational conflict in dorsomedial prefrontal cortex was reduced in the NPD group.

Collectively, low generosity in NPD appears to arise from reduced consideration of prosocial motives,

which obviates motivational conflict with selfish motives and entails reduced activity in neural conflict

monitoring systems. Yet, our data also indicate that NPD is not marked by an absolute indifference to

others' needs. This points to potentials for improving interpersonal relationships, effectively supporting

the well-being of patients and their peers.

Introduction

Narcissistic personality disorder (NPD) is characterized by a pervasive pattern of grandiosity, need for admiration, and lack of empathy[1]. Relationships in patients with NPD are described as largely superficial and serving as a means for self-esteem regulation, rather than being motivated by mutuality and genuine interest in the other person[1]. Most of these descriptive interpersonal features derive from clinical observations. Laboratory studies that prove these concepts using experimental designs are lacking[2]. The aim of this study was to investigate selfish behavior, a key interpersonal pattern in NPD, on a behavioral and neural level.

"Narcissism" can be conceptualized as a dimensional personality trait including grandiose and vulnerable features[3,4]. Grandiose features are also reflected in the descriptive DSM-5 criteria set of NPD[1] and include an exaggerated sense of self-importance and need for admiration. Vulnerable features include traits such as anger, shame, and insecurity[3]. A pervasive motif of grandiose narcissism is to maintain an exaggerated self-esteem, which is reflected in social behavior[5,6]. Non-clinically grandiose individuals overestimate their abilities[7], tend to be extraverted[8], and display interpersonal skills like charmingness[9,10], which presumably facilitates making new acquaintances[6,11,12]. In the long run, however, non-clinical grandiosity predicts more detrimental outcomes[5,12] reflected in antagonistic and defensive behavior[9,13,14], vindictiveness[15,16], and dysfunctional relationships[17,18]. Non-clinical narcissistic vulnerability similarly relates to defensiveness and self-enhancement[10], but also to emotional instability, low extraversion[19], low self-esteem[20], and aggression[21]. Most behavioral studies on "narcissism" are performed in non-clinical samples using the Narcissism Personality Inventory (NPI) that mainly captures grandiose features of the construct[21], making the transferability of these data into clinical samples disputable[22]. Pathological narcissism likely represents a singular dimension, with NPD as an extreme[23] that is linked to reduced empathy[24,25], and distress of close others[26].

In healthy subjects, laboratory studies have shown that higher trait narcissism predicts higher individual acquisitiveness but diminishes the group's total harvest in common goods games[27]. Similarly,

trait vulnerability predicts reduced generosity, lower sensitivity to potential punishment for low

generosity, and increased retaliation[28]. This coheres with classic findings that non-clinical grandiosity

predicted aggression, especially following self-esteem threats[29]. Yet, when prosocial motives gain

salience through empathizing or perspective taking, both selfishness[30–32] and aggression[31,33]

diminish in healthy individuals, and the arising arbitration between selfish and prosocial motives entails

slower reaction times (RTs)[34–37]. Habitually selfish/prosocial individuals respond quickly when behaving

selfishly/prosocially, whereas deviations from default behavior are slower[36]. Hence, to the extent that

individuals inform their behavior by reconciling opposing interpersonal motives, experimentally induced

conflict between such motives should slow down responses and potentially shift them away from default

tendencies[34,36].

Regarding NPD, this evidence from non-clincal samples stimulates divergent hypotheses: Either,

pronounced selfish behavior, as also frequently observed in non-clinical narcissism[27–29], could go along

with low motivational conflict if selfish motives trump prosocial motives. Alternatively, if interpersonal

deliberation in NPD entails motives like being perceived as likeable or attractive[9,10], achievable for

instance through generous actions, eventually selfish behavior should demand to first resolve conflict with

co-existing prosocial motives. In the first hypothesis, low motivational conflict should be reflected in fast

RTs, whereas the second hypothesis would predict slower RTs[34,36].

Neurally, motivational conflict should relate to activity in areas of medial prefrontal cortex, such

as dorsomedial prefrontal cortex (dmPFC), anterior cingulate cortex (ACC), and the supplementary motor

complex (SMC, including supplementary motor area (SMA), and preSMA)[38]. These regions are commonly

associated with cognitive control[39,40], which also holds in the social domain[41–43]. Further, since

ventromedial prefrontal cortex (vmPFC) is associated with subjective value computations of competing

choice options[44] and action outcomes[45], activity in this region may also contribute to reduced

prosocial decision making in NPD. Moreover, the vmPFC tracks state self-esteem[46,47], social inequality

assessment[48,49], and guides prosocial behavior in concert with the temporoparietal junction

(TPJ)[37,50,51]. The TPJ includes regions at the border of temporal and parietal lobes including superior

temporal sulcus (STS) and angular gyrus (AG)[52], implicated in higher-level social-cognitive

processes[53,54] such as moral judgement[55] or perspective taking[30]. Last, the anterior insula (AI) is

involved in interoceptive awareness[56], empathic processes[30,57], detection of social norm

violations[42]. Together with ACC, it serves the neural processing of interpersonal motives[58].

Here, patients diagnosed with NPD and a group of healthy control subjects (CTL) were scanned

using functional magnetic resonance imaging (fMRI) during a prosocial decision-making task. On each trial,

participants had two options to split a monetary endowment between themselves and an unknown

recipient. The splits systematically varied to manipulate the level of conflict between selfish and prosocial

motives. Additionally, some trials allowed punishing the recipient by taking money from them. Concluding

from previous studies in non-clinical samples with narcissistic traits [27–29], we expected lower generosity

in the group of patients diagnosed with NPD than in CTL group, reflecting their decision making to be

eventually governed by selfish motives. Additionally, we expected activity in neural conflict monitoring

systems (i.e. dmPFC, ACC, SMC)[39] and differential RT patterns[34,36] to track the motivational conflict

implied in the offer, potentially allowing to differentiate patients with NPD from CTL subjects.

                                                                     Methods

    Sample

        A total of N=42 participants were included in the study. Participants belonging to the NPD group

    were recruited from Charité - Universitätsmedizin Berlin, Campus Benjamin Franklin, and were both

    inpatients and outpatients, but were included in the study only after being discharged from treatment.

    CTL group participants were recruited by announcements in the press and via publicly placed

    advertisements. The study was approved by the local ethics committee at Charité Berlin (EA4/092/10).

    Participants gave written informed consent and received monetary compensation for participating in the

    study. Structured Clinical Interviews for DSM IV axes I and II Disorders (SCID-I; SCID-II; 60,61; German

    Versions: 62,64) were administered to all participants by a trained clinical psychologist to validate NPD

    diagnosis and assess comorbidity (Supplementary Table 1). Potential participants for the CTL group were

    excluded if completion of SCID-I or SCID-II indicated the presence of any mental disorder. As assessed in

    prior studies of our group, interrater reliability (κ=0.80) and internal consistency (Cronbach's α=0.86; sum

    of criteria) of SCID-II NPD diagnoses was acceptable [63]. Five participants were excluded from the analysis

    due to various reasons (Supplementary Methods), leaving a total of 37 participants (CTL: n=19, NPD: n=18)

    in the final analysis sample.

    Procedure

        Before the study, a set of paper-pencil questionnaires were sent to all participants (Supplementary

    Table 1). Upon arrival at the MRI facility, all participants were briefed about the study protocol and gave

    written informed consent. They were informed that inside the MRI they were going to repeatedly decide

    how to share an endowment of 1000 points (equivalent to 10 €) between themselves and another person

    (henceforth: recipient) by choosing between two alternative splits of this endowment. To ensure

    interpersonal relevance of the task, they were informed that after task completion, a number of their

    choices would be selected randomly, defining how much money would be paid to them and to the next

study participant (i.e. the recipient). Additionally, participants received money based on the choices of one

other participant who was assigned randomly. All participants were tested alone and never met any other

participant.

Experimental Task

Stimulus presentation was performed using the *Presentation* software package (Neurobehavioral

Systems, Albany, CA). On each trial, one option was displayed on the left and the other on the right side

of the screen, with the amounts for the participant and the recipient on the top and the bottom,

respectively (Figure 1). Participants chose by pressing a button with either the index or the middle finger.

In a baseline (BL) condition, both options assigned precisely 500 points to each person. In five other

conditions the level of conflict between selfish and prosocial motives was manipulated by systematically

splitting the endowment, with one option favoring the participant more strongly (selfish option) than the

other option (prosocial option). In one condition, options were very similar, inducing low motivational

conflict (low conflict, LC). On other trials, one option assigned a high payoff to the participant, leaving little

to the recipient, whereas the other option proposed a fairer split (medium conflict, MC). Still other trials

did not allow to resolve motivational conflict this way, because avoiding selfishness required to choose an

extremely prosocial option (high conflict, HC). Further, one condition allowed participants to increase their

gain by subtracting money from the recipient, or to resort to a fairer split (medium conflict punishment,

MCP). Finally, one condition also allowed participants to increase their payoff at the recipient's cost, but

the alternative would be to choose an extremely prosocial option (high conflict punishment, HCP). On each

trial, following 500 ms fixation, a choice had to be made within the 3000 ms display of the options,

otherwise the trial was coded as invalid. Each trial onset was triggered by an MRI pulse, and the inter-trial-

interval (ITI) was jittered between 500 and 6500 ms in steps of 2000 ms (further information in

Supplementary Methods).

Statistical analyses of behavioral data

Statistical analyses of behavioral data were performed using JASP[64], jamovi[65], and the

p.adjust() function in R[66]. We analyzed percentages of prosocial choices and median RTs for the different

experimental conditions using repeated measures analysis of variance (ANOVA) and nonparametric tests,

as appropriate. Since both choice options in the BL condition offered the same amount of 500 points to

the player and the recipient, no selfish or prosocial option was defined and BL choice data were not

analyzed. However, RTs from the BL condition were analyzed and it served as an explicit baseline for the

analysis of fMRI data. To test the effects of increasing motivational conflict on behavior and neural activity,

we compared HC trials against the average of LC and MC trials (LC&MC), because the latter both allowed

to resolve the conflict with the same pro-social option in which the participant still received more payoff

but amounts were rather similar (i.e. 550 and 450, respectively). This was not the case in the HC condition,

in which the pro-social option implied a relative loss for the participant and gain for the recipient (200 and

800, respectively). Last, to test the main effect of possible punishment, we averaged MCP and HCP trials

(PUN) and compared this to the average of MC and HC trials (noPUN).

FMRI data acquisition and preprocessing

Echo-planar images (EPIs) were acquired on a Magnetom TrioTim Syngo (Siemens, Munich,

Germany). The entire run consisted of 600 EPIs (37 ascending slices; 20% gap; 3*3*3 mm voxels; TR=2000

ms; TE=30 ms; FA=70°; FoV=192; GRAPPA=2). Additionally, a high-resolution T1-weighted MPRAGE image

(176 ascending slices; 50% gap; 1*1*1 mm voxels; TR=1900 ms; TE=2.52 ms; FA=9°; FoV=256; GRAPPA=2)

was acquired to improve spatial normalization. All fMRI data were preprocessed and analyzed using SPM

12[67] in Matlab R2019b (The Mathworks, Natick, Massachusetts, USA). EPIs were corrected for

acquisition delay (slice timing), with the middle slice as reference, and for head motion (realignment). The

MPRAGE image was coregistered to the mean EPI obtained from realignment and segmented using the

unified segmentation algorithm implemented in SPM. EPIs were normalized to a common stereotactic

reference frame (MNI) space using the transformation parameters obtained from segmentation, and

smoothed using an 8 mm full-width-at-half-maximum isotropic Gaussian kernel.

Statistical analysis of fMRI data

All fMRI data were analyzed using SPM12[67], using a two-stage mixed effects model. The first-

level general linear model (GLM) contained six regressors of interest, one for the BL condition and one for

each of the Choice conditions, plus the six rigid-body movement parameters obtained from realignment

as regressors of no interest, to account for variance related to head motion, and a single constant. At the

second level, we constructed a full-factorial ANOVA model, which included the contrast images of each

Choice condition against the BL condition (e.g. HC>BL), separated by group, resulting in the within-subjects

factor *condition* with five levels, and the between-subjects factor *group* (CTL, NPD).

To test for within-subject effects of RTs and prosocial choices, respectively, we set up two

additional GLMs for each participant. Within each GLM, we modelled BL trials separately from Choice

trials, and parametrically modulated, in this order, the latter with dummy predictors that indicated

whether medium (MC, MCP) or high (HC, HCP) conflict trials were presented, and whether or not a given

trial was a PUN trial (i.e., MCP, HCP). Last, we included either a predictor indicating the participant's choice

(prosocial=1; selfish=-1), or the RT on each trial (log-transformed and z-scored). Parametric modulators

were orthogonalized. Additional regressors modelled invalid trials and head motion. One participant in the

CTL group and three participants in the NPD group never chose the prosocial option and were therefore

excluded from these analyses.

Region-of-Interest Analysis

Since our primary focus was to investigate how motivational conflict engages regions associated

with conflict monitoring, we used Neurosynth[68] to perform an automated meta-analysis of functional

neuroimaging studies by searching for the term "conflict" and obtaining the association test map

thresholded at $p<.01$ (false-discovery-rate-corrected; 337 studies). This map was then limited to include

only clusters of 5 or more contiguous voxels and used as a region-of-interest, largely covering dorsal ACC,

SMC, and aspects of bilateral AI (henceforth: *conflict mask*).

<p style="text-align:center">**Results**</p>

Participants in NPD Group Acted More Selfishly, but Avoided Punishing Recipients

Across all Choice conditions, the NPD group (*MD*=9.80%, IQR=24.13%, *min*=0.0%, *max*=97.81%) chose the prosocial option less often than the CTL group (*MD*=45.20%, IQR=34.74%, *min*=0.0%, *max*=86.41%; Mann-Whitney-U test: *W*=260.5, one-sided *p*=.003, rank-biserial correlation=0.52, 90% *CI*=[0.26;0.72]; Figure 2 & Table 1).

When motivational conflict was low, the CTL group behaved more prosocially than the NPD group (LC&MC, CTL group: *MD*=51.11%, *IQR*=59.53; NPD group: *MD*=3.97%, *IQR*=24.58; *W*=266.5, one-sided *p*=0.002, rank-biserial correlation=0.56, 90%, CI=[0.30;0.74]). When conflict increased, prosocial choices diminished in both groups (difference: LC&MC minus HC; CTL group: *MD*=29.69%, *IQR*=46.40, *W*=171.00, *p*<.001; NPD group: *MD*=1.20%, *IQR*=5.43, *W*=100.5, *p*=.003), and this effect was larger in the CTL group (*W*=288.50, two-sided *p*<.001, rank-biserial correlation=0.69, 95% *CI*=[0.43;0.84]). However, under HC, the CTL group still behaved more prosocially than the NPD group (CTL: *MD*=15.22%, *IQR*=25.65%; NPD: *MD*=0.00%, *IQR*=22.23%; *W*=229, one-sided, Holm-corrected *p*=0.046, rank-biserial correlation=0.34, 90% *CI*=[0.04;0.59]). Thus, the difference in prosociality between the groups was reduced, but still present, when prosocial behavior was very disadvantageous for the participant.

Across groups, prosociality increased when selfishness required to punish the recipient (difference PUN minus noPUN, CTL: *MD*=26.37%, *IQR*=48.80, *W*=158, one-sided *p*<.001, rank-biserial correlation=0.85, 90% *CI*=[0.67;0.93]; NPD: *MD*=1.63%, *IQR*=32.15, *W*=87, one-sided *p*=.016, rank-biserial correlation=0.66, 95% *CI*=[0.28;0.86]). This increase was descriptively, but not significantly, larger in the CTL than in the NPD group (*W*=219, one-sided *p*=.074, rank-biserial correlation=0.28, 90% *CI*=[-0.03;0.54]). Hence, having to punish others for one's own benefit led to more prosocial behavior.

Responses in NPD Group were Fast, but Slowed Down for Prosocial Decisions

RTs did not generally differ between groups (ANOVA with factors trial type [BL, Choice] and group [CTL,

NPD]: $F(1,35)=3.39$, $p=.074$, $\eta_p^2=.09$). However, there was a significant main effect of trial type

($F(1,35)=37.15$, $p<.001$, $\eta_p^2=.52$), that was modulated by group (trial type*group interaction: $F(1,35)=11.09$,

$p=.002$, $\eta_p^2=.24$; Figure 2). Post-hoc tests showed that the CTL group responded slower during Choice trials

($t(54.2)=3.21$, two-sided $p_{Holm}=.009$), but there was no difference during BL trials ($t(54.2)=0.02$, two-sided

$p_{Holm}=.987$). Moreover, motivational conflict slowed RTs in the CTL group (Choice>BL: $t(35)=6.76$,

$p_{Holm}<.001$), but not in the NPD group ($t(35)=1.93$, $p_{Holm}=.186$). Thus, both groups responsed comparably

fast unless interpersonal motives needed to be reconciled, which slowed responding only in the CTL group.

Median RTs did not differ between selfish and prosocial decisions across groups (ANOVA with

factors decision [selfish, prosocial] and group [CTL, NPD]: no main effect of decision: $F(1,31)=4.03$, $p=.054$,

$\eta_p^2=.12$; factor group: $F(1,31)=2.91$, $p=.098$, $\eta_p^2=.09$). However, the effect of decision differed between

groups (decision*group: $F(1,32)=8.56$, $p=.006$, $\eta_p^2=.22$; Figure 3), showing that the CTL group was slower

than the NPD group during selfish choices (CTL>NPD: $t(49.2)=2.93$, $p_{Holm}=.026$; Figure 3), but not during

prosocial choices (CTL>NPD: $t(49.2)=0.03$, $p_{Holm}=1.000$). Moreover, RTs within the CTL group did not differ

between decisions ($t(31)=0.68$, $p_{Holm}=1.000$), but the NPD group was faster for selfish than for prosocial

decisions ($t(31)=-3.34$, $p_{Holm}=.013$), which was not due to group-specific effects of the experimental

conditions (Supplementary Table S4).

RTs were insensitive to level of motivational conflict across groups (ANOVA with factors level of

conflict [LC&MC, HC] and group [CTL, NPD]: $F(1,35)=1.86$, $p=.181$, $\eta_p^2=.05$). Yet, we found a main effect of

group ($F(1,35)=8.53$, $p=.006$, $\eta_p^2=.20$), that was driven by the HC condition (CTL>NPD: $t(35)=3.21$,

$p_{Holm}=.017$). The interaction of both factors was not significant ($F(1,35)=3.50$, $p=.070$, $\eta_p^2=.09$). Thus, when

prosociality was disadvantageous for the participants, responses in the CTL group were slower than in the

NPD group.

Last, the option to punish did not impact reaction times across groups (ANOVA with factors

punishment [noPUN, PUN] and group [CTL, NPD]: $F(1,35)=0.00$, $p=.983$, $\eta_p^2=.00$). Again, the CTL group

responded slower than the NPD group (main effect of group: $F(1,35)=8.30$, $p=.007$, $\eta_p^2=.19$), which was due

to noPUN trials (CTL>NPD: $t(43.2)=3.06$, $p_{Holm}=.023$). No significant interaction was found

(punishment*group: $F(1,35)=1.06$, $p=.311$, $\eta_p^2=.03$).

Conflict-Related Neural Activity During Prosocial Decision Making Was Reduced in NPD Group

Across groups, presence of motivational conflict (Choice>BL) elicited activity in right dorsal midcingulate

cortex (MNI x,y,z: 6,30,42; $T=5.79$, $k=68$, $p_{FWE}=.001$; whole-brain), right dorsolateral prefrontal cortex

(dlPFC; 46,34,22; $T=6.45$, $k=146$, $p_{FWE}<.001$), bilateral parietal cortex (right: 28,-60,44; $T=9.32$, $k=1833$,

$p_{FWE}<.001$; left: -26,-70,32; $T=9.63$, $k=1349$, $p_{FWE}<.001$), and further regions (see Supplementary

Information for full activation coordinates). This effect was stronger for the CTL than the NPD group in left

dorsomedial prefrontal cortex (dmPFC; -6,30,38; $T=3.86$, $k=29$, $p_{FWE}=.023$) and left preSMA (-4,12,52;

$T=4.44$, $k=10$, $p_{FWE}=.003$; -corrected inside the *conflict* mask, Figure 4), while the opposite comparison (i.e.

Choice>BL * NPD>CTL) did not yield any significant results, indicating that considering potential

interpersonal consequences is linked to conflict-related neural activity particularly in the CTL group.

Contrasting HC against the average of LC and MC (HC>LC&MC) to test for effects of increasing

motivational conflict on neural activity did not yield any significant effects or interactions.

When selfish behavior required punishing others (PUN>noPUN), activity increased in right AG (34,-

68,52; $T=5.35$, $k=73$, $p_{FWE}=.006$; whole-brain), right inferior temporal gyrus (ITG; 54,-56,-10; $T=5.24$, $k=19$,

$p_{FWE}=.009$), and right superior parietal cortex (26,-72,54; $T=4.84$, $k=1$, $p_{FWE}=.043$). The effect in right ITG

was significant when performing this analysis in the CTL group alone (48,-56,-8; $T=5.20$, $k=33$, $p_{FWE}=.004$;

whole-brain), which also showed significantly stronger activations for PUN vs noPUN trials in left AI (-

32,18,-14; $T=5.25$, $k=14$, $p_{FWE}=.009$), right inferior frontal gyrus (50,20,16; $T=4.89$, $k=3$, $p_{FWE}=.037$), and right

middle temporal gyrus (62,-44,2; $T=5.26$, $k=35$, $p_{FWE}=.008$). However, despite no significant effects in the

NPD group, PUN>noPUN did not differ between groups. Hence, when selfish behavior was possible only

by punishing others, this induced stronger activity in brain regions like right AG and ITG, which was driven

by the CTL group.

Opportunities to Punish Increased Conflict-Related Neural Activity Across Groups

To predict task-related neural activity with individual differences in prosociality, we regressed the first-

level Choice>BL contrast images on the participants' mean percentages of prosocial choices while

controlling for group differences. Across groups, higher prosociality predicted increasing activity in right

dlPFC (36,38,14; T=5.80, k=3, p_{FWE}=.040; whole-brain), but no group differences were found. Regressing

the contrast HC>LC&MC on the corresponding difference in prosociality (i.e., HC minus LC&MC) did not

yield any significant effects. However, increasing prosociality from noPUN to PUN trials predicted

corresponding increases in neural activity in right SMA (6,6,48; T=6.16, k=3, p_{FWE}=.018; whole-brain) across

groups. Thus, higher overall prosociality predicted neural activity in right dlPFC, and increased prosociality

in order not to punish others was linked to increased SMA activity across groups.

Since participants in the NPD group showed low overall prosociality and responded more slowly

during prosocial than during selfish choices, we expected stronger trial-by-trial conflict-related neural

activity for prosocial choices in the NPD than in the CTL group[34,36,39], but no such effect was found.

Reaction Times Predicted Conflict-Related Neural Activity Within and Across Participants

Last, under motivational conflict, slower participants showed higher activity in right mid cingulate gyrus

(Choice>BL: 4,32,36; T=5.37, k=53, p_{FWE}=.001; corrected inside the *conflict* mask), left AI (-38,18,2; T=5.87,

k=22, p_{FWE}<.001), and left SMA (-4,14,50; T=5.37, k=111, p_{FWE}=.001). Similarly, slower RTs due to increasing

conflict were associated with higher neural activity in SMA (HC>LC&MC: 2,16,44; T=6.56, k=224, p_{FWE}<.001;

corrected inside the *conflict* mask), and left (-32,16,2; T=5.9, k=16, p_{FWE}<.001) and right AI (40,16,0; T=4.73,

k=6, p_{FWE}=.008). Additionally, slower trial-by-trial RTs predicted neural activity in ACC, bilateral AI, and

lateral prefrontal and parietal cortices, and an exploratory *F*-test indicated that trial-by-trial RTs were less

predictive of mid cingulate neural activity in the NPD than in the CTL group (-6,26,36; *F*=18.05, *k*=1,

p_{FWE}=.048; corrected inside the *conflict* mask).

Discussion

In the present study we investigated how prosocial decision making and underlying neural processes in

patients diagnosed with NPD are shaped by conflicting interpersonal motives. Compared to control

subjects, the NPD group behaved less prosocially, responded faster, and showed less neural activity in

regions associated with conflict monitoring such as the SMA and cingulate cortex[38,39]. When

prosociality was disadvantageous for the participants, it diminished in both groups, although control

subjects still behaved more prosocially than the NPD group. Both groups were similarly affected by the

requirement to punish others for one's own benefit, which shifted behavior away from selfishness, and

increased neural activity in right AG and ITG. Last, when participants in the NPD group acted prosocially,

thus deviating from their selfish mode of behavior, this slowed responding to a level comparable to the

CTL group.

Our finding of low generosity in participants diagnosed with NPD is well in line with existing

literature from non-clinical narcissistic samples[27,28]. We hypothesized that this may either result from

more readily available tendencies towards selfishness[35] or, alternatively, from cognitively effortful

arbitration between conflicting interpersonal motives; i.e. either to selfishly increase one's own gain or to

behave prosocially in order to be seen in a good light by others[9,10]. Our findings speak to the former

hypothesis, as the mainly selfish responses in the NPD group were generally fast, which was paralleled by

comparably low conflict-related neural activity[38,39], and rare prosocial choices were considerably

slower. Participants in the CTL group more readily utilized the fairness information provided in each trial

to guide their decisions, causing motivational conflict and the need to engage in cognitive control[34,39].

In sum, our data nicely coincide with earlier research showing that deviations from selfish or prosocial

habits are marked by slowing RTs[36]. Thus, using response times one may reveal individual differences in

social value orientations[35], with participants in the NPD group being more strongly oriented toward

selfishness.

Notably, behavior in the NPD group was not fully insensitive to its potential social consequences.

Similar to the CTL group, participants in the NPD group acted more selfishly when prosociality was believed

to be highly detrimental, and acted more prosocially when their own benefit was conditional on punishing

others. Prosocial behavior can be shaped by inducing prosocial motives[58], and explicit instructions to

consider others' perspectives can increase prosocial responding in non-clinical narcissistic individuals[69].

Although we did not provide such explicit instructions, our task was effective for pitting selfish against

prosocial motives. Notably, recipients were anonymous, which is known to impact behavior in prosocial

decision games[70], and did not have an option to reciprocate, invalidating any potential economic

repercussions of unfair behavior. Despite these circumstances, patients diagnosed with NPD showed some

responsiveness to others' potential needs, underlining that pathological narcissism does not equate to an

absolute incapacity, but perhaps a reduced propensity to act prosocially[71]. This finding points to the

potential of empathy interventions to increase perspective taking[72] for increasing prosocial

behavior[58], which could be reflected in neural activity in networks mediating interpersonal

motivations[32,58,71]. Such interventions could entail practicing to actively consider how one's social

behavior affects others, and to imagine how others will respond during interpersonal interactions. In

consequence, such therapeutic interventions could elicit higher prosociality and more favorable

evaluations from others, stabilizing self-esteem[46,73] and overall well-being[74].

Our findings also speak to neural underpinnings of prosociality more generally. Precisely, neural

activity in SMA, which is linked to cognitive control of behavior[38], was higher for participants who

favored prosociality over punishing others for their own benefit. When such punishment was possible, this

increased activity in AG, which may reflect more specifically social-cognitive processes like moral

reasoning[55], or domain-general cognitive processes like attentional reorienting[75] that may also be

engaged during social behavior. In addition, ITG was activated when punishment was possible, which has

previously been associated with fairness considerations[42]. Last, participants with higher overall levels of

prosociality showed increased activity in dlPFC, which was also responsive to the general presence of

conflicting interpersonal motives, alongside other regions of the cognitive control network (i.e. dmPFC, AI,

ACC, SMA, and parietal cortex).

So far, few studies have investigated the neural correlates of interpersonal behavior in NPD using

experimental paradigms in combination with fMRI. By doing so, our data contribute to a more detailed

understanding of NPD in terms of impairments in interpersonal functioning, as introduced by DSM-5,

section 3 [1]. Collectively, our data indicate that selfish behavior in NPD relies on diminished consideration

of prosocial motives, which obviates the need to resolve conflict with selfish tendencies. However, this

may reflect a reduced propensity, not ability, to act prosocially[69,71]. Potentially, this reflects differences

between NPD and, for instance, antisocial personality disorder, which could be tested explicitly in future

studies. In terms of treatment implications, our results point to possibilities for enhancing prosocial

behavior and thereby improving social relationships and well-being in NPD. Regarding social cognition and

interpersonal behavior more broadly, our findings support the notion that these functions rely on domain-

general control processes [38–40], that draw on contextual social information to decide between generous

or selfish behavior[43].

Acknowledgements

We would like to thank Sylvia Richter and Thomas Münte for helpful discussions during

development of the task and Michael Scholz for his help with the optimization of the trial sequence for the

fMRI task. We further thank Jürgen Baudewig, Ricardo Heydenblut, and Christian Kainz for assistance with

MRI data acquisition. Research leading to this publication has been funded by the Deutsche

Forschungsgemeinschaft (Sören Krach; KR 3803/9-1; Aline Vater and Stefan Roepke; EXC 302; Björn Schott;

SFB 779, TP A08).

Disclosures

The authors declare no conflicts of interest.

Supplementary Information

Supplementary information is available at MP's website.

References

1. American Psychiatric Association (2013): *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. Washington, DC: American Psychiatric Association. https://doi.org/10.1176/appi.books.9780890425596

2. Roepke S, Vater A (2014): Narcissistic personality disorder: An integrative review of recent empirical data and current definitions. *Curr Psychiatry Rep* 16. https://doi.org/10.1007/s11920-014-0445-0

3. Cain NM, Pincus AL, Ansell EB (2008): Narcissism at the crossroads: Phenotypic description of pathological narcissism across clinical theory, social/personality psychology, and psychiatric diagnosis. *Clin Psychol Rev* 28: 638–656.

4. Koepernik T, Jauk E, Kanske P (2021): Lay theories of grandiose and vulnerable narcissism. *Curr Psychol*. https://doi.org/10.1007/s12144-020-01296-w

5. Morf CC, Rhodewalt F (2001): Unraveling the paradoxes of narcissism: A dynamic self-regulatory processing model. *Psychol Inq* 12: 177–196.

6. Campbell WK, Foster JD (2007): The Narcissistic Self: Background, an Extended Agency Model, and Ongoing Controversies. In: Sedikides C, Spencer J, editors. *The Self*. New York: Psychology Press, pp 115–138.

7. Farwell L, Wohlwend-Lloyd R (1998): Narcissistic Processes: Optimistic Expectations, Favorable Self-Evaluations, and Self-Enhancing Attributions. *J Pers* 66: 65–83.

8. Paulhus DL, Williams KM (2002): The Dark Triad of personality: Narcissism, Machiavellianism, and psychopathy. *J Res Pers* 36: 556–563.

9. Back MD, Küfner ACP, Dufner M, Gerlach TM, Rauthmann JF, Denissen JJA (2013): Narcissistic admiration and rivalry: Disentangling the bright and dark sides of narcissism. *J Pers Soc Psychol* 105: 1013–1037.

10. Hart W, Adams J, Burton KA, Tortoriello GK (2017): Narcissism and self-presentation: Profiling grandiose and vulnerable Narcissists' self-presentation tactic use. *Pers Individ Dif* 104: 48–57.

11. Campbell WK, Brunell AB, Finkel EJ (2006): Narcissism, Interpersonal Self-Regulation, and Romantic

Relationships: An Agency Model Approach. In: Vohs KD, Finkel EJ, editors. *Self and Relationships:*

Connecting Intrapersonal and Interpersonal Processes. New York, NY: Guilford Press, pp 57–83.

12. Paulhus DL (1998): Interpersonal and intrapsychic adaptiveness of trait self-enhancement: A mixed

blessing? *J Pers Soc Psychol* 74: 1197–1208.

13. Stucke TS, Sporer SL (2002): When a grandiose self-image is threatened: Narcissism and self-concept

clarity as predictors of negative emotions and aggression following ego-threat. *J Pers* 70: 509–532.

14. Ferriday C, Vartanian O, Mandel DR (2011): Public but not private ego threat triggers aggression in

narcissists. *Eur J Soc Psychol* 41: 564–568.

15. Dickinson KA, Pincus AL (2003): Interpersonal Analysis of Grandiose and Vulnerable Narcissism. *J Pers*

Disord 17: 188–207.

16. Fatfouta R, Gerlach TM, Schröder-Abé M, Merkl A (2015): Narcissism and lack of interpersonal

forgiveness: The mediating role of state anger, state rumination, and state empathy. *Pers Individ Dif*

75: 36–40.

17. Wurst SN, Gerlach TM, Dufner M, Rauthmann JF, Grosz MP, Küfner ACP, *et al.* (2017): Narcissism and

romantic relationships: The differential impact of narcissistic admiration and rivalry. *J Pers Soc*

Psychol 112: 280–306.

18. Sauls D, Zeigler-Hill V, Vrabel JK, Lehtman MJ (2019): How do narcissists get what they want from

their romantic partners? The connections that narcissistic admiration and narcissistic rivalry have

with influence strategies. *Pers Individ Dif* 147: 33–42.

19. Miller JD, Hoffman BJ, Gaughan ET, Gentile B, Maples J, Keith Campbell W (2011): Grandiose and

Vulnerable Narcissism: A Nomological Network Analysis. *J Pers* 79: 1013–1042.

20. Crowe ML, Edershile EA, Wright AGC, Campbell WK, Lynam DR, Miller JD (2018): Development and

validation of the Narcissistic Vulnerability Scale: An adjective rating scale. *Psychol Assess* 30: 978–

983.

21. Pincus AL, Ansell EB, Pimentel CA, Cain NM, Wright AGC, Levy KN (2009): Initial Construction and

Validation of the Pathological Narcissism Inventory. *Psychol Assess* 21: 365–379.

22. Vater A, Schröder-Abé M, Ritter K, Renneberg B, Schulze L, Bosson JK, Roepke S (2013): The

Narcissistic Personality Inventory: A Useful Tool for Assessing Pathological Narcissism? Evidence

From Patients With Narcissistic Personality Disorder. *J Pers Assess* 95: 301–308.

23. Aslinger EN, Manuck SB, Pilkonis PA, Simms LJ, Wright AGC (2018): Narcissist or narcissistic?

Evaluation of the latent structure of narcissistic personality disorder. *J Abnorm Psychol* 127: 496–

502.

24. Ritter K, Dziobek I, Preißler S, Rüter A, Vater A, Fydrich T, *et al.* (2011): Lack of empathy in patients

with narcissistic personality disorder. *Psychiatry Res* 187: 241–247.

25. Schulze L, Dziobek I, Vater A, Heekeren HR, Bajbouj M, Renneberg B, *et al.* (2013): Gray matter

abnormalities in patients with narcissistic personality disorder. *J Psychiatr Res* 47: 1363–1369.

26. Miller JD, Campbell WK, Pilkonis PA (2007): Narcissistic personality disorder: relations with distress

and functional impairment. *Compr Psychiatry* 48: 170–177.

27. Campbell WK, Bush CP, Brunell AB, Shelton J (2005): Understanding the social costs of narcissism:

The case of the tragedy of the commons. *Personal Soc Psychol Bull* 31: 1358–1368.

28. Böckler A, Sharifi M, Kanske P, Dziobek I, Singer T (2017): Social decision making in narcissism:

Reduced generosity and increased retaliation are driven by alterations in perspective-taking and

anger. *Pers Individ Dif* 104: 1–7.

29. Bushman BJ, Baumeister RF (1998): Threatened Egotism, Narcissism, Self-Esteem, and Direct and

Displaced Aggression: Does Self-Love or Self-Hate Lead to Violence. *J Pers Soc Psychol* 75: 219–229.

30. Tusche A, Böckler A, Kanske P, Trautwein F-M, Singer T (2016): Decoding the Charitable Brain:

Empathy, Perspective Taking, and Attention Shifts Differentially Predict Altruistic Giving. *J Neurosci*

36: 4719–4732.

31. Paciello M, Fida R, Cerniglia L, Tramontano C, Cole E (2013): High cost helping scenario: The role of

empathy, prosocial reasoning and moral disengagement on helping behavior. *Pers Individ Dif* 55: 3–

7.

32. Gallo S, Paracampo R, Müller-Pinzler L, Severo MC, Blömer L, Fernandes-Henriques C, *et al.* (2018):

The causal role of the somatosensory cortex in prosocial behaviour. *Elife* 7: 1–31.

33. Carlo G, Mestre MV, Samper P, Tur A, Armenta BE (2010): Feelings or cognitions? Moral cognitions

and emotions as longitudinal predictors of prosocial and aggressive behaviors. *Pers Individ Dif* 48:

872–877.

34. Evans AM, Dillon KD, Rand DG (2015): Fast But Not Intuitive, Slow But Not Reflective: Decision

Conflict Drives Reaction Times in Social Dilemmas. *J Exp Psychol Gen* 144: 951–966.

35. Cornelissen G, Dewitte S, Warlop L (2011): Are Social Value Orientations Expressed Automatically?

Decision Making in the Dictator Game. *Personal Soc Psychol Bull* 37: 1080–1090.

36. Yamagishi T, Matsumoto Y, Kiyonari T, Takagishi H, Li Y, Kanai R, Sakagami M (2017): Response time

in economic games reflects different types of decision conflict for prosocial and proself individuals.

Proc Natl Acad Sci 114: 6394–6399.

37. Hutcherson CA, Bushong B, Rangel A (2015): A Neurocomputational Model of Altruistic Choice and Its

Implications. *Neuron* 87: 451–462.

38. Nachev P, Kennard C, Husain M (2008): Functional role of the supplementary and pre-supplementary

motor areas. *Nat Rev Neurosci* 9: 856–869.

39. Botvinick MM, Cohen JD, Carter CS (2004): Conflict monitoring and anterior cingulate cortex: An

update. *Trends Cogn Sci* 8: 539–46.

40. Bonini F, Burle B, Liegeois-Chauvel C, Regis J, Chauvel P, Vidal F (2014): Action Monitoring and Medial

Frontal Cortex: Leading Role of Supplementary Motor Area. *Science (80-)* 343: 888–891.

41. Cutler J, Campbell-Meiklejohn D (2019): A comparative fMRI meta-analysis of altruistic and strategic

decisions to give. *Neuroimage* 184: 227–241.

42. Feng C, Luo YJ, Krueger F (2015): Neural signatures of fairness-related normative decision making in

the ultimatum game: A coordinate-based meta-analysis. *Hum Brain Mapp* 36: 591–602.

43. Zaki J, Hennigan K, Weber J, Ochsner KN (2010): Social cognitive conflict resolution: Contributions of

domain-general and domain-specific neural systems. *J Neurosci* 30: 8481–8488.

44. Bartra O, McGuire JT, Kable JW (2013): The valuation system: A coordinate-based meta-analysis of

BOLD fMRI experiments examining neural correlates of subjective value. *Neuroimage* 76: 412–427.

45. Stolz DS, Müller-Pinzler L, Krach S, Paulus FM (2020): Internal control beliefs shape positive affect and

associated neural dynamics during outcome valuation. *Nat Commun* 11: 1230.

46. Will G-J, Rutledge RB, Moutoussis M, Dolan RJ (2017): Neural and computational processes

underlying dynamic changes in self-esteem. *Elife* 6: 1–21.

47. Eisenberger NI, Inagaki TK, Muscatell K a., Byrne Haltom KE, Leary MR (2011): The Neural Sociometer:

Brain Mechanisms Underlying State Self-esteem. *J Cogn Neurosci* 23: 3448–3455.

48. Tricomi E, Rangel A, Camerer CF, O'Doherty JP (2010): Neural evidence for inequality-averse social

preferences. *Nature* 463: 1089–1091.

49. Aoki R, Matsumoto M, Yomogida Y, Izuma K, Murayama K, Sugiura A, *et al.* (2014): Social equality in

the number of choice options is represented in the ventromedial prefrontal cortex. *J Neurosci* 34:

6413–6421.

50. Strombach T, Weber B, Hangebrauk Z, Kenning P, Karipidis II, Tobler PN, Kalenscher T (2014): Social

discounting involves modulation of neural value signals by temporoparietal junction. *Proc Natl Acad*

Sci 112: 1619–1624.

51. Park SQ, Kahnt T, Dogan A, Strang S, Fehr E, Tobler PN (2017): A neural link between generosity and

happiness. *Nat Commun* 8: 15964.

52. Schurz M, Tholen MG, Perner J, Mars RB, Sallet J (2017): Specifying the brain anatomy underlying

temporo-parietal junction activations for theory of mind: A review using probabilistic atlases from

different imaging modalities. *Hum Brain Mapp* 38: 4788–4805.

53. Decety J, Lamm C (2007): The role of the right temporoparietal junction in social interaction: How

low-level computational processes contribute to meta-cognition. *Neuroscientist* 13: 580–593.

54. Saxe R, Kanwisher N (2003): People thinking about thinking people: The role of the temporo-parietal

junction in "theory of mind." *Neuroimage* 19: 1835–1842.

55. Young L, Camprodon JA, Hauser M, Pascual-Leone A, Saxe R (2010): Disruption of the right

temporoparietal junction with transcranial magnetic stimulation reduces the role of beliefs in moral

judgments. *Proc Natl Acad Sci* 107: 6753–6758.

56. A.D. Craig (2002): How do you feel? Interoception: the sense of the physiological condition of the

body. *Nat Rev Neurosci* 3: 655–666.

57. Kanske P, Böckler A, Trautwein FM, Singer T (2015): Dissecting the social brain: Introducing the

EmpaToM to reveal distinct neural networks and brain-behavior relations for empathy and Theory

of Mind. *Neuroimage* 122: 6–19.

58. Hein G, Morishima Y, Leiberg S, Sul S, Fehr E (2016): The brain's functional network architecture

reveals human motives. *Science (80-)* 351: 84–90.

59. First M, Spitzer R, Gibbon M, Williams JBW (1996): *Structured Clinical Interview for DSM-IV Axis I*

Disorders (SCID-I). New York: New York State Psychiatric Institute, Biometrics Research Department.

60. First MB, Gibbon M, Spitzer RL, Williams JBW, Benjamin L (1997): *Structured Clinical Interview for*

DSM–IV Personality Disorders (SCID–II). Washington, DC: American Psychiatric Press.

61. Wittchen HU, Zaudig M, Fydrich T (1997): *SKID-I - Strukturiertes Klinisches Interview Für DSM–IV–TR -*

Achse I [Structured Clinical Interview for DSM–IV–TR - Axis I]. Göttingen: Hogrefe.

62. Fydrich T, Renneberg B, Schmitz B, Wittchen H-U (1997): *SKID-II - Strukturiertes Klinisches Interview*

Für DSM–IV Achse II: Persönlichkeitsstörungen [SCID II, Structured Clinical Interview for DSM–IV Axis

II: Personality Disorders]. Göttingen: Hogrefe.

63. Ritter K, Vater A, Rüsch N, Schröder-Abé M, Schütz A, Fydrich T, *et al.* (2014): Shame in patients with

narcissistic personality disorder. *Psychiatry Res* 215: 429–437.

64. JASP Team (2019): JASP (Version 0.11.1).

65. The jamovi project (2020): jamovi [no. 1.2]. Retrieved from https://www.jamovi.org

66. R Core Team (2013): R: A language and environment for statistical computing. Vienna, Austria,

Austria: R Foundation for Statistical Computing.

67. Ashburner J, Barnes G, Chen C, Daunizeau J, Moran R, Henson R, *et al.* (, February): SPM12 Manual.

https://doi.org/10.1111/j.1365-294X.2006.02813.x

68. Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD (2011): Large-scale automated synthesis

of human functional neuroimaging data. *Nat Methods* 8: 665–670.

69. Hepper EG, Hart CM, Sedikides C (2014): Moving Narcissus: Can Narcissists Be Empathic? *Personal*

Soc Psychol Bull 40: 1079–1091.

70. Eckel CC, Grossman PJ (1996): Altruism in anonymous dictator games. *Games Econ Behav* 16: 181–

191.

71. Meffert H, Gazzola V, Den Boer JA, Bartels AAJ, Keysers C (2013): Reduced spontaneous but relatively

normal deliberate vicarious representations in psychopathy. *Brain* 136: 2550–2562.

72. Weisz E, Zaki J (2017): Empathy-Building Interventions: A Review of Existing Work and Suggestions for

Future Directions. In: Seppala EM, Simon-Thomas E, Brown SL, Worline MC, Cameron CD, Doty JR,

editors. *The Oxford Handbook of Compassion Science.* New York: Oxford University Press, pp 205–

218.

73. Leary MR (2005): Sociometer theory and the pursuit of relational value: Getting to the root of self-

esteem. *Eur Rev Soc Psychol* 16: 75–111.

74. Morelli SA, Lee IA, Arnn ME, Zaki J (2015): Emotional and instrumental support provision interact to

predict well-being. *Emotion* 15: 484–493.

75. Seghier ML (2013): The angular gyrus: Multiple functions and multiple subdivisions. *Neuroscientist*

19: 43–61.

    Figure legends

Figure 1. Structure of the experimental task. **Top.** Structure of a single trial. Following a jittered fixation

(500-6500 ms), a trial was initialized for 500 ms and followed by a 3000 ms display of the two possible

splits of the endowment of 1000 points (equivalent to 10€). Participants chose between options with a

button press. Responses slower than 3000 ms were discarded. Player A is the participant and Player B the

recipient. **Bottom.** Overview of all experimental conditions, with mean number of points assigned to each

player and the preset maximum deviation from the mean assignment (max +/-). The split assigning most

money to the participant was defined to be the selfish options, whereas the prosocial option was the one

splitting the endowment in a relatively more beneficial way for the recipient. Note that due to this

definition, the prosocial option did not necessarily assign more money to the recipient than to the

participant. There were 12 different, pseudorandomized trial sequences that were designed to optimize

BOLD efficiency. Thus, the number of trials in each sequence varied between 223 and 229 (see text for

details).

Figure 2. Behavioral data. **a.** Percentages of prosocial choices for the CTL group and NPD group overall

(left), for trials with low and medium (LC&MC) vs high motivational conflict (HC; middle), and separately

for trials without (noPUN: MC, HC), and with punishment (PUN: MCP, HCP). **b.** Median reaction times in

seconds for BL and Choice trials (left), for LC&MC trials and HC trials (middle), and separately for PUN and

noPUN trials. Where shown, small crossed bars represent interaction between group (CTL, NPD) and factor

on x-axis. Central bars of boxplots show median, lower and upper box borders show 25^{th} and 75^{th}

percentile and whiskers end at last data point within 1.5 times the interquartile range from lower or upper

box border.

Figure 3. Reaction times separated by decision. Small crossed bar represents interaction between group

(CTL, NPD) and decision made (selfish, prosocial). Small crossed bars represents interaction between group

(CTL, NPD) and choice (selfish, prosocial). Central bars of boxplots show median, lower and upper box

borders show 25th and 75th percentile and whiskers end at last data point within 1.5 times the interquartile

range from lower or upper box border.

Figure 4. Neural activity during the experimental task. **a.** Neural activity in regions associated with the term

conflict (green outline) including dorsomedial prefrontal cortex (dmPFC) and supplementary motor area

(SMA) was higher for the contrast Choice>BL (p<.05, whole-brain FWE-corrected; displayed at p<.0001,

uncorrected). This effect was larger in the CTL group than in the NPD group, visualized in the bar plots.

Neural activation data in bar plots are based on mean parameter estimate across all voxels in dmPFC and

SMA, respectively, in which the interaction of Choice>BL * CTL>NPD was significant at p<.05, small-volume

FWE-corrected inside the *conflict* mask). **b.** Neural activity in right angular gyrus (AG) and right inferior

temporal gyrus (ITG) was higher for PUN than for noPUN trials, but did not differ between groups. Bar

plots show mean parameter estimates across all voxels in right AG and right ITG in which the contrast

PUN>noPUN was significant at p<.05, whole-brain FWE-corrected (displayed at p<.005, uncorrected). Inset

in bottom right corner depicts activation in right AG, rotated around the y axis by 45 degrees. Errorbars

are +/- 1 standard error of the mean.

Tables

Table 1. Behavioral data by group and experimental condition

Choice behavior	CTL			NPD							
Condition	median % prosocial	[25th, 75th percentile]		median % prosocial	[25th,75th percentile]		W	one-sided p-value[1]	Rank-Biserial Correlation	90% *CI* for Rank-Biserial Correlation	
										Lower	Upper
Overall	45.20	35.30	70.03	9.80	1.38	25.51	260.5	.003	0.52	0.26	0.72
LC	53.49	17.58	79.18	4.65	0	28.33	246	.046	0.44	0.15	0.66
MC	53.33	29.35	88.04	2.30	0	19.99	273.5	.005	0.60	0.36	0.77
HC	15.22	4.35	30.00	0.00	0	22.23	229	.046	0.34	0.04	0.59
MCP	95.65	45.65	100.00	18.22	0	85.87	242	.046	0.42	0.13	0.64
HCP	79.17	37.47	97.83	7.03	0	79.19	245	.046	0.43	0.15	0.65

Reaction times	CTL		NPD						95% *CI* for *d*	
Condition	*Mean* (ms)	*SD*	*Mean* (ms)	*SD*	*t*	*df*	two-sided p-value[1]	*d*	Lower	Upper
Overall	1138.63	231.72	951.50	218.88	2.52	35	.016	0.83	0.15	1.50
BL	889.08	166.62	887.89	187.96	0.02	35	.984	0.01	-0.64	0.65
LC	1180.98	262.65	1060.97	283.77	1.34	35	.571	0.44	-0.22	1.09
MC	1164.63	301.72	950.72	275.30	2.25	35	.124	0.74	0.07	1.40
HC	1310.61	362.42	976.31	258.75	3.21	35	.017	1.06	0.36	1.74
MCP	1096.95	234.73	993.14	359.65	1.05	35	.606	0.34	-0.31	1.00
HCP	1393.84	414.36	1007.19	374.47	2.97	35	.027	0.98	0.29	1.66

Note. CTL=Control group. NPD=Group diagnosed with Narcissistic Personality Disorder. CI=confidence interval. LC=Low Conflict. MC=Medium Conflict. HC=High Conflict. MCP=Medium Conflict Punishment. HCP=High Conflict Punishment.
[1]p-values for individual conditions (i.e. not "Overall") are Holm-corrected.

 Figures

baseline trials			mean	max +/-	mean	max +/-
baseline	BL	participant	500	0	500	0
		recipient	500		500	

non baseline trials			selfish option		prosocial option	
trialtype			mean	max +/-	mean	max +/-
low conflict	LC	participant	600	20	550	20
		recipient	400		450	
medium conflict	MC	participant	800	60	550	20
		recipient	200		450	
high conflict	HC	participant	800	60	200	60
		recipient	200		800	
medium conflict punishment	MCP	participant	1200	60	550	20
		recipient	-200		450	
high conflict punishment	HCP	participant	1200	60	200	60
		recipient	-200		800	

622   Figure 1

624   Figure 2

626   Figure 3

628   Figure 4