Title: Long-term Exposures to Air Pollutants Affect FeNO in Children: A Longitudinal Study

Authors: Yue Zhang, PhD[1,2,3], Sandrah P. Eckel, PhD[4], Kiros Berhane, PhD[5], Erika Garcia, PhD[4], Patrick Muchmore, PhD[4], Noa Ben-Ari Molshatzki, PhD[4], Edward B. Rappaport, MS[4], William S. Linn, MA[4], Rima Habre, ScD[4], Frank D. Gilliland, MD, PhD [4]

Affiliations:
[1] Department of Internal Medicine, University of Utah, Salt Lake City, UT, USA;
[2] Department of Family and Preventive Medicine, University of Utah, Salt Lake City, UT, USA;
[3] Veteran Affairs Salt Lake City Health Care System, Salt Lake City, UT, USA;
[4] Department of Preventive Medicine, University of Southern California, Los Angeles, CA, USA;
[5] Department of Biostatistics, Columbia University, New York, NY, USA;

Corresponding author contact information:
Yue Zhang
Division of Epidemiology
Department of Internal Medicine
University of Utah
295 Chipeta Way
Salt Lake City, UT, 84018
E-mail: zhang.yue@hsc.utah.edu
Tel: 801-213-3735
Fax: 801-581-3623

Funding: National Institute of Health, Environmental Protection Agency, Hastings Foundation.

ABSTRACT

Fractional exhaled nitric oxide (FeNO) is a marker of airway inflammation shown to be responsive to short-term air pollution exposures; however, effects of long-term exposures are uncertain. Using longitudinal assessments of FeNO and air pollutant exposures, we aimed to determine whether FeNO is a marker for chronic effects of air pollution exposures after accounting for short-term exposures effects.

FeNO was assessed up to six times 2004-2012 in 3607 schoolchildren from 12 communities in the Southern California Children's Health Study. Within-community long-term ambient air pollution exposures ($PM_{2.5}$, PM_{10}, NO_2, O_3) were represented by differences between community-specific annual averages and the eight-year average spanning the study period. Linear mixed-effect models estimated within-participant associations of annual average air pollution with current FeNO, controlling for previous FeNO, prior seven-day average pollution, potential confounders, and community-level random intercepts. We considered effect modification by sex, ethnicity, asthma, and allergy at baseline.

We found FeNO was positively associated with annual average air pollution, after accounting for short-term exposures. One standard deviation higher annual $PM_{2.5}$ and NO_2 exposures ($PM_{2.5}$:2.0 $\mu g/m^3$; NO_2:2.7 ppb) were associated, respectively, with 4.6% (95%CI:2.3%-6.8%) and 6.5% (95%CI:4.1%-8.9%) higher FeNO. These associations were larger among females. We found little evidence supporting association with PM_{10} or O_3.

Annual average $PM_{2.5}$ and NO_2 levels were associated with FeNO in schoolchildren, adding new evidence that long-term exposure affects FeNO beyond the well-documented short-term effects. Longitudinal FeNO measurements may be useful as an early marker of chronic respiratory effects of long-term $PM_{2.5}$ and NO_2 exposures in children.

Key messages
- We show strong evidence that long-term exposures to air pollutants affect FeNO, independent of the well-documented associations with short-term exposures to air pollution
- Longitudinal FeNO measurements may be useful as an early marker of chronic respiratory effects of long-term air pollution exposures in children.

Capsule summary
Annual average $PM_{2.5}$ and NO_2 were associated with FeNO in schoolchildren, adding new evidence that long-term exposure affects FeNO beyond the well-documented short-term effects.

Key words:
FeNO, pollution, particulate matter, nitrogen dioxide, children

Abbreviations
ATS: American Thoracic Society
BMI: body mass index
CHS: Southern California Children's Health study
EC: elemental carbon
FeNO: Fractional exhaled nitric oxide
NO: Nitric oxide
NO_2: Nitrogen dioxide
O_3: Ozone
$PM_{2.5}$: particulate matter with an aerodynamic diameter of 2.5 or less
PM_{10}: particulate matter with an aerodynamic diameter of 10 μm or less
SHS: secondhand tobacco smoke

INTRODUCTION

Breathing polluted air has a multitude of impacts on human health, many of which are thought to be driven by inflammatory mechanisms. The primary route of exposure is inhalation, thus the respiratory system is a major target organ for toxic effects as it receives the largest dose. The fractional concentration of exhaled nitric oxide (FeNO) is a marker of airway inflammation which has been shown to be useful for the assessment of acute respiratory effects of short-term exposure (hours to weeks) to ambient air pollutants,[1-3] including particulate matter with an aerodynamic diameter of 2.5 or 10 μm or less ($PM_{2.5}$ and PM_{10}, respectively), black smoke, elemental carbon (EC), nitrogen dioxide (NO_2), and ultrafine particles. In the Southern California Children's Health study (CHS), we have reported positive cross-sectional associations of short-term average PM_{10}, $PM_{2.5}$, and ozone (O_3) exposures in the week(s) before FeNO measurement with FeNO.[4]

Currently, there is limited and inconsistent evidence for effects of longer-term air pollution exposures on FeNO. For example, a number of cross-sectional studies have related FeNO to either location-based proxies for long-term exposures to traffic/industrial activity or long-term (seasonal to annual average) ambient pollution. Some of these studies found significant positive associations[5-7] (two with adjustment for short-term exposures[8, 9]) but others did not.[10-15] An early analysis of CHS data over a short follow-up period is one of the few studies to date relating longitudinally assessed FeNO to long-term ambient air pollution exposures. Using two assessments of FeNO approximately one year apart in ~1200 schoolchildren, we found that increased annual average $PM_{2.5}$ or NO_2 was associated with increased within-participant FeNO, after adjusting for short-term exposures (in the prior week).[16]

We hypothesize that FeNO is affected by both long-term and short-term exposures to ambient air pollution. In this study, the estimand of interest is long-term ambient air pollution exposure effect, and short-term exposure is treated as nuisance parameter. We examined the full longitudinal CHS data—which includes up to six repeated FeNO assessments in more than 3000 schoolchildren—to investigate whether long-term (i.e., annual average) ambient air pollution exposures are associated with within-participant changes in FeNO, after accounting for short-term (i.e., weekly average) exposures.

MATERIAL AND METHODS

Study Population

Participants in this study were part of the most recent southern California Children's Health Study (CHS) cohort, which started enrollment in 2002-2003 from kindergarten and first grade classrooms. Detailed information on study design, participant recruitment, and data collection has been reported previously.[17] This analysis includes 3607 school children aged 6-17 years from 12 communities in Southern California, who had FeNO assessed up to 6 times in approximately annual visits from 2004-2005 through 2011-2012. Ethical approval of the protocol was obtained from the University of Southern California institutional review board (approval # HS-13-00150, continuing review date April 17, 2019). Informed assent was obtained for each child, and informed consent from a parent or guardian. The research protocol was reviewed and approved by the University of Southern California Health Campus Institutional Review Board.

FeNO measurement

FeNO was measured at schools following the American Thoracic Society (ATS) guidelines using
an offline technique at the first two visits (2004-2005 and 2005-2006 school years) and online
technique in the subsequent four visits. For offline measurement, breath was collected in
aluminized Mylar bags using commercial breath sampling kits (Sievers Division, GE Analytical
Instruments, Boulder, Colorado, USA). Each child took 2-3 initial deep breaths through the kit's
NO scrubber followed by a near-vital-capacity breath and immediate controlled exhalation
through the sampling kit. The collected bag samples were measured in central laboratory for
FeNO level at targeted 100 ml/sec flow rate using a standard chemiluminescent analyzer system
(Sievers Division). For online measurement, FeNO was measured directly using EcoMedics
CLD-88-SP analyzers with DeNOx accessories to provide NO-free inhaled air (EcoPhysics Inc,
Ann Arbor, MI, USA/Duernten, Switzerland). Each child was asked to provide at least three
maneuvers at the target 50 ml/sec flow rate. The plateau concentration for each maneuver was
taken from the 3 second plateau with the minimum NO coefficient of variation. FeNO was
represented by the mean of two or three ATS acceptable plateaus that differed by less than 15%
or 1 *ppb* where all ATS acceptable plateaus were less than 10 *ppb*. Offline FeNO data were
converted to estimates of FeNO as would be measured online at 50 ml/sec expiratory flow. The
conversion was performed using a statistical model determined in the post hoc comparison study,
which involved 362 children measured on and off line in the same testing session. We exclude
observations with reported use of inhaled corticosteroids. Details of FeNO data collection,
quality control, and conversion procedures have been reported previously.[18-20]
149
Air Pollution Data
Daily 24-hour average $PM_{2.5}$, PM_{10}, NO_2, O_3 (10am-6pm only), and temperature data were
obtained for the study period from central monitoring sites in each community operated by local
air pollution agencies in conformance with US Environmental Protection Agency (EPA)
requirements. For each pollutant, community-specific study period averages were computed (a
seven-year average, from 2004/05-2011/12). For each pollutant and each participants' FeNO test
date, annual average pollution was calculated based on the 12 months preceding the test date in
that participant's community. Weekly averages were calculated based on the 7 days preceding
the test date. Air pollution data was interpolated to fill in data gaps on days with missing
exposure using modeled predictions based on data from nearby monitors. While such processes
could be potential sources of bias, previous sensitivity analyses on CHS data that limited use to
complete data have not altered main study findings.[4] The primary "long-term" exposure of
interest for this study was the annual within-community fluctuation, defined as the difference
between annual averages and the seven year average concentration for each community. We
refer to seven day average as "short-term" exposures hereafter.
165
Other Covariates
Race/ethnicity, medication use, current wheeze status, exposure to secondhand tobacco smoke
(SHS), asthma history, and allergy history were determined from annual self-administered
questionnaires. Questionnaires were completed by parents from baseline to year 2006-2007, and
by children from follow-up year 2007-2008 onwards. Questionnaires were collected at the school
visit during which the child's FeNO test was performed. Ambient air nitric oxide was measured
at the time of FeNO collection. Trained technicians measured height and weight at every FeNO
test date following a standardized protocol. Age- and sex-specific body mass index (BMI)

percentiles were calculated based on the Centers for Disease Control and Prevention BMI growth
charts[21].
176
Statistical Analysis
FeNO had a right-skewed distribution, so we used geometric mean and geometric standard
deviation to quantify central tendency and dispersion and performed all analyses with natural
log-transformed FeNO. In descriptive data analyses, we summarized the characteristics of the
study participants, compared the distribution of baseline year log FeNO by these characteristics,
and also summarized the distribution of the community-specific seven-year pollutant averages
and the annual within-community fluctuations. We also calculated Pearson correlation
coefficients of annual within-community fluctuations between pairs of pollutants.
185
We estimated linear mixed effect models to examine the association of log FeNO with annual
within-community fluctuations in air pollution, adjusting for short-term exposures, confounders,
and design variables as well as log FeNO at the previous visit. More details on the model
specification could be found in the Supplementary appendix. Potential confounders included sex,
race/ethnicity, and respiratory allergy at baseline, as well as time-varying measures of asthma
status, medication use, current wheeze, secondhand tobacco smoke exposure, recent respiratory
illness, concurrent room air nitric oxide, month of FeNO test, and average temperature in the
week preceding the FeNO test. To assess confounding, we used the analytic change-in-estimate
($\geq 10\%$) criterion. We also evaluated whether long-term exposure associations were modified by:
baseline asthma, baseline allergy, sex, and ethnicity. In sensitivity analyses, the final models
were refitted by: (a) removing the five communities which contributed to only the first four (of
six) FeNO visits, (b) removing one community at a time, (c) adding a community-level random
slope on long-term exposure to investigate potential heterogeneity by community, (d) subsetting
to only online FeNO collection (the last four of the six visits), or (e) accounting for the serial
correlation of FeNO observations from same participant by using an autoregressive within-
participant level random component instead of controlling for previous visit log FeNO. In all
models, missing data were assumed to be missing at random. All analyses were conducted using
the statistical package R version 3.5.1[22]. Hypothesis tests were performed under a 0.05
significance level and a two-sided alternative.
205
RESULTS
The 3607 children, with mean age 9 at baseline, were evenly distributed by sex (male: 49.7%),
primarily Hispanic (56.3%) or non-Hispanic White (33.2%), 41.5% had allergies, 15.1% had
asthma, and 3% had reported a recent respiratory illness (Table 1). At baseline, the geometric
mean FeNO was 12.1 ppb (geometric SD: 1.9 ppb) and baseline FeNO was statistically
significantly associated with most participant characteristics. Specifically, boys on average had
significantly higher FeNO than girls (p-value=0.015). FeNO differed by race/ethnicity (p-value
<0.001), with African American children having higher FeNO than the other race/ethnicity
groups. Children with respiratory conditions, such as asthma, wheeze, allergy, and recent
respiratory illness, had statistically significantly higher FeNO than those without (p-values all
<0.001). FeNO was positively associated with age and height (p-value <0.001), but not BMI (p-
value=0.831). FeNO varied by month of collection (p-value<0.001), with the lowest geometric
means in cooler months (December – March).
219

Table 1: Summary of Participant Characteristics and FeNO at Baseline (N=3607)

Variable	Category	N (%)	Geometric mean FeNO (GSD)[*]	p-value[†]
Sex	Male	1792 (49.7)	12.4 (2.0)	0.015
	Female	1815 (50.3)	11.9 (1.9)	
Race/ethnicity	Non-Hispanic White	1193 (33.2)	11.4 (1.8)	<0.001
	Hispanic	2022 (56.3)	12.4 (1.9)	
	African American	122 (3.4)	15.5 (2.2)	
	Asian	173 (4.8)	11.2 (1.9)	
	Other	82 (2.3)	14.2 (2.1)	
ETS	Yes	174 (5.2)	13.1 (2.0)	0.181
	No	3207 (94.8)	12.0 (1.9)	
Allergy	Yes	1292 (41.5)	13.6 (2.0)	<0.001
	No	1820 (58.5)	11.3 (1.8)	
Age	<=9	2443 (67.7)	11.2 (1.8)	<0.001
	>9	1164 (32.3)	14.3 (2.0)	
Height	<=132cm	1947 (54.3)	11.4 (1.8)	<0.001
	>132cm	1638 (45.7)	13.1 (2.0)	
Asthma	Yes	546 (15.1)	16.3 (2.3)	<0.001
	No	3060 (84.9)	11.5 (1.8)	
Medication Use	Yes	299 (9.4)	17.2 (2.3)	<0.001
	No	2896 (90.6)	11.6 (1.8)	
Wheeze	Yes	368 (10.9)	16.6 (2.2)	<0.001
	No	2995 (89.1)	11.6 (1.9)	
BMI	<=75%	1803 (50.3)	12.1 (1.9)	0.831
	>75%	1781 (49.7)	12.1 (1.9)	
Month	Jan	319 (8.8)	10.8 (2.0)	<0.001
	Feb	554 (15.4)	11.0 (1.8)	
	Mar	678 (18.8)	11.7 (1.9)	
	Apr	629 (17.4)	13.6 (1.9)	
	May	323 (9.0)	14.4 (1.9)	
	Jun	367 (10.2)	13.7 (1.9)	
	Oct	117 (3.2)	13.1 (1.8)	
	Nov	250 (6.9)	12.1 (1.8)	
	Dec	370 (10.3)	10.1 (2.0)	
Room air NO	<=5	1831 (50.8)	13.3 (1.9)	<0.001
	>5	1775 (49.2)	11.0 (1.9)	
Respiratory Illness	Yes	82 (3.0)	15.4 (2.1)	<0.001
	No	2631 (97)	11.8 (1.9)	

[*]: Results based on geometric means (GM) and geometric standard deviations (GSD) due to skewness of FeNO data.
[†]: p-values are based on t-tests for comparisons of natural log-transformed FeNO levels across categories (not

Community-specific seven-year average air pollution concentrations ranged approximately two
to three-fold across the 12 communities (Table 2). Most communities experienced a downward
trend in annual average $PM_{2.5}$ and NO_2 over the study period (Figure 1), which was a major
source of variation in the annual within-community fluctuations for these pollutants. The
standard deviation of annual within-community fluctuations was 2.0 $\mu g/m^3$ for $PM_{2.5}$, 6.7
$\mu g/m^3$ for PM_{10}, 2.7 ppb for NO_2, and 2.5 ppb for O_3 (Table 2). As shown in Table 3, $PM_{2.5}$ and
PM_{10} had the highest correlation in seven-year averages across-communities (Pearson's R=0.81),
while $PM_{2.5}$ and NO_2 had the largest correlation in annual within-community fluctuations
(Pearson's R=0.64) and annual within-community fluctuations of O_3 were negatively correlated
with the other pollutants. Log FeNO was correlated with log FeNO at the previous visit
(Pearson's R=0.65), indicating moderately strong first order autocorrelation.

Table 2: Distribution of the 7-year Average Air Pollutant Concentrations and Annual Within-Community Fluctuation across 12 Communities

Pollutant[#]	7-year Average		Annual Within-Community Fluctuation	
	Mean (SD)	Min-Max	Mean (SD)	Min-Max
$PM_{2.5}$, $\mu g/m^3$	12.9 (3.8)	6.8-19.4	0.1 (2.0)	-3.8-4.8
PM_{10}, $\mu g/m^3$	32.0 (8.3)	22.5-52.7	2.1 (6.7)	-10.2-23.3
NO_2, ppb	16.7 (6.3)	6.7-25.6	-0.1 (2.7)	-7.2-5.7
O_3, ppb	45.1 (8.4)	30.7-56.6	-0.1 (2.5)	-6.1-7.7

[#]: All pollutants are 24 hr average, except for ozone (10am–6pm average).

238
239
240

Table 3: Pearson Correlation Coefficients of Annual Within-Community Fluctuation in Air Pollution (lower triangular matrix) and 7-year average air pollutant concentrations (upper triangular matrix).

Pollutant[#]	$PM_{2.5}$	PM_{10}	NO_2	O_3
$PM_{2.5}$	1	0.81	0.75	0.2
PM_{10}	0.55	1	0.29	0.31
NO_2	0.64	0.37	1	-0.02
O_3	-0.41	-0.55	-0.43	1

[*]: P-value < 0.05

[#]: All pollutants are 24 hr average, except for ozone (10am–6pm average).

241
242

Figure 1: Community-specific annual average air pollution during the study period.

Note: 1=Alpine; 2=Lake Elsinore; 3=Lake Gregory; 6=Long Beach; 7=Mira Loma; 8=Riverside; 9=San Dimas;

11=Santa Maria; 12=Upland; 13=Glendora; 14=Anaheim; 15=San Bernardino; 16=Santa Barbara. Study

Year 3=2004-2005; Study Year 4=2005-2006; Study Year 5=2006-2007; Study Year 6=2007-2008; Study

Year 7=2008-2009; Study Year 8=2009-2010; Study Year 9=2010-2011; Study Year 10=2011-2012; The dot

points are the annual average air pollution level prior to the dates of FeNO measurement, and triangle

points are the annual average air pollution level at each study year. The curve lines are the fitted trends
of annual average air pollution level across the study years for each community.
Long-term exposures to $PM_{2.5}$ and NO_2 were positively associated with within-person changes in
FeNO, after adjustment for covariates and short-term exposure (Table 4). A one SD (2.0 μg/m^3)
increase in the annual within-community fluctuation of $PM_{2.5}$ was associated with a 4.6% within-
participant increase in FeNO (95% CI: 2.3%-6.8%), controlling for covariates including FeNO at
the previous visit and seven-day average $PM_{2.5}$. Similarly, a one SD (2.7 ppb) increase in the
annual within-community fluctuation of NO_2 was associated with a 6.5% within-participant
increase in FeNO (95% CI: 4.1%-8.9%), controlling for covariates. We found no evidence for an
association of long-term PM_{10} and FeNO. Long-term O_3 was negatively associated with FeNO in
the primary analysis, but this association was attenuated and no longer statistically significant in
sensitivity analyses (Tables 1S and 2S). All models were adjusted for sex, race, respiratory
allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness,
room air nitric oxide, month, temperature, and short-term exposure of air pollution.
264

Table 4: Adjusted* percent difference in FeNO associated with a one standard deviation
(SD)‡ increase in long-term air pollution exposures†

Pollutant	Percent difference in FeNO (95% CI)
$PM_{2.5}$	4.55 (2.33, 6.82)
PM_{10}	0.63 (-1.88, 3.21)
NO_2	6.46 (4.08, 8.9)
O_3	-2.62 (-4.49, -0.71)

*Models adjusted for: sex, race, respiratory allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness, room air nitric oxide, month, temperature, prior 7 day average of the same pollutant, and natural log-transformed FeNO at the previous visit.
†Long-term exposures are annual within-community fluctuations, calculated as the difference between annual air pollutant concentration and seven-year average concentration.
‡SDs are: 2.0 μg/m^3 for $PM_{2.5}$, 6.7 μg/m^3 for PM_{10}, 2.7 ppb for NO_2 and 2.5 ppb for O_3.

265
266
There was evidence that the associations of long-term $PM_{2.5}$ and NO_2 with FeNO varied by sex,
with larger estimated associations in females (interaction p-values: 0.007 for $PM_{2.5}$ and 0.066 for
NO_2) (Table 5). In sensitivity analyses (Supplementary Table 1), we still observed statistically
significant positive associations of long-term $PM_{2.5}$ and NO_2 with FeNO after: (a) removing the
five communities followed only through four years, (b) removing one community at a time
(Supplementary Table 2), and (c) taking out the adjustment for short-term air pollution, though
we did observe some heterogeneity in the associations.
274

Table 5: Interaction Tests[*] on the Effects of Annual Within-Community Fluctuation in Air Pollutions[#] on FeNO level

Effect Modifier		$PM_{2.5}$ % diff/Unit (95% CI)	PM_{10} % diff/Unit (95% CI)	NO_2 % diff/Unit (95% CI)	O_3 % diff/Unit (95% CI)
Sex	Female	3.1 (1.8,4.3)	0.2 (-0.2,0.6)	2.7 (1.8,3.6)	-1.0 (-1.9,-0.2)
	Male	1.4 (0.1,2.6)	-0.1 (-0.5,0.4)	1.8 (0.8,2.9)	-1.1 (-2.0,-0.1)
	Interaction P-value	0.007	0.143	0.066	0.957
Asthma	No	2.4 (1.2 3.5)	0.1 (-0.3,0.5)	2.4 (1.5,3.2)	-1.0 (-1.8,-0.2)
	Yes	1.1 (-1.1,3.4)	-0.2 (-0.9,0.5)	2.3 (0.7,3.9)	-1.1 (-2.8,0.5)
	Interaction P-value	0.249	0.288	0.919	0.899
Allergy	No	2.6 (1.3,3.9)	0.1 (-0.3,0.6)	2.5 (1.5,3.5)	-1.3 (-2.2,-0.4)
	Yes	2.1 (0.7,3.6)	0.1 (-0.4,0.5)	2.1 (1.1,3.2)	-1.5 (-2.5,-0.5)
	Interaction P-value	0.491	0.753	0.471	0.731
Hispanic	No	2.4 (1.0,3.7)	-0.1 (-0.5,0.4)	2.2 (1.3,3.1)	-1.4 (-2.3,-0.6)
	Yes	2.2 (1.0,3.4)	0.2 (-0.2,0.6)	2.6 (1.5,3.7)	-0.6 (-1.5,0.4)
	Interaction P-value	0.799	0.232	0.420	0.083

[*]: Annual Within-community Fluctuation: The difference between annual air pollutant concentration and 7-year average concentration (within-communities). All models adjusted for sex, race, respiratory allergy, asthma, medication use, wheeze, secondhand tobacco smoking, recent respiratory illness, room air nitric oxide, month, temperature, and air pollution level in the prior week.
[#]: All pollutants are 24 hr average, except for ozone (10am–6pm average).

276
277

DISCUSSION

In a large, prospective cohort study of schoolchildren in 12 Southern California communities, we found statistically significant positive associations of FeNO—assessed longitudinally over an eight-year period—with within-community fluctuations in annual average $PM_{2.5}$ and NO_2 after adjusting for prior week exposure, previous visit FeNO, and confounders. This provides the strong evidence that long-term exposures to air pollutants affect FeNO, independent of the well-documented associations with short-term exposures to air pollution.

Results of this longitudinal study (using all six FeNO visits in the CHS) are consistent with a preliminary longitudinal analysis of CHS data (using only the first two online FeNO assessments),[16] which had found *within-participant* associations of long-term $PM_{2.5}$ and NO_2 (but not PM_{10} or O_3) with FeNO while adjusting for short-term exposure. The preliminary CHS study had targeted within-participant changes in FeNO using a different but complementary statistical approach in which the change in FeNO was modeled as a function of the change in annual average air pollution.[16] When studying FeNO, it is important to consider within-participant changes over time from longitudinally collected data, given the considerable unexplained across-participant heterogeneity in FeNO typically observed in cross-sectional studies (e.g., $R^2 < 0.3$).[23] Indeed, in our data we observed moderately strong autocorrelation in

FeNO indicating relatively stable within-person FeNO. As an additional example, in 158
asthmatic adults, the intraclass correlation of FeNO over one year was 0.73, with low baseline
FeNO tending to stay low and higher baseline FeNO tending to stay high, but with more
variability.[24] To our knowledge, the only other study relating longitudinally assessed FeNO to
long-term air pollution exposures found positive but non-statistically-significant associations of
FeNO (measured twice over six months in 130 children) with seasonal black carbon levels at
subjects' residence (estimated by land-use regression), even after adjusting for short-term
exposures (2 hour, 24 hour, or 7 day average black carbon from central site monitors).[25] This
study's statistical analysis did not fully target within-participant associations since it used linear
mixed models with a participant-level random intercept but did not decompose time-varying
exposures (X_{ij}) into within- and between-person terms (i.e., $X_{ij}-\bar{X}_i.$ and $\bar{X}_i.$).[26]
A number of cross-sectional studies have related FeNO to long-term (seasonal to annual average)
ambient pollution or location-based proxies for long-term exposures to traffic/industrial activity.
By design, these studies can estimate only between-participant associations of exposures with
FeNO and hence are subject to the limitations of this approach. Some cross-sectional studies
provide evidence supporting long-term exposure effects. In 241 healthy, full-term infants,
prenatal average NO_2 from a central monitoring site was associated with increased FeNO
measured during sleep at five weeks of age, even after adjusting for mean NO_2 in the 2 days
prior to the FeNO test or residential distance to major road.[9] Two studies relating FeNO to
proximity to industrial activity found FeNO to be elevated in children living closer to oil
refinery/petrochemical activity[6] and in children attending schools closer to a steel mill,[7] though
neither study adjusted for short-term air pollution exposures. Three studies[5, 8, 27] found a similar
magnitude positive association between a proxy measure for traffic exposure (length of road in a
circular buffer around the home), with the strongest associations for the smallest buffers around
the home. While the length of road traffic exposure proxy likely represents both long- and short-
term exposures, one of these studies adjusted for short-term exposure (central site 24 hour
average ambient pollution) and found the length of road association to be robust to the
adjustment,[8] suggesting a long-term exposure effect. Other cross-sectional studies have found no
evidence for long-term exposure effects, perhaps due to exposure metrics relying on spatial
variation which are not sensitive to traffic sources on smaller, local roads (e.g., land-use
regression predicted seasonal or annual averages,[14, 15, 25] traffic counts,[10, 13] or distance to major
road[11, 13]) or too few locations with high/low exposure to separate the source of interest from
potential confounders.[12]
Many study communities observed declines in annual average $PM_{2.5}$ and NO_2 during the study
period. These declines were a key source of the variation in our long-term exposure metric
(within-community annual fluctuations, calculated as the difference between annual averages
and the seven-year average concentration for each community). Our standard interpretation of
the observed positive associations as "percent *increase* in FeNO per SD *increase* in exposure"
(e.g., 6.5% increase in FeNO (95% CI: 4.1%-8.9%) per 2.7 ppb increase in long-term NO_2),
could alternatively be reformulated as a "percent *decrease* in FeNO per SD *decrease* in
exposure". For example, scaled to an observed declines, in Glendora annual average NO_2
decreased by 11.3 ppb over the study period, and we estimated a 30% *decrease* in FeNO (95%
CI: 18.2% to 42.9%) per 11.3 ppb *decrease* in long-term NO_2). A number of studies have used
natural or designed experiments to demonstrate that FeNO decreased following decreasing
(primarily short-term) exposures. For example, following the adoption of clean fuel/technology
school buses, 10-30% reductions in FeNO were observed in 275 schoolchildren.[28] Following

dramatic drops in air pollution in response to anti-air pollution measures enacted for the Beijing
Olympics, FeNO decreased ~57% in 125 health young adults (and subsequently rose 135% after
the games)[3] and ~27% in 36 children.[29] Within-participant FeNO also declined in a group of 42
retired teachers relocated from a higher air pollution city to a lower pollution city for 9 days[30]
and in 37 untreated allergic children moved from an urban to a rural environment for 7 days,
with the reductions in FeNO restricted to participants with the highest FeNO at baseline in the
urban environment.[31]
Several additional results in our study merit further discussion. We found evidence for larger
effects of long-term $PM_{2.5}$ and NO_2 in females as compared to males (p=0.007 and 0.07,
respectively), but no evidence for effect modification by asthma or allergy. In the preliminary
CHS analysis of long-term air pollution effects on FeNO, we had found no evidence for effect
modification by sex.[16] Sex/gender differences have been observed previously in air pollution
epidemiology analyses, generally with larger effects for boys in early life and girls/women in
later childhood and older age.[32] We also found a statistically significant negative association for
long-term O_3, which was not robust to our various sensitivity analyses. Within-community
annual average O_3 was negatively correlated with $PM_{2.5,}$ PM_{10}, and NO_2 and unlike those
pollutants, there were trends of increasing O_3 during the study period. Finally, the association of
long-term NO_2 with FeNO was larger in magnitude when leaving out one community (Upland)
or including a community-level random slope. Upland had the highest annual average NO_2 at the
beginning of the study period, and one of the largest declines over the study period.
This study has several strengths, including: a large ethnically diverse population-based cohort of
schoolchildren; prospective longitudinal assessment of FeNO (up to 6 measurements over an
eight year period); a Southern California study location with substantial within- and between-
community variation in ambient air pollution exposures during the study period, representative
of the range observed nationally in the United States; as well as a study design and statistical
analyses approach that allowed us to target within-person associations of long-term air pollution
and FeNO (automatically controlling for time-constant confounders) while adjusting for short-
term air pollution exposures.
This study has several limitations. The method for FeNO assessment changed during the study
period, as technology improved (offline assessment used for the first two FeNO visits and online
assessment used for the next four FeNO visits). Although the offline values were converted to
predicted online values based on a published internally-developed prediction model with an $R^2 =$
0.94 [33], there remains a possibility that this change in breath collection technique may have
impacted our effect estimates. In a sensitivity analysis restricting to only online FeNO visits, the
association of long-term NO_2 with FeNO remained similar and statistically significant (though
with wider confidence intervals because of smaller sample size) and the association of long-term
$PM_{2.5}$ with FeNO was still positive, but attenuated and no longer statistically significant. A
limitation of our study design is that FeNO was assessed in only the most recent CHS cohort,
allowing only for investigation of associations with within-community fluctuations in annual
average air pollution. We were unable to investigate associations of within-community
fluctuations in multi-year averages. Previous high impact CHS publications have taken
advantage of multiple CHS cohorts (with staggered recruitment over time from the same
communities) to demonstrate that, in the same communities, declines in long-term (multi-year)
average air pollution have been associated with improved lung function growth,[34] reduced
bronchitic symptoms,[35] and reduced asthma incidence.[36] However, the most recent CHS cohort

was still recruited from multiple cities, with a wide range of annual average ambient air pollution
levels at baseline and differing trends over the course of the study period.
388
389 In conclusion, our findings provide the strongest evidence to date that long-term $PM_{2.5}$ and NO_2
390 exposures affect within-participant FeNO, independent of the effects of short-term exposures,
391 even during a study period with declining pollution levels. Longitudinal FeNO measurements
392 may be useful as an early marker of chronic respiratory effects of long-term air pollution
393 exposures in children.
394

Acknowledgements
This work was supported by the Office of the Director of the National Institutes of Health
(UG3OD023287 and UG3OD023249); the National Institutes of Environmental Health Sciences
(grant # R01ES023262, R01ES027860); the Southern California Environmental Health Sciences
Center (grant # P30ES007048) funded by the National Institute of Environmental Health
Sciences; the Children's Environmental Health Center (grant #s P01ES009581, R826708-01 and
RD831861-01) funded by the National Institute of Environmental Health Sciences and the
Environmental Protection Agency; the Maternal and Developmental Risks from Environmental
and Social Stressors (MADRES) Center (grant #s P50ES026086, 83615801-0) funded by the
National Institute of Environmental Health Sciences, the National Institute for Minority Health
and Health Disparities and the Environmental Protection Agency and the Hastings Foundation.
We are indebted to the school principals, teachers, students and parents in each of the study
communities for their cooperation and especially to the members of the field team for their
efforts.

REFERENCES

1.   La Grutta S, Ferrante G, Malizia V, Cibella F, Viegi G. Environmental effects on fractional exhaled nitric oxide in allergic children. J Allergy (Cairo) 2012; 2012:916926.

2.   Scarpa MC, Kulkarni N, Maestrelli P. The role of non-invasive biomarkers in detecting acute respiratory effects of traffic-related air pollution. Clin Exp Allergy 2014; 44:1100-18.

3.   Huang W, Wang G, Lu SE, Kipen H, Wang Y, Hu M, et al. Inflammatory and oxidative stress responses of healthy young adults to changes in air quality during the Beijing Olympics. Am J Respir Crit Care Med 2012; 186:1150-9.

4.   Berhane K, Zhang Y, Linn WS, Rappaport EB, Bastain TM, Salam MT, et al. The effect of ambient air pollution on exhaled nitric oxide in the Children's Health Study. Eur Respir J 2011; 37:1029-36.

5.   Eckel SP, Berhane K, Salam MT, Rappaport EB, Linn WS, Bastain TM, et al. Residential traffic-related pollution exposures and exhaled nitric oxide in the children's health study. Environ Health Perspect 2011; 119:1472-7.

6.   Rusconi F, Catelan D, Accetta G, Peluso M, Pistelli R, Barbone F, et al. Asthma symptoms, lung function, and markers of oxidative stress and inflammation in children exposed to oil refinery pollution. J Asthma 2011; 48:84-90.

7.   Acat M, Aydemir Y, Turğut M, Çörtük M, Çakar M, Yaşar Z, et al. Comparison of the Fractional Exhaled Nitric Oxide Levels in Adolescents at Three Schools Located Three Different Distances from a Large Steel Mill. Can Respir J   2017; 2017.

8.   Dales R, Wheeler A, Mahmud M, Frescura AM, Smith-Doiron M, Nethery E, et al. The influence of living near roadways on spirometry and exhaled nitric oxide in elementary schoolchildren. Environ Health Perspect 2008; 116:1423-7.

9.   Latzin P, Röösli M, Huss A, Kuehni CE, Frey U. Air pollution during pregnancy and lung function in newborns: a birth cohort study. Eur Respir J 2009; 33:594-603.

10.   Cakmak S, Mahmud M, Grgicak-Mannion A, Dales RE. The influence of neighborhood traffic density on the respiratory health of elementary schoolchildren. Environ Int 2012; 39:128-32.

11.   Brown MS, Sarnat SE, DeMuth KA, Brown LAS, Whitlock DR, Brown SW, et al. Residential proximity to a major roadway is associated with features of asthma control in children. PLoS One 2012; 7:e37044.

12.   Spencer-Hwang R, Soret S, Ghamsary M, Rizzo N, Baum M, Juma D, et al. Gender Differences in Respiratory Health of School Children Exposed to Rail Yard–Generated Air Pollution: The ENRRICH Study. J Environ Health 2016; 78:8.

13.   Graveland H, Van Roosbroeck SA, Rensen WM, Brunekreef B, Gehring U. Air pollution and exhaled nitric oxide in Dutch schoolchildren. Occup Environ Med 2010.

14.   Liu C, Flexeder C, Fuertes E, Cyrys J, Bauer C-P, Koletzko S, et al. Effects of air pollution on exhaled nitric oxide in children: Results from the GINIplus and LISAplus studies. Int J Hyg Environ Health 2014; 217:483-91.

15.   Yang A, Janssen NA, Brunekreef B, Cassee FR, Hoek G, Gehring U. Children's respiratory health and oxidative potential of PM2. 5: the PIAMA birth cohort study. Occup Environ Med 2016; 73:154-60.

16. Berhane K, Zhang Y, Salam MT, Eckel SP, Linn WS, Rappaport EB, et al. Longitudinal
effects of air pollution on exhaled nitric oxide: the Children's Health Study. Occup
Environ Med 2014; 71:507-13.
17. McConnell R, Berhane K, Yao L, Jerrett M, Lurmann F, Gilliland F, et al. Traffic,
susceptibility, and childhood asthma. Environ Health Perspect 2006; 114:766-72.
18. Linn WS, Rappaport EB, Berhane KT, Bastain TM, Salam MT, Gilliland FD. Extended
exhaled nitric oxide analysis in field surveys of schoolchildren: a pilot test. Pediatr
Pulmonol 2009; 44:1033-42.
19. Linn WS, Rappaport EB, Eckel SP, Berhane KT, Zhang Y, Salam MT, et al. Multiple-
Flow Exhaled Nitric Oxide, Allergy, and Asthma in a Population of Older Children.
Pediatr Pulmonol 2013.
20. Islam T, Urman R, Gauderman WJ, Milam J, Lurmann F, Shankardass K, et al. Parental
stress increases the detrimental effect of traffic exposure on children's lung function. Am
J Respir Crit Care Med 2011; 184:822-7.
21. 2009. CfDCaPCASpftCgc. A SAS porgram for the CDC growth charts. . 2009.
22. Team RDC. R: a language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. 2014.
23. Garcia-Marcos P, Soriano-Pérez M, Perez-Fernández V, Valverde-Molina J. Exhaled
nitric oxide in school children: Searching for the lost variability. Allergol immunopathol
2016; 44:206-13.
24. Silkoff P, Laviolette M, Singh D, FitzGerald J, Kelsen S, Backer V, et al. Longitudinal
stability of asthma characteristics and biomarkers from the Airways Disease Endotyping
for Personalized Therapeutics (ADEPT) study. Respir Res 2016; 17:43.
25. De Prins S, Dons E, Van Poppel M, Panis LI, Van de Mieroop E, Nelen V, et al. Airway
oxidative stress and inflammation markers in exhaled breath from children are linked
with exposure to black carbon. Environ Int 2014; 73:440-6.
26. Enders CK, Tofighi D. Centering predictor variables in cross-sectional multilevel models:
a new look at an old issue. Psychological methods 2007; 12:121.
27. Holguin F, Flores S, Ross Z, Cortez M, Molina M, Molina L, et al. Traffic-related
exposures, airway function, inflammation, and respiratory symptoms in children. Am J
Respir Crit Care Med 2007; 176:1236-42.
28. Adar SD, D'Souza J, Sheppard L, Kaufman JD, Hallstrand TS, Davey ME, et al.
Adopting clean fuels and technologies on school buses. Pollution and health impacts in
children. Am J Respir Crit Care Med 2015; 191:1413-21.
29. Lin W, Huang W, Zhu T, Hu M, Brunekreef B, Zhang Y, et al. Acute respiratory
inflammation in children and black carbon in ambient air before and during the 2008
Beijing Olympics. Environ Health Perspect 2011; 119:1507.
30. Hao Y, Zhang G, Han B, Xu X, Feng N, Li Y, et al. Prospective evaluation of respiratory
health benefits from reduced exposure to airborne particulate matter. Int J Environ Health
Res 2017; 27:126-35.
31. Renzetti G, Silvestre G, D'Amario C, Bottini E, Gloria-Bottini F, Bottini N, et al. Less air
pollution leads to rapid reduction of airway inflammation and improved airway function
in asthmatic children. Pediatrics 2009; 123:1051-8.
32. Clougherty JE. A growing role for gender analysis in air pollution epidemiology.
Environmental health perspectives 2009; 118:167-76.

33. Linn WS, Berhane KT, Rappaport EB, Bastain TM, Avol EL, Gilliland FD. Relationships
of online exhaled, offline exhaled, and ambient nitric oxide in an epidemiologic survey of
schoolchildren. J Expo Sci Environ Epidemiol 2009; 19:674-81.
34. Gauderman WJ, Urman R, Avol E, Berhane K, McConnell R, Rappaport E, et al.
Association of improved air quality with lung development in children. N Engl J Med
2015; 372:905-13.
35. Berhane K, Chang C-C, McConnell R, Gauderman WJ, Avol E, Rapapport E, et al.
Association of changes in air quality with bronchitic symptoms in children in California,
1993-2012. JAMA 2016; 315:1491-501.
36. Garcia E, Berhane KT, Islam T, McConnell R, Urman R, Chen Z, et al. Association of
changes in air quality with incident asthma in children in California, 1993-2014. Jama
2019; 321:1906-15.