Strong and Significant Associations of Single Nucleotide Variants (SNVs) in the Promoter and 3'-Untranslated Region (3'-UTR) of *Vascular Endothelial Growth Factor* (*VEGF*) Gene with Head and Neck Cancers

Sadia Ajaz [1*], Rabbia Muneer[1], Aisha Siddiqa[2], Muhammad Ali Memon[2], Sadaf Firasat[3], Aiysha Abid[3], Shagufta Khaliq[4]

[1.] Dr. Panjwani Center for Molecular Medicine and Drug Research (PCMD), International Center for Chemical and Biological Sciences (ICCBS), University of Karachi, Karachi-75270, Pakistan.

[2.] Atomic Energy Medical Centre (AEMC), Jinnah Postgraduate Medical Centre (JPMC), Karachi, Pakistan.

[3.] Centre for Human Genetics and Molecular Medicine, Sindh Institute of Urology and Transplantation (SIUT), Karachi, Pakistan.

[4.] Department of Human Genetics and Molecular Biology, University of Health Sciences, Lahore, Pakistan.

[*]Corresponding Author:
Sadia Ajaz, *Ph. D.*
E-mail: sadiaajaz.pcmd@iccs.edu

Strong and Significant Associations of Single Nucleotide Variants (SNVs) in the Promoter and 3'-Untranslated Region (3'-UTR) of *Vascular Endothelial Growth Factor* (*VEGF*) Gene with Head and Neck Cancers

Abstract

Vascular Endothelial Growth Factor (VEGF) has a potent role in tumorigenesis and metastasis. *VEGF* gene is highly polymorphic. Specific variants have been associated with risk and disease progression in disorders with microvascular involvement such as diabetes mellitus and cancer. However, no study has reported association analysis of common single nucleotide variants (SNVs) in the promoter and 3' untranslated region (3'-UTR) with head and neck cancers (HNCs). The present study addresses this gap. It investigates the association of two SNVs, -2578 C/A (rs699947) in the promoter region and +936 C/T (rs3025039) in 3'-UTR of the *VEGF* gene, with risk of HNCs and tumour characteristics.

After the ethical approval and informed consent, 323 participants were enrolled in the study. In the case-control component of the study, there were 121 HNC patients and 202 controls. Demographic details and medical history were recorded. Peripheral blood samples (10ml) were collected in ACD-coated vacutainers. DNA isolation was carried out by organic-solvent method. PCR-RFLP methods were optimized for the genotyping of selected SNVs. The protocol was validated by Sanger sequencing. The susceptibility association was determined by using genetic models applying Hardy-Weinberg equilibrium. Statistical analyses included Chi-squared test of independence or Fisher's Exact test with significance at p-value <0.05, and odds ratios with 95% confidence interval.

VEGF -2578 A-allele, A-carrier genotype (dominant model), and AA genotype (co-dominant model) were significantly asso*c*iated with protective effect against HNCs (*p-values* <0.01, <0.042, and < 0.009, respectively). The odds ratios (95% CI) were: 0.651 (0.469 - 0.904), 0.613 (0.381 – 0.985), and 0.393 (0.193 - 0.804), respectively. The association analysis of *VEGF* +936 C/T polymorphism showed that T-allele, CT (over-dominant model), and T-carrier (dominant model) genotypes were significantly associated with the risk for head and neck cancers (*p-values* < 0.047, < 0.037, and < 0.038, respectively). The odds ratios (95% CI) were 1.882 (1.001 – 3.536); 2.060 (1.035 – 4.102); and 2.023 (1.032 – 3.966), respectively.

Additionally, *VEGF* +936 CT and T-carrier genotypes showed significant associations with higher tumour grade (*p-value* <0.029, and <0.037). The odds ratios (95% CI) were 10.00 (1.435 – 72.806) and 9.00 (1.294 – 62.594), respectively.

VEGF -2578 C/A and +936 C/T may be further investigated as susceptibility genetic markers for HNCs. Specifically, -2578 A-variant has a protective effect, whereas +936 T-variant increases the risk. Furthermore, +936 CT and T-carrier genotypes are also associated with higher tumour grade. The present study is the first positive association analysis report of both selected SNVs with HNCs in any population.

INTRODUCTION:

VEGF is a potent mitogen and a critical pro-angiogenic factor [1-3]. Increased VEGF expression is associated with tumourigenesis [4, 5] and metastasis [6]. VEGF concentrations, and hence the function, may vary depending upon the sequence variations in the encoding gene [7].

 The *VEGF* gene, is located on chromosome 6p21.1. It is 16kb in size. It is one of the most polymorphic genes with more than 100 reported genetic variations [8]. *VEGF* gene expression variations have been shown in association with the following SNVs: -1154 G/A (rs1570360), -

2578 C/A (rs699947) and -1498 C/T (rs833061) polymorphisms in the promoter region, -634 G/C (rs2010963) in the 5' untranslated region (UTR) and +936 C/T (rs3025039) in the 3' UTR [7, 9-10]. These SNVs have been associated with certain epithelial cancers in different populations.

Head and neck cancers (HNCs) show high incidence in South-Asia [11]. However, no data is available for the contribution of variations in critical genes like *VEGF* to HNCs in this region.

In order to address the paucity of data, the present study performs the association analysis of two frequent SNV in the *VEGF* gene, rs699947 (or -2578 C/A) and rs3025039 (or +936 C/T) with HNC cases from Pakistan.

MATERIALS AND METHODS:

Ethics Statement

The protocol is approved by the ethical review committees of the participating institutions. These include the independent ethics review committee (IEC) of the International Center for Chemical and Biological Sciences (ICCBS), University of Karachi, Karachi, Pakistan [ICCBS/IEC-016-BS/HT-2016/Protocol/1.0], the Ethics Review Committee (ERC) of the Atomic Energy Medical Centre (AEMC), Jinnah Postgraduate Medical Centre (JPMC), Karachi, Pakistan [Admin-3 (257)/2016], and the ERC, Sindh Institute of Urology and Transplantation (SIUT), Karachi, Pakistan. All the participants signed a written informed-consent form prior to sampling.

Study Population, Demographic, and Tumour Data Collection:

The present study comprised two components: case-control study design to assess the contribution of selected SNVs to HNC susceptibility in the local population, and a cross-sectional study design to evaluate the association of particular SNVs with tumour progression .

The participants belonged to Southern-Pakistan. The sampling for cases was carried out at AEMC, JPMC, Karachi, Pakistan from July 2016 – July 2017, while the samples from controls had been collected earlier at SIUT, Karachi, Pakistan from March 2009 - November 2012.

All cases were biopsy-proven primary HNC patients. Information regarding tumour site, histopathology, tumour size, tumour grade, and TNM classification were recorded from the pathology reports in the patients' medical files. For controls, blood samples from individuals in the age range of 40 – 60 years were selected. In particular, the controls visited SIUT and had no personal history of any cancer at the time of sampling as determined from their medical records.

Sample Collection

The present study comprised 323 participants, with 121 cases and 202 controls. In case of patients 8-10ml, while for controls 4-5ml of blood samples were collected in ACD coated vacutainers and either processed immediately or stored at 4°C until DNA extraction.

DNA Extraction and Spectrophotometric Analysis

DNA was extracted using standard phenol-chloroform method [12]. Quality of the extracted DNA and concentration were determined spectrophotometrically. The samples had OD_{260}/OD_{280} ratio between 1.7 – 1.9. No DNA fragmentation was observed in agarose gel run for quality control purpose

Genotyping

Genotyping for *VEGF* -2578C/A (rs699947) and +936C/T (rs3025039) were carried out by Polymerase Chain Reaction (PCR) followed by Restriction Fragment Length Polymorphism (RFLP) at Dr. Panjwani Center for Molecular Medicine and Drug Research (PCMD), ICCBS, University of Karachi, Karachi, Pakistan.

Genotyping accuracy was validated by sequencing selected samples for the detected polymorphic variants by Sanger sequencing, commercially.

PCR-RFLP based genotyping assays have been described previously [13, 14].

Genotyping was confirmed by second blind evaluation. Overall evaluation of concordance rates was 100% for both *VEGF* -2578C/A and +936C/T polymorphisms. The call rates of genotyping reproducibility were also 100% for randomly selected samples.

Validation of Genotyping Assays:

The genotyping assays were further validated by Sanger sequencing commercially.

Statistical Analysis

The demographic, clinical, and laboratory data were entered, coded, and processed with SPSS® software (version 19). The selected SNPs were tested for Hardy-Weinberg equilibrium (HWE) [14]. The p-values >0.05 showed that the genotypes were in HWE.

Associations between each SNP, HNC risk, and clinico-pathological features were analyzed by Chi-squared test of independence or Fisher's exact test, where appropriate, with requisite degrees of freedom. Statistical significance was selected at p-values < 0.05. Odds ratios with 95% confidence interval were calculated to quantify the strength of genetic association.

RESULTS:

Demographic Comparison and Clinical Data:

The demographics of 121 cases and 202 controls were analyzed. The mean ages of cases and controls were 50.06 ± 1.29 and 46.36 ± 0.36, respectively

The clinico-pathological features of the patients are presented in table 1.

Table 1. Histopathological features of the HNC tumours of the cases

S.No	Parameter*	N or value %
1	**Gender (n=108)**	
1a	Male	75 (69.4%)

1b	Female	33(30.6%)
2	**Tumor Site (n=100)**	
2a	Buccal cavity/oral cavity	38(38%)
2b	Cheek	19(19%)
2c	Tongue	14(14%)
2d	Larynx	8(8%)
2e	Pharynx	7(7%)
2f	Lip	5(5%)
2g	Unspecified parts of mouth	5(5%)
2h	Nasal cavity	2(2%)
2i	Glottis	1(1%)
2j	Salivary gland	1(1%)
3	**Histological Sub-type (n=102)**	
3a	SCC	92(90.2%)
3ai	MD SCC	45(44.11%)
3aii	WD SCC	21(20.6%)
3aiii	PD SCC	7(6.86%)
3aiv	Well to MD SCC	6(5.88%)
3b	Sarcomas	3(2.95%)
3c	Basal cell carcinoma	2(1.96%)
3d	Adenoid cystic carcinoma	2(1.96%)
3e	Others	2(1.96%)
3f	Acinic cell carcinoma	1(0.98%)
4	**Tumor size (n=80)**	
4b	☐5cm	48(60%)
4a	≥5cm	32(40%)
5	**Tumor stage (n=74)**	
5a	Stage (III-IV)	61(82.44%)
5b	Stage (I-II)	13(17.56%)
6	**Tumor Grade (n=61)**	
6a	Low grade (1-2)	56(91.8%)
6b	High grade(3-4)	5(8.2%)

SCC=Squamous cell carcinoma, MD SCC= Moderately differentiated Squamous cell carcinoma, WD SCC= Well differentiated Squamous cell carcinoma, PD SCC= Poorly differentiated Squamous cell carcinoma

*missing data

Significant Protective Effect of *VEGF* (rs699947) -2578 A-Allele, A-carrier Genotype (Dominant Model), and AA-genotype (Co-dominant Model) Against HNCs:

The representative gels for amplification and restriction digestion are shown in Figures 1 and 2, respectively. Validation by Sanger sequencing is shown in the inset of figure 2.

Figure 1: Amplification of *VEGF* -2578C/A on 2.5% agarose gel with 100 bp ladder as a DNA marker. Ethidium bromide was used for staining.

Figure 2: Genotyping of *VEGF* -2578C/A digested with *BglII* enzyme. 325 bp DNA band showed CC genotype, combination of 123 bp, 202 bp, and 325 bp DNA band showed CA genotype, while 202 bp and 123 bp DNA band represented AA genotype.100 bp ladder was used as a marker.

Figure 2 (inset): Electropherogram of Sanger sequencing for *VEGF* -2578C/A polymorphism (arrows represent the site of polymorphism). A-Homozygous variant (A/A), B-Heterozygous variant (C/A), C-Homozygous wild type (C/C).

The genotypes for *VEGF* -2578C/A polymorphism (CC, CA, and AA) were in Hardy-Weinberg equilibrium for both cases and controls (HNC patients: χ^2=0.13, 2 df, p >0.05; Controls: χ^2=0.6, 2 df, p >0.05).

The genotypic and allelic frequencies of *VEGF* -2578C/A polymorphism are given in the table 2. As shown in the table, a significant protective effect of A-allele (OR.: 0.65; 95%CI: 0.47 – 0.90), A-carrier and AA genotypes [OR.: 0.61 (95% CI: 0.38 – 0.98; OR: 0.39 (95% CI 0.19 – 0.80)] were observed against HNCs.

Table 2. Distribution of the *VEGF* -2578C/A genotypes and allele frequencies (with standard errors) in controls and cases (HNC patients). Chi-squared test and odds-ratios (95% Confidence Interval) for the risk of HNC.

VEGF -2578C/A Polymorphism	Controls (n = 202)	HNC patients (n = 121)	Chi-square (p-value)	Odds-ratios (OR) (95% Confidence Interval)
Genotypes	N (%)	N (%)		
CC	58 (28.7%)	48 (39.7%)	Ref.	Ref.
CA	101 (50%%)	59 (48.8%)	1.88 (0.17)	0.71 (0.43– 1.16)
AA	43 (21.3%)	14 (11.6%)	6.75 (0.009)**	0.393 (0.193 – 0.804)**
A-Carrier	144 (71.3%)	73 (60.3%)	4.12 (0.04)*	0.613 (0.381 – 0.985)*
Allele Frequencies				
p[C]	0.54$\pm$0.04	0.64$\pm$0.04	Ref.	Ref.
q[A]	0.46$\pm$0.04	0.36$\pm$0.04	6.62 (0.01)**	0.651 (0.469 - 0.904)*

*significant; **highly significant

Significant Risk Association of *VEGF* +936 T-Allele, CT (Over-dominant Model), and T-carrier (Dominant Model) Genotypes with HNCs:

The representative gels for amplification and restriction digestion are shown in Figures 3 and 4, respectively. Validation by Sanger sequencing is shown in the inset of figure 4.

Figure 3 Amplification of *VEGF* +936C/T on 2.5% agarose gel with 100 bp ladder as a DNA marker. Ethidium bromide was used for staining.

Figure 4. Genotyping of *VEGF* +936C/T digested with *NlaIII* enzyme. A-DNA band of 208 bp represented CC genotype, while 86 bp, 122 bp, and 208 bp DNA bands showed CT genotype, B-122 bp and 86 bp DNA band depicted TT genotype.100 bp ladder was used as a marker.

Figure 4 (inset): Electropherogram of Sanger sequencing for *VEGF* +936C/T SNP (arrows represent the site of polymorphism). A-Wildtype homozygous (C/C), B-Variant heterozygous (C/T), C-Variant homozygous (T/T).

The chi-squared test of independence showed that *VEGF* +936C/T polymorphism was significantly associated with the risk for HNCs (Table 3). The frequencies of T-allele were 13% and 7% in cases and controls, respectively. The respective frequencies of CT genotypes were 19% and 10.3%, whereas those of T-carrier genotypes were 19.8% and 10.9% among cases and controls, respectively. As shown in table 3, chi-squared test of independence showed a significant increased risk for HNCs in CT and T-carrier genotypes.

Table 3. Distribution of the *VEGF* +936C/T genotypes and allele frequencies (with standard errors) in controls and cases (HNC patients). Chi-square test and odds-ratios (95% Confidence Interval) for the risk of HNC.

VEGF +936C/T Polymorphism	Controls (n = 156)	HNC patients (n = 121)	Chi-square (p-value)/Fisher's Exact Test (p-value)	Odds-ratios (OR) (95% Confidence Interval)
Genotypes	N (%)	N (%)		
CC	139 (89.1%)	97 (80.2%)	Ref.	Ref.
CT	16 (10.3%)	23 (19.0%)	4.35 (0.037)*	2.06 (1.03– 4.10)*
TT	01 (0.6%)	01 (0.8%)	0.06 (1.00) ¶	1.43 (0.09 – 23.19)
T-Carrier	24 (10.9%)	24 (19.8%)	4.32 (0.038)*	2.02 (1.03 – 3.96)*
Allele Frequencies				
p[C]	0.94±0.02	0.87±0.03	Ref.	Ref.
q[T]	0.06±0.02	0.13±0.03	3.96 (0.047)*	1.882 (1.001– 3.536)*

*significant result; ¶Fisher's Exact Test;

Significant Association of *VEGF* +936 CT, and T-carrier Genotypes with Tumour Grade in HNCs:

The association analysis of *VEGF* -2578C/A and +936C/T polymorphism with clinico-pathological features of HNC cases is shown in table 4. Significant associations were observed between CT- and T-carrier genotypes with tumour grade in HNCs. The odds ratios (95% CI) were 10.29 (1.45 – 72.81) and 9 (1.29 – 62.59), respectively.

Table 4. Chi-squared test for the association of -2578 C/A and +936 C/T alleles with tumour characteristics in HNCs.

VEGF POLYMORPHISMS	Tumour Size (<=5 vs >5) χ^2 value	Tumour Stage (T1-T2 vs T3-T4) χ^2 value	Tumour Grade (G1-G2 vs G3-G4) χ^2 value
-2578C (AA vs AC/CC)	0.722	0.892	0.792
-2578C (AA vs CC)	0.615	0.039	0.61
-2578C (AC vs CC)	0.42	0.471	0.459
+936 (CC/CT vs TT)	0.604	0.358	**0.037****
+936 (CC vs TT)	0.09	0.6	0.838
+936 (CC vs CT)	0.652	0.246	**0.029****

**statistically significant

DISCUSSION

Cancer is a complex disorder. It has a number of etiologies. The underlying causes could be environmental (e.g. UV exposure), occupational (e.g. chemicals, radioactive materials and asbestos), life style related factors (e.g. tobacco, alcohol) and biological factors including age or inherited genetic defects, certain mutation, and functional polymorphisms in putative genes. VEGF plays an important role in tumour development and progression among different epithelial cancers [16].

Genetic variations in the regulatory region or the gene itself influence gene expression. Thus, these variations are potential genetic markers in disease susceptibility, progression, and/or treatment response [17]. In the highly polymorphic *VEGF* gene, SNVs in the regulatory regions, i.e. promoter, 5' UTR, and 3' UTR have been linked with variations in the gene expression [16, 18-19]. These polymorphisms have been associated with carcinomas including breast [20], melanoma [21], lung [22], kidney [13], and prostate [23] cancers.

Here, we report significant associations of two SNVs in the *VEGF* gene, -2578C/A in the promoter and +936C/T in 3' (UTR), with HNCs. To the best of our knowledge, this is the first such report from South Asia.

Among the eligible published reports from other parts of the world (i.e. studies which fulfill STROBE criteria) [24], two studies each have reported association of *VEGF* -2578 C/A [25, 26] and +936 C/T with HNCs risk [25, 27].

In case of -2578C/A, the present study reports strong and significant protective effect of -2578 A-allele, A-carrier, and AA genotypes against HNCs. The earlier published reports are from Caucasian populations, one from Germany [25] and the other from Serbia [26]. In contrast to the present study, the German study reports increased risk for oral squamous cell carcinomas (OSCC) with -2578CA genotype, while Serbian study does not report independent association of this SNV with HNCs. It is pertinent to mention here that the power of the present study for *VEGF* -2578C/A is more than 90%.

In 3' UTR, +936C/T was not linked with risk for OSCCs in Taiwanese population [27] However, in the German study +936 T allele was associated with increased risk [25], which is in agreement with the present report. The frequency of T allele in Caucasian populations is much higher (approx. 23%) than the South-Asian populations (approx. 10%) [9]. The frequency is further reduced to 7% in PJL (Punjabis in Lahore)-1000 genomes project, and the present study.

Role of VEGF polymorphisms in modulating the VEGF expression has been reported with varying results [28, 29]. For *VEGF* -2578C/A, the correlation of AA genotype with lower VEGF levels has been shown experimentally [30,31]. In agreement, another study reports that C-allele is associated with higher VEGF levels [32]. In contrast one study reports the co-relation of A allele to higher VEGF mRNA levels [33].

The possible mechanism for the observed association may be explained by binding of GATA-2 and MZF-1 transcription factors to VEGF.

The -2578C/A SNP is positioned in the GATA-2 binding site. The -2578 A allele is likely to reduce the binding specificity. The -2578 C allele may produce a binding site GATA-2, that ultimately permits activation of *VEGF* promoter via promoter binding of transcription factor GATA-2. Therefore, the -2578 C allele may increase VEGF levels and promotes angiogenesis as well as tumor cells proliferation. Whereas, the -2578 A allele can reduce the transcription by decreasing the binding affinity to this region [34].

VEGF promoter region containing -2578C/A polymorphism has a binding site for the Myeloid Zinc Finger-1 (MZF-1). It was reported that *VEGF* promoter region at -2549 position had 18 bp fragment deletion/insertion (D/I) that has an association with -2578 SNP. Individuals with the -2578A allele had 18 bp insertion, while those with the -2578C allele did not have this D/I. This D/I polymorphism of *VEGF* has an association with increased transcriptional activity [35].

In oral cancers, various studies have reported the contribution of *VEGF* with conflicting results. While, several studies found no association between *VEGF* expression and angiogenesis in oral cancers [34]. *VEGF* -2578C/A has been correlated with nasopharyngeal carcinomas. The -2578 CC homozygotes have higher VEGF levels in cells as compared to -2578 AA individuals. It has an association with diseases in which inflammation and angiogenesis play important role. This polymorphism showed the association of C allele with NPC in Tunisian population [34]. In Caucasian population, it has correlation with breast cancer [35]. While, the -2578 A allele has significant association with increased risk of renal cell carcinomas in Pakistani population [13].

In *VEGF* +936C/T polymorphism, T allele has been associated with higher risk for HNCs [36-39]. In *VEGF* +936C/T polymorphism, T allele has been associated with lower VEGF levels [38-

39]. In the present study, we propose two possible mechanisms for the observed data: first, that lower VEGF expression may contribute to the risk, but not to tumour progression in HNCs, which require increased VEGF levels. Secondly, the allele could be in linkage disequilibrium with another locus, which increases the risk for HNCs. Indeed, +1451 C/T (rs3025040) in 3'UTR, located in the miR-199a or miR-199b binding site exhibits strong linkage disequilibrium (r^2 = 1.0) with +936C/T in Chinese population [37].

The C allele at +936 in 3′-UTR of *VEGF* has an activator enhancer-binding protein 4 (AP-4) binding site that is absent in case of +936 T allele [40]. An AP-4 DNA binding site (CAGCTG) is present in the 3′-UTR of *VEGF*. SNP in this region reduces the AP-4 binding to DNA in their DNA-binding domain that ultimately affect *VEGF* expression [41].

Most of the published literature presents the association of these polymorphisms with tumour properties such as invasion, stage or grade. The present study also reports significant association of the polymorphism in 3'UTR with tumour grade.

Furthermore, the discrepancies in association with serum concentration levels may result from differences in the health and disease status of the participants. Thus, well designed studies, clearly describing the characteristics of participants are recommended.

In conclusion, to the best of our knowledge, the present study is the first report of strong and significant association of *VEGF* -2578 C/A and +936 C/T polymorphisms with HNCs in South-Asia. This molecular investigation shall aid in establishing genotypic markers for risk and prognosis of HNCs

ACKNOWLEDGEMENTS:

The authors would like to thank Institution core facilities (PCMD/ICCBS), AEMC, JPMC staff

and all the participants of the study for their co-operation.

CONFLICT OF INTEREST:

None to declare.

FUNDING DISCLOSURE:

None to declare.

REFERENCES:

1. Arcondeguy T, Lacazette E, Millevoi S, Prats H, Touriol C. VEGF-A mRNA processing, stability and translation: a paradigm for intricate regulation of gene expression at the post-transcriptional level. Nucleic Acids Res. 2013 Jul 12;41(17):7997-8010.
2. Genetic Home Reference. Retrieved [02.05.2018] from https://ghr.nlm.nih.gov/gene/VEGFA#location
3. Koutras A, Kotoula V, Fountzilas G. Prognostic and predictive role of vascular endothelial growth factor polymorphisms in breast cancer. Pharmacogenomics. 2015 Jan;16(1):79-94.
4. Ghias K. Role of VEGF in head and neck squamous cell carcinomas. J Pioneer Med Sci. 2014 Jul;4(4):163-4.
5. Zhao SF, Yang XD, Lu MX, Sun GW, Wang YX, Zhang YK, Pu YM, Tang EY. Prognostic significance of VEGF immunohistochemical expression in oral cancer: a meta-analysis of the literature. Tumor Biol. 2013 Oct 1;34(5):3165-71
6. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. Cell. 2011 Mar 4;144(5):646-74.
7. Kämmerer PW, Al□Nawas B, Kalkan S, Liese J, Fruth K, Frerich B, Brieger J. Angiogenesis□related prognosis in patients with oral squamous cell carcinoma—role of the VEGF+ 936 C/T polymorphism. J Oral Pathol Med. 2015 Jul;44(6):429-36.
8. Ensembl 2018. Retrieved [02.05.2018] from https://www.ensembl.org/Homo_sapiens/Gene/Variation_Gene/Table?db=core;g=ENSG00000112715;r=6:43770184-43786487
9. Formento JL, Etienne-Grimaldi MC, Francoual M, Pagès G, Onesto C, Formento P, Chamorey E, Dassonville O, Poissonnet G, Milano G. Influence of the VEGF-A 936C> T germinal polymorphism on tumoral VEGF expression in head and neck cancer. Pharmacogenomics. 2009 Aug;10(8):1277-83.

10. Makni L, Stayoussef M, Ghazouani E, Mezlini A, Almawi WY. Distinct association of VEGF-A polymorphisms with laryngeal and nasopharyngeal cancer. Meta Gene. 2016 Dec 31;10:90-4.
11. Fitzmaurice, 2018
12. Sambrook, J. & Russell, D.W. Molecular cloning: a laboratory manual. 2001. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).
13. Ajaz S, Khaliq S, Abid A, Hassan AS, Hashmi A, Sultan G, et al. Association of a single-nucleotide polymorphism in the promoter region of the VEGF gene with the risk of renal cell carcinoma. Genet Test Mol Biomarkers. 2011 Sep 1;15(9):653-7.
14. Ajaz, S., Siddiqa, A., Ali, S. M., Firasat, S., Abid, A., Khaliq, S., & Memon, M. A. (2019). Association analysis of the VEGF gene variants with breast cancer susceptibility and tumor characteristics in an indigenous population: A pilot study. Gene Reports, 16, 100447.
15. http://www.oege.org/software/hwe-mr-calc.shtml
16. Lee SH, Jeong D, Han YS, Baek MJ. Pivotal role of vascular endothelial growth factor pathway in tumor angiogenesis. Annals of surgical treatment and research. 2015 Jul 1;89(1):1-8.
17. Dong PP. Association of vascular endothelial growth factor expression and polymorphisms with the risk of gestational diabetes mellitus. Journal of Clinical Laboratory Analysis. 2018 Oct 23:e22686.
18. Paradowska-Gorycka A, Pawlik A, Romanowska-Prochnicka K, Haladyj E, Malinowski D, Stypinska B, Manczak M, Olesinska M. Relationship between VEGF gene polymorphisms and serum VEGF protein levels in patients with rheumatoid arthritis. PloS one. 2016 Aug 11;11(8):e0160769.
19. Koukourakis MI, Papazoglou D, Giatromanolaki A, Bougioukas G, Maltezos E, Sivridis E. VEGF gene sequence variation defines VEGF gene expression status and angiogenic activity in non-small cell lung cancer. Lung cancer. 2004 Dec 1;46(3):293-8.
20. Shevchenko AV, Konenkov VI, GARBUKOV EY, Stakheeva MN. Polymorphism of regulatory regions of growth factor gene of vascular endothelium VEGFA in breast cancer patients with family history. Problems in oncology. 2012 Dec 15;58(6):773-6.
21. Howell WM, Bateman AC, Turner SJ, Collins A, Theaker JM. Influence of vascular endothelial growth factor single nucleotide polymorphisms on tumour development in cutaneous malignant melanoma. Genes and immunity. 2002 Jun;3(4):229
22. Li Y, Liang J, Liu X, Liu H, Yin B, Xiao J, Bi Z. Correlation of polymorphisms of the vascular endothelial growth factor gene and the risk of lung cancer in an ethnic Han group of North China. Experimental and therapeutic medicine. 2012 Apr 1;3(4):673-6.
23. Martinez-Fierro ML, Garza-Veloz I, Rojas-Martinez A, Ortiz-Lopez R, Castruita-de la Rosa C, Ortiz-Castro Y, Lazalde-Ramos BP, Cervantes-Villagrana AR, Castañeda-Lopez ME, Gomez-Guerra L, Delgado-Enciso I. Positive association between vascular endothelial growth factor (VEGF)-2578 C/A variant and prostate cancer. Cancer Biomarkers. 2013 Oct 31;13(4):235-41.
24. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, Strobe Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology

(STROBE) Statement: guidelines for reporting observational studies. International journal of surgery. 2014 Dec 1;12(12):1495-9.

25. Kämmerer PW, Toyoshima T, Eletr S, Kämmerer P, Kuhr K, Al□Nawas B, Brieger J. Single nucleotide polymorphisms of the vascular endothelial growth factor gene associated with incidence of oral squamous cell carcinoma. Journal of oral pathology & medicine. 2010 Nov;39(10):786-92.

26. Cheng, C. Y., Chang, C. S., Liu, C. J., & Kao, S. Y. (2008). Vascular endothelial growth factor 936 C/T polymorphism is associated with vascular invasion in oral squamous cell carcinoma. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology, 106(1), 79-84.

27. Supic G, Jovic N, Zeljic K, Kozomara R, Magic Z. Association of VEGF-A genetic polymorphisms with cancer risk and survival in advanced-stage oral squamous cell carcinoma patients. Oral oncology. 2012 Nov 1;48(11):1171-7.

28. Etienne□Grimaldi MC, Formento P, Degeorges A, Pierga JY, Delva R, Pivot X, et al. Prospective analysis of the impact of VEGF□A gene polymorphisms on the pharmacodynamics of bevacizumab□based therapy in metastatic breast cancer patients. Br J Clin Pharmacol. 2011 Jun;71(6):921-8.

29. Jain L, Vargo CA, Danesi R, Sissung TM, Price DK, Venzon D, et al. The role of vascular endothelial growth factor SNPs as predictive and prognostic markers for major solid tumors. Mol Cancer Ther. 2009 Sep 15:1535-7163.

30. Schneider BP, Radovich M, Miller KD. The role of vascular endothelial growth factor genetic variability in cancer. Clin Cancer Res. 2009 Aug 25:1078-0432.

31. Schneider BP, Wang M, Radovich M, Sledge GW, Badve S, Thor A, et al. Association of vascular endothelial growth factor and vascular endothelial growth factor receptor-2 genetic polymorphisms with outcome in a trial of paclitaxel compared with paclitaxel plus bevacizumab in advanced breast cancer: ECOG 2100. J Clin Oncol. 2008 Oct 1;26(28):4672.

32. Shahbazi M, Fryer AA, Pravica V, Brogan IJ, Ramsay HM, Hutchinson IV, et al. Vascular endothelial growth factor gene polymorphisms are associated with acute renal allograft rejection. J Am Soc Nephrol. 2002 Jan 1;13(1):260-4

33. Zang J, Hu Y, Xu X, Ni J, Yan D, Liu S, et al. Elevated serum levels of vascular endothelial growth factor predict a poor prognosis of platinum-based chemotherapy in non-small cell lung cancer. Onco Targets Ther. 2017;10:409.

34. Nasr, H. B., Chahed, K., Bouaouina, N., and Chouchane, L. (2008). Functional vascular endothelial growth factor -2578C/A polymorphism in relation to nasopharyngeal carcinoma risk and tumor progression. Clinica Chimica Acta, 395(1-2), 124-129.

35. Yang, B., Cross, D. F., Ollerenshaw, M., Millward, B. A., and Demaine, A. G. (2003). Polymorphisms of the vascular endothelial growth factor and susceptibility to diabetic microvascular complications in patients with type 1 diabetes mellitus. *Journal of Diabetes and its Complications, 17*(1), 1-6.

36. Yapijakis, C., Vairaktaris, E., Vassiliou, S., Vylliotis, A., Nkenke, E., Nixon, A., Neukam, F. (2007). The low *VEGF* production allele of the+ 936C/T polymorphism is strongly associated with increased risk for oral cancer. *Journal of Cancer Research and Clinical Oncology, 133*(10), 787-791.

37. Zhao J, Bai Y, Jin L, Weng Y, Wang Y, Wu H, Li X, Huang Y, Wang S. A functional variant in the 3□-UTR of VEGF predicts the 90-day outcome of ischemic stroke in Chinese patients. PloS one. 2017 Feb 24;12(2): e0172709.

38. Doi K, Noiri E, Nakao A, Fujita T, Kobayashi S, Tokunaga K. Functional polymorphisms in the vascular endothelial growth factor gene are associated with development of end-stage renal disease in males. J Am Soc Nephrol. 2006 Mar 1;17(3):823-30.

39. Krippl P, Langsenlehner U, Renner W, Yazdani□Biuki B, Wolf G, Wascher TC, et al. A common 936 C/T gene polymorphism of vascular endothelial growth factor is associated with decreased breast cancer risk. Int J Cancer. 2003 Sep 10;106(4):468-71.

40. Renner, W., Kotschan, S., Hoffmann, C., Obermayer-Pietsch, B., and Pilger, E. (2000). A common 936 C/T mutation in the gene for vascular endothelial growth factor is associated with vascular endothelial growth factor plasma levels. *Journal of Vascular Research, 37*(6), 443-448.

41. Cao, J., Tang, M., Li, W. L., Xie, J., Du, H., Tang, W. B., Li, Y. (2009). Upregulation of activator protein-4 in human colorectal cancer with metastasis. *International Journal of Surgical Pathology, 17*(1), 16-21.