Manuscript title: Characterizing all-cause excess mortality patterns during COVID-19 pandemic

in Mexico

Sushma Dahal, MPH[1*], Juan M. Banda, PhD[2], Ana I. Bento, PhD[3], Kenji Mizumoto,
PhD[1,4,5], Gerardo Chowell, PhD[1]

[1] Department of Population Health Sciences, School of Public Health, Georgia State University,

Atlanta, GA, USA

[2] Department of Computer Science, College of Arts and Sciences, Georgia State University,

Atlanta, GA, USA

[3] Department of Epidemiology and Biostatistics, School of Public Health, Indiana University

Bloomington, Indiana, USA

[4] Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University

Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan

[5] Hakubi Center for Advanced Research, Kyoto University, Yoshidahonmachi, Sakyo-ku, Kyoto,

Japan

* Corresponding author:

Email: sdahal2@student.gsu.edu (SD)

Abstract

Background: Low testing rates and delays in reporting hinder the estimation of the mortality burden associated with the COVID-19 pandemic. During a public health emergency, estimating all cause excess deaths above an expected level of death can provide a more reliable picture of the mortality burden. Here, we aim to estimate the absolute and relative mortality impact of COVID-19 pandemic in Mexico.

Methods: We obtained weekly mortality time series due to all causes for Mexico, and by gender, and geographic region from 2015 to 2020. We also compiled surveillance data on COVID-19 cases and deaths to assess the timing and intensity of the pandemic and assembled weekly series of the proportion of tweets about 'death' from Mexico to assess the correlation between people's media interaction about 'death' and the rise in pandemic deaths. We estimated all-cause excess mortality rates and mortality rate ratio increase over baseline by fitting Serfling regression models and forecasted the total excess deaths for Mexico for the first four weeks of 2021 using the generalized logistic growth model.

Results: We estimated the all-cause excess mortality rate associated with the COVID-19 pandemic in Mexico in 2020 at 26.10 per 10,000 population, which corresponds to 333,538 excess deaths. Males had about 2-fold higher excess mortality rate (33.99) compared to females (18.53). Mexico City reported the highest excess death rate (63.54) and RR (2.09) compared to rest of the country (excess rate=23.25, RR=1.62). While COVID-19 deaths accounted for only 38.64% of total excess deaths in Mexico, our forecast estimate that Mexico has accumulated a total of ~61610 [95% PI: 60003, 63216] excess deaths in the first four weeks of 2021. Proportion

of tweets was significantly correlated with the excess mortality (ρ=0.508 [95% CI: 0.245, 0.701],

p-value=0.0004).

Conclusion: The COVID-19 pandemic has heavily affected Mexico. The lab-confirmed COVID-

19 deaths accounted for only 38.64% of total all cause excess deaths (333,538) in Mexico in

2020. This reflects either the effect of low testing rates in Mexico, or the surge in number of

deaths due to other causes during the pandemic. A model-based forecast indicates that an

average of 61,610 excess deaths have occurred in January 2021.

Keywords: All-cause mortality, COVID-19, Excess mortality, Mexico, Pandemic, 2020

Background

SARS-CoV-2 continues to spread unabated in many parts of the world. In particular, Latin

American countries are being heavily affected by the COVID-19 pandemic with a total of

22,467,574 cases including 709,062 deaths as of March 12, 2021 [1]. Mexico, one of the highly

populated countries in the world with approximately 42% of the people living in poverty [2],

documented the first imported COVID-19 case on 27th February, 2020 and currently ranks 3rd in

the world in terms of numbers of COVID-19 reported deaths, with a total of 192,488 recorded

deaths (7.33% of total deaths globally) as of March 12, 2021 [1]. Delays in the implementation

of social distancing interventions, mixed reactions towards the stay-at-home order

recommendations, and phased reopening of the country have facilitated sustained transmission of

COVID-19 in Mexico [3].

Mexico has one of the lowest per-capita COVID-19 testing rates in the world with about 17 tests

per 1000 people in total [4]. The low testing rates, compounded by reporting delays, hinders the

estimation of the mortality burden associated with the COVID-19 pandemic based on

surveillance data alone. Instead, a more reliable picture of the effect of COVID-19 pandemic on

mortality can be derived by estimating excess deaths above a baseline or expected level of death

[5, 6]. These estimates can provide information about the deaths that are directly or indirectly

attributed to the pandemic [6]. Indeed, some deaths could be misclassified as COVID-19 deaths,

or some could be occurring in the context of overburdened health care systems. Thus, tracking

all-cause mortality in near real time can help assess whether excess deaths are occurring during a

specific period of time and spatial area [6].

Here we report our estimates of the absolute and relative mortality impact of the COVID-19

pandemic in Mexico using cyclical Serfling regression models together with publicly available

weekly all-cause mortality data from 2015 to 2020 by gender and for Mexico City and other

areas of Mexico. Further, we collected and analyzed weekly twitter data from Mexico about

'deaths' during the COVID-19 pandemic in correlation with the excess all-cause death rate and

COVID-19 death rate. We supplemented our analyses with social media data from Twitter,

which has been found useful to interpret epidemiological trends [7]. In prior work, Google

Trends, Wikipedia searches, and Twitter data have been used for predicting COVID-19 deaths

[8]. Taking this data fusion approach even further, other researchers have used Twitter data,

alongside smart thermometer data, up-to-date clinician search logs as well as Apple and Cuebiq

mobility indices to build near-real time early warning systems for COVID-19 [9].

We also generated predictions of excess mortality for the first 4 weeks of January 2021 using

generalized logistic growth model [10].

Methods

Data: We obtained weekly all-cause death counts based on epidemiological weeks for Mexico

which were also stratified by gender and geographic region from January to December 2020 as

well as for the preceding 5 years (2015-2019) in order to establish a baseline mortality level [11].

We accessed publicly available weekly mortality data available from National Institute of

Statistics and Geography (INEGI) for the years from 2015 to 2018, and data available from

National Population Registry (RENAPO) for the years 2019 and 2020 [11]. The last week of

December 2020, also includes first 2 days of January 2021. To gauge the timing and relative

intensity of the pandemic in Mexico, we examined surveillance data characterizing the weekly

number of laboratory-confirmed COVID-19 cases and deaths, which were obtained from the

official website of the Mexican Ministry of Health through the Directorate General of

Epidemiology [12]. Population size estimates used to calculate mortality rates were obtained

from National Population Council (CONAPO) of Mexico [13].

Statistical analysis: To investigate and quantify the mortality pattern associated with the

COVID-19 pandemic in Mexico, we estimated excess all-cause mortality rates per 10,000

population at the national level and for Mexico City, and other areas of Mexico and by gender.

The excess death rate corresponds to the overall mortality rate above a seasonal baseline of the

expected mortality rates in the absence of the COVID-19 pandemic using standard statistical

methods [5, 14-17].

Definition of pandemic periods and excess mortality estimation: We estimated the baseline

mortality level by fitting cyclical Serfling regression models to all-cause deaths in non-COVID-

19 period, after excluding data from March to December 2020. We included a combination of

linear terms with sine and cosine terms describing time trend and seasonal change respectively,

which is described by the following equation [5, 15]:

Weekly death rates$(t) = a + \alpha \times t + \beta*\sin(2 \times \pi/52.17 \times t) + \gamma*\cos(2 \times \pi/52.17 \times t),$ (1)

Where, a is the intercept and t represents the epidemic week.

Once a weekly baseline and 95% CI were established, periods of COVID-19 pandemic were

defined as the weeks in 2020 where the observed all-cause mortality rate exceeded the upper

95% confidence limit of the baseline mortality level. The same pandemic period was used for

estimating the total excess mortality rate for entire Mexico, Mexico City, Mexico excluding

Mexico City, and gender specific excess mortality rates using established methodology [5, 14-

17]. Excess all-cause mortality rate was defined as the difference between the observed and

model adjusted baseline mortality rates for each week constituting the pandemic period.

Negative excess mortality estimates were replaced by zeros in our analyses. Overall pandemic

excess mortality attributed to all cause for total population, each gender group, Mexico City, and

Mexico excluding Mexico City was calculated by summing the excess death rates across the

pandemic weeks in 2020 [14, 16]. We also calculated the rate ratio (RR), the ratio of observed

all-cause mortality rate during pandemic period to the model predicted baseline mortality level in

the absence of COVID-19 for the given group.

Twitter data analysis: We used a clean version of the publicly available dataset of tweets version

42 [18], the clean version of this dataset removes all re-tweets, keeping only directly initiated

posts by users. We filtered all tweets, by removing all other languages via their ISO 639-1

language code, to only keep the tweets in Spanish (es) and those that originated from Mexico via

its country code MX. Additionally, we removed tweets from news agencies and bot accounts.

We used the following terms to subset the tweets per day: "muerto, muerta, fallecio, murio,

deceso, fallecimiento, defunción, óbito, expiración, defuncion, obito, expiracion, perdio la vida,

sin vida". In English, these terms reflect the meanings "dead, deceased, died, death, expiration,

lost life, lifeless". We collected a total of 1,223,096 and 32,423,282 unique tweets reflecting

death and total tweets respectively from March to December 2020. Next, we overlayed the curve

of proportion of tweets on death out of total tweets in a given week over the mortality rate curve

to inspect the relationship between the mortality rate and the proportion of tweets. We also

calculated correlation coefficients between proportion of weekly tweets and the weekly excess

death rate and the weekly COVID-19 death rate.

Short term forecast of excess deaths: We used generalized logistic growth model (GLM) to

predict excess deaths during the first four weeks of 2021. GLM characterizes epidemic growth

by estimating (i) the intrinsic growth rate, r (ii) a dimensionless "deceleration of growth"

parameter, p and (iii) the final epidemic size, k_0. The deceleration parameter modulates the

epidemic growth patterns including the sub-exponential growth ($0 < p < 1$), constant incidence (p

$=0$) and exponential growth dynamics ($p = 1$). The GLM model is given by the following

differential equation:

$$\frac{dC(t)}{dt} = rC^p(t)\left(1 - \frac{C(t)}{k_0}\right) \qquad (2)$$

Where, $\frac{dC(t)}{dt}$ describes the incidence (of excess deaths) over time t and the cumulative number of

excess deaths at time t is given by $C(t)$. To forecast the number of excess deaths during the first

4 weeks of 2021, we calibrated the model using weekly excess deaths during the last six weeks

of 2020 (week starting from November 22 to the week starting from December 27, 2020). We

utilized parametric bootstrapping approach with a Poisson error structure [10].

Results

Between March 1, 2020 and January 2, 2021, as of surveillance data updated on February 26,

2021, a total of 1,364,557 laboratory-confirmed COVID-19 cases, and 128,886 COVID-19-

related deaths were captured by the epidemiological surveillance system in Mexico. The national

daily series of new cases and deaths due to COVID-19 are shown in Figure 1. The number of

cases rapidly rises from April to July followed by a downward trend which again takes off from

mid-September. A similar temporal pattern can also be gleaned from the time series of COVID-

19 related deaths.

Figure 1. Daily series of new laboratory-confirmed COVID-19 cases and deaths in Mexico, from

March 1, 2020- January 2, 2021

Out of total 44 weeks from March 1, 2020 to January 2, 2021, 38 weeks starting from week 16

(April 12-18, 2020) had the excess death rate greater than 0. The excess death rate peaked on

week 29 (July 12-18, 2020) with the excess death rate of 1.01 per 10,000 population, and on

week 53 (December 27, 2020-January 2, 2021) with the excess death rate of 1.06 per 10,000

population. The weekly timeseries of all-cause mortality rate per 10,000 population in Mexico is

shown in (Figure 2). We found that peaks in all-cause death rates aligned with the peaks in

COVID-19 laboratory-confirmed death rates captured by the surveillance system. The curve of

weekly proportion of tweets from Mexico about death is overlaid with the mortality rate curve in

Figure 2.

Figure 2. Mortality rate per 10,000 population, Mexico, 2015-2020. The red line is the weekly

all-cause death rate. COVID-19 death rate curve is shown in yellow. Dotted lines highlight 2020

COVID-19 pandemic period. The Serfling seasonal regression model baseline (*black curve*) and

corresponding upper limit of the 95% confidence interval of the baseline (*green curve*) are also

shown. The weekly frequency of tweets about death is shown by blue curve. Excess all-cause

mortality rate is the difference between the observed and model adjusted baseline mortality rates

for each week where observed total all-cause mortality rate exceeded the upper 95% confidence

limit of the baseline.

Twitter trends show engagement of people in Mexico with the hashtag terms (Figure 2). The

trend of tweets with the hashtag term related to deaths (explained in the methods section) for the

baseline years of 2018 and 2019 shows a sharp increase in the twitter chatter about deaths during

the pandemic compared to the baseline years. There was a relatively weak but statistically

significant correlation of the weekly proportion of tweets with the weekly excess mortality rate

ρ=0.508 [95% CI: 0.245, 0.701], p-value=0.0004, and the weekly COVID-19 mortality rate

ρ=0.526 [95% CI: 0.268, 0.714], p-value=0.0002, in the study period

The weekly timeseries of all-cause mortality rate per 10,000 population in the country of Mexico

by gender are displayed in Figure 3.

Figure 3. Mortality rate per 10,000 by gender, Mexico. Excess all-cause mortality rate is the

difference between the observed and model adjusted baseline mortality rates for each week

where observed total all-cause mortality rate exceeded the upper 95% confidence limit of the

baseline in the country.

Similarly, the weekly timeseries of all-cause mortality rate per 10,000 population for Mexico

City and for the rest of Mexico are shown in Figure 4.

Figure 4. Mortality rate for Mexico City and Mexico excluding Mexico City. Excess all-cause

mortality rate is the difference between the observed and model adjusted baseline mortality rates

for each week where observed total all-cause mortality rate exceeded the upper 95% confidence

limit of the baseline in the country.

In Table 1, we present the estimates of all-cause excess mortality rate per 10,000 population and

the rate ratio estimates for each studied group, including the estimates at the national level. We

estimated an excess death rate at 26.10 per 10,000 population in Mexico from March 1 to

January 2, 2021. This corresponds to 333,538 excess deaths during the pandemic period. In the

same period, a total of 128,886 lab-confirmed COVID-19 deaths corresponds to 38.64% of the

total estimated excess deaths.

Table 1. Estimates and their uncertainty for all-cause excess mortality rates per 10,000

population and RR during COVID-19 pandemic, Mexico, March 1, 2020- January 2, 2021.

	All-cause excess death rate per 10,000 population [95% CI]	Rate ratio* [95% CI]	Total number of all-cause excess deaths [95% CI]	Deaths due to COVID-19 (% of total number of all-cause excess deaths)
Mexico	26.10 [23.33-28.87]	1.67 [1.56-1.80]	333,538 [298,139-368,936]	128,886 (38.64%)
Mexico City	63.54 [58.15-68.92]	2.09 [1.91-2.30]	57,304 [52,443-62,157]	16,127 (28.14%)
Mexico excluding Mexico City	23.25 [20.58-25.93]	1.62 [1.51-1.75]	276,149 [244,436-307,980]	112,759 (40.83%)
Male, Mexico	33.99 [30.91-37.08]	1.76 [1.64-1.89]	212,667 [193,397-232,001]	81,489 (38.32%)
Female, Mexico	18.53 [15.97-21.08]	1.56 [1.45-1.69]	120,861 [104,164-137,494]	47,397 (39.22%)

*calculated as the ratio of total observed death rate to total baseline death rate during the
pandemic period*

The excess mortality rate in Mexico City (63.54) was about 2.7-fold higher than the rest of the

Mexico (23.25) (proportion test, p-value <0.0001). Interestingly, COVID-19 deaths in Mexico

City accounted for only 28.14% % of the total estimated excess deaths in Mexico City, compared

to 40.83% in the rest of Mexico. Excess mortality rate among males nearly doubled the rate

among females (proportion test, p-value <0.0001, and the proportion of COVID-19 deaths out of

total excess deaths was similar, 38.32% among males and 39.22% among females (proportion

test, p-value <0.0001).

Our estimates of both the absolute and relative excess mortality rate as measured by the rate ratio

of observed vs baseline mortality rate was highest for Mexico City (2.09) compared to other

areas of Mexico (1.62) and for males (1.76). The rate ratio (RR) at the national level was

estimated at 1.67.

Finding from the average weekly forecast of excess deaths generated from the GLM model

calibrated to weeks starting from November 22 to December 27, 2020 shows that Mexico had a

total of ~61610 excess deaths in first four weeks of 2021 (Figure 5).

Figure 5. Model-based forecast of excess number of deaths for first 4 weeks of 2021, Mexico.

Blue circles are the estimates of excess mortality rate and model fit based on generalized logistic

growth model are shown by the black line. The red dashed lines represent the upper and lower

bound of 95% prediction interval. The vertical dashed black line denotes the end of calibration

period and start of forecasting period.

Discussion

Monitoring the excess mortality rate during the course of a pandemic is one of the key

approaches for evaluating pandemic mortality impact [19]. In this study we characterized the

excess mortality impact during COVID-19 pandemic in Mexico from March 1, 2020 to January

2, 2021. The pandemic was associated with an excess mortality rate of 26.10 per 10,000

population (a total of 333,538 excess deaths). Further, COVID-19 laboratory-confirmed deaths

comprised only 38.64% of total excess deaths during the studied period. Our findings indicate

that the COVID-19 pandemic has exerted a particularly devastating mortality burden on the

Mexican population. We found that the all-cause excess-death rate among males was twice as

high as the excess death rate among women in Mexico. This finding is in line with the previous

studies, indicating that more men die from COVID-19 than women [20-22]. Several factors such

as differences in the prevalence of comorbidities [23] as well as risk behaviors such as smoking

and drinking [24], frequency of hand washing [25-27] and delays in health care seeking [22]

could be contributing to a higher risk of COVID-19 death among males.

We found that both the all-cause excess death rate and the rate ratio were the highest in Mexico

City, compared to rest of the country. A previous study reported that Mexico City was the most

affected area during 2009-10 A/H1N1 influenza pandemic in Mexico [28]. Mexico City is one

of the most crowded cities in the world [29] and has been significantly affected by air pollution

for decades [30]. Prior work has identified high population density [31, 32] and long-term

exposure to ambient air pollution [33, 34] as significant predictors of COVID-19 death. Besides,

the long term exposure to ambient air pollution, and living in overcrowded setting is not random

and might interact with other social determinants of health [34] such as poverty, unemployment,

and lack of healthcare access that increase the risk of death during natural disasters.

The fraction of COVID-19 attributed excess deaths was lower in Mexico (38.64%) compared to

more than 65% in the USA [19, 35], and Germany [36]. From March to May 2020, the number

of all cause excess deaths in the US was only 28% higher than the official record of COVID-19

deaths for that period [19]. Subsequently, from March 15, 2020 to January 30, 2021 an estimated

527,500 excess deaths occurred in the USA of which 83.3% were attributed to COVID-19 [37].

In developed countries like Germany where the COVID-19 pandemic management has been

considered a success story, the estimated excess number of deaths during the first wave of

pandemic was lower than the reported number of COVID-19 deaths (+8071 estimated excess

deaths vs. 8674 reported COVID-19 deaths) [36]. The low proportion of COVID-19 deaths out

of total excess deaths could be driven by low COVID-19 testing rates in the country [4], delays

in reporting COVID-19 deaths [38], diagnostic delays for fatal conditions such as cancer [39],

issues related to the sensitivity of reverse transcriptase-PCR (RTPCR) test [40], disruption of

routine health care owing to the collapse of the health system, or intentional choices of not

visiting health facilities due to fear of contracting the virus, exacerbating the effects on the health

of vulnerable groups [41]. Moreover, the pandemic triggered a mental health crisis that has given

rise to an increase in self-harm and suicide [42, 43]. Our forecast of excess deaths during the first

four weeks of 2021 showed an increasing trend which was in line with the high morbidity trend

of COVID-19 cases surrounding that time period [44].

Our Twitter signal indicate an increasing trend in the Twitter chatter about deaths from mid-

January 2020 that peaked in mid-February 2020. This increase coincided with a series of events

including the declaration of the novel coronavirus outbreak in Wuhan as a public health

emergency of international concern (PHEC) (January 30, 2020), and the WHO-China joint

mission of experts from different countries to inform planning on next steps in the response to

COVID-19 outbreak (16-24 February, 2020) [45]. Following a short period of decline, the

Twitter chatter showed a substantial increase, just as the trend in lab-confirmed COVID-19

deaths started to rise in Mexico. However, it substantially declined during the next few months

probably due to pandemic fatigue. This normalization of the pandemic also reflects people's

beliefs of the government signaling that the pandemic was subsiding during the second half of

2020 [46, 47].

Several factors could explain the large number of all-cause excess deaths in Mexico (333,538).

First, Mexico has a high burden of non-communicable diseases. In 2019, the top 5 leading causes

of deaths were ischemic heart disease, diabetes, chronic kidney disease, cirrhosis, and stroke

[48]. These comorbidities have been found to be associated with severe outcomes including

death due to COVID-19 [49, 50]. Therefore, COVID-19 pandemic in a country like Mexico with

high prevalence of chronic diseases, as well as with a health system struggling with absenteeism

and health worker infections might have led to this alarming number of excess deaths [51, 52]. It

is worth noting that, Mexico has the highest number of health worker deaths due to the COVID-

19 pandemic (~1400 deaths) in the world [53-55].

The COVID-19 pandemic has a higher all cause excess mortality compared to 2009 A/H1N1

pandemic in Mexico [56] as shown in Table 2. Past work has reported estimates of respiratory

excess mortality in Mexico City during the 1918-19 influenza pandemic [57]. Our all-cause

excess mortality rate of 63.54 for Mexico City was less than the estimated respiratory excess

mortality of 72.90 per 10,000 population for the three waves of 1918 influenza pandemic in

Mexico City [57] (table 2).

Table 2. Comparison of excess death rate and rate ratio across different pandemics in Mexico

and Mexico City

Pandemic	Place	Mortality outcome	Time period	Excess death rate per 10000 population [95% CI]	Rate ratio
COVID-19 pandemic	Mexico	All cause	March 1, 2020 to January 2, 2021	26.10 [23.33, 28.87]	2.09
A/H1N1 pandemic [56]	Mexico	All cause	April 2009- April 2010	2.46 [1.95-2.96]	NA
COVID-19 pandemic	Mexico City	All cause	March 1, 2020 to January 2, 2021	63.54 [58.15, 68.92]	2.09 [1.91, 2.30]
1918 influenza pandemic in [57]	Mexico City	Respiratory	April 1918 to March 1920	72.90	NA
			April -May, 1918	6.60	1.2
			October to December, 1918	47.00	7.0
			February to March, 1920	19.30	2.6

Our study has several limitations. As excess death rates will be strongly different among

subgroups (it is quite high among the elderly, and those with underlying diseases), overall

estimate is affected by structure of the population. A detailed data on death certificate with age

and underlying diseases information will provide more accurate estimates. Likewise, we can not

rule out the possibility of negative excess deaths following this elevated period of excess

mortality due to new innovations on vaccination and treatment which might prevent serious

complications and deaths, or due to the reduction of the vulnerable populations such as the

elderly during the initial pandemic years. Similarly, the COVID-19 deaths data that we have used

might be underestimated because of different factors such as very low testing rates in Mexico,

and misclassification of COVID-19 deaths. Further studies are needed to shed light on the extent

of deaths directly attributable to COVID-19 and those that are related to other causes.

Conclusion

Our estimate of all-cause excess mortality rate at 26.10 per 10,000 population during COVID-19

pandemic in Mexico provides a reliable estimate of the mortality impact of COVID-19 in a hard-

hit Latin American country with a low testing rate. As more refined mortality data becomes

available on different sub-groups of the population, further studies on excess mortality could

elucidate the mortality impact of the COVID-19 pandemic in Mexico. Our findings indicate that

Mexico has been disproportionately affected by the COVID-19 pandemic.

Declarations

Ethics approval and consent to participate: Not Applicable

Consent for publication: Not Applicable

Availability of data and material: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interest: The authors declare that they have no competing interest.

Funding: GC is partially supported by NSF grant No.'s 2026797, 2034003, and NIH R01 GM 130900. KM is supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI [grant 20H03940] and Japan Science and Technology Agency (JST) J-RAPID [grant JPMJJR2002].

Author Contributions: Conceptualization, G.C., and S.D.; methodology, G.C., S.D., J.M.B., K.M.; validation, G.C.; formal analysis, S.D.; investigation, S.D.; resources, G.C., S.D., J.M.B.; writing—original draft preparation, S.D.; writing, review and editing, S.D., G.C., J.M.B., A.I.B, K.M. All authors have read and approved the final version of the manuscript.

Acknowledgements: Not Applicable

References

1.   **WHO Coronavirus (COVID-19) dashboard** [https://covid19.who.int/]

2.   **Number of people living in poverty in Mexico between 2008 and 2018 (in millions)**

[https://www.statista.com/statistics/1039479/mexico-people-living-poverty/]

3.   Tariq A, Banda JM, Skums P, Dahal S, Castillo-Garsow C, Espinoza B, Brizuela NG,

Saenz RA, Kirpich A, Luo R: **Transmission dynamics and forecasts of the COVID-19**

pandemic in Mexico, March 20-November 11, 2020. *medRxiv* 2021.

4.   Ortega RP: **Mexico's coronavirus czar faces criticism as COVID-19 surges** In:

Science. December 9 2020. [https://www.sciencemag.org/news/2020/12/mexico-s-

coronavirus-czar-faces-criticism-covid-19-surges]

5.   Serfling RE: **Methods for current statistical analysis of excess pneumonia-influenza**

deaths. *Public health reports* 1963, **78**(6):494.

6.   CDC: **Excess Deaths Associated with COVID-19**

[https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm]

7.   Mavragani A, Gkillas K: **COVID-19 predictability in the United States using Google**

Trends time series. *Scientific reports* 2020, **10**(1):1-12.

8.   O'Leary DE, Storey VC: **A Google–Wikipedia–Twitter Model as a Leading Indicator**

of the Numbers of Coronavirus Deaths. *Intelligent Systems in Accounting, Finance and*

Management 2020, **27**(3):151-158.

9.   Kogan NE, Clemente L, Liautaud P, Kaashoek J, Link NB, Nguyen AT, Lu FS, Huybers

P, Resch B, Havas C: **An early warning approach to monitor COVID-19 activity with**

multiple digital traces in near real time. *Science Advances* 2021, **7**(10):eabd6989.

10.   Yan P, Chowell G: **Quantitative methods for investigating infectious disease**

outbreaks, vol. 70: Springer; 2019.

11.   Government of Mexico: **Excess mortality in Mexico**

[https://coronavirus.gob.mx/exceso-de-mortalidad-en-mexico/]

12.   Government of Mexico: **Historical databases COVID-19: published on 26 February**

2021 [https://www.gob.mx/salud/documentos/datos-abiertos-bases-historicas-direccion-

general-de-epidemiologia?idiom=es]

13.   CONAPO: **Projections of the Population of the Municipalities of Mexico, 2015-2030**

14.   Chowell G, Simonsen L, Flores J, Miller MA, Viboud C: **Death patterns during the**

1918 influenza pandemic in Chile. *Emerging infectious diseases* 2014, **20**(11):1803.

15.   Chowell G, Viboud C, Simonsen L, Miller MA, Acuna-Soto R, Díaz JMO, Martínez-

Martín AF: **The 1918–19 influenza pandemic in Boyaca, Colombia**. *Emerging*

infectious diseases 2012, **18**(1):48.

16.   Dahal S, Jenner M, Dinh L, Mizumoto K, Viboud C, Chowell G: **Excess mortality**

patterns during 1918–1921 influenza pandemic in the state of Arizona, USA. *Annals*

of epidemiology 2018, **28**(5):273-280.

17.   Viboud C, Eisenstein J, Reid AH, Janczewski TA, Morens DM, Taubenberger JK: **Age-**

and sex-specific mortality associated with the 1918–1919 influenza pandemic in

Kentucky. *The Journal of infectious diseases* 2013, **207**(5):721-729.

18.   **A large-scale COVID-19 Twitter chatter dataset for open scientific research -- an**

international collaboration [https://arxiv.org/abs/2004.03688]

19.   Weinberger DM, Chen J, Cohen T, Crawford FW, Mostashari F, Olson D, Pitzer VE,

Reich NG, Russi M, Simonsen L: **Estimation of excess deaths associated with the**

389   **COVID-19 pandemic in the United States, March to May 2020**. *JAMA Internal*

390   *Medicine* 2020, **180**(10):1336-1344.

391   20.   Bhopal SS, Bhopal R: **Sex differential in COVID-19 mortality varies markedly by**

392   **age**. *Lancet (London, England)* 2020.

393   21.   Gebhard C, Regitz-Zagrosek V, Neuhauser HK, Morgan R, Klein SL: **Impact of sex and**

394   **gender on COVID-19 outcomes in Europe**. *Biology of Sex Differences* 2020, **11**(1):1-

395   13.

396   22.   Griffith DM, Sharma G, Holliday CS, Enyia OK, Valliere M, Semlow AR, Stewart EC,

397   Blumenthal RS: **Men and COVID-19: a biopsychosocial approach to understanding**

398   **sex differences in mortality and recommendations for practice and policy**

399   **interventions**. *Preventing chronic disease* 2020, **17**:E63.

400   23.   James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, Abbastabar H, Abd-

401   Allah F, Abdela J, Abdelalim A: **Global, regional, and national incidence, prevalence,**

402   **and years lived with disability for 354 diseases and injuries for 195 countries and**

403   **territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study**

404   **2017**. *The Lancet* 2018, **392**(10159):1789-1858.

405   24.   Cai H: **Sex difference and smoking predisposition in patients with COVID-19**. *The*

406   *Lancet Respiratory Medicine* 2020, **8**(4):e20.

407   25.   Suen LK, So ZY, Yeung SK, Lo KY, Lam SC: **Epidemiological investigation on hand**

408   **hygiene knowledge and behaviour: a cross-sectional study on gender disparity**. *BMC*

409   *Public Health* 2019, **19**(1):401.

26. Rebecca Onion: **The long history of the hand-washing gender gap**. In: *Slate*. February 15, 2020. [https://slate.com/technology/2020/02/women-hand-washing-more-than-men-why-coronavirus.html]

27. Alyson Krueger: **Where Women Are Ahead of Men: Hand Washing** In: *The New York Times*. March 17, 2020. [https://www.nytimes.com/2020/03/17/us/women-men-hand-washing-coronavirus.html]

28. Sarria-Guzmán Y, Bernal J, Biase MD, Muñoz-Arenas LC, González-Jiménez FE, Mosso C, León-Lorenzana AD, Fusaro C: **Using demographic data to understand the distribution of H1N1 and COVID-19 pandemics cases among federal entities and municipalities of Mexico**. *PeerJ* 2021.

29. Diamond M: **The 20 most crowded citites on Earth**. In: *INSIDER*. July 8, 2019. [https://www.insider.com/most-crowded-places-on-earth-2017-10]

30. Romo R: **Mexico City's extreme air pollution, in photos**. In: *CNN*. May 17, 2019. [https://www.cnn.com/2019/05/17/americas/mexico-city-pollution-in-photos-intl]

31. Roy S, Ghosh P: **Factors affecting COVID-19 infected and death rates inform lockdown-related policymaking**. *PloS one* 2020, **15**(10):e0241165.

32. Rosenthal BM: **Density Is New York City's Big 'Enemy' in the Coronavirus Fight** In: *The New York Times.* [https://www.nytimes.com/2020/03/23/nyregion/coronavirus-nyc-crowds-density.html]

33. Wu X, Nethery RC, Sabath M, Braun D, Dominici F: **Air pollution and COVID-19 mortality in the United States: Strengths and limitations of an ecological regression analysis**. *Science advances* 2020, **6**(45):eabd4049.

34.   Benmarhnia T: **Linkages between air pollution and the health burden from COVID-**

19: methodological challenges and opportunities. *American journal of epidemiology*

2020, **189**(11):1238-1243.

35.   Rossen LM, Branum AM, Ahmad FB, Sutton P, Anderson RN: **Excess deaths**

associated with COVID-19, by age and race and ethnicity—United States, January

26–October 3, 2020. *Morbidity and Mortality Weekly Report* 2020, **69**(42):1522.

36.   Stang A, Standl F, Kowall B, Brune B, Böttcher J, Brinkmann M, Dittmer U, Jöckel K-H:

Excess mortality due to COVID-19 in Germany. *Journal of Infection* 2020, **81**(5):797-

801.

37.   Josh Katz, Denise Lu, Sanger-Katz M: **527,000 More U.S. Deaths Than Normal Since**

Covid-19 Struck In.; Feb 24, 2021.

38.   Gutierrez E, Rubli A, Tavares T: **Delays in death reports and their implications for**

tracking the evolution of COVID-19. *Available at SSRN 3645304* 2020.

39.   Maringe C, Spicer J, Morris M, Purushotham A, Nolte E, Sullivan R, Rachet B,

Aggarwal A: **The impact of the COVID-19 pandemic on cancer deaths due to delays**

in diagnosis in England, UK: a national, population-based, modelling study. *The*

lancet oncology 2020, **21**(8):1023-1034.

40.   Tahamtan A, Ardebili A: **Real-time RT-PCR in COVID-19 detection: issues affecting**

the results. *Expert review of molecular diagnostics* 2020, **20**(5):453-454.

41.   Roberton T, Carter ED, Chou VB, Stegmuller AR, Jackson BD, Tam Y, Sawadogo-

Lewis T, Walker N: **Early estimates of the indirect effects of the COVID-19**

pandemic on maternal and child mortality in low-income and middle-income

countries: a modelling study. *The Lancet Global Health* 2020, **8**(7):e901-e908.

42.   Sahoo S, Bharadwaj S, Parveen S, Singh AP, Tandup C, Mehra A, Chakrabarti S, Grover

S: **Self-harm and COVID-19 Pandemic: An emerging concern–A report of 2 cases**

from India. *Asian journal of psychiatry* 2020.

43.   Kawohl W, Nordt C: **COVID-19, unemployment, and suicide**. *The Lancet Psychiatry*

2020, **7**(5):389-390.

44.   **Mexico Coronavirus Map and Case Count**. In: *The New York Times* March 14, 2021.

[https://www.nytimes.com/interactive/2020/world/americas/mexico-coronavirus-

cases.html]

45.   WHO: **Archived: WHO Timeline-COVID-19** [https://www.who.int/news/item/27-04-

2020-who-timeline---covid-19]

46.   Dominguez P: **Coronavirus está controlado y a la baja en todo el país: secretario de**

Salud In: *La Mañanera.* 10/11/2020. [https://www.milenio.com/politica/covid-19-

pandemia-controlada-32-entidades-ssa]

47.   Espino LA: **Opinion: La verdad sobre la pandemia está en disputa en México**. In:

The Washington Post July 8, 2020. [https://www.washingtonpost.com/es/post-

opinion/2020/07/08/la-verdad-sobre-la-pandemia-esta-en-disputa-en-mexico/]

48.   IHME: **Mexico** [http://www.healthdata.org/mexico]

49.   Jordan RE, Adab P, Cheng K: **Covid-19: risk factors for severe disease and death**. In.:

British Medical Journal Publishing Group; 2020.

50.   Kim SJ, Bostwick W: **Social vulnerability and racial inequality in COVID-19 deaths**

in Chicago. *Health education & behavior* 2020, **47**(4).

51.   **The Lancet: Latest global disease estimates reveal perfect storm of rising chronic**

diseases and public health failures fuelling COVID-19 pandemic

[http://www.healthdata.org/news-release/lancet-latest-global-disease-estimates-reveal-

perfect-storm-rising-chronic-diseases-and]

52.   Dyer O: **Covid-19: Mexico acknowledges 50 000 more deaths than official figures**

show. In.: British Medical Journal Publishing Group; 2020.

53.   Kahn C: **Mexico Records The Highest Number Of Health Care Worker Deaths**

From COVID-19. In: *NPR*. 2020. [https://www.npr.org/2020/09/17/914103034/mexico-

records-the-highest-number-of-health-care-worker-deaths-from-covid-19]

54.   **Mexico, Demographic data as of July 1, 2020, economic data for 2019**

[https://www.census.gov/popclock/world/mx]

55.   **Covid: Mexico passes 100,000 coronavirus deaths**. In: *BBC News*. 2020.

[https://www.bbc.com/news/world-latin-america-55011840]

56.   Charu V, Chowell G, Palacio Mejia LS, Echevarría-Zuno S, Borja-Aburto VH, Simonsen

L, Miller MA, Viboud C: **Mortality burden of the A/H1N1 pandemic in Mexico: a**

comparison of deaths and years of life lost to seasonal influenza. *Clinical infectious*

diseases 2011, **53**(10):985-993.

57.   Chowell G, Viboud C, Simonsen L, Miller MA, Acuna-Soto R: **Mortality patterns**

associated with the 1918 influenza pandemic in Mexico: evidence for a spring herald

wave and lack of preexisting immunity in older populations. *The Journal of infectious*

diseases 2010, **202**(4):567-575.