Genome sequencing and analysis of an emergent SARS-CoV-2 variant characterized by multiple spike protein mutations detected from the Central Visayas Region of the Philippines

Francis A. Tablizo[1], Kenneth M. Kim[1], Carlo M. Lapid[1], Marc Jerrone R. Castro[1], Maria Sofia L. Yangzon[1], Benedict A. Maralit[2], Marc Edsel C. Ayes[3], Eva Maria Cutiongco-de la Paz[4,7], Alethea R. De Guzman[5], Jan Michael C. Yap[1], Jo-Hannah S. Llames[2], Sheila Mae M. Araiza[2], Kris P. Punayan[2], Irish Coleen A. Asin[2], Candice Francheska B. Tambaoan[2], Asia Louisa U. Chong[2], Karol Sophia Agape R. Padilla[2], Rianna Patricia S. Cruz[1], El King D. Morado[1], Joshua Gregor A. Dizon[1], Razel Nikka M. Hao[6], Arianne A. Zamora[5], Devon Ray Pacial[5], Juan Antonio R. Magalang[5], Marissa Alejandria[7], Celia Carlos[7], Anna Ong-Lim[7], Edsel Maurice Salvaña[7], John Q. Wong[7], Jaime C. Montoya[7], Maria Rosario Singh-Vergeire[7] and Cynthia P. Saloma[4,7]*

* To whom correspondence should be addressed. Email address: cpsaloma@up.edu.ph

[1] Core Facility for Bioinformatics, Philippine Genome Center, University of the Philippines System

[2] DNA Sequencing Core Facility, Philippine Genome Center, University of the Philippines System

[3] Clinical Genomics Laboratory, Philippine Genome Center, University of the Philippines System

[4] Philippine Genome Center, University of the Philippines System

[5] Epidemiology Bureau, Department of Health, Philippines

[6] Disease Prevention and Control Bureau, Department of Health, Philippines

[7] Inter-Agency Task Force on Emerging Infectious Diseases (IATF) Task Force on COVID-19 Variants, Department of Health, Philippines

ABSTRACT

The emergence of SARS-CoV-2 variants of concern such as the B.1.1.7, B.1.35 and the P.1 have prompted calls for governments worldwide to increase their genomic biosurveillance efforts. Globally, quarantine and outbreak management measures have been implemented to stem the introduction of these variants and to monitor any emerging variants of potential clinical significance domestically. Here, we describe the emergence of a new SARS-CoV-2 lineage, mainly from the Central Visayas region of the Philippines. This emergent variant is characterized by 13 lineage-defining mutations, including the co-occurrence of the E484K, N501Y, and P681H mutations at the spike protein region, as well as three additional radical amino acid replacements towards the C-terminal end of the said protein. A three-amino acid deletion at positions 141 to 143 (LGV141_143del) in the spike protein was likewise seen in a region preceding the 144Y deletion found in the B.1.1.7 variant. A single amino acid replacement, K2Q, at the N-terminus of ORF8 was also shared by all 33 samples sequenced. The mutation profile of this new virus variant warrants closer investigation due to its potential public health implications. The current distribution of this emergent variant in the Philippines and its transmission are being monitored and addressed by relevant public health agencies to stem its spread in nearby islands and regions in the country.

INTRODUCTION

On March 11, 2020, the coronavirus disease 2019 (COVID-19), caused by the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), was declared by the World Health Organization (WHO) as a global pandemic, causing over 115 million infections and over 2.5 million deaths as of March 2, 2021. The global uncontrolled spread of SARS-CoV-2 has given rise to various viral lineages circulating worldwide, with at least three (3) variants being flagged as variants of concern.

Early in December 2020, the United Kingdom reported the emergence of the B.1.1.7 variant

(*20I/501Y.V1 Variant of Concern (VOC) 202012/0)*, defined by 17 amino acid changes, including eight (8)

mutations in the spike protein (Rambaut et al., 2020). In addition to the B.1.1.7, two (2) more variants of

concern were reported: the B.1.351 lineage (South African Variant *20H/501Y.V2*) and the more recent

P.1 lineage (*20J/501Y.V3*) of Brazilian origin. All three lineages of concern share the spike N501Y

mutation, whereas the P.1 and B.1.351 lineages additionally share two (2) other mutations in the spike

protein (K417N/T and E484K) (COVID-19 Genomics UK Consortium [CoG-UK], 2021). Even though the set

of mutations/deletions found in the B.1.1.7, B.1.351, and P.1 lineages appear to have emerged

independently, these viruses have all been associated with a rapid increase in cases in their respective

places of origin (Faria et al., 2021; Rambaut et al., 2020a; Tegally et al., 2020). The WHO has encouraged

countries globally to increase routine systematic sequencing of SARS-CoV-2 viruses to better understand

SARS-CoV-2 transmission and to monitor possible emergence of new variants (World Health

Organization [WHO], 2021).

The Philippines is among the countries adversely hit by the COVID-19 pandemic, with over 580,000

confirmed cases and over 12,000 deaths attributed to the disease as of March 2, 2021. Although border

restrictions are in place to prevent the spread of the virus, local and international travel is still ongoing,

primarily for returning overseas Filipinos (ROFs) from various countries around the world. The

announcement of the UK variant last December 2020 prompted the government to conduct a

nationwide genomic surveillance program to detect SARS-CoV-2 variants, particularly among ROFs in

their ports of entry, as well as from local case clusters in different regions of the archipelago.

In this study, we report the detection of 33 Philippine SARS-CoV-2 cases that were found to share 13

lineage-defining mutations that may have potential biological significance, including the co-occurrence

of the E484K, N501Y, and P681H amino acid replacements at the spike protein region. The mutation

profile observed for this set of viruses has not been described in currently known SARS-CoV-2 lineages of

concern and were detected primarily from the Central Visayas Region of the country during a period

that coincides with a recorded surge in COVID-19 cases in the area.

RESULTS

The Philippine Department of Health, through its surveillance efforts, monitors and analyzes COVID-19

case data since the start of the pandemic. A decline in cases were seen after cases peaked in August of

2020. From January 21, the Department saw an increase in cases particularly in key areas in the country,

including Central Visayas. While an increase in cases was not unexpected given the increased mobility,

crowding, and gatherings during the holiday season, the cases in Central Visayas continued to rise as

other areas started to plateau around February (**Figure 1**). To further enrich the surveillance data

gathered from this area, samples from Central Visayas were prioritized for whole genome sequencing of

the virus. In total, we sequenced 60 samples from this region.

We report here the detection of 33 Philippine SARS-CoV-2 infections caused by viruses initially classified

to be under lineage B.1.1.28, but were found to have a co-occurrence of the spike protein E484K,

N501Y, and P681H mutations. The samples were collected from January 16 to February 02, 2021 and

most were among the sequenced samples from Central Visayas, with a few others from laboratory

submissions from the National Capital Region (NCR) and Davao Region. **Table 1** shows the breakdown of

sample counts per region of laboratory facility.

The three (3) co-occurring mutations have all been identified as significant SNPs in lineages of concern.

The E484K and N501Y co-occur in the P.1 lineage (Brazilian Variant) and the B.1.351 lineage (South

African Variant). In addition, N501Y and P681H are both defining single nucleotide polymorphisms

(SNPs) for the B.1.1.7 lineage (UK Variant) (CoG-UK, 2021). However, a profile of other high-frequency

mutations in the Philippine samples indicate that they do not share any of the other defining SNPs from

any of these lineages (**Figure 2**) (Faria et al., 2021; Rambaut et al., 2020a; Tegally et al., 2020). All the

samples also carry the globally prevalent spike D614G mutation.

Apart from the spike protein mutations previously associated with SARS-CoV-2 lineages of concern, ten

additional amino acid replacements were also observed in the 33 Philippine cases, including the

following: L3201P, D3681E, L3930F and P4715L in the ORF1ab polyprotein; E1092K, H1101Y and V1176F

in the spike protein; K2Q in the ORF8 protein product; and R203K and G204R in the nucleocapsid

protein. Three (3) other mutations were found in 32 of the 33 Philippine samples, which include the

D1554G and A5692V amino acid replacements at the ORF1ab polyprotein, as well as a three-amino acid

deletion at the spike protein at positions 141 to 143 (LGV141_143del). Five (5) synonymous mutations

were also seen: F924F, S2433S and N3928N in ORF1ab; and spike protein G593G and S875S. A list of the

commonly observed mutations for this viral group is shown in **Table 2** and the designated signature

mutations are also indicated. The significance of these mutations, particularly the combination of

mutations in the spike and other regions of the viral genome, on the transmissibility, pathogenicity, and

immunogenicity of this emergent variant remains to be studied.

Among the three known lineages of concern (*i.e.* B.1.1.7, P.1, B.1.351), the B.1.1.7 viruses have been the

most frequently detected so far in the Philippines, majority of which were collected from the main ports

of entry among ROFs and in a cluster of cases in the Cordillera Administrative Region (CAR), north of the

capital Manila. A handful of the B.1.351 viruses have also been detected, but none of the P.1 lineage as

of this writing. Infections from SARS-CoV-2 viruses belonging to the B.1.1.28 lineage appear to be more

widespread, with cases detected in six (6) different regions. However, the majority of the B.1.1.28 cases

in Central Visayas exhibited the signature mutations described for the emergent variant, designated

here as PHL-B.1.1.28 (**Figure 3A**).

Phylogenetic tree reconstruction of the depicted local SARS-CoV-2 cases, together with representative

sequences of the B.1.1.7, B.1.1.28, and P.1 lineages obtained from GISAID, shows that PHL-B.1.1.28

viruses form a distinct monophyletic clade that is separate from other lineages (**Figure 3B**).

DISCUSSION

In this study, we detected 33 Philippine SARS-CoV-2 infections, majority of which were collected from

laboratory facilities in Central Visayas, National Capital Region, and Davao Region (**Table 1**), caused by

viruses initially classified under lineage B.1.1.28, but with the co-occurrence of the E484K, N501Y, and

P681H spike protein mutations. These three (3) mutations have all been identified as defining mutations

in lineages that have attracted global concern with evidence of biological significance.

The E484K mutation, which is a defining SNP of the B.1.351 and P.1 lineages, has been identified as an

immune escape mutation associated with a loss of neutralizing activity by vaccine-elicited antibodies

(Collier et al., 2021). The N501Y mutation, which co-occurs with E484K in the B.1.351 and P.1 lineages

and is also found in B.1.1.7 viruses, is associated with higher transmissibility and increased binding

affinity to the ACE2 receptor (Leung et al., 2021; Luan et al., 2021). The P681H mutation on the other

hand that is found in the B.1.1.7 lineage, is immediately adjacent to the furin cleavage site at the S1/S2

boundary of the spike protein.   The interaction of this site with furin and the ACE2 receptor is

implicated in viral transmission and systemic infection (Hasan et al., 2020; Wang et al, 2020). However,

none of the other defining SNPs for these lineages are found in the Philippine samples, suggesting that

these three (3) mutations may have arisen independently. All the samples also harbor the globally

prevalent D614G mutation, which has been previously associated with increased transmissibility and

higher viral loads (Hou et al., 2020; Lorenzo-Redondo et al., 2020).

The mutation profile (**Figure 2**) for this group of viruses further indicates that they have a characteristic

combination of mutations not observed in other known SARS-CoV-2 lineages of concern. Apart from

E484K, N501Y, D614G, and P681H, we found three (3) other amino acid replacements located towards

the C-terminal end of the spike protein in all of the 33 samples: E1092K, H1101Y, and V1176F (**Table 2**).

Although much is unknown about the actual effects of these mutations in terms of viral transmissibility,

pathogenicity, and its immunogenicity, they all represent radical amino acid replacements with possible

implications in protein structure and activity — E1092K being a change from a negatively charged

residue to a positively charged one; the H1101Y is a change from a positively charged residue to a

hydrophobic one, and the V1176F from a smaller aliphatic residue to a relatively larger aromatic one.

Interestingly, a single amino acid replacement (K2Q) at the N-terminal end of the ORF8 protein product

was also observed in all samples, which presents a possible significant change from a positively charged

residue to a polar uncharged one.

Moreover, we found a deletion of three (3) amino acids at the spike protein residues 141 to 143

(LGV141_143del) in 32 of the 33 samples included in this report. This deletion is adjacent to the tyrosine

residue deletion at the spike protein position 144 (Y144del), which is one of the hallmark mutations of

viruses in the B.1.1.7 lineage. Deletions in this region of the spike protein, including Δ144/145 and Δ141-

144, have been previously shown to alter the protein's antigenicity, with one particular monoclonal

antibody (4A8 MAb) against SARS-CoV-2 unable to detect mutants harboring the said deletions in a Vero

E6 (human kidney epithelial) cell line experiment (McCarthy et al., 2021).

Phylogenetic reconstruction of the 33 Philippine samples with representative samples from the B.1.1.7

and B.1.1.28 lineages, including the P.1 lineage which is derived from B.1.1.28, indicate that they

represent an emergent lineage within the global phylogenetic tree (**Figure 3B**). The phylogeny reveals

that these samples, referred to here as PHL-B.1.1.28, cluster into a well-defined monophyletic clade

(red) that diverges from the B.1.1.28 lineage (yellow). Notably, the tree topology shows that this clade

does not derive from the P.1 lineage, indicating that the E484K and N501Y mutations in samples may

represent recurrent mutations at these sites, rather than inheritance from the latter. The tree also

indicates a recent common ancestor for PHL-B.1.1.28, consistent with the emergence of a novel SARS-

CoV-2 variant.

The geographic distribution of the variant described here further supports its position to be separate

from the B.1.1.28 lineage. While other samples assigned to B.1.1.28 in the Philippines are relatively

abundant in other regions and rare in Central Visayas, the reverse is true for samples bearing the

signature mutations of PHL-B.1.1.28 (**Figure 3A**), suggesting that the emergence of this variant took

place within a geographically separate area of the Philippines centered in Cebu Island. This observation

is particularly notable because the Central Visayas region has been experiencing a sharp spike in COVID-

19 cases since early January 2021 according to statistics compiled by the Republic of the Philippines

Department of Health (**Figure 1**). One possible cause of this spike could be the increased mobility of

people post-holiday season or the emergence of a novel variant with higher transmissibility during this

period, consistent with the collection dates for the described samples in late January to early February.

Although the earliest collection (January 16, 2021) recorded for this group of viruses was from a sample

submitted by a testing laboratory situated at the NCR, we note that hardly any samples were collected

from Central Visayas for sequencing prior to January 31, highlighting the fact that much still remains

hidden about the evolutionary history of this emergent variant.

CONCLUSIONS

Presented in this report are several lines of evidence suggesting the emergence of a novel variant of

SARS-CoV-2 in the Philippines meriting further investigation, represented by 33 samples collected in

areas mainly from Central Visayas. First, all of these samples, collectively referred to here as PHL-

B.1.1.28, have three (3) spike protein mutations that are considered as defining SNPs for other lineages

of concern, and are associated with changes to viral behavior such as increased transmissibility or

antigenic escape. Second, they exhibit a characteristic mutation profile not found in other lineages,

comprising 13 signature non-synonymous mutations including others in the spike protein that are also

implicated in altered protein structure or activity. Third, the samples cluster into a distinct monophyletic

clade within the global tree that diverges from B.1.1.28, similar to the P.1 lineage. And fourth, these

samples were collected mainly at a time and location that was experiencing a sharp spike in COVID-19

cases, raising the possibility of a more transmissible SARS-CoV-2 variant as a potential cause.

METHODOLOGY

Nasopharyngeal swabs from SARS-CoV-2 infected individuals were collected and subjected to

automated RNA extraction using the PANAMAX™48 Automated Nucleic Acid Extraction Unit with

GenAmplify™ BeaDxMax™ Viral DNA/RNA Extraction Kit. Enrichment and sequencing of the extracted

SARS-CoV-2 genetic material were subsequently performed using the Illumina COVIDSeq workflow. The

resulting sequence reads were mapped to the reference SARS-CoV-2 genome sequence (NCBI Accession

No. NC_045512.2) using minimap2 v2.17 (Li, 2018) with preset parameters for accurate genomic short-

read mapping (-x sr). The output mapping file was further processed using Samtools v1.10 (Li et al.,

2009) to fill the mate coordinates and insert sizes field (fixmate), sort according to reference coordinates

(sort), and remove read duplicates (markdup). Mapping and coverage statistics were obtained using

Samtools (flagstat and coverage) as well.

The consensus sequence generation and intrahost variant calling were primarily done using the tool iVar

v1.3 (Grubaugh et al., 2019) with default parameters. Briefly, primer clipping and quality trimming was

implemented using the ivar trim function. Intrahost variants were then detected using a combination of

Samtools mpileup and ivar variants, with the parameters described in the iVar manual page

(https://andersen-lab.github.io/ivar/html/manualpage.html). Following the same manual, the detected

intrahost variants were used to identify and exclude sequence reads associated with mismatched primer

indices using the getmasked and removereads functions of iVar. The consensus sequence assemblies

were then generated using Samtools mpileup and ivar consensus commands with default parameters

(minimum quality = 20, minimum coverage = 10x).

From the consensus sequence assemblies, SARS-CoV-2 lineages were assigned using the tool Pangolin

v2.3.0 (https://github.com/cov-lineages/pangolin; Rambaut et al., 2020b). Variants relative to the

generated consensus sequences were also identified using MUMmer v4.0 (Kurtz et al., 2004) as

implemented in the annotation tool RATT (Otto et al., 2011). Summary mutation tables and plots were

generated using in-house scripts written in Python.

Phylogenetic trees were produced by initially generating a multiple sequence alignment of the local

SARS-CoV-2 consensus sequences using the tool MAFFT v7.407 (Katoh et al., 2002), supplemented with

publicly available sequences from the EpiCoV database of the Global Initiative for Sharing All Influenza

Data (GISAID) platform (Shu and McCauley, 2017). The resulting alignment was automatically trimmed

using trimAl (Capella-Gutierrez et al., 2009). Phylogenetic trees were then reconstructed using a local

instance of the Nextstrain analysis platform (Hadfield et al., 2018).

REFERENCES

Capella-Gutierrez, S., Silla-Martinez, J., & Gabaldon, T. (2009) trimAl: a tool for automated alignment

trimming in large-scale phylogenetic analyses. Bioinformatics. 25(15):1972-1973.

Collier, D., De Marco, A., Ferreira, I., Meng, B., Datir, R., Walls, A. C., ... & CITIID-NIHR BioResource

COVID-19 Collaboration. (2021). SARS-CoV-2 B.1.1.7 sensitivity to mRNA vaccine-elicited,

convalescent and monoclonal antibodies. medRxiv. doi:

https://doi.org/10.1101/2021.01.19.21249840

COVID-19 Genomics UK Consortium (CoG-UK). (2021) COG-UK report on SARS-CoV-2 Spike mutations of

interest in the UK 15[th] January 2021. Available from: https://www.cogconsortium.uk/wp-

content/uploads/2021/01/Report-2_COG-UK_SARS-CoV-2-Mutations.pdf.

Faria, N. R., Claro, I. M., Candido, D., Moyses Franco, L. A., Andrade, P. S, Coletti, T. M., ... on behalf of

CADDE Genomic Network. (2021) Genomic characterisation of an emergent SARS-CoV-2 lineage

in Manaus: preliminary findings. Virological. Available from: https://virological.org/t/genomic-

characterisation-of-an-emergent-sars-cov-2-lineage-in-manaus-preliminary-findings/586.

Grubaugh, N. D., Gangavarapu, K., Quick, J., Mtterson, N. L., de Jesus J. G., Main, B. J., ... & Andersen,

K.G. (2019). An amplicon-based sequencing framework for accurately measuring intrahost virus

diversity using PrimalSeq and iVar. Genome Biology. 20(8). https://doi.org/10.1186/s13059-018-

1618-7.

Hadfield, J., Megill, C., Bell, S. M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P., Bedford, T., &

Neher, R. A. (2018) Nextstrain: real-time tracking of pathogen evolution. Bioinformatics.

34(23):4121-4123.,

Hasan, A., Paray, B. A., Hussain, A., Qadir, F. A., Attar, F., Aziz, F. M., ... & Falahati, M. (2020). A review on

the cleavage priming of the spike protein on coronavirus by angiotensin-converting enzyme-2 and

furin. *Journal of Biomolecular Structure and Dynamics*, 1-9.

Hou, Y. J., Chiba, S., Halfmann, P., Ehre, C., Kuroda, M., Dinnon, K. H., ... & Baric, R. S. (2020). SARS-CoV-2

D614G variant exhibits efficient replication ex vivo and transmission in vivo. *Science*, *370*(6523),

1464-1468.

Katoh, K., Rozewicki, J., & Yamada, K. D. (2002) MAFFT: a novel method for rapid multiple sequence

alignment based on fast Fourier transform. Nucleic Acids Research. 30(14):3059-3066.

Kurtz, S., Philippy, A., Delcher, A. D., Smoot, M., Shumway, M., Antonescu, C., & Salzberg, S. L. (2004)

Versatile and open software for comparing large genomes. Genome Biology. 5: R12.

Leung, K., Shum, M. H., Leung, G. M., Lam, T. T., & Wu, J. T. (2021). Early transmissibility assessment of

the N501Y mutant strains of SARS-CoV-2 in the United Kingdom, October to November 2020.

Eurosurveillance, *26*(1), 2002106.

Li, H. (2018) Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34(18): 3094-3100.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R.

(2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25(16):2078-9.

Lorenzo-Redondo, R., Nam, H. H., Roberts, S. C., Simons, L. M., Jennings, L. J., Qi, C., ... & Ozer, E. A.

(2020). A clade of SARS-CoV-2 viruses associated with lower viral loads in patient upper airways.

EBioMedicine, *62*, 103112.

Luan, B., Wang, H., & Huynh, T. (2021). Molecular Mechanism of the N501Y Mutation for Enhanced

Binding between SARS-CoV-2's Spike Protein and Human ACE2 Receptor. *bioRxiv*, 2021-01. doi:

https://doi.org/10.1101/2021.01.04.425316

McCarthy, K. R., Rennick, L. J., Nambuli, S., Robinson-McCarthy, L. R., Bain, W. G., Haidar, G., & Duprex,

W. P. (2021) Recurrent deletions in the SARS-CoV-2 spike glycoprotein drive antibody escape.

Science. eabf6950. DOI: 10.1126/science.abf6950.

Otto, T. D., Dillon, G. P., Degrave, W. S., & Berriman, M. (2011) RATT: Rapid Annotation Transfer Tool.

Nucleic Acids Research. 39(9): e57.

Rambaut, A., Loman, N., Pybus, O., Barclay, W., Barrett, J., Carabelli, A., Connor, T., Robertson, D. L.,

Volz, E., on behalf of COVID-19 Genomics Consortium UK. (2020a) Preliminary genomic

characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of spike

mutations. Virological. Available from: https://virological.org/t/preliminary-genomic-

characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-

mutations/563.

Rambaut, A., Homes, E.C., O'Toole, A., Hill, V., McCrone, J. T., Ruis, C., du Plessis, L. & Pybus, O. G.

(2020b) A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic

epidemiology. Nature Microbiology. 5: 1403-1407.

Tegally, H., Wilkinson, E., Giovanetti, M., Iranzadeh, A., Fonseca, V., Giandhari, J., ... & de Oliveira, T.

(2020) Emergence and rapid spread of a new severe acute respiratory syndrome-related

coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. medRxiv. doi:

https://doi.org/10.1101/2020.12.21.20248640.

Wang, Q., Qiu, Y., Li, J. Y., Zhou, Z. J., Liao, C. H., & Ge, X. Y. (2020). A unique protease cleavage site

predicted in the spike protein of the novel pneumonia coronavirus (2019-nCoV) potentially related

to viral transmissibility. *Virologica Sinica*, 1-3.

World Health Organization (WHO). (2021) Genomic sequencing of SARS-CoV-2: a guide to

implementation for maximum impact on public health. Available from:

https://www.who.int/publications/i/item/9789240018440.

ACKNOWLEDGEMENT

318 This study was funded by the Republic of the Philippines Department of Health, the Philippine Council

319 for Health Research and Development of the Department of Science and Technology, and the University

320 of the Philippines System. The authors would also like to thank the contributing institutions to the

321 Philippine Genomic Biosurveillance Network for their logistical support.

COMPETING INTERESTS

325 The authors declare no competing interests.

ETHICS APPROVAL

329 The methods used in this study has been approved by the Single Joint Research Ethics Board of the

330 Department of Health, Republic of the Philippines, with approval code SJREB-2021-11, as part of a larger

331 research program entitled "A retrospective study on the national genomic surveillance of COVID-19

332 transmission in the Philippines by SARS-CoV-2 genome sequencing and bioinformatics analysis".

DATA AVAILABILITY

Genome sequences of the 33 Philippine SARS-CoV-2 samples reported here are deposited at the EpiCoV^{TM} database of the Global Initiative for Sharing All Influenza Data (GISAID) with accession codes EPI_ISL_1122426 to EPI_ISL_ 1122458. The acknowledgement table for other GISAID sequences used in this study can be found in **Supplementary Table S1**.

340 **LIST OF TABLES**

341 **Table 1.** Regions in the Philippines where the emergent SARS-CoV-2 variant was detected.

Region of Laboratory Facility	# Cases	Date of Collection	Remarks
Central Visayas	29	January 31 – February 02, 2021	Mostly from the Island of Cebu
National Capital Region	3	January 16 – February 01, 2021	Earliest recorded collection
Davao Region	1	January 23, 2021	

344 **Table 2.** Mutations observed in at least 32 of the 33 Philippine SARS-CoV-2 samples comprising an

345 emergent virus variant. The column "Signature" signifies the mutations found in this group of viruses

346 that are likely to have functional implications or are not prevalent in other lineages.

Affected Gene	Nucleotide Change	Amino Acid Change	# Cases	Signature
Non-synonymous Mutations				
ORF1ab	4962A>G	D1554G	32	Yes
ORF1ab	9867T>C	L3201P	33	Yes
ORF1ab	11308C>A	D3681E	33	Yes
ORF1ab	12053C>T	L3930F	33	Yes
ORF1ab	14408C>T	P4715L	33	
ORF1ab	17339C>T	A5692V	32	Yes
S	21981_21989del	LGV141_143del	32	Yes
S	23012G>A	E484K	33	Yes
S	23063A>T	N501Y	33	Yes
S	23403A>G	D614G	33	
S	23604C>A	P681H	33	Yes
S	24836G>A	E1092K	33	Yes
S	24863C>T	H1101Y	33	Yes
S	25088G>T	V1176F	33	Yes
ORF8	27897A>C	K2Q	33	Yes
N	28881_28882GG>AA	R203K	33	
N	28883G>C	G204R	33	
Synonymous Mutations				
ORF1ab	3037C>T	F924F	33	
ORF1ab	7564C>T	S2433S	33	
ORF1ab	12049C>T	N3928N	33	
S	23341T>C	G593G	33	
S	24187T>A	S875S	33	

348   **LIST OF FIGURES**

351   **Figure 1. Epidemic curve in Central Visayas, Philippines.**   The figure above shows the distribution of

352   reported COVID-19 confirmed cases from the Central Visayas region as to their date of onset of illness as

353   of 02 March 2021. A steep rise in cases have been reported since the January 21, 2021.

Figure 2. Mutation profile of 33 Philippine SARS-CoV-2 samples comprising an emergent viral lineage.

A total of 14 amino acid replacements were observed in all samples (labeled in red), including seven (7) spike protein mutations. Among the spike protein mutations, four (4) have been previously associated with lineages of concern (*i.e.*, E484K, N501Y, D614G, and P681H) while three (3) additional replacements were observed towards the C-terminal region of the protein (*i.e.*, E1092K, H1101Y, and V1176F). Interestingly, a single amino acid replacement at the N-terminus of ORF8 (*i.e.*, K2Q) was also found in all samples. Three (3) other mutations were seen in 32 of the 33 samples (labeled in green), including a three-amino acid deletion at the spike protein positions 141 to 143. Lastly, five (5) synonymous mutations (labeled in gray) were also detected in all of the cases. For comparison, the defining mutations for the P.1, B.1.1.7, and B.1.351 lineages of concern are marked below in red, blue, and green boxes, respectively. None of the samples exhibit mutation profiles similar to any of the three (3) known lineages of concern.

Figure 3. Geographic distribution and phylogenetic clustering of an emergent SARS-CoV-2 lineage in

the Philippines designated as PHL-B.1.1.28. (A) Geographic distribution of viruses under lineages

B.1.1.28 (Yellow), B.1.1.7 (Purple), P.1 (Orange) and the emergent variant PHL-B.1.1.28 (Red) in the

Philippines. Most of the samples detected to have the signature mutations of PHL-B.1.1.28 were from

the Central Visayas region. Note that viruses from lineage P.1 have not been detected in the country to

date. **(B)** Phylogenetic tree of B.1.1.28, B.1.1.7, P.1, and PHL-B.1.1.28 sequences showing that the

emergent variant forms a monophyletic clade separate from other lineages included in the analysis. The

B.1.1.7 and B.1.1.28 clades include samples collected both globally and in the Philippines.

388