Title Page:

Demonstrate, observe, assist perform (DOAP) versus structured educational video (SEV) in imparting standard skill in male urinary bladder catheterisation.

Authors:

Name	Designation & Department	Mobile *	E-mail id
1. Dr Santosh Balakrishnan	Associate Professor of Surgery	9747058458	santosurg@gmail.com
2. Dr Lijo Paul	Senior Resident: Surgery	9965067539	drlijopaul@gmail.com
3. Dr Minu N Rajan	Junior Resident: Surgery	9400634782	minunrajan1994@gmail.com
4. Dr Sherin A Arthungal	Junior Resident: Surgery	8722684015	sherinalias92@gmail.com

*(India country code +91 to be prefixed)

Institution of affiliation of all authors:

Department of Surgery, MOSC Medical College Hospital, Kolenchery. Ernakulam. Kerala. India. 682311

Corresponding Author: Dr Lijo Paul

Funding: Self

Conflict of Interest: None declared

IEC & IRB Approval: IRB Approval Date: 29/11/2019 IEC Protocol Number: MOSC/IEC/398/2019

MESH Key words: Medical education, Competency-Based Education, Simulation

Abbreviations:

SEV (Structured Educational Video); DOAP (Demonstrate, Observe, Assist, Perform); CRRI (Compulsory Residential Rotating Internship); MBC (Male bladder catheterisation); IMG (Indian Medical Graduate)

Abstract:

Introduction: Conventional teaching and learning methods have been seen to fail to assure achievement of competencies in male bladder catherization in the Indian Medical Graduate (IMG) with wide variation noted in competencies. This could be remedied by introduction of Structured training methods.

Aims and Objective: The study aimed to investigate DOAP method of training in a Skill Lab against training through a Structured Educational Video (SEV) with the objective of comparing their efficacy in training Final MBBS Students in the psychomotor skill of performing male bladder catheterisation.

Material & Methods: Final MBBS students fulfilling selection criteria were randomly allocated into two comparable groups. One group underwent Skill lab training using DOAP method while the other group underwent training using a SEV by the same instructor. CRRI interns, regularly performing MBC at work by virtue of conventional training, with 6-8 months experience formed a control group. All participants underwent assessment of skill in MBC by skill lab OSCE evaluation, by assessors who were blinded to the participant's method of training. Data was recorded and analysed using standard statistical software. Trial evaluation from the trial groups was obtained using Survey monkey tool.

Observation: There was no statistically significant difference in the ability of DOAP group or SEV group in being able to safely perform MBC though a higher level of confidence was expressed with their training by DOAP group. Both trial groups statistically outperformed the control group.

Conclusion: Structured training assures competence. Video-assisted Training produces comparable results though DOAP method is preferred by students. A combination of the techniques may facilitate optimal training.

Manuscript

Introduction:

Wide variation is observed in skills in performing common bedside clinical procedure of male bladder catheterisation among final phase MBBS students trained through conventional training and assessment [1,2]. Conventional teaching and learning methods may not allow assurance of achievement of competencies in the Indian Medical Graduate IMG. Introduction of structured training methods could offer a reliable method of facilitation of learning and assured skill acquisition.

Aim:

The team aimed to investigate the suitability of Demonstrate, observe, assist perform (DOAP) method of training in a Skill Lab simulator compared with demonstration using a Structured Educational Video (SEV) in imparting psychomotor skill training in a bedside procedure to medical students.

Objective:

The objective of this trial was to evaluate the efficacy of training by DOAP in comparison to training through a SEV in acquiring skill in performing male bladder catheterisation among Final phase MBBS Students

Material & Methods:

The study was performed as a mandated research project as part of the 1-year Advanced course in Medical Education administered by the National Medical Council through its nodal centre in Kottayam, Kerala. The total duration of the trial was therefore restricted by course regulation to 6 months between conceptualisation & presentation of results as a poster to the course peer group & faculty.

The study was designed as a randomised comparative study conducted in the investigator's institution. A SEV was created mirroring the steps and instructions conveyed by the DOAP training session. This SEV was approved through peer review by four consultant surgeons, one of whom was a consultant Urologist. The DOAP session & OSCE assessment of all groups was conducted in a simulation lab with procedure demonstrated and assessed using a standard male catheterisation trainer model. The group being trained through a SEV were trained by being shown the video twice in the seminar room in the presence of the trainer who led the training session and also reinforced the steps by verbal repetition between two successive viewings of the SEV.

The larger population of the study are Final phase MBBS Students of Ernakulam district in Kerala, India.

Sampling method: All Final phase MBBS students fulfilling selection criteria were randomly assigned a trial number by blinded chit pick-up for anonymity with a securely held key for verification if necessary. They were divided into two equal groups based on random selection and grouping again by blinded chit pick-ups. Compulsory Residential Rotating Internship (CRRI) interns were recruited through their random allocation to the department as per their rotation, to serve as a control group. The interns posted to the department for the duration of the study would have completed between six to eight months of CRRI with regular opportunity at work to perform MBC. They represent a comparable group who have just completed final phase MBBS and have learnt the procedure through the conventional apprenticeship method.

Inclusion Criteria:

1. Medical Students from Final Phase who have not had skill training in male bladder catheterisation who consent to participate
2. CRRI interns posted to the department of surgery subject to their consent.

Exclusion Criteria:

1. Medical students who have had structured skill training through any means in bladder catheterisation.
2. CRRI interns who had any structured training in bladder catheterisation other than traditional apprenticeship.
3. CRRI interns from the senior additional batch.
4. Subjects who have Allergy to latex.

Sample size:

All final phase MBBS Students who fulfil inclusion criteria were recruited. The number of participants was limited by the maximum number of students fulfilling criteria in the institution over the permitted study period.

Intervention:

Institutional review board and Ethical Committee clearance and written informed consent from all participants was taken. The DOAP group underwent structured training using the DOAP method in batches of not more than 6 subjects with demonstration and observation as a group and individual opportunity to assist and perform. The group being trained through a SEV was trained in batches of 15, as limited by the size of the seminar room available for screening the video. They were shown the video two times in the presence of the trainer who verbally reinforced the steps between viewings, to match the visual and auditory exposure through DOAP. A group of 35 CRRI interns posted in our department in numbers to reasonably match the test groups, act as controls trained by conventional apprenticeship method of training to verify the validity and impact of either of the study methods of training

over current practice as recommended by members of the institutional research committee. Participants in all groups were individually assessed and scored using an existing validated OSCE assessment form [3] for their skill in male catheterisation by assessors blinded to the intervention in question. The OSCE form assessed completion of 26 vital steps of the procedure. Eight vital steps were allocated two marks while all others had one mark making up a total score of 34. Following assessments, an online programme satisfaction evaluation questionnaire was administered to each participant. Debrief to subjects was provided only after collection of evaluation questionnaire. In the interest of fairness, a crossover training was provided by sending the video to all in the DOAP group and DOAP training was offered to all in the SEV group after the assessments and online evaluation was complete in order to avoid bias in response.

Data Collection Method and Analysis:

OSCE valuation scores in whole numbers for all groups were compiled in excel sheets. SPSS software was used for data analysis. The institutional bio-statistician aided the analysis in view of the authors' limited skill with the software tool. $P<0.05$ was considered as statistically significant. OSCE scores were summarised using appropriate tests of significance using these tools. Evaluation response received from both groups using Survey Monkey online survey tool, was analysed for difference in levels of agreement on various elements of the training experience with their allocated training method after they completed the OSCE assessment. Data was analysed using MS Excel.

Results:

All trial intervention groups; DOAP group (n=39), SEV group (n=39) and the control group participants (n=35) successfully completed catheterisation in the Simulator model in the skills lab. OSCE scores were awarded by assessors blinded to the interventions. Weight given to each of the 26 elements in the OSCE assessment as well as total scores out of 34 were recorded on proformas and compiled on MS Excel. The scores were found to follow normal distribution (Kolmogorov-Smirnov test) in control, DOAP and SEV groups. (**Fig 1**) Means and standard deviations were calculated. Initial One-way Anova was performed using SPSS software to assess significant difference in the mean scores for each of the three groups. Fischer's exact test showed (F value=65.53; $P<0.001$) significant difference between the three groups (**Table 1**).

In keeping with the objective of the study a Post Hoc Test (Tukey-HSD) performed to compare the DOAP and SEV groups was done. There was no significant difference between the mean scores of the two groups ($P=0.92$). There was significant difference however between SEV and DOAP groups individually against the control group (**Table 2**).

Assessment of the feedback response on SurveyMonkey showed a statistically significant difference in the level of agreement with all elements of training surveyed favouring the DOAP group (**Table 3**). They reported higher levels of satisfaction with the method of training, felt it reinforced elements of risk, safety and caution involved in performing the procedure and also strongly recommended it as a modality of training for trainees at their level. The video group reported lower levels of satisfaction with the confidence instilled by the training process in performing the procedure and were less convinced of the suitability of video assisted training as a method of psychomotor training for trainees at their level of expertise.

In a sub-analysis 8 critical elements of the procedure assessed in the OSCE (Questions 3, 6,10,14, 17, 18, 22, 26) were compiled (**Table 4**). Failure to perform each critical element of the procedure was compared between DOAP and SEV groups. It was observed, that

contrary to the respondent's feedback, the SEV group actually outperformed the DOAP group in the elements of the OSCE (Q3,10,17,18) that pertained to safety and risk mitigation in the procedure. They performed poorly in the steps (Q6,14, 22 & 26) that pertained to pre-procedure hand wash, adequate use of lubricant anaesthetic gel, post procedure replacement of prepuce to normal position and documentation in the notes. The difference failed to achieve statistical significance in any except Q22, pertaining to repositioning of prepuce post procedure, which favoured of the DOAP group.

Despite the significant lower level of confidence expressed by the SEV group, there was no statistically significant difference in their ability to perform the procedure safely vis-à-vis the DOAP group as seen by their OSCE score assessment.

Each of the intervention groups outperformed the Control group with the difference reaching statistical significance.

Inherent Bias & Redressal:

1. Researcher's interest in Structured competency-based training:

   • Excluded by recruiting two trial blinded assessors for each group assessment.

2. Repetition bias for DOAP group:

   • Minimised by limiting feedback to after completing OSCE assessment and submitting evaluation questionnaire.

   • SEV group shown video two times in presence of trainer to match visual impact with opportunity to discuss steps with trainer to match opportunity afforded to DOAP group.

Discussion:

The training of a medical graduate requires development of almost every domain of learning. Most core competencies require the application of more than one skill domain. The move towards competency based medical education (CBME) globally had highlighted this need and also the challenges inherent to ensuring achievement of these competencies [1]. Wide variability in procedural skills have been observed among junior doctors even after adoption of CBME raising questions about their readiness to undertake roles expected of them [2].

Simulation based training provides an evidence-based solution to imparting skills to the modern trainee in the current practise backdrop of reduced availability of trainee patient contact time as a result of reduced inpatient stay for most treatments and reduced tolerance for training related morbidity [4]. Skill labs allow repeated training and assessment in a safe environment till achievement of skills in objectively demonstrated before advancing to the clinical settings. Peyton's 4 step technique (DOAP) is an effective method of psychomotor skill training. This technique has been shown to be significantly effective in procedures involving multiple sequential steps [5,6].

Many studies have confirmed the positive significant benefit of adding multimedia-based training to traditional text-based training [7]. Students have also shown grater satisfaction with the use of video assisted learning in comparison with traditional methods of skill training through demonstrations [8]. No significant difference was noted in procedural skill when video-based teaching was compared to live demonstrations for training in orthodontal procedures [8]. Students report personal comfort, availability of media for review and clarification and scope for personal visualisation and reflection as possible reasons for this preference [8,9].

The present study showed that, though the subjects trained through DOAP reported greater satisfaction and confidence with their training as compared to those trained through the use of a SEV, there was no significant difference in their OSCE performance scores in the skill lab. There was no statistical difference even in the analysis of 8 critical steps selected from

the OSCE questionnaire apart from the step requiring the subject to declare the need to protract the prepuce post catheterisation in uncircumcised individuals where the difference reached statistical significance favouring the DOAP group. The DOAP group also had a numerical advantage over the SEV group in performing documentation of the procedure, though the difference didn't reach statistical significance.

This marginal advantage could be accounted to the simulator model featuring a circumcised penis and the need to document the procedure being conveyed only as a closing statement by the narrator in the SEV rather than a scene of this being done.

The lack of visual impact of both these steps could have resulted in failure on the part of the subject to register the verbal suggestion impressing the need to protract the prepuce post catheterisation in uncircumcised males and ensure documentation of the procedure in the case notes. Including visuals of these steps being performed in the training video could potentially level the field.

Video based training thus appears to be as effective a tool when compared to the DOAP method for training in simple bedside clinical procedures.

There is evidence that structured training improves performance in healthcare professionals in procedural, communication and clinical examination skills over learning by apprenticeship and informal workplace-based learning [10,11].

The present study findings also show a clear statistical advantage in the OSCE scores achieved by subjects trained through either method of structured training (DOAP or SEV) over the subjects in the control group who were trained by the conventional way through apprenticeship and unstructured work-based supervised training.

Conclusion:

Structured training appears to be a key element in implementing CBME to ensure development of competencies. Personalised training through DOAP in small groups appears to be the ideal method of imparting training in common bedside clinical procedures but needs more time and faculty hours. Training using Structured educational videos with the presence and reinforcement by a trainer allows training of larger groups and appears to be equally effective.

References:

1. Vogel D, Harendza S; Basic practical skills teaching and learning in undergraduate medical education–a review on methodological evidence. GMS Journal for Medical Education 2016, August 15; Vol.33(4); Doc 64.

2. Bugaj T J, Nikendei C, Groener J B, Stiepak J, Huber J, Möltner A, Herzog W, Koechel A. Ready to run the wards? – A descriptive follow-up study assessing future doctors' clinical skills. BMC Med Educ. 2018 Nov 12;18(1):257.

3. University of California Irvine School of Medicine Medical Education Simulation Center. http://sites.uci.edu/medsim/files/2015/04/FOLEY-CATHETER-INSERTION-CHECKLIST.pdf

4. Scalese RJ, Obeso VT, Issenberg SB. Simulation Technology for Skills Training and Competency Assessment in Medical Education. J Gen Intern Med. 2008 Jan;23 Suppl 1:46-9

5. Balafoutas D, Joukhadar R, Kiesel M, Häusler S, Loeb S, Woeckel A, Herr D. The Role of Deconstructive Teaching in the Training of Laparoscopy. JSLS. 2019 Apr-Jun;23(2). pii: e2019.00020. doi: 10.4293/JSLS.2019.00020

6. Gradl-Dietsch G, Lübke C, Horst K, Simon M, Modabber A, Sönmez TT, Münker R, Nebelung S, Knobe M. Peyton's four-step approach for teaching complex spinal manipulation techniques – a prospective randomized trial. BMC Med Educ. 2016 Nov 3;16(1):284.

7. Buch SV, Treschow FP, Svendsen JB, Worm BS. Video- or text-based e-learning when teaching clinical procedures? A randomized controlled trial. Adv Med Educ Pract. 2014 Aug 16;5:257-62

8. Botelho MG, Gao X, Jagannathan N. A qualitative analysis of students' perceptions of videos to support learning in a psychomotor skills course. Eur J Dent Educ. 2019 Feb;23(1):20-27.

9. Alqahtani ND, Al-Jewair T, Al-Moammar K, Albarakati SF, ALkofide EA. Live demonstration versus procedural video: a comparison of two methods for teaching an orthodontic laboratory procedure. BMC Med Educ. 2015 Nov 4;15:199. (doi: 10.1186/s12909-015-0479-y)

10. Jünger J, Schäfer S, Roth C, Schellberg D, Friedman Ben-David M, Nikendei C. Effects of basic clinical skills training on objective structured clinical examination performance. Med Educ. 2005 Oct;39(10):1015-20.

11. Rebuck D, Dzyngel B, Khan K, Kesten RN, Chapman KR. The effect of structured versus conventional inhaler education in medical housestaff. J Asthma. 1996;33(6):385-93.

Acknowledgment:

Figures:

Fig 1: Histogram of OSCE results: Normal Distribution

Tables:

Table 1: Three-way Anova Comparison

OSCE Score Results			Anova	
Group	n= Numbers	Mean OSCE Score	F Value	P-Value.
Control	35	17.11		
DOAP	39	25.23	65.528	<0.001
SEV	39	24.92		

Table 2: One Way Anova: Tukey HSD Post Hoc Intergroup Comparison

One Way Anova with Tukey HSD Post Hoc Inter Group Comparison			
Tukey HSD Post Hoc Comparison of Groups	Mean Difference	P Value	Statistical Significance of difference
DOAP v/s SEV	0.308	0.917	No significant difference
Control v/s DOAP	-8.116	<0.001	Significant favouring DOAP
Control v/s SEV	-7.809	<0.001	Significant favouring SEV

Table 3: Analysis of Study subject's Trial experience feedback (Surveymonkey®).

Questions presented	Groups	Numbers in Strong agreement	Percentage in Strong Agreement	P- Value: Significant if P< 0.05
1. Captivating & Promotes interest	DOAP	25	73.5	0.002
	SEV	12	34.3	
2. Clear explanation of Procedure	DOAP	25	73.5	0.007
	SEV	14	40	
3. Elements of risk and safety conveyed	DOAP	18	52.9	0.045
	SEV	9	25.7	
4. Confidence afforded to perform	DOAP	17	50	<0.001
	SEV	4	11.4	
5. Suitability of teaching method	DOAP	26	76.5	<0.001
	SEV	5	14.3	
6. Whether they would recommend this method	DOAP	25	73.5	<0.001
	SEV	3	8.6	
7. Overall satisfaction with teaching method	DOAP	26	76.5	<0.001
	SEV	7	20	
Degree of affirmative agreement with the teaching method was significantly higher in the DOAP Group with a P value of <0.05 in every element.				

Table 4. Subgroup analysis: Scores in 8 critical steps of procedure

Critical Steps assessed by OSCE	Fail / Pass	DOAP	SEV	P- Value Significant if P<0.05
Q3: Taking Consent	Fail	7	4	0.520
	Pass	32	35	
Q6: Hand washing pre-procedure	Fail	1	5	0.200
	Pass	38	34	
Q10: Clean by no-touch technique	Fail	6	3	0.480
	Pass	33	36	
Q14: Insertion of adequate anaesthetic gel	Fail	1	3	0.620
	Pass	38	36	
Q17: No-touch catheter insertion	Fail	2	0	0.490
	Pass	37	39	
Q18: Confirming Urine flow	Fail	11	5	0.092
	Pass	28	34	
Q22: Retraction of prepuce post procedure*	Fail	17	28	0.012
	Pass	22	11	
Q26: Documentation	Fail	6	10	0.260
	Pass	33	29	
* Bias Correction Q22 discounted because MBC simulator had a circumcised penile shaft & hence could not be reliably assessed on OSCE.				