

1 Estimating the impact of reopening schools on the reproduction number
2 of SARS-CoV-2 in England, using weekly contact survey data

3 *James D Munday^{*1a}, Christopher I Jarvis^{*1b}, Amy Gimma^{1c}, Kerry LM Wong^{1d}, Kevin van*

4 *Zandvoort^{1e}, CMMID COVID-19 Working Group, Sebastian Funk^{1f}, W. John Edmunds^{1g}*

5 *1 Centre for Mathematical Modelling of Infectious Disease, London School of Hygiene and*
6 *Tropical Medicine.*

7 *a: James.Munday@lshtm.ac.uk (Corresponding Author)*

8 *b: Christopher.Jarvis@lshtm.ac.uk*

9 *c: Amy.Gimma@lshtm.ac.uk*

10 *d: Kerry.Wong@lshtm.ac.uk*

11 *e: Kevin.Van-Zandvoort@lshtm.ac.uk*

12 *f: Sebastian.Funk@lshtm.ac.uk*

13 *g: John.Edmunds@lshtm.ac.uk*

14 **Abstract**

15 *Background*

16 Schools have been closed in England since the 4th of January 2021 as part of the national
17 restrictions to curb transmission of SARS-CoV-2. The UK Government plans to reopen
18 schools on the 8th of March. Although there is evidence of lower individual-level
19 transmission risk amongst children compared to adults, the combined effects of this with
20 increased contact rates in school settings are not clear.

21 *Methods*

22 We measured social contacts when schools were both open or closed, amongst other
23 restrictions. We combined these data with estimates of the susceptibility and infectiousness
24 of children compared with adults to estimate the impact of reopening schools on the
25 reproduction number.

26 *Results*

27 Our results suggest that reopening all schools could increase R from an assumed baseline
28 of 0.8 to between 1.0 and 1.5, or to between 0.9 and 1.2 reopening primary or secondary
29 schools alone.

30 *Conclusion*

31 Our results suggest that reopening schools is likely to halt the fall in cases observed in
32 recent months and risks returning to rising infections, but these estimates rely heavily on the
33 current estimates or reproduction number and the current validity of the susceptibility and
34 infectiousness profiles we use.

35 **Keywords:** School closure, SARS-CoV-2, COVID-19, Social Contacts, Reproduction
36 Number, CoMix

37 Introduction

38 School closures have been implemented in many countries as part of a broader response to
39 the COVID-19 pandemic [1]. It is well established that children are at low risk of
40 hospitalisation and death as a direct result of infection [2, 3]. Despite this lower risk, there is
41 concern that allowing transmission amongst younger age-groups increases risk of infection
42 in adults, who are at substantially higher risk. The role of schools in transmission is
43 therefore an important question. On the 4th of January 2021, a third national lockdown was
44 announced in England to curb transmission of SARS-CoV-2 [4]. This included the closure of
45 schools, a measure the UK government plans to reverse on the 8th of March.

46 The direct and indirect impact of school closures and eventual reopening is still unclear.
47 There is mixed evidence around the role of schools in community transmission. Existing
48 studies of transmission within schools have wide ranging results [5–7]. Other work
49 demonstrates an increased prevalence amongst school-aged-children when schools return
50 [8, 9] and a higher risk of infections entering households through children than adults.
51 However, the evidence that schools drive transmission in the community remains scarce [10,
52 11]. A particular challenge for many analyses is bias resulting from the age-dependence in
53 case ascertainment due to varying rates of asymptomatic infection [12]. This challenge is
54 then further complicated by changes in epidemiology due to the emergence of new variants
55 [13].

56 The potential change in transmission of SARS-CoV-2 upon reopening schools predominantly
57 depends on a combination of two factors. Firstly, the age-specific risk of transmission upon
58 contact. Secondly, the likely increased rate of contact between members of the population
59 due to school reopening. Multiple studies aimed at understanding the relative transmission
60 risk associated with children indicate lower susceptibility [14–16] and some indicate lower
61 infectiousness [14]. However, evidence of lower transmission risk amongst children alone is
62 insufficient to quantify the impact of reopening schools. There is a need to combine the

63 estimates of reduced susceptibility and infectiousness with age specific contact patterns in
64 this age-group social contacts amongst school-aged-children.

65 There is abundant evidence that children's contacts increase when schools are open,
66 presenting opportunities for increased infectious disease transmission which is well
67 documented in other pathogens such as influenza [17]. Nonetheless, it is important to
68 capture how these contacts vary under the specific conditions presented during the current
69 pandemic response, where social distancing and other mitigations are in effect within
70 schools.

71 CoMix is a large-scale comprehensive social contact survey which has collected data on
72 social contacts in the UK on a weekly basis since the 24th of March 2020 [18]. In this paper,
73 we estimate the impact of opening schools on the reproduction number in England, by
74 combining social-contact data collected during periods where schools were open and closed
75 [18] with estimates of age-stratified susceptibility and infectiousness [14–16].

76 **Methods**

77 **CoMix Data**

78 CoMix is a longitudinal behavioural survey, launched on the 24th of March 2020. The sample
79 is broadly representative of the UK adult population with data collected from approximately
80 2000 individuals per week. Participants are invited to respond to the survey once every two
81 weeks. We collected weekly data by running two alternating panels. Parents complete the
82 survey on behalf of children (17 years old or younger). Participants record direct,
83 face-to-face contacts made on the previous day, specifying certain characteristics for each
84 contact including the age and sex of the contact, whether contact was physical (skin-to-skin
85 contact), and where the contact occurred (e.g. at home, work, while undertaking leisure
86 activities, etc). Further details have been published elsewhere [18]. The contact survey is
87 based on an approach developed for the POLYMOD contact survey [19]. We provide a brief
88 descriptive analysis of the contacts recorded during the November and January lockdown
89 periods by age group and geographical region.

90 **Constructing contact matrices and estimating reproduction number**

91 We constructed age-stratified contact matrices for nine age-groups (0-4, 5-11, 12-17, 18-29,
92 30-39, 40-49, 50-59, 60-69, and 70+). Participants did not report exact ages of contacts, we
93 therefore sampled from the reported age-group with a weighting consistent with contacts
94 reported in the POLYMOD survey. We fitted a truncated negative binomial model to calculate
95 the mean contacts between each participant and contact age-groups. To ensure reciprocity
96 in contacts, we multiplied the matrix by population size vector for England, using United
97 Nations World Population Prospects data [20], before taking the cross-diagonal mean and
98 then dividing by the same population vector again.

99 **Profiles of Age-dependent transmission risk**

100 We consider five age-dependent susceptibility and infectiousness profiles (Table 1):

101 The first profile (i) assumed equal susceptibility and infectiousness in all age groups. This is
102 unlikely to reflect reality but provides an upper limit as a reference point to compare the other
103 profiles.

104 For the second profile (ii) we used results from a mathematical modelling study by Davies et.
105 al [14]. which estimated relative susceptibility and clinical fraction in 9 age groups. The work
106 also reports estimates of 50% infectiousness of sub-clinical cases and reports clinical
107 fraction by age. We used this to calculate infectiousness per age group further detailed in
108 Table 1.

109 The third profile (iii), was based on analyses of household transmission patterns from the
110 Office for National Statistics (ONS) Community Infection Study [15]; 50% susceptibility in
111 children relative to adults but equal infectiousness.

112 For the fourth profile (iv), we performed a meta-analysis of prevalence studies included in a
113 systematic review by Viner et al [16]. We used a random effects model based on the data
114 from Figure 4 of their paper. This resulted in 64% (51% - 81%, 95% confidence interval [CI])
115 susceptibility in children relative to adults, we assumed equal infectiousness between
116 children and adults [16];

117 For the fifth profile (v), we used an independent estimate of relative susceptibility in children
118 (31%, see results section), quantified by comparing reproduction numbers estimated from
119 CoMix data and using a well-established time-series method developed by Abbott et. al [21],
120 which uses a time-series of cases to determine the instantaneous reproduction number
121 under an assumed generation interval and infection to reporting delay distribution.

122 **Table 1** Susceptibility and infectiousness profiles taken from Davies et.al.[14], ONS reports
 123 and Viner et al[16]

Study	Age groups	Susceptibility	Infectiousness	Clinical Fraction
Davies et al ¹	0-4	0.4 (0.25, 0.57)	0.61	0.29 (0.18, 0.44)
	5-10	0.4 (0.25, 0.57)	0.61	0.29 (0.18, 0.44)
	11-17	0.4 (0.27, 0.53)	0.61	0.21 (0.12, 0.31)
	18-29	0.79 (0.59, 0.96)	0.64	0.27 (0.18, 0.38)
	30-39	0.86 (0.69, 0.98)	0.67	0.33 (0.24, 0.43)
	40-49	0.80 (0.61, 0.96)	0.70	0.40 (0.28, 0.52)
	50-59	0.82 (0.63, 0.97)	0.75	0.49 (0.37, 0.60)
	60-69	0.88 (0.70, 0.99)	0.82	0.63 (0.49, 0.76)
	70+	0.74 (0.56, 0.90)	0.85	0.69 (0.57, 0.82)
		Susceptibility	Infectiousness	
ONS ²	0-4	0.5 (0.35, 0.75)	1.0 (0.7, 1.5)	
	5-10	0.5 (0.35, 0.75)	1.0 (0.7, 1.5)	
	11-17	0.5 (0.35, 0.75)	1.0 (0.7, 1.5)	
	18-29	1.0	1.0	
	30-39	1.0	1.0	
	40-49	1.0	1.0	
	50-59	1.0	1.0	
	60-69	1.0	1.0	
	70+	1.0	1.0	
		Susceptibility	Infectiousness	
Viner et al ³	0-4	0.64 (0.51, 0.81)	1.0 (assumed)	
	5-10	0.64 (0.51, 0.81)	1.0 (assumed)	
	11-17	0.64 (0.51, 0.81)	1.0 (assumed)	
	18-29	1.0	1.0	
	30-39	1.0	1.0	
	40-49	1.0	1.0	
	50-59	1.0	1.0	
	60-69	1.0	1.0	
	70+	1.0	1.0	
		Susceptibility	Infectiousness	
CoMix fit	0-4	0.31 (0.30, 0.31)	1.0	
	5-10	0.31 (0.30, 0.31)	1.0	
	11-17	0.31 (0.30, 0.31)	1.0	
	18-29	1.0	1.0	
	30-39	1.0	1.0	
	40-49	1.0	1.0	
	50-59	1.0	1.0	
	60-69	1.0	1.0	
	70+	1.0	1.0	

¹ 95% Credible Intervals

² Approximate results inferred from plot in[15] unknown quantification of uncertainty

³ 95% Confidence Interval

124 Inferring age dependent transmission risk using CoMix data

125 We established independent estimates of susceptibility and infectiousness in children
126 relative to adults. We did this by comparing estimates of R using CoMix contact data with
127 estimates of the time-varying reproduction number in England calculated using case data
128 [21]. To capture the change in contact rates as schools returned in September 2020

129 We calculated a reproduction number resulting from two-weekly rolling contact matrices \mathbf{C}_t
130 and assumed relative susceptibility and infectiousness vectors \mathbf{s} and \mathbf{i} to be:

$$R = r \text{Eig}(\mathbf{C}_t \circ (\mathbf{i} \otimes \mathbf{s})) \quad (1)$$

131 We simplified \mathbf{s} and \mathbf{i} such that adult age-groups (18+) were 1.0 and child age groups were
132 equal, s and i . We inferred s and r , keeping i at 1.0, by fitting our estimates using maximum
133 likelihood estimation to those calculated using the EpiNow2 package [21]. We assumed
134 gamma distributed uncertainty in the time-varying estimates which we parameterised using
135 the mean μ_{rt} and standard deviation σ_{rt} of these estimates over each survey period used to
136 calculate CoMix derived eigenvalues.

$$R \sim \text{Gamma}(\mu_{R_t}, \sigma_{R_t}) \quad (2)$$

137 To show the likelihood surface of relative susceptibility and infectiousness, we calculated the
138 likelihood of a range of combinations of i and s while fitting r .

139 We fitted over 2 periods of time. Firstly, between 27th July and 10th October to most clearly
140 capture the impact of schools returning in the summer whilst minimising issues related to
141 gradual acquisition of natural immunity. Second, We fitted over a longer period of time
142 incorporating data from 10th June.

143 We omitted data at the end of August in both fits due to a short spike in reproduction number
144 estimates, which we believe resulted from large numbers of imported cases from
145 recreational travel. We further omitted two weeks in July when contacts were not recorded

146 for children. We assessed sensitivity to the fitted period, by using a range of fitting options
147 (Figure S4).

148 **Evaluating the impact of reopening schools on Reproduction Number**

149 We created contact matrices using CoMix data collected during the second lockdown, (5th
150 November to 2nd December 2020) to represent contacts during a lockdown with schools
151 open. We used data from 5th to 18th of January 2021 for contacts during a lockdown with
152 schools closed (Supplementary Figures, Figure S1). We constructed further synthetic
153 contact matrices representing opening primary or secondary schools by replacing the
154 contacts of 5-10 year-olds (primary) and 11-17 year-olds (secondary) in the 'schools open'
155 contact matrix (second lockdown), with those from the 'schools closed' contact matrix (third
156 lockdown) (Supplementary Figures, Figure S2).

157 Since the basic reproduction number scales linearly with the dominant eigenvalue of a matrix
158 of effective contact [22], the ratio of the eigenvalues of two effective contact matrices
159 provides a relative change in reproduction number between the three scenarios considered.

160 In the case where infectiousness and susceptibility are equal in all age groups, the effective
161 contact matrix is proportional to the contact matrix itself. Under the scenarios where we
162 assumed infectiousness and susceptibility vary with age, we converted measured contact
163 matrices to effective contact matrices by taking the outer product of the estimated age
164 stratified infectiousness profile and susceptibility profile vectors and calculating the
165 eigenvalues of the Hadamard product of the resulting matrix and the contact matrices.

166 To demonstrate the potential impact of reopening schools, we estimated the relative increase
167 (k) in reproduction number (R) by calculating the ratio of dominant eigenvalues of the
168 effective contact matrix associated with the respective reopening scenario and from the
169 current lockdown period.

$$k = \frac{Eig(\mathbf{C}_{Scenario} \circ (\mathbf{i} \otimes \mathbf{s}))}{Eig(\mathbf{C}_{LD3} \circ (\mathbf{i} \otimes \mathbf{s}))} \quad (3)$$

170 We also calculated how R varies from baseline values between 0.7 and 1.0, from official UK
171 estimates of the reproduction number from [23].

172 **Results**

173 **Descriptive analysis**

174 Adults' contacts were similar when comparing both periods of national lockdown, this is
175 consistent across all settings and regions. Although children's contacts at home were similar
176 between the two periods, contacts at school and other locations were consistently higher in
177 lockdown 2 than lockdown 3. Contacts were very similar between lockdowns in all age-group
178 combinations other than those between children (Figure 1). For participants under 18
179 years-old, the mean number of contacts that were also under 18 years-old was between 6.3
180 (3.9 - 9.0, 90% CI) and 16.7 (13.1 - 20.4, 90% CI) across the regions of England during the
181 November Lockdown. Such contacts were highest in South East, South West and Yorkshire
182 and Humber and lowest in London. The mean number of contacts between children reduced
183 to between 1.8 (1.3 - 2.5, 90% CI) and 2.6 (1.9 - 3.3, 90% CI) during the January Lockdown.

184 **Estimating susceptibility in children relative to adults using CoMix data.**

185 Fitting the R estimates from CoMix data to time-varying R estimates over a period from 27th
186 July to 10th October we estimated susceptibility of 44% (43.5% - 0.45.4%, 95% CI) in
187 children relative to adults (Figure 2, A & C), consistent with profiles ii and iii. When we fitted
188 from the 10th June to 10th October, 2020, we estimated 31% (29.8% - 31.4%, 95% CI)
189 relative susceptibility in children compared to adults (Figure 2, B & D), near the lower range
190 of ONS and Davies et al estimates. We chose to apply the second estimate as the fifth
191 susceptibility profile (v) to represent this lower bound (Table 1) and present fits to other date
192 ranges in the supplementary material (Supplementary Figures, Figure S4).

193 **Evaluation of the impact of reopening schools**

194 Incorporating estimates of differential susceptibility and infectiousness of children compared
195 with adults (profiles ii - v), full school reopening increased R by a factor of between 1.3 and
196 1.9 times the baseline value across the four profiles used (including 90% CI range) (Figure

197 3, Table 2). This would result in an increase of R from 0.8 to above 1.0 for these four profiles.
 198 Partial school reopening resulted in smaller increases in R from 0.8 to between 0.9 and 1.2.
 199 **Table 2** Expected resultant R if schools were reopened for different baseline values of R
 200 reported as median (90% CI)

Susceptibility/ Infectiousness	Attendance	Baseline R			
		0.7	0.8	0.9	1.0 (Scale factor)
1. Equal	Both	1.6 (1.5 - 1.6)	1.8 (1.7 - 1.9)	2.0 (1.9 - 2.1)	2.2 (2.1 - 2.3)
	Primary	1.1 (1.0 - 1.1)	1.2 (1.2 - 1.3)	1.4 (1.3 - 1.5)	1.5 (1.4 - 1.6)
	Secondary	1.1 (1.0 - 1.2)	1.3 (1.2 - 1.3)	1.4 (1.3 - 1.5)	1.6 (1.5 - 1.7)
2. Davies et al	Both	1.1 (1.0 - 1.1)	1.2 (1.1 - 1.3)	1.4 (1.3 - 1.4)	1.5 (1.4 - 1.6)
	Primary	0.9 (0.8 - 0.9)	1.0 (0.9 - 1.0)	1.1 (1.1 - 1.2)	1.2 (1.2 - 1.3)
	Secondary	0.9 (0.8 - 0.9)	1.0 (1.0 - 1.1)	1.1 (1.1 - 1.2)	1.3 (1.2 - 1.3)
3. ONS	Both	1.1 (1.1 - 1.2)	1.3 (1.2 - 1.3)	1.4 (1.4 - 1.5)	1.6 (1.5 - 1.7)
	Primary	0.9 (0.8 - 0.9)	1.0 (1.0 - 1.1)	1.1 (1.1 - 1.2)	1.3 (1.2 - 1.3)
	Secondary	0.9 (0.9 - 1.0)	1.0 (1.0 - 1.1)	1.2 (1.1 - 1.2)	1.3 (1.3 - 1.4)
4. Viner et al	Both	1.3 (1.2 - 1.3)	1.4 (1.4 - 1.5)	1.6 (1.5 - 1.7)	1.8 (1.7 - 1.9)
	Primary	0.9 (0.9 - 1.0)	1.1 (1.0 - 1.1)	1.2 (1.1 - 1.3)	1.3 (1.3 - 1.4)
	Secondary	1.0 (0.9 - 1.0)	1.1 (1.1 - 1.2)	1.2 (1.2 - 1.3)	1.4 (1.3 - 1.4)
5. CoMix fit	Both	0.9 (0.9 - 1.0)	1.1 (1.0 - 1.1)	1.2 (1.2 - 1.3)	1.4 (1.3 - 1.4)
	Primary	0.8 (0.8 - 0.9)	0.9 (0.9 - 1.0)	1.1 (1.0 - 1.1)	1.2 (1.1 - 1.2)
	Secondary	0.8 (0.8 - 0.9)	1.0 (0.9 - 1.0)	1.1 (1.0 - 1.1)	1.2 (1.2 - 1.3)

201 When we assumed equal infectiousness and susceptibility between all age groups (profile i),
 202 reopening schools resulted in more substantial relative changes in R . Full school reopening
 203 increased R by a factor of between 2.1 and 2.3 (Figure 3, Table 2), resulting in an increase
 204 of R to roughly 1.7-1.9 from a baseline of 0.8 (Table 2). Partial re-opening increased R from
 205 0.8 to 1.2-1.3 (Figure 3). We included these estimates for completeness but stress that
 206 assuming that children are equally infectious and susceptible as adults is not compatible
 207 with results from previous studies or our own estimates (Figure 2).

208 Discussion

209 The potential impact of reopening schools on transmission of SARS-CoV-2 is uncertain.

210 Although there have been many attempts to quantify the relative susceptibility and
211 infectiousness of children and adults, these estimates need to be assessed alongside rates
212 of contact to give an indication of the overall risk of transmission in any given setting. We
213 combined social contact data from a large-scale survey in England during two periods of
214 national lockdown, one with schools open and the other with schools closed, with estimates
215 of relative susceptibility of children and adults. We used these data to quantify the potential
216 impact of reopening schools on reproduction number.

217 Whereas adults' contacts were generally similar between the two periods of lockdown, there
218 was markedly higher contact between children during the November lockdown, when
219 schools were open. We observed the change in contacts at school but also in other contacts
220 outside of the home. Increased contact outside of school and home settings includes
221 contacts in wrap around care, which would be expected to rise, however it could also
222 indicate reduced overall adherence amongst children when attending schools physically.

223 The differences in contacts suggest that reopening all schools is highly likely to increase R
224 above 1.0, from an assumed current value 0.8. Reopening primary or secondary is likely to
225 increase R above 1.0. This would be expected to stop or reverse the fall in cases that has
226 been observed since January 2021 [24]. The risk of cases increasing following the reopening
227 of schools is highly dependent on the current value of R . Although cases of the current
228 dominant variant (B.1.1.7) appeared to be increasing whilst national lockdown was still in
229 place in November [10, 13], the latest national serology surveys suggest that immunity levels
230 have substantially increased across the UK [24], resultant from both infections and the
231 national COVID-19 vaccination program. These changes in overall immunity should be
232 reflected in the current estimates of R , but these estimates are lagged due to delays in
233 reporting [25].

234 In November, when schools were open, there was substantial variation in contacts between
235 children by region. We have not presented regional estimates of the impact of reopening
236 schools on R due to low numbers of observations between the lower-level age-group
237 aggregation used in the construction of contact matrices, however the variation in mean
238 contacts points to potential geographical variation in the impact of reopening schools, which
239 may be lower in London than other parts of the country.

240 There are a number of important limitations to this work: Contacts in different settings likely
241 contribute differently to transmission, but we assumed all contacts make equal contributions
242 to transmission, as these differences are not well quantified in the context of control
243 measures. If contacts at school are lower risk than those outside of school the impact of
244 reopening schools would be lower. The age-stratified susceptibility profile is likely to change
245 over time as natural immunity is acquired in the population. The profiles we used each reflect
246 a single point in time. Changes in the relative immunity in children would alter the relative
247 impact of school contacts on overall transmission. We assume adult contacts revert to those
248 observed when all schools were open, which is conservative, in reality, particularly for partial
249 reopening scenarios, adult contacts may not fully return to the same levels. Furthermore,
250 there may also be differences in adherence to restrictions between the two lockdowns,
251 unrelated to school closure. However, the change in adults' contacts between the two
252 periods was relatively small. The proportion of children in school varied over time due to
253 exclusion-based control measures during the autumn, though the proportion attending
254 school remained high during the November lockdown (Supplementary Figures, Figure S3).
255 Contacts of children are reported by parents, which may impact their reliability, particularly in
256 school, where parents are unlikely to witness students' behaviour, it is unclear whether this
257 would lead to systematic bias in reporting either more or fewer contacts.

258 Our work evaluates the impact of reopening schools on the reproduction number in England,
259 which gives an indication of how transmission may be affected. However, there are other
260 factors that reopening schools may introduce, such as the potential for children's contact at

261 school to provide routes of transmission between households, facilitating long chains of
262 transmission that would be otherwise impossible[26]. We are not able to capture these
263 network effects in this analysis, however they may play an important role in the change in
264 epidemiology between school closure and reopening. Second, there is evidence for lower
265 prevalence in primary school than secondary schools [8]. Our framework has not captured
266 these differences suggesting there may be additional factors that reduce the impact of
267 reopening primary schools relative to secondary schools. Furthermore, additional
268 management strategies such as mass testing of school children, may serve to reduce the
269 risk that a contact in a school results in infection beyond those implemented last year.
270 Importantly, with the recent emergence of new variants, particularly B.1.1.7 [27], the baseline
271 R will depend on proportions of these variants as well as contact patterns. Furthermore,
272 these proportions are likely to change, potentially altering the implications of reopening
273 schools.

274 Our results suggest reopening schools is likely to increase R close to or above 1.0, which
275 would stop the decrease in cases observed in recent months. However, precise estimates
276 rely heavily on the baseline values of R and the profiles of susceptibility, generally assuming
277 lower susceptibility and no greater infectiousness in children relative to adults.

278 **List of abbreviations**

279 CI Confidence Interval

280 ONS Office for National Statistics

281 UK United Kingdom

282 **Declarations**

283 **Ethics approval and consent to participate**

284 Participation in this opt-in study was voluntary, and all analyses were carried out on
285 anonymised data. The study and method of informed consent was approved by the ethics
286 committee of the London School of Hygiene & Tropical Medicine Reference number 21795.

287 **Consent to publish**

288 Not applicable

289 **Availability of data and materials**

290 Although it is not possible to share the contact survey data used to generate the contact
291 matrices used in this analysis. The analysis code and contact matrices used are available in
292 an online repository here: https://github.com/jdmunday/CoMix_schools_reopening

293 **Competing interests**

294 None

295 **Funding**

296 CoMix is funded by the EU Horizon 2020 Research and Innovations Programme - project
297 EpiPose (Epidemic Intelligence to Minimize COVID-19's Public Health, Societal and
298 Economical Impact, No 101003688) and by the Medical Research Council (Understanding
299 the dynamics and drivers of the COVID-2019 epidemic using real-time outbreak analytics
300 MC_PC 19065).

301 The following funding sources are acknowledged as providing funding for the named
302 authors. Elrha R2HC/UK FCDO/Wellcome Trust/This research was partly funded by the
303 National Institute for Health Research (NIHR) using UK aid from the UK Government to
304 support global health research. The views expressed in this publication are those of the
305 author(s) and not necessarily those of the NIHR or the UK Department of Health and Social
306 Care (KvZ). This project has received funding from the European Union's Horizon 2020
307 research and innovation programme - project EpiPose (101003688: AG, WJE).
308 FCDO/Wellcome Trust (Epidemic Preparedness Coronavirus research programme
309 221303/Z/20/Z: KvZ). This research was partly funded by the Global Challenges Research
310 Fund (GCRF) project 'RECAP' managed through RCUK and ESRC (ES/P010873/1: CIJ).
311 NIHR (PR-OD-1017-20002: WJE). UK MRC (MC_PC_19065 - Covid 19: Understanding the
312 dynamics and drivers of the COVID-19 epidemic using real-time outbreak analytics: WJE).
313 Wellcome Trust (210758/Z/18/Z: JDM, SFunk). Department of Health and Social Care
314 School Infection Study (PHSEZU7510) (JDM, WJE). No funding (KW).

315 The following funding sources are acknowledged as providing funding for the working group
316 authors. BBSRC LIDP (BB/M009513/1: DS). This research was partly funded by the Bill &
317 Melinda Gates Foundation (INV-001754: MQ; INV-003174: KP, MJ, YL; INV-016832: SRP;
318 NTD Modelling Consortium OPP1184344: CABP, GFM; OPP1139859: BJQ; OPP1183986:
319 ESN; OPP1191821: MA). BMGF (INV-016832; OPP1157270: KA). EDCTP2
320 (RIA2020EF-2983-CSIGN: HPG). ERC Starting Grant (#757699: MQ). This project has
321 received funding from the European Union's Horizon 2020 research and innovation
322 programme - project EpiPose (101003688: KP, MJ, PK, RCB, YL). FCDO/Wellcome Trust
323 (Epidemic Preparedness Coronavirus research programme 221303/Z/20/Z: CABP). This
324 research was partly funded by the Global Challenges Research Fund (GCRF) project
325 'RECAP' managed through RCUK and ESRC (ES/P010873/1: TJ). HDR UK
326 (MR/S003975/1: RME). HPRU (This research was partly funded by the National Institute for
327 Health Research (NIHR) using UK aid from the UK Government to support global health
328 research. The views expressed in this publication are those of the author(s) and not

329 necessarily those of the NIHR or the UK Department of Health and Social Care200908:
330 NIB). MRC (MR/N013638/1: NRW). Nakajima Foundation (AE). NIHR (16/136/46: BJQ;
331 16/137/109: BJQ, FYS, MJ, YL; Health Protection Research Unit for Modelling Methodology
332 HPRU-2012-10096: TJ; NIHR200908: AJK, RME; NIHR200929: FGS, MJ, NGD;
333 PR-OD-1017-20002: AR). Royal Society (Dorothy Hodgkin Fellowship: RL; RP\EA\180004:
334 PK). UK DHSC/UK Aid/NIHR (PR-OD-1017-20001: HPG). UK MRC (MC_PC_19065 - Covid
335 19: Understanding the dynamics and drivers of the COVID-19 epidemic using real-time
336 outbreak analytics: NGD, RME, SC, TJ, YL; MR/P014658/1: GMK). Authors of this research
337 receive funding from UK Public Health Rapid Support Team funded by the United Kingdom
338 Department of Health and Social Care (TJ). UKRI Research England (NGD). Wellcome Trust
339 (206250/Z/17/Z: AJK, TWR; 206471/Z/17/Z: OJB; 208812/Z/17/Z: SC, SFlasche;
340 210758/Z/18/Z: JH, KS, SA, SRM). No funding (AMF, AS, CJVA, DCT, JW, KEA, YWDC).

341 **Authors contributions**

342 JDM, CIJ, WJE conceived of and planned the analysis; JDM and CIJ performed the main
343 analysis with input from WEJ and SF; SF provided estimates of time-varying reproduction
344 number; CIJ, KvZ, and WEJ designed the CoMix contact survey, CIJ, AG, KW, and KvZ
345 cleaned and managed the contact survey data; All authors wrote and reviewed the
346 manuscript. The CMMID COVID-19 Working Group provided discussion and comments.

347 **Acknowledgements**

348 The authors wish to thank Dr Thomas House for his support with interpretation of the ONS
349 susceptibility estimates. We also thank members of SPI-M for their useful discussion which
350 helped shape the final version of this work. We would like to thank the team at Ipsos, who
351 have been excellent in running the survey, collecting the data and allowing for the CoMix
352 study to be implemented rapidly. Finally, we thank Katie Collis for proof reading and
353 excellent discussions.

354 *The following authors were part of the Centre for Mathematical Modelling of Infectious Disease*
355 *COVID-19 Working Group. Each contributed in processing, cleaning and interpretation of data,*
356 *interpreted findings, contributed to the manuscript, and approved the work for publication: Yang Liu,*
357 *Joel Hellewell, Nicholas G. Davies, C Julian Villabona-Arenas, Rosalind M Eggo, Akira Endo, Nikos I*
358 *Bosse, Hamish P Gibbs, Carl A B Pearson, Fiona Yueqian Sun, Mark Jit, Kathleen O'Reilly, Yalda*
359 *Jafari, Katherine E. Atkins, Naomi R Waterlow, Alicia Rosello, Yung-Wai Desmond Chan, Anna M*
360 *Foss, Billy J Quilty, Timothy W Russell, Stefan Flasche, Simon R Procter, William Waites, Rosanna C*
361 *Barnard, Adam J Kucharski, Thibaut Jombart, Graham Medley, Rachel Lowe, Fabienne Krauer,*
362 *Damien C Tully, Kiesha Prem, Jiayao Lei, Oliver Brady, Frank G Sandmann, Sophie R Meakin, Kaja*
363 *Abbas, Gwenan M Knight, Matthew Quaife, Mihaly Koltai, Sam Abbott, Samuel Clifford.*

364 **Additional Files**

365 Supplementary Figures

366 **Figure Captions**

367 **Figure 1. Contacts in the national lockdown periods in November (Lockdown 2) and**
368 **January (Lockdown 3). A)** the distribution of the number of reported contacts in Home,
369 Work, School and Other locations for Adult (> 17 years old) and Child (<= 17 years old)
370 participants. **B)** Mean contacts reported between Children and Adults in each region of
371 England. Error bars show the 90% CI (bootstrapped, 1000 samples).



372 **Figure 2: R estimates using CoMix data fit to time-varying reproduction number**
373 **estimates based on the time series of cases [21].** Transformed likelihood for different
374 combinations of relative susceptibility and infectiousness based on data from **A)** August to
375 October and **B)** June to October and the corresponding R estimates in **C)** and **D)**
376 respectively. 90% CI of the estimates are shown by Grey rectangles for CoMix and the red
377 ribbon for the time-varying reproduction number estimates from case data, red bars show
378 their mean for the CoMix survey periods. Grey shaded areas indicate fitted periods.







379 **Figure 3: The impact of reopening schools on the reproduction number. A)** the relative
380 increase in R (the ratio of dominant eigenvalues between contact matrices for each
381 reopening scenario and that for current contact patterns) under different estimates of the age
382 profile of susceptibility and infectiousness. **B)** The estimated R after reopening schools
383 (points, 90% CI bars) from baseline R of 0.7, 0.8, 0.9 and 1.0 (vertical line). Dashed vertical
384 lines show R = 1.0.

385 References

- 386 1. PHSM. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/phsm>.
387 Accessed 4 Mar 2021.
- 388 2. O'Driscoll M, Ribeiro Dos Santos G, Wang L, Cummings DAT, Azman AS, Paireau J, et al.
389 Age-specific mortality and immunity patterns of SARS-CoV-2. *Nature*. 2021;590:140–5.
- 390 3. Poletti P, Tirani M, Cereda D, Trentini F, Guzzetta G, Marziano V, et al. Age-specific
391 SARS-CoV-2 infection fatality ratio and associated risk factors, Italy, February to April 2020.
392 *Euro Surveill*. 2020;25. doi:10.2807/1560-7917.ES.2020.25.31.2001383.
- 393 4. Prime Minister's Office, Street 10 Downing. Prime Minister announces national lockdown.
394 GOV.UK. 2021.
395 <https://www.gov.uk/government/news/prime-minister-announces-national-lockdown>.
396 Accessed 22 Jan 2021.
- 397 5. Ismail SA, Saliba V, Lopez Bernal J, Ramsay ME, Ladhani SN. SARS-CoV-2 infection and
398 transmission in educational settings: a prospective, cross-sectional analysis of infection
399 clusters and outbreaks in England. *Lancet Infect Dis*. 2021;21:344–53.
- 400 6. Fontanet A, Grant R, Tondeur L, Madec Y, Grzelak L, Cailleau I, et al. SARS-CoV-2
401 infection in primary schools in northern France: A retrospective cohort study in an area of
402 high transmission. doi:10.1101/2020.06.25.20140178.
- 403 7. Stein-Zamir C, Abramson N, Shoob H, Libal E, Bitan M, Cardash T, et al. A large
404 COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020.
405 *Eurosurveillance*. 2020;25. doi:10.2807/1560-7917.es.2020.25.29.2001352.
- 406 8. Office for National Statistics. Coronavirus (COVID-19) Infection Survey, UK. Office for
407 National Statistics; 2020.
408 <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurvey/pilot/13november2020>. Accessed 4 Mar
409 2021.
- 411 9. Riley S, Ainslie KEC, Eales O, Walters CE, Wang H, Atchison C, et al. High prevalence of
412 SARS-CoV-2 swab positivity and increasing R number in England during October 2020:
413 REACT-1 round 6 interim report. medRxiv. 2020;:2020.10.30.20223123.
- 414 10. Mensah AA, Sinnathamby M, Zaidi A, Coughlan L, Simmons R, Ismail SA, et al.
415 SARS-CoV-2 infections in children following the full re-opening of schools and the impact of
416 national lockdown: prospective, national observational cohort surveillance, July-December
417 2020, England. *J Infect*. 2021. doi:10.1016/j.jinf.2021.02.022.
- 418 11. Southall E, Holmes A, Hill EM, Atkins BD, Leng T, Thompson RN, et al. An analysis of
419 school absences in England during the Covid-19 pandemic. *bioRxiv*. 2021.
420 doi:10.1101/2021.02.10.21251484.
- 421 12. Flasche S, Edmunds WJ. The role of schools and school-aged children in SARS-CoV-2
422 transmission. *The Lancet infectious diseases*. 2021;21:298–9.
- 423 13. Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated
424 transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*. 2021.
425 doi:10.1126/science.abg3055.
- 426 14. Davies NG, Klepac P, Liu Y, Prem K, Jit M, CMMID COVID-19 working group, et al.



- 427 Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nat Med.*
428 2020;26:1205–11.
- 429 15. Scientific Advisory Group for Emergencies. TFC: Children and transmission, 4 November
430 2020. 2020.
431 <https://www.gov.uk/government/publications/tfc-children-and-transmission-4-november-2020>.
432 Accessed 22 Jan 2021.
- 433 16. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, et al.
434 Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With
435 Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr.* 2020.
436 doi:10.1001/jamapediatrics.2020.4573.
- 437 17. Eames KTD, Tilston NL, Edmunds WJ. The impact of school holidays on the social
438 mixing patterns of school children. *Epidemics.* 2011;3:103–8.
- 439 18. Jarvis CI, Van Zandvoort K, Gimma A, Prem K, CMMID COVID-19 working group,
440 Klepac P, et al. Quantifying the impact of physical distance measures on the transmission of
441 COVID-19 in the UK. *BMC Med.* 2020;18:124.
- 442 19. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts
443 and mixing patterns relevant to the spread of infectious diseases. *PLoS Med.* 2008;5:e74.
- 444 20. World Population Prospects - Population Division - United Nations.
445 <https://population.un.org/wpp/>. Accessed 6 Mar 2021.
- 446 21. Abbott S, Hellewell J, Thompson RN, Sherratt K, Gibbs HP, Bosse NI, et al. Estimating
447 the time-varying reproduction number of SARS-CoV-2 using national and subnational case
448 counts. *Wellcome Open Res.* 2020;5:112.
- 449 22. Wallinga J, Teunis P, Kretzschmar M. Using data on social contacts to estimate
450 age-specific transmission parameters for respiratory-spread infectious agents. *Am J*
451 *Epidemiol.* 2006;164:936–44.
- 452 23. The R value and growth rate in the UK.
453 <https://www.gov.uk/guidance/the-r-number-in-the-uk>. Accessed 10 Feb 2021.
- 454 24. Office for National Statistics. Coronavirus (COVID-19) Infection Survey, UK. Office for
455 National Statistics; 2021.
456 <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurveypilot/12february2021>. Accessed 4 Mar
457 2021.
- 459 25. Gostic KM, McGough L, Baskerville EB, Abbott S, Joshi K, Tedijanto C, et al. Practical
460 considerations for measuring the effective reproductive number, R_t . *PLoS Comput Biol.*
461 2020;16:e1008409.
- 462 26. Munday JD, Sherratt K, Meakin S, Endo A, Pearson CAB, Hellewell J, et al. Implications
463 of the school-household network structure on SARS-CoV-2 transmission under different
464 school reopening strategies in England. *medRxiv.* 2020.
465 <https://www.medrxiv.org/content/10.1101/2020.08.21.20167965v1>.
- 466 27. Public Health England. Investigation of novel SARS-CoV-2 variant: Variant of Concern
467 202012/01. 2020.
468 [https://www.gov.uk/government/publications/investigation-of-novel-sars-cov-2-variant-variant](https://www.gov.uk/government/publications/investigation-of-novel-sars-cov-2-variant-variant-of-concern-20201201)
469 [-of-concern-20201201](https://www.gov.uk/government/publications/investigation-of-novel-sars-cov-2-variant-variant-of-concern-20201201). Accessed 26 Jan 2021



A**B****C****D**

A

Study: 1. Equal 2. Davies et al 3. ONS 4. Viner et al 5. CoMix fit


**B**