The plasmablast response to SARS-CoV-2 mRNA vaccination is dominated by non-neutralizing antibodies and targets both the NTD and the RBD

Fatima Amanat[1,2], Mahima Thapa[3], Tinting Lei[3], Shaza M. Sayed Ahmed[3], Daniel C. Adelsberg[1], Juan Manuel Carreno[1], Shirin Strohmeier[1], Aaron J. Schmitz[3], Sarah Zafar[1], Julian Q Zhou[4], Willemijn Rijnink[1], Hala Alshammary[1], Nicholas Borcherding[3], Ana Gonzalez Reiche[5], Komal Srivastava, Emilia Mia Sordillo[6], Harm van Bakel[5,7], The Personalized Virology Initiative[#], Jackson S. Turner[3], Goran Bajic[1], Viviana Simon[1,8,9], Ali H. Ellebedy[3,10,11], Florian Krammer[1]

Affiliations

[1]*Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[2]*Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[3]*Department of Pathology & Immunology, Washington University School of Medicine, St. Louis, MO, 63110, USA*
[4]*AbCellera Biologics Inc., Vancouver, BC V5Y 0A1, Canada*
[5]*Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[6]*Department of Pathology, Molecular and Cell Based Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[7]Icahn Institute for Data Science and Genomic Technology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[8]*Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[9]*The Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA*
[10]*Andrew M. and Jane M. Bursky Center for Human Immunology and Immunotherapy Programs, Washington University School of Medicine, Saint Louis, MO 63110, USA*
[11]*Center for Vaccines and Immunity to Microbial Pathogens, Washington University School of Medicine, Saint Louis, MO 63110, USA*

[#]Personalized Virology Initiative (PVI) Study group/Consortium (in alphabetical order): Dr. Bulbul Ahmed, Dr. Deena Altman, Angela Amoako, Mahmoud Awawda, Katherine Beach, Carolina Bermúdez-González, Rachel Chernet, Lily Eaker, Shelcie Fabre, Emily. D. Ferreri, Daniel Floda, Charles Gleason, Dr. Giulio Kleiner, Dr. Denise Jurczyszak, Julia Matthews, Wanni Mendez, Dr. Lubbertus CF Mulder, Jose Polanco, Kayla Russo, Ashley Salimbangon, Dr. Miti Saksena, Amber S. Shin , Levy Sominsky, Sayahi Suthakaran, Dr. Ania Wajnberg

Corresponding authors
Goran Bajic, goran.bajic@mssm.edu
Viviana Simon, viviana.simon@mssm.edu
Ali H. Ellebedy, ellebedy@wustl.edu
Florian Krammer, florian.krammer@mssm.edu

Summary

In this study we profiled vaccine-induced polyclonal antibodies as well as plasmablast derived mAbs from individuals who received SARS-CoV-2 spike mRNA vaccine. Polyclonal antibody responses in vaccinees were robust and comparable to or exceeded those seen after natural infection. However, the ratio of binding to neutralizing antibodies after vaccination was greater than that after natural infection and, at the monoclonal level, we found that the majority of vaccine-induced antibodies did not have neutralizing activity. We also found a co-dominance of mAbs targeting the NTD and RBD of SARS-CoV-2 spike and an original antigenic-sin like backboost to seasonal human coronaviruses OC43 and HKU1. Neutralizing activity of NTD mAbs but not RBD mAbs against a clinical viral isolate carrying E484K as well as extensive changes in the NTD was abolished, suggesting that a proportion of vaccine induced RBD binding antibodies may provide substantial protection against viral variants carrying single E484K RBD mutations.

Introduction

Understanding of the innate and adaptive immune responses to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has progressed rapidly since the beginning of the coronavirus disease 2019 (COVID-19) pandemic (Carvalho et al., 2021). Polyclonal antibody responses against the spike protein of the virus in serum, and to a lesser degree also at mucosal surfaces, have been well characterized with respect to their kinetics, binding capacity and functionality (Grandjean et al., 2020; Isho et al., 2020; Iyer et al., 2020; Ripperger et al., 2020; Seow et al., 2020; Wajnberg et al., 2020). Similarly, encouraging data have been published on both the plasmablast response and the memory B-cell response induced by SARS-CoV-2 infection (Dan et al., 2021; Gaebler et al., 2020; Guthmiller et al., 2021; Huang et al., 2021; Robbiani et al., 2020; Rodda et al., 2021; Wilson et al., 2020). The immune responses to SARS-CoV-2 vaccination, including to mRNA-based vaccines, are less well studied since these vaccines have only become available in the last months of 2020 (Baden et al., 2020; Polack et al., 2020). However, understanding vaccine-induced immunity is of high importance given the goal to achieve immunity for most people through vaccination, rather than as a consequence of infection.

The receptor binding domain (RBD) of the SARS-CoV-2 spike is an important target for serological and B-cell studies because it directly interacts with the cellular receptor angiotensin converting enzyme 2 (ACE2) mediating host cell entry (Letko et al., 2020; Wrapp et al., 2020). Antibodies binding to the RBD can potently block attachment of the virus to ACE2 and thereby neutralize the virus (Barnes et al., 2020). As a consequence, RBD-based vaccines are in development in addition to full length spike-based vaccines (Krammer, 2020). Analyses of the B-cell responses to the spike generally focus on the RBD and on cells sorted with RBD baits introducing an inherent bias by omitting non-RBD targets (Cao et al., 2020; Gaebler *et al.*, 2020; Robbiani *et al.*, 2020; Weisblum et al., 2020). This is also true for B cells and monoclonal antibodies (mAbs) isolated from vaccinated individuals (Wang et al., 2021). However, other epitopes within the spike protein, notably the N-terminal domain (NTD) but also S2, do harbor neutralizing epitopes (Chi et al., 2020; Liu et al., 2020; McCallum et al., 2021b; Song et al., 2020). In fact, the NTD is heavily mutated in the three most prominent variants of concern (VOCs, B.1.1.7, B.1.351 and P.1 (Davies et al., 2021; Faria et al., 2021; Tegally et al., 2020)). Here, we studied the unbiased plasmablast response to SARS-CoV-2 mRNA-based vaccination and report several new findings. First, we document that RBD and NTD co-dominate as B-cell targets on the viral spike protein, highlighting the importance of the NTD. We also report the first vaccine-induced NTD mAbs. In addition, we show that the majority of mAbs isolated are non-neutralizing, which is reflective of the higher binding to neutralization ratios found in serum after vaccination compared to natural infection. Finally, data from plasmablasts suggest that, at least, some of the vaccine-induced response is biased by pre-existing immunity to human β-coronaviruses.

Results

The polyclonal antibody response to mRNA vaccination exceeds titers seen in convalescent individuals but is characterized by a high ratio of non-neutralizing antibodies

In late 2020, six adult participants of an ongoing observational study received mRNA-based SARS-CoV-2 vaccines (**Suppl. Table 1**). Blood from these individuals (termed V1-V6) was collected at several time points including before vaccination (for 4/6), after the first vaccination and at several time points after the second vaccination. We examined their immune responses to recombinant spike protein and RBD in enzyme-linked immunosorbent assays (ELISA), in comparison to those of 30 COVID-19 survivors (**Figure 1A and 1B, Suppl. Table 1**). The sera from convalescent individuals were selected based on their anti-spike titers and grouped into three groups (low +: n=8; moderate ++: n=11; and high +++: n=11, based on the antibody titer measured in the Mount Sinai's CLIA laboratory (Wajnberg *et al.*, 2020)), in order to facilitate identifying different features that may track with the strength of the antibody response. Five out of six vaccinees produced anti-spike and anti-RBD responses that were, at the peak, markedly higher than responses observed even in the high titer convalescent group while one vaccinee (V4) produced titers comparable to the high titer group. Notably, the antibody response peaked one week after the second vaccine dose, followed by a decline in titers over the following weeks as expected from an antibody response to vaccination. Interestingly, anti-RBD antibody titers seemed to decline faster than anti-spike antibody titers, which appeared to be more stable over time. We also measured neutralizing antibody titers using authentic SARS-CoV-2 and found a similar trend with all vaccinees displaying high titers, even though V4 responded with delayed kinetics (**Figure 1C**). Importantly, although at the peak response, the vaccine group mounted neutralization titers that fell in the upper range for the high convalescent group, they did not exceed that group markedly. This finding prompted us to calculate the proportions of spike binding to neutralizing antibodies. For the convalescent group, we found that individuals with lower titers had a higher proportion of binding to neutralizing antibodies than high responding convalescent individuals (**Figure 1D**). When determined at the time of peak response, the vaccinees had the highest proportion of binding to neutralizing antibody titers, indicating an immune response focused on non-neutralizing antibodies or an induction of less potent neutralizing antibodies in general (or both). These proportions remained stable over time with the ratio of binding to neutralizing antibodies in vaccinated individuals being significantly higher than those observed for any of the three convalescent groups (p = 0.0004, 0.0002 and 0.0041 for the three groups respectively; **Suppl. Figure 1**). We also investigated the spike binding to RBD binding ratio and found no difference to convalescent individuals except a general trend towards proportionally less RBD binding over time in the vaccinees (**Suppl. Figure 1**).

mRNA vaccination induces a modest but measurable immune response to seasonal β-coronavirus spike proteins

It has been reported that SARS-CoV-2 infection induces an original antigenic sin-type immune response against human coronaviruses (hCoVs) to which the majority of the human population has pre-existing immunity (Aydillo et al., 2020; Song *et al.*, 2020). Here, we explored whether this phenomenon is also induced by SARS-CoV-2 mRNA vaccination. Antibody titers in four vaccinees against spike protein from α-coronaviruses 229E and NL63 were detectable at the pre-vaccination time point, but did not increase substantially post-vaccination (**Figure 1E-F**; for V5 and V6 no pre-vaccination serum was available). However, titers against the spike proteins of β-coronaviruses OC43 and HKU1 increased substantially in these four vaccinees after vaccination (**Figure 1G-H)**. Thus, vaccination with mRNA SARS-

140   CoV-2 spike also boosts immune responses against seasonal β-coronavirus spike proteins in a manner
141   reminiscent of that reported for natural infection with SARS-CoV-2.

Plasmablast response to SARS-CoV-2 mRNA vaccination targets both the RBD and the NTD

In order to characterize the B-cell response to vaccination in an unbiased manner, plasmablasts
were single-cell sorted from blood specimens obtained from three individuals (V3, V5 and V6) one week
after the booster immunization (**Suppl. Figure 2**). All mAbs were generated from single-cell sorted
plasmablasts and probed for binding to recombinant SARS-CoV-2 spike protein. Twenty-one (40 mAbs
were screened, with 28 being clonally unique, **Suppl. Table 2**) spike-reactive mAbs were isolated from V3,
six (82 screened, 20 unique) from V5 and fifteen (84 screened, 24 unique) from V6 (**Figure 2A**). Using
recombinant spike, RBD, NTD and S2 proteins, we mapped the domains to which these mAbs bind.
Interestingly, only a minority of these antibodies recognized RBD (24% for V3, 47% for V6 and no RBD
binders were identified for V5) (**Figure 2B and 2E**). A substantial number of the isolated mAbs bound to
NTD including 14% for V3, 33% for V5 and 33% for V6 (**Figure 2C and 2E**). These data indicate that RBD
and NTD are co-dominant in the context of mRNA-induced plasmablast response. The epitopes for the
majority of the remaining spike binding mAbs, 52% for V3, 50% for V5 and 20% for V6, mapped to S2
(**Figure 2D and 2E**). Only three mAbs were not accounted for in terms of binding target (two for V3 and
one for V5, **Figure 2E**).

The majority of isolated mAbs from SARS-CoV2 vaccinees are non-neutralizing

All antibodies were tested for neutralizing activity against the USA-WA1/2020 strain of SARS-CoV-
2. Only a minority of the binding antibodies, even those targeting the RBD, showed neutralizing activity
(**Figure 2F and 2G**). For V3, only one (an RBD binder) out of 21 mAbs (5%) displayed neutralizing activity
(**Figure 2G**). For V5, a single NTD antibody neutralized authentic SARS-CoV-2 (17%) (**Figure 2G**). The
highest frequency of neutralizing antibodies was found in V6 (34%) with one RBD neutralizer and four NTD
neutralizers (**Figure 2G**). Interestingly, the highest neutralizing potency was found in mAb PVI.V5-6, an
NTD binder followed by PVI.V6-4, an RBD binder.

We also tested all the antibodies for reactivity to the spike proteins of the four hCoVs 229E, NL63,
HKU1 and OC43. No antibody binding to the spike proteins of α-coronaviruses 229E and NL63 was found
but we identified five mAbs (including three from V3, one from V5 and one from V6) that bound, to varying
degrees, to the spike of OC43, which, like SARS-CoV-2, is a β-coronavirus (**Figure 2H**). Three mAbs showed
strong binding (PVI.V3-8, PVI.V3-12 and PVI.V6-1), while PVI.V3-17 showed an intermediate binding
phenotype and PVI-V5-1 bound very weakly. Three of these mAbs also showed binding to the spike of
HKU1, another β-coronavirus. Of these, PVI.V6-1 showed only very weak binding while PVI.3-8 and PVI.3-
12 had low minimal binding concentrations (MBCs) indicating higher affinity (**Figure 2I**).

The spike-reactive plasmablast response is dominated by IgG1+ cells and is comprised of a mixture of
cells with low and high levels of somatic hypermutation (SHM)

Single-cell RNA sequencing (scRNAseq) was performed on bulk sorted plasmablasts from the
three vaccinees (V3, V5, V6) to comprehensively examine the transcriptional profile, isotype distribution
and somatic hypermutation (SHM) of vaccine-induced plasmablasts. We analyzed 4,584, 3,523 and 4,461
single cells from subjects V3, V5, and V6, respectively. We first verified the identity of sequenced cells as
plasmablasts through the combined expression of B cell receptors (BCRs) (**Figure 3A**) and that of the

canonical transcription as well as other factors essential for plasma cell differentiation, such as *PRDM1*,
XBP1 and *MZB1* (**Figure 3B**). To identify vaccine-responding B cell clones among the analyzed
plasmablasts, we used scRNAseq to also analyze gene expression and V(D)J libraries from the sorted
plasmablasts and clonally matched the BCR sequences to those from which spike-specific mAbs had been
made. Using this method, we recovered 332, 7 and 1,384 BCR sequences from the scRNAseq data that are
clonally related to the spike-binding mAbs derived from subjects V3, V5 and V6, respectively (**Figure 3C**).
It is important to note here that we were not able to recover clonally related sequences for all of the mAbs
that we cloned and expressed from each of the three vaccinees.

We next examined the isotype and IgG subclass distribution among the recovered sequences.
IgG1 was by far the most dominant isotype in the three vaccinees (**Figure 3D**). Finally, we assessed the
level of somatic hypermutation (SHM) among the mAbs-related sequences from the three subjects. We
used the SHM levels observed in human naïve B cells and seasonal influenza virus vaccination-induced
plasmablasts that were previously published for comparison (Turner et al., 2020). Spike-reactive
plasmablasts from V3 and V6 but not V5 had accumulated SHM at levels that are significantly greater than
those observed with naïve B cells (**Figure 3E, left panel**). Strikingly, the SHM among V6 plasmablasts was
equivalent to those observed after seasonal influenza virus vaccination (**Figure 3E, left panel**). We
reasoned that the high level of SHM among spike-reactive plasmablasts may be derived from those
targeting conserved epitopes that are shared with human β-coronaviruses. Indeed, we found that the
SHM level among clones that are related to cross-reactive mAbs was significantly higher than their non-
cross-reactive counterparts (**Figure 3E, right panel**).

Competition of RBD binding neutralizing mAbs with ACE2 and affinity of variant RBDs for human ACE2

Two mAbs were identified as neutralizing and binding to RBD. We wanted, therefore, to test if
they competed with ACE2 for RBD binding. Concentration-dependent competition was indeed observed
for both mAbs demonstrating that inhibition of ACE2 binding is the mechanism of action of the two mAbs
(**Figure 4**). Since we prepared RBD proteins of viral variants of concern for analysis of antibody binding
(see below), we also wanted to assess the affinity of each variant RBD for human ACE2. Using biolayer
interferometry (BLI), we measured association and dissociation rates of the N501Y RBD mutant (B.1.1.7
carries that mutation as its sole RBD mutation), Y453F, as found in mink isolates (Larsen et al., 2021),
N439K, which is found in some European clades (Thomson et al., 2021), a combination of Y453F and
N439K, E484K (part of B.1.351 and P.1) as well as for the B.1.351 and the P.1 RBDs for a recombinant
version of human ACE2 (**Figure 4A, 4B and 4D**). Almost all of the single and double mutations in RBD tested
increased affinity to human ACE2. Specifically, N501Y and Y453F combined with N439K increased affinity
for human ACE2 by 5-fold (**Figure 4D, Suppl. Figure 3**). In contrast, E484K on its own decreased affinity by
4-fold. Of note, the B.1.351 RBD affinity for ACE2 was comparable to that of the wild-type RBD. These data
were confirmed using an ELISA-based method which showed the same trends (**Suppl. Figure 4**).

Binding profiles of polyclonal serum and mAbs to RBDs carrying mutations found in viral variants of
concern

Next, we assessed binding of sera from vaccinated individuals, COVID-19 survivors and mAbs
derived from plasmablasts to variant RBDs. Our panel of RBDs includes published mAb escape mutants,
RBD mutants detected by the Mount Sinai Hospital's Pathogen Surveillance Program in patients seeking
care at the Mount Sinai Health System in NYC as well as mutations found in viral variants of interest and
variants of concern (Baum et al., 2020; Greaney et al., 2021; Larsen *et al.*, 2021; Thomson *et al.*, 2021;

Weisblum *et al.*, 2020). Serum from convalescent individuals showed strong fluctuations depending on
the viral variant (**Figure 5A**). In general, single mutants E406Q, E484K and F490K exerted the biggest
impact on binding. However, complete loss of binding was rare and 2-4-fold reduction in binding was more
common. Interestingly, almost all sera bound better to N501Y RBD (B.1.1.7) than to wild-type RBD
(average 129% compared to wild type). Conversely, the B.1.351 RBD caused, on average, a 39% reduction
in binding. The impact was slightly lower for the P.1 RBD (average 70% binding compared to wild-type).
For sera from the six vaccinated individuals, however, the highest reduction seen was only two-fold for
E406Q, N440K, E484K and F490K (**Figure 5B**). Of note, the vaccinees' later samples (V1=d89, V2=d102,
V3=d47, V4=d48, V5=49 and V6=48) were assayed to allow for some affinity maturation. The highest
reduction observed for E484K, F484A, B.1.351 and P.1 were also approximately two-fold but this did not
apply to all six vaccinees. Some vaccinees maintained binding levels against these RBDs at levels
comparable to wild-type RBD.

RBD binding mAbs were also tested for binding to the same variants. In general, mAbs maintained
binding levels within 2-fold of the binding seen with the wild-type RBD with some exceptions. In fact, for
most mAbs, no impact on binding was observed (**Figure 5C**) with the exception of PVI.V3-9, which lost
binding to the RBD carrying F486A. Although there was a negative impact on binding of several mAbs to
the B.1.351 variant, binding was almost unaffected by the mutations in the P.1 variant RBD. Only one
mAb, PVI.V6-4, showed a drop in binding to P.1.

Escape of an NTD and E484K mutant virus from polyclonal post-vaccination serum is negligible but
NTD mutations significantly impact the neutralizing activity of NTD binding mAbs

Through the Mount Sinai Hospital's Pathogen Surveillance Program, we had access to the SARS-
CoV-2 isolate PV14252 (Clade 20C, Pango lineage B.1) that featured two mutations (W64R, L141Y) and
one deletion (Δ142-145) in the NTD as well as the E484K mutation in the RBD (**Figure 5D**). To determine
the susceptibility of this virus variant to neutralization by post-vaccination serum, we performed
microneutralization assays. Wild-type SARS-CoV-2 and PV14252 were tested in parallel to ensure that the
assay setup for both viruses allowed comparison. We found a relatively minor impact when testing
polyclonal sera from vaccinees for neutralizing activity (**Figure 5E**). The activity of sera from V2, V5 and V6
slightly increased while the activity for V1, V3 and V4 decreased. Next, we tested the seven neutralizing
mAbs that we isolated from plasmablasts. Consistent with their binding profiles in the variant RBD ELISA,
the two RBD mAbs neutralized both viruses with comparable efficiency (**Figure 5F**). In fact, the activity of
PVI.V3-9 increased slightly (**Figure 5F**). In stark contrast, all five anti-NTD antibodies completely lost
neutralizing activity against PV14252 due to mutations present in the NTD of this viral isolate.

B.1.1.7 and B.1.351 partially escape from plasmablast derived neutralizing antibodies

We also tested the neutralizing activity of the two RBD and the five NTD antibodies against the
variants of concern B.1.1.7 and B.1.351 (**Figure 5G**). Both variants contain deletions as well as mutations
in the NTD. In addition, B.1.1.7 carries the N501Y RBD mutation and B.1.351 carries N417K, E484K and
N501Y mutations in the RBD (**Figure 4 A and B**). The two RBD binding antibodies lost no (PVI.V6-4) or little
(PVI.V3-9) neutralizing activity against B.1.1.7. However, PVI.V3-9 lost all activity against B.1.351 and the
remaining neutralizing activity of PVI.V6-4 was low (but measurable). All but one (PVI.V6-11)) NTD mAbs
lost neutralizing activity against B.1.1.7 and all of them lost neutralizing activity against B.1.351 once more
highlighting the importance of changes in the NTD on the antibody activity.

Discussion

Our knowledge of B-cell responses to SARS-CoV-2 mRNA vaccination remains incomplete. We
urgently need information about the nature of polyclonal vaccine-induced responses as well as unbiased,
in depth analyses of plasmablast responses. Our data provide important new insights into these responses
in comparison with immune responses to natural infection. Indeed, SARS-CoV-2 infection results in a very
heterogeneous antibody response to the spike protein in terms of antibody quantity. In contrast, mRNA
vaccination appears to induce a high antibody response of relatively homogenous titers. However, we
also found that vaccinees generate more non-neutralizing antibodies than COVID-19 survivors resulting in
a lower ratio of neutralizing to binding antibodies. These data were already apparent in the early phase
clinical trials but remained unrecognized at the time (Walsh et al., 2020). Interestingly, low titer
convalescent serum had the highest relative amount of neutralizing antibodies, whereas the proportion
of binding antibodies was increased in sera with higher measured antibody titers. The majority of
plasmablasts sampled after vaccination do, in fact, produce non-neutralizing antibodies. Two recent
studies have performed a similarly unbiased plasmablast analysis for individuals naturally infected with
SARS-CoV-2 (Cho et al., 2021; Huang *et al.*, 2021). Of course, the antibody response after SARS-CoV-2
infection is not only targeting the spike protein but several other proteins expressed by the virus. When
accounting for spike binding only, these studies report proportions of 44% and 25% neutralizing antibodies
(Cho *et al.*, 2021; Huang *et al.*, 2021). While plasmablast analysis is in general not quantitative (e.g. one
clone per clonotype is selected etc.) our analysis of post-vaccination plasmablasts found a lower number
of neutralizing antibodies (17%).

Future studies are needed to reveal the role of non-neutralizing antibodies in SARS-CoV-2 immune
protection. Indeed, antibody functions other than neutralization have been shown to correlate with
protection (Bartsch et al., 2021; Gorman et al., 2021; Schäfer et al., 2021). The importance of absolute
antibody titers and not ratios is underscored by the fact that post-vaccination neutralization titers were
equal to or exceeded the titers found in the high responder convalescent group.

Of the four seasonal CoVs that are widely circulating in humans, β-coronaviruses OC43 and HKU1 have
higher homology to SARS-CoV-2 spike. Vaccinated individuals mounted a response to spike proteins from
OC43 and HKU1 but not to α-coronaviruses 229E and NL63. This phenomenon resembles the immune
imprinting described in influenza virus immunology and has already been shown for natural infection with
SARS-CoV-2 where a 'backboost' to β-coronaviruses was also found (Aydillo *et al.*, 2020; Song *et al.*, 2020).
A few of the mAbs isolated in our study had, indeed, such a cross-reactive phenotype. It remains unclear
whether these antibodies, which target mostly S2 epitopes, contribute to protection against SARS-CoV-2,
OC43 or HKU1 infection. However, the cross-reactive epitopes of mAbs that do bind SARS-CoV-2, HKU1
and OC43 spikes could provide the basis for future pan-β-coronavirus vaccines. While it is likely the case
that the B-cells producing these mAbs come from recall responses and were initially induced by human β-
coronaviruses (which is supported by serology and of course the extensive SHM that the mAbs show),
they could hypothetically also be *de novo* induced antibodies. While this is probably not the case, we
cannot exclude this possibility with our current data.
302

303   Another interesting point we noted is the co-dominance of RBD and NTD. Previous analyses of B-
304 cell responses to SARS-CoV-2 mRNA vaccination focused on cells baited by labeled RBD (Wang *et al.*,
305 2021). We, in contrast, took an unbiased approach to sort and clone plasmablasts in an antigen-agnostic

manner. We found similar levels of NTD and RBD binders with many mAbs binding to epitopes outside the RBD and the NTD. In one vaccinee not a single RBD binding mAbs was isolated with the caveats that the overall number of mAbs derived from that individual were low and their polyclonal serum antibody responses included RBD recognition. These data suggest that the NTD, which also harbors neutralizing epitopes, is - at least - as important as the RBD and warrants as much attention. In fact, five out of seven neutralizing antibodies isolated in this study bound to the NTD and only two targeted the RBD. Recent studies analyzing the plasmablast response after natural infection have found a similar co-dominance of RBD and NTD (Cho *et al.*, 2021; Huang *et al.*, 2021) with one study reporting 59 mAbs targeting the RBD, 64 targeting the NTD and 46 binding outside of RBD and NTD and the second study finding 10 RBD mAbs, 13 non-RBD S1 binding mAbs (strongly suggesting NTD binding) and 9 mAbs targeting S2. Further characterization of the mAbs obtained in our study showed a complete loss of neutralization against an authentic, replication-competent variant virus that harbored extensive changes in the NTD. All NTD mAbs also lost neutralizing activity against B.1.351 and all but one lost activity against B.1.1.7. These observations may explain why a reduction in neutralization against the viral variant of concern B.1.1.7 is seen in some studies despite the fact the N501Y substitution in the RBD of this variant does not significantly impact binding and neutralizing activity (Emary et al., 2021). The key role of NTD as target for antibodies has recently also been shown using memory B cell derived mAbs (McCallum et al., 2021a).

In addition, we assessed the impact of different RBD mutations on affinity towards human ACE2. Interestingly, N501Y increased the affinity by five-fold. This increase in receptor binding affinity may contribute to the higher infectivity of B.1.1.7, which carries this mutation in its RBD. In contrast, introduction of E484K reduced the affinity by 4-fold which may explain why virus variants carrying only the E484K mutation have rarely spread efficiently, although viruses carrying E484K have been detected since the fall of 2020 in a handful of patients receiving care at the Mount Sinai Health System and have also been reported in immunocompromised patients (Choi et al., 2020). It is tempting to speculate that the N501Y mutation enables the acquisition of E484K without a fitness loss. In fact, the B.1.351 RBD, which carries N501Y and E484K (as well as N417K) showed binding to hACE2 that was similar to wild- type RBD. Recently, B.1.1.7 variant strains carrying E484K, in addition to N501Y, have been isolated in the UK (PHE, 2021), providing evidence for the hypothesis that N501Y enables acquisition of mutations in the RBD that may be detrimental to receptor binding. However, recent expansion of B.1.526, a lineage also featuring E484K but without N501Y in New York City, suggests that this fitness loss may be overcome by other, yet uncharacterized, changes in the virus as well (Annavajhala et al., 2021; Lasek-Nesselquist et al., 2021). Interestingly, binding of convalescent sera to the N501Y RBD was also increased, suggesting that changes that increase affinity for the receptor may also increase affinity of a set of antibodies that may mimic the receptor.

We also noted that the two neutralizing antibodies against the RBD showed some reduced binding to a mutant RBD carrying the E484K mutation while having similar or even increased neutralizing potency against a variant virus carrying the E484K mutation as the only change in its RBD. The reduced affinity of the E484K variant RBD for hACE2 could render the virus more susceptible to RBD binding mAbs. Thus, an antibody binding to the RBD may just be more effective in interfering with a low affinity as compared to a high affinity RBD-hACE2 interaction. Increased affinity as an escape mechanism for viruses has been described in the past (Hensley et al., 2009; O'Donnell et al., 2012) and the converse mechanism could be at play here.

Whether or not the current vaccines will provide effective protection against circulating and emerging viral variants of concern is an important question which has gathered a lot of attention in early 2021. Our data indicate that reduction in binding to the E484K and B.1.351 variant RBDs was minor (often only 2-fold) compared to reported reduction in neutralization (which ranges from 6-8 fold to complete loss of neutralization (Cele et al., 2021; Wibmer et al., 2021; Wu et al., 2021)). Although not tested here, it is likely that the reduction in binding to full length spike is even lower, given the many epitopes on the spike other than NTD and RBD. The maintenance of binding to a large degree observed in this study suggests that viral variants will have a minor impact on serological assays which are currently in wide use for medical, scientific and public health reasons. Binding, non-neutralizing antibodies have also been shown to have a protective effect in many viral infections (Asthagiri Arunkumar et al., 2019; Dilillo et al., 2014; Saphire et al., 2018) and may be a factor in the substantial residual protection seen in the Johnson & Johnson and Novavax vaccine trials against B.1.351 in South Africa (Shinde et al., 2021). Production of non-neutralizing antibodies may also play a role in protection by mRNA vaccines after the first dose, as it is substantial and occurs during a time when neutralizing antibody titers are either very low or absent (Baden *et al.*, 2020; Dagan et al., 2021; Polack *et al.*, 2020). Finally, although some antibodies may lose neutralizing activity due to reduced affinity, they do still bind. Furthermore, B cells with these specificities potentially could undergo affinity maturation after exposure to a variant virus or a variant spike-containing vaccine, leading to high affinity antibodies to variant viruses of concern.

In summary, we demonstrate that the antibody responses to SARS-CoV-2 mRNA vaccination comprise a large proportion of non-neutralizing antibodies and are co-dominated by NTD and RBD antibodies. The NTD portion of the spike represents, thus, an important vaccine target. Since all viral variants of concern are heavily mutated in this region, these observations warrant further attention to optimize SARS-CoV-2 vaccines. Finally, broadly cross-reactive mAbs to β-coronavirus spike proteins are induced after vaccination, and suggest a potential development path for a pan- β-coronavirus vaccine.

Acknowledgements

We would like to thank the study participants for their generosity and willingness to participate in longitudinal COVID19 research studies. None of this work would be possible without their contributions. We would like to thank Dr. Randy A. Albrecht for oversight of the conventional BSL3 biocontainment facility, which makes our work with live SARS-CoV-2 possible. We are also grateful for Mount Sinai's leadership during the COVID-19 pandemic. We want to especially thank Drs. Peter Palese, Carlos Cordon-Cardo, Dennis Charney, David Reich and Kenneth Davis for their support. We would also like to thank Bassem Mohamed and Wooseob Kim for their help with preparing the scRNAseq libraries. This work was partially funded by the NIAID Collaborative Influenza Vaccine Innovation Centers (CIVIC) contract 75N93019C00051, NIAID Center of Excellence for Influenza Research and Surveillance (CEIRS, contract # HHSN272201400008C and HHSN272201400006C), NIAID grants U01AI141990 and U01AI150747, by the generous support of the JPB Foundation and the Open Philanthropy Project (research grant 2020-215611 (5384); and by anonymous donors. J.S.T. was supported by NIAID 5T32CA009547.

Conflict of interest statement

The Icahn School of Medicine at Mount Sinai has filed patent applications relating to SARS-CoV-2 serological assays and NDV-based SARS-CoV-2 vaccines which list Florian Krammer as co-inventor. Viviana Simon is also listed on the serological assay patent application as co-inventors. Mount Sinai has spun out a company, Kantaro, to market serological tests for SARS-CoV-2. Florian Krammer has consulted for Merck and Pfizer (before 2020), and is currently consulting for Pfizer, Seqirus and Avimex. The Krammer laboratory is also collaborating with Pfizer on animal models of SARS-CoV-2. Ali Ellebedy has consulted for InBios and Fimbrion Therapeutics (before 2021) and is currently a consultant for Mubadala Investment Company. The Ellebedy laboratory received funding under sponsored research agreements that are unrelated to the data presented in the current study from Emergent BioSolutions and from AbbVie.

Materials and methods.

Human subjects and specimen collection. The study protocols for the collection of clinical specimens from individuals with and without SARS-CoV-2 infection by the Personalized Virology Initiative were reviewed and approved by the Mount Sinai Hospital Institutional Review Board (IRB-16-16772; IRB-16-00791; IRB-20-03374). All participants provided written informed consent prior to collection of specimen and clinical information. All specimens were coded prior to processing and analysis. An overview of the characteristics of the vaccinees as well as the study participants with and without COVID-19 is provided in **Suppl. Table 1**. The vaccinees received two doses of the Pfizer mRNA vaccine.

Whole blood was collected via phlebotomy in serum separator tubes (SST) or ethylenediaminetetraacetic acid (EDTA) tubes. Serum was collected after centrifucation as per manufacturers' instructions. Peripheral blood mononuclear cells (PBMCs) isolation was performed by density gradient centrifugation using SepMate tubes (Stemcell) according to manufacturers' instructions. PBMCs were cryo-preserved and stored in liquid nitrogen until analysis.

Recombinant proteins. All recombinant proteins were produced using Expi293F cells (Life Technologies). Receptor binding domain (RBD) and spike protein of SARS-CoV-2 (GenBank: MN908947.3) was cloned into a mammalian expression vector, pCAGGS as described earlier (Amanat et al., 2020b; Stadlbauer et al., 2020). RBD mutants were generated in the pCAGGS RBD construct by changing single residues using mutagenesis primers. All proteins were purified after transient transfections with each respective plasmid. Six-hundred million Expi293F cells were transfected using the ExpiFectamine 293 Transfection Kit and purified DNA. Supernatants were collected on day four post transfection, centrifuged at 4,000 g for 20 minutes and finally, the supernatant was filtered using a 0.22 um filter. Ni-NTA agarose (Qiagen) was used to purify the protein via gravity flow and proteins were eluted as previously described (Amanat *et al.*, 2020b; Stadlbauer *et al.*, 2020). The buffer was exchanged using Amicon centrifugal units (EMD Millipore) and all recombinant proteins were finally re-suspended in phosphate buffered saline (PBS). Proteins were also run on a sodium dodecyl sulphate (SDS) polyacrylamide gels (5–20% gradient; Bio-Rad) to check for purity (Amanat et al., 2018; Margine et al., 2013). Plasmids to express recombinant spike proteins of 229E, HKU1, NL63 and OC43 were generously provided by Dr. Barney Graham (Pallesen et al., 2017). NTD and S2 proteins were purchased from SinoBiologics.

ELISA. Ninety-six well plates (Immulon 4 HBX; Thermo Scientific) were coated overnight at 4°C with recombinant proteins at a concentration of 2 ug/ml in PBS (Gibco; Life Technologies) and 50 uls/well. The next day, the coating solution was discarded. One hundred uls per well of 3% non-fat milk prepared in PBS

(Life Technologies) containing 0.01% Tween-20 (TPBS; Fisher Scientific) was added to the plates to block
the plates for 1 hour at room temperature (RT). All serum dilutions were prepared in 1% non-fat milk
prepared in TPBS. All serum samples were diluted 3-fold starting at a dilution of 1:50. After the blocking
step, serum dilutions were added to the respective plates for two hours at RT. Next, plates were washed
thrice with 250 uls/well of TPBS to remove any residual primary antibody. Secondary antibody solution
was prepared in 1% non-fat milk in TPBS as well and 100 uls/well was added to the plates for 1 hour at
RT. For human samples, anti-human IgG conjugated to horseradish peroxidase (HRP) was used at a
dilution of 1:3000 (Millipore Sigma; catalog #A0293). For mouse samples, anti-mouse IgG conjugated to
HRP was used at the same dilution (Rockland antibodies and assays; catalog #610-4302). Specifically, a
mouse anti-histidine antibody (Takara; catalog #631212) was used as a positive control to detect proteins
with a hexa-histidine tag. Once the secondary incubation was done, plates were again washed thrice with
250uls/well of TPBS. Developing solution was made in 0.05M phosphate-citrate buffer at pH 5 using o-
phenylenediamine dihydrochloride tablets (Sigma-Aldrich; OPD) at a final concentration of 0.04 mg/ml.
One hundred uls/well of developing solution was added to each plate for exactly 10 minutes after which
the reaction was halted with addition of 50 uls/well of 3M hydrochloric acid (HCl). Plates were read at an
optical density of 490 nanometers using a Synergy 4 (BioTek) plate reader. Eight wells on each plate
received no primary antibody (blank wells) and the optical density in those wells was used to assess
background. Area under the curve was calculated by deducting the average of blank values plus 3 times
standard deviation of the blank values. All data was analyzed in Graphpad Prism 7. This protocol has been
described in detail earlier (Bailey et al., 2019; Wohlbold et al., 2015).

Purified monoclonal antibodies were used at a concentration of 30 ug/ml and then subsequently diluted
3-fold. Purified monoclonal antibodies were only incubated on the coated plates for an hour. The
remaining part of the protocol was the same as above (Amanat et al., 2020a; Wohlbold et al., 2016).

Bio-layer Interferometry Binding Experiments. Bio-layer Interferometry (BLI) experiments were
performed using the BLItz system (fortéBIO, Pall Corporation). Recombinant human Fc fusion ACE2
(SinoBiological) was immobilized on an anti-human IgG Fc biosensor, and RBDs were then applied to
obtain binding affinities. Single-hit concentrations were tested at 5.8 µM for binding. All measurements
were repeated in subsequent independent experiments. K_D values were obtained through local fit of the
curves by applying a 1:1 binding isotherm model using vendor-supplied software. All experiments were
performed in PBS pH 7.4 and at room temperature.
461
hACE2 competition interferometry experiments, Interferometry experiments were performed using a
BLItz instrument (fortéBIO, Sartorius). Polyhistidine-tagged Fabs were immobilized on Ni-NTA biosensors
at 10 µg/ml and SARS-CoV-2 RBD was supplied as analyte at 5µM alone or pre-mixed with hACE2-Fc at
different concentrations. Maximal signal at association (Rmax) was used to plot the concentration-
dependent competition with hACE2. All experiments were performed in PBS at pH 7.4 and at room
temperature.
468
RBD-hACE2 ELISA. 25ng of hACE2-Fc fusion protein expressed in HEK293 cells were adhered to high-
capacity binding, 96 well-plates (Corning) overnight in PBS. Plates were blocked with 5% BSA in PBS
containing Tween-20 (PBS-T) for 1hr at room temperature (RT). Blocking solution was discarded and 5-
fold dilutions of 6xHis-tagged RBDs in PBS were added to wells and incubated for 1hr at RT. Plates were
then washed three times with PBS-T. Anti-polyhistidine IgG-Biotin (Abcam) in PBS-T was added to each
and incubated for 1hr at RT. Plates were then washed three times with PBS-T. Streptavidin-HRP (Abcam)
in PBS-T was added to each and incubated for 1hr at RT. Plates were then washed three times with PBS-T

Plates were developed using 1-Step Ultra TMB substrate (ThermoFisher), stopped with sulfuric acid and
immediately read using a plate reader at 450nm. Data were plotted using Prism 9 (GraphPad Software)
and affinities determined by applying a nonlinear regression model.
479

Viruses and cells. Vero.E6 cells (ATCC #CRL-1586) cells were maintained in culture using Dulbecco's
Modified Eagles Medium (DMEM, Gibco) which was supplemented with 10% fetal bovine serum (FBS,
Corning) and antibiotics solution containing 10,000 units/mL of penicillin and 10,000 µg/mL of
streptomycin (Pen Strep, Gibco)(10). Wild type SARS-CoV-2 (isolate USA-WA1/2020), hCoV-19/South
Africa/KRISP-K005325/2020 (B.1.351, BEI Resources NR-54009) and hCoV-19/England/204820464/2020
(B.1.1.7, BEI Resources NR-54000) were grown in cells for 3 days, the supernatant was clarified by
centrifugation at 4,000 g for 5 minutes and aliquots were frozen at -80°C for long term use. The viruses
were subjected to deep sequencing to ensure that no mutations had taken place in culture. The polybasic
cleavage site changed to WRAR in the B.1.351 variant virus during cultivation in cell culture (as known for
this virus at BEI Resources) and no other unexpected mutations occurred. A primary virus isolate,
PV14252, bearing mutations and deletions in the spike was obtained by incubating 200 uls of viral
transport media from the nasopharyngeal swab with Vero.E6 cells. The sequence of the passage 2 viral
isolate was identical to the sequence obtained directly from the clinical specimen. Sequencing was
performed on the Illumina platform as described previously (Gonzalez-Reiche et al., 2020). Both
replication competent viruses were used to test serum from study participants and antibodies for
neutralization activity.
496

Neutralization assay. Twenty-thousand cells in 100 uls per well were seeded on sterile 96-well cell culture
plates one day prior to the neutralization assay. In general, cells were used at 90% confluency to perform
the assay. All serum samples were heat-inactivated to eliminate any complement activity. Serial dilutions
of serum samples were made in 1X minimal essential medium (MEM; Life Technologies) starting at a
dilution of 1:20. All work with authentic SARS-CoV-2 (isolate USA-WA1/2020 and PV14252) was done in a
biosafety level 3 (BSL3) laboratory following institutional biosafety guidelines and has been described in
much greater detail earlier (Amanat *et al.*, 2020b; Amanat et al., 2020c). Six hundred median cell culture
infectious doses (TCID$_{50}$s) of authentic virus (USA-WA1/2020 and PV14252) was added to each serum
dilution and virus-serum mixture was incubated together for 1 hour inside the biosafety cabinet. Media
from the cells was removed and 120 uls of the virus-serum mixture was added onto the cells for 1 hour at
37°C. After one hour, the virus-serum mixture was removed and 100 uls of each corresponding dilution
was added to every well. In addition, 100uls of 1X MEM was also added to every well. Cells were incubated
for 48 hours at 37°C after which the media was removed and 150 uls of 10% formaldehyde (Polysciences)
was added to inactivate the virus. For assay control, remdesivir was used against both the wild type virus
as well as the patient isolate. After 24 hours, cells were permeabilized and stained using an anti-
nucleoprotein antibody 1C7 as discussed in detail earlier (Amanat *et al.*, 2020b; Sun et al., 2020).

Cell sorting and flow cytometry. Staining for sorting was performed using cryo-preserved PBMCs in 2%
FBS and 2 mM ethylenediaminetetraacetic acid (EDTA) in PBS (P2). Cells were stained for 30 min on ice
with CD20-Pacific Blue (2H7, 1:400), Zombie Aqua, CD71-FITC (CY1G4, 1:200), IgD-PerCP-Cy5.5 (IA6-2,
1:200), CD19-PE (HIB19, 1:200), CD38-PE-Cy7 (HIT2, 1:200), and CD3-Alexa 700 (HIT3a, 1:200), all
BioLegend. Cells were washed twice, and single plasmablasts (live singlet CD19$^+$ CD3$^-$ IgD^{lo} CD38$^+$ CD20$^-$
CD71$^+$) were sorted using a FACSAria II into 96-well plates containing 2 µL Lysis Buffer (Clontech)
supplemented with 1 U/µL RNase inhibitor (NEB) and immediately frozen on dry ice, or bulk sorted into
PBS supplemented with 0.05% BSA and processed for single cell RNAseq.

Monoclonal antibody (mAb) generation. Antibodies were cloned as described previously (Wrammert et
al., 2011). Briefly, VH, Vκ, and Vλ genes were amplified by reverse transcription-PCR and nested PCR

reactions from singly sorted plasmablasts using primer combinations specific for IgG, IgM/A, Igκ, and Igλ
from previously described primer sets (Smith et al., 2009) and then sequenced. To generate recombinant
antibodies, restriction sites were incorporated via PCR with primers to the corresponding heavy and light
chain V and J genes. The amplified VH, Vκ, and Vλ genes were cloned into IgG1 and Igκ expression vectors,
respectively, as described previously (Nachbagauer et al., 2018; Wrammert et al., 2008). Heavy and light
chain plasmids were co-transfected into Expi293F cells (Gibco) for expression, and antibody was purified
with protein A agarose (Invitrogen).

Single-cell RNAseq library preparation and sequencing. Bulk-sorted plasmablasts were processed using
the following 10× Genomics kits: Chromium Next GEM Single Cell 5' Kit v2 (PN-1000263); Library
Construction Kit (PN-1000190); Chromium Next GEM Chip K Single Cell Kit (PN-1000286); Chromium Single
Cell Human BCR Amplification Kit (PN- 1000253), and Dual Index Kit TT Set A (PN-1000215). The cDNAs
were prepared after GEM generation and barcoding, followed by GEM RT reaction and bead cleanup
steps. Purified cDNA was amplified for 10–14 cycles before cleaning with SPRIselect beads. Then, samples
were evaluated on a 4200 TapeStation (Agilent) to determine cDNA concentration. B-cell receptor (BCR)
target enrichments were performed on full-length cDNA. Gene expression and enriched BCR libraries were
prepared as recommended by the Chromium Next GEM Single Cell 5' Reagent Kits v2 (Dual Index) user
guide, with appropriate modifications to the PCR cycles based on the calculated cDNA concentration. The
cDNA libraries were sequenced on Novaseq S4 (Illumina), targeting a median sequencing depth of 50,000
and 5,000 read pairs per cell for gene expression and BCR libraries, respectively.

Single cell RNAseq analysis. Single-cell RNA sequencing and BCR sequencing data was processed using
Cell Ranger v5.0 and the GRCh38-2020 version of the human genome provided by the manufacturer. Total
recovered cells by RNA sequencing were V3: 6,608, V5: 5,256, and V6: 6,325 with a mean of 90.64% read
mapped to the genome. Count matrices were processed in R (v4.0.2) using the Seurat (v3.2.2) R package
(Stuart et al., 2019). Cells were filtered for percentage of mitochondrial genes less than 15% and number
features less than 4,000. The three specimen sequencing runs were integrated using log-normalized count
values and canonical correlation approach (Stuart *et al.*, 2019) with 2,000 variable features. The resulting
single-cell object underwent principal component analysis and the top 30 principal components were used
for uniform manifold approximation and projection and identifying neighbors. Clustering was performed
using a resolution of 0.6. The integrated RNA sequencing object included 12,568 cells with V3: 4,584, V5:
3,523, and V6: 4,461 cells. The filtered contig annotation output of Cell Ranger vdj were loaded into R and
processed using the scRepertoire (v1.1.3) R package (Borcherding et al., 2020). Clonotypes were assigned
using igraph (v1.2.6) network analysis of components generated from CDR3 sequences greater than or
equal to 0.85 normalized Levenshtein distance. Percent of cells expressing genes along the UMAP
embedding was visualized using the schex (v1.3.0) R package. For mutation analysis, heavy chains of mAbs
and single-cell BCRs first underwent V(D)J gene annotation using IgBLAST (v1.14.0) (Ye et al., 2013) with
human reference (release 201931-4) from the international ImMunoGeneTics information system (IMGT)
(Giudicelli et al., 2005) and then parsing using Change-O (v0.4.6) (Gupta et al., 2015). Mutation frequency
was calculated, as described in (Turner *et al.*, 2020), using the "calcObservedMutations" function from
SHazaM (v.1.0.2) (Gupta *et al.*, 2015) and by counting the number of nucleotide mismatches from the
germline sequence in the heavy chain variable segment leading up to the complementary-determining
region 3 (CDR3), while excluding the first 18 positions that could be error-prone due to the primers used
for generating the mAb sequences.

Structure visualization and statistical analysis. Structural figures were modeled and rendered in Pymol (The PyMOL Molecular Graphics System, Version 2.4 Schrödinger, LLC). Statistical analysis was performed in GraphPad Prism using a one-way ANOVA with correction for multiple comparisons.

References

Amanat, F., Duehr, J., Huang, C., Paessler, S., Tan, G.S., and Krammer, F. (2020a). Monoclonal Antibodies with Neutralizing Activity and Fc-Effector Functions against the Machupo Virus Glycoprotein. J Virol *94*. 10.1128/JVI.01741-19.

Amanat, F., Duehr, J., Oestereich, L., Hastie, K.M., Ollmann Saphire, E., and Krammer, F. (2018). Antibodies to the Glycoprotein GP2 Subunit Cross-React between Old and New World Arenaviruses. mSphere *3*. 10.1128/mSphere.00189-18.

Amanat, F., Stadlbauer, D., Strohmeier, S., Nguyen, T.H.O., Chromikova, V., McMahon, M., Jiang, K., Arunkumar, G.A., Jurczyszak, D., Polanco, J., et al. (2020b). A serological assay to detect SARS-CoV-2 seroconversion in humans. Nat Med *26*, 1033-1036. 10.1038/s41591-020-0913-5.

Amanat, F., White, K.M., Miorin, L., Strohmeier, S., McMahon, M., Meade, P., Liu, W.C., Albrecht, R.A., Simon, V., Martinez-Sobrido, L., et al. (2020c). An In Vitro Microneutralization Assay for SARS-CoV-2 Serology and Drug Screening. Curr Protoc Microbiol *58*, e108. 10.1002/cpmc.108.

Annavajhala, M.K., Mohri, H., Zucker, J.E., Sheng, Z., Wang, P., Gomez-Simmonds, A., Ho, D.D., and Uhlemann, A.C. (2021). A Novel SARS-CoV-2 Variant of Concern, B.1.526, Identified in New York. medRxiv. 10.1101/2021.02.23.21252259.

Asthagiri Arunkumar, G., Ioannou, A., Wohlbold, T.J., Meade, P., Aslam, S., Amanat, F., Ayllon, J., García-Sastre, A., and Krammer, F. (2019). Broadly Cross-Reactive, Nonneutralizing Antibodies against Influenza B Virus Hemagglutinin Demonstrate Effector Function-Dependent Protection against Lethal Viral Challenge in Mice. J Virol *93*. 10.1128/JVI.01696-18.

Aydillo, T., Rombauts, A., Stadlbauer, D., Aslam, S., Abelenda-Alonso, G., Escalera, A., Amanat, F., Jiang, K., Krammer, F., Carratala, J., and García-Sastre, A. (2020). Antibody Immunological Imprinting on COVID-19 Patients. medRxiv, 2020.2010.2014.20212662. 10.1101/2020.10.14.20212662.

Baden, L.R., El Sahly, H.M., Essink, B., Kotloff, K., Frey, S., Novak, R., Diemert, D., Spector, S.A., Rouphael, N., Creech, C.B., et al. (2020). Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med. 10.1056/NEJMoa2035389.

Bailey, M.J., Broecker, F., Duehr, J., Arumemi, F., Krammer, F., Palese, P., and Tan, G.S. (2019). Antibodies Elicited by an NS1-Based Vaccine Protect Mice against Zika Virus. mBio *10*. 10.1128/mBio.02861-18.

Barnes, C.O., West, A.P., Huey-Tubman, K.E., Hoffmann, M.A.G., Sharaf, N.G., Hoffman, P.R., Koranda, N., Gristick, H.B., Gaebler, C., Muecksch, F., et al. (2020). Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal Common Epitopes and Recurrent Features of Antibodies. Cell *182*, 828-842.e816. 10.1016/j.cell.2020.06.025.

Bartsch, Y.C., Fischinger, S., Siddiqui, S.M., Chen, Z., Yu, J., Gebre, M., Atyeo, C., Gorman, M.J., Zhu, A.L., Kang, J., et al. (2021). Discrete SARS-CoV-2 antibody titers track with functional humoral stability. Nat Commun *12*, 1018. 10.1038/s41467-021-21336-8.

Baum, A., Fulton, B.O., Wloga, E., Copin, R., Pascal, K.E., Russo, V., Giordano, S., Lanza, K., Negron, N., Ni, M., et al. (2020). Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. Science *369*, 1014-1018. 10.1126/science.abd0831.

Borcherding, N., Bormann, N.L., and Kraus, G. (2020). scRepertoire: An R-based toolkit for single-cell immune receptor analysis. F1000Res *9*, 47. 10.12688/f1000research.22139.2.

Cao, Y., Su, B., Guo, X., Sun, W., Deng, Y., Bao, L., Zhu, Q., Zhang, X., Zheng, Y., Geng, C., et al. (2020).
Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-Throughput Single-Cell Sequencing
of Convalescent Patients' B Cells. Cell *182*, 73-84.e16. 10.1016/j.cell.2020.05.025.
Carvalho, T., Krammer, F., and Iwasaki, A. (2021). The first 12 months of COVID-19: a timeline of
immunological insights. Nat Rev Immunol *21*, 245-256. 10.1038/s41577-021-00522-1.
Cele, S., Gazy, I., Jackson, L., Hwa, S.-H., Tegally, H., Lustig, G., Giandhari, J., Pillay, S., Wilkinson, E.,
Naidoo, Y., et al. (2021). Escape of SARS-CoV-2 501Y.V2 variants from neutralization by convalescent
plasma. medRxiv, 2021.2001.2026.21250224. 10.1101/2021.01.26.21250224.
Chi, X., Yan, R., Zhang, J., Zhang, G., Zhang, Y., Hao, M., Zhang, Z., Fan, P., Dong, Y., Yang, Y., et al. (2020).
A neutralizing human antibody binds to the N-terminal domain of the Spike protein of SARS-CoV-2.
Science *369*, 650-655. 10.1126/science.abc6952.
Cho, H., Gonzales-Wartz, K.K., Huang, D., Yuan, M., Peterson, M., Liang, J., Beutler, N., Torres, J.L., Cong,
Y., Postnikova, E., et al. (2021). Ultrapotent bispecific antibodies neutralize emerging SARS-CoV-2
variants. bioRxiv, 2021.2004.2001.437942. 10.1101/2021.04.01.437942.
Choi, B., Choudhary, M.C., Regan, J., Sparks, J.A., Padera, R.F., Qiu, X., Solomon, I.H., Kuo, H.H., Boucau,
J., Bowman, K., et al. (2020). Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host.
N Engl J Med *383*, 2291-2293. 10.1056/NEJMc2031364.
Dagan, N., Barda, N., Kepten, E., Miron, O., Perchik, S., Katz, M.A., Hernán, M.A., Lipsitch, M., Reis, B.,
and Balicer, R.D. (2021). BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. N
Engl J Med. 10.1056/NEJMoa2101765.
Dan, J.M., Mateus, J., Kato, Y., Hastie, K.M., Yu, E.D., Faliti, C.E., Grifoni, A., Ramirez, S.I., Haupt, S.,
Frazier, A., et al. (2021). Immunological memory to SARS-CoV-2 assessed for up to 8 months after
infection. Science *371*. 10.1126/science.abf4063.
Davies, N.G., Abbott, S., Barnard, R.C., Jarvis, C.I., Kucharski, A.J., Munday, J.D., Pearson, C.A.B., Russell,
T.W., Tully, D.C., Washburne, A.D., et al. (2021). Estimated transmissibility and impact of SARS-CoV-2
lineage B.1.1.7 in England. Science, eabg3055. 10.1126/science.abg3055.
Dilillo, D.J., Tan, G.S., Palese, P., and Ravetch, J.V. (2014). Broadly neutralizing hemagglutinin stalk-
specific antibodies require FcγR interactions for protection against influenza virus in vivo. Nat Med.
10.1038/nm.3443.
Emary, K.R.W., Golubchik, T., Aley, P.K., Ariani, C.V., Angus, B., Bibi, S., Blane, B., Bonsall, D., Cicconi, P.,
Charlton, S., et al. (2021). Efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine against SARS-CoV-2 variant of
concern 202012/01 (B.1.1.7): an exploratory analysis of a randomised controlled trial. Lancet *397*, 1351-
1362. 10.1016/S0140-6736(21)00628-0.
Faria, N.R., Claro, I.M., Candido, D., Franco, L.A.M., Andrade, P.S., Coletti, T.M., Silva, C.A.M., Sales, F.C.,
Manula, E.R., Aguiar, R.S., et al. (2021). <h1 data-topic-id="586" style="font-size: 1.7511em; margin: 0px
0px 4px; font-family: Helvetica, Arial, sans-serif; width: 999px; line-height: 1.2; overflow-wrap: break-
word; color: rgb(34, 34, 34); background-color: rgb(255, 255, 255);">Genomic characterisation of an
emergent SARS-CoV-2 lineage in Manaus: preliminary findings.
Gaebler, C., Wang, Z., Lorenzi, J.C.C., Muecksch, F., Finkin, S., Tokuyama, M., Ladinsky, M., Cho, A.,
Jankovic, M., Schaefer-Babajew, D., et al. (2020). Evolution of Antibody Immunity to SARS-CoV-2.
bioRxiv. 10.1101/2020.11.03.367391.
Giudicelli, V., Chaume, D., and Lefranc, M.P. (2005). IMGT/GENE-DB: a comprehensive database for
human and mouse immunoglobulin and T cell receptor genes. Nucleic Acids Res *33*, D256-261.
10.1093/nar/gki010.
Gonzalez-Reiche, A.S., Hernandez, M.M., Sullivan, M.J., Ciferri, B., Alshammary, H., Obla, A., Fabre, S.,
Kleiner, G., Polanco, J., Khan, Z., et al. (2020). Introductions and early spread of SARS-CoV-2 in the New
York City area. Science. 10.1126/science.abc1917.

Gorman, M.J., Patel, N., Guebre-Xabier, M., Zhu, A., Atyeo, C., Pullen, K.M., Loos, C., Goez-Gazi, Y.,
Carrion, R., Tian, J.-H., et al. (2021). Collaboration between the Fab and Fc contribute to maximal
protection against SARS-CoV-2 in nonhuman primates following NVX-CoV2373 subunit vaccine with
Matrix-M™ vaccination. bioRxiv, 2021.2002.2005.429759. 10.1101/2021.02.05.429759.
Grandjean, L., Saso, A., Torres, A., Lam, T., Hatcher, J., Thistlethwayte, R., Harris, M., Best, T., Johnson,
M., Wagstaffe, H., et al. (2020). Humoral Response Dynamics Following Infection with SARS-CoV-2.
medRxiv, 2020.2007.2016.20155663. 10.1101/2020.07.16.20155663.
Greaney, A.J., Starr, T.N., Gilchuk, P., Zost, S.J., Binshtein, E., Loes, A.N., Hilton, S.K., Huddleston, J.,
Eguia, R., Crawford, K.H.D., et al. (2021). Complete Mapping of Mutations to the SARS-CoV-2 Spike
Receptor-Binding Domain that Escape Antibody Recognition. Cell Host Microbe *29*, 44-57.e49.
10.1016/j.chom.2020.11.007.
Gupta, N.T., Vander Heiden, J.A., Uduman, M., Gadala-Maria, D., Yaari, G., and Kleinstein, S.H. (2015).
Change-O: a toolkit for analyzing large-scale B cell immunoglobulin repertoire sequencing data.
Bioinformatics *31*, 3356-3358. 10.1093/bioinformatics/btv359.
Guthmiller, J.J., Stovicek, O., Wang, J., Changrob, S., Li, L., Halfmann, P., Zheng, N.Y., Utset, H., Stamper,
C.T., Dugan, H.L., et al. (2021). SARS-CoV-2 Infection Severity Is Linked to Superior Humoral Immunity
against the Spike. mBio *12*. 10.1128/mBio.02940-20.
Hensley, S.E., Das, S.R., Bailey, A.L., Schmidt, L.M., Hickman, H.D., Jayaraman, A., Viswanathan, K.,
Raman, R., Sasisekharan, R., Bennink, J.R., and Yewdell, J.W. (2009). Hemagglutinin receptor binding
avidity drives influenza A virus antigenic drift. Science *326*, 734-736. 10.1126/science.1178258.
Huang, K.A., Tan, T.K., Chen, T.H., Huang, C.G., Harvey, R., Hussain, S., Chen, C.P., Harding, A., Gilbert-
Jaramillo, J., Liu, X., et al. (2021). Breadth and function of antibody response to acute SARS-CoV-2
infection in humans. PLoS Pathog *17*, e1009352. 10.1371/journal.ppat.1009352.
Isho, B., Abe, K.T., Zuo, M., Jamal, A.J., Rathod, B., Wang, J.H., Li, Z., Chao, G., Rojas, O.L., Bang, Y.M., et
al. (2020). Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19
patients. Sci Immunol *5*. 10.1126/sciimmunol.abe5511.
Iyer, A.S., Jones, F.K., Nodoushani, A., Kelly, M., Becker, M., Slater, D., Mills, R., Teng, E., Kamruzzaman,
M., Garcia-Beltran, W.F., et al. (2020). Dynamics and significance of the antibody response to SARS-CoV-
2 infection. medRxiv. 10.1101/2020.07.18.20155374.
Krammer, F. (2020). SARS-CoV-2 vaccines in development. Nature *586*, 516-527. 10.1038/s41586-020-
2798-3.
Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., Zhang, Q., Shi, X., Wang, Q., Zhang, L., and Wang, X.
(2020). Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. Nature
581, 215-220. 10.1038/s41586-020-2180-5.
Larsen, H.D., Fonager, J., Lomholt, F.K., Dalby, T., Benedetti, G., Kristensen, B., Urth, T.R., Rasmussen, M.,
Lassaunière, R., Rasmussen, T.B., et al. (2021). Preliminary report of an outbreak of SARS-CoV-2 in mink
and mink farmers associated with community spread, Denmark, June to November 2020. Euro Surveill
26. 10.2807/1560-7917.ES.2021.26.5.210009.
Lasek-Nesselquist, E., Lapierre, P., Schneider, E., George, K.S., and Pata, J. (2021). The localized rise of a
B.1.526 SARS-CoV-2 variant containing an E484K mutation in New York State. medRxiv,
2021.2002.2026.21251868. 10.1101/2021.02.26.21251868.
Letko, M., Marzi, A., and Munster, V. (2020). Functional assessment of cell entry and receptor usage for
SARS-CoV-2 and other lineage B betacoronaviruses. Nat Microbiol *5*, 562-569. 10.1038/s41564-020-
0688-y.
Liu, L., Wang, P., Nair, M.S., Yu, J., Rapp, M., Wang, Q., Luo, Y., Chan, J.F., Sahi, V., Figueroa, A., et al.
(2020). Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike. Nature *584*, 450-
456. 10.1038/s41586-020-2571-7.

Margine, I., Palese, P., and Krammer, F. (2013). Expression of functional recombinant hemagglutinin and
neuraminidase proteins from the novel H7N9 influenza virus using the baculovirus expression system. J
Vis Exp, e51112. 10.3791/51112.
McCallum, M., De Marco, A., Lempp, F.A., Tortorici, M.A., Pinto, D., Walls, A.C., Beltramello, M., Chen,
A., Liu, Z., Zatta, F., et al. (2021a). N-terminal domain antigenic mapping reveals a site of vulnerability for
SARS-CoV-2. Cell. 10.1016/j.cell.2021.03.028.
McCallum, M., Marco, A., Lempp, F., Tortorici, M.A., Pinto, D., Walls, A.C., Beltramello, M., Chen, A., Liu,
Z., Zatta, F., et al. (2021b). N-terminal domain antigenic mapping reveals a site of vulnerability for SARS-
CoV-2. bioRxiv. 10.1101/2021.01.14.426475.
Nachbagauer, R., Shore, D., Yang, H., Johnson, S.K., Gabbard, J.D., Tompkins, S.M., Wrammert, J.,
Wilson, P.C., Stevens, J., Ahmed, R., et al. (2018). Broadly Reactive Human Monoclonal Antibodies
Elicited following Pandemic H1N1 Influenza Virus Exposure Protect Mice against Highly Pathogenic H5N1
Challenge. J Virol *92*. 10.1128/JVI.00949-18.
O'Donnell, C.D., Vogel, L., Wright, A., Das, S.R., Wrammert, J., Li, G.M., McCausland, M., Zheng, N.Y.,
Yewdell, J.W., Ahmed, R., et al. (2012). Antibody pressure by a human monoclonal antibody targeting
the 2009 pandemic H1N1 virus hemagglutinin drives the emergence of a virus with increased virulence
in mice. mBio *3*. 10.1128/mBio.00120-12.
Pallesen, J., Wang, N., Corbett, K.S., Wrapp, D., Kirchdoerfer, R.N., Turner, H.L., Cottrell, C.A., Becker,
M.M., Wang, L., Shi, W., et al. (2017). Immunogenicity and structures of a rationally designed prefusion
MERS-CoV spike antigen. Proc Natl Acad Sci U S A *114*, E7348-E7357. 10.1073/pnas.1707304114.
PHE (2021). Investigation of novel SARS-CoV-2 variant Variant of Concern 202012/01 Technical briefing
5.
Polack, F.P., Thomas, S.J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L., Pérez Marc, G.,
Moreira, E.D., Zerbini, C., et al. (2020). Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. N
Engl J Med. 10.1056/NEJMoa2034577.
Ripperger, T.J., Uhrlaub, J.L., Watanabe, M., Wong, R., Castaneda, Y., Pizzato, H.A., Thompson, M.R.,
Bradshaw, C., Weinkauf, C.C., Bime, C., et al. (2020). Orthogonal SARS-CoV-2 Serological Assays Enable
Surveillance of Low-Prevalence Communities and Reveal Durable Humoral Immunity. Immunity *53*, 925-
933.e924. 10.1016/j.immuni.2020.10.004.
Robbiani, D.F., Gaebler, C., Muecksch, F., Lorenzi, J.C.C., Wang, Z., Cho, A., Agudelo, M., Barnes, C.O.,
Gazumyan, A., Finkin, S., et al. (2020). Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. Nature *584*, 437-442. 10.1038/s41586-020-2456-9.
Rodda, L.B., Netland, J., Shehata, L., Pruner, K.B., Morawski, P.A., Thouvenel, C.D., Takehara, K.K.,
Eggenberger, J., Hemann, E.A., Waterman, H.R., et al. (2021). Functional SARS-CoV-2-Specific Immune
Memory Persists after Mild COVID-19. Cell *184*, 169-183.e117. 10.1016/j.cell.2020.11.029.
Saphire, E.O., Schendel, S.L., Fusco, M.L., Gangavarapu, K., Gunn, B.M., Wec, A.Z., Halfmann, P.J.,
Brannan, J.M., Herbert, A.S., Qiu, X., et al. (2018). Systematic Analysis of Monoclonal Antibodies against
Ebola Virus GP Defines Features that Contribute to Protection. Cell *174*, 938-952.e913.
10.1016/j.cell.2018.07.033.
Schäfer, A., Muecksch, F., Lorenzi, J.C.C., Leist, S.R., Cipolla, M., Bournazos, S., Schmidt, F., Maison, R.M.,
Gazumyan, A., Martinez, D.R., et al. (2021). Antibody potency, effector function, and combinations in
protection and therapy for SARS-CoV-2 infection in vivo. J Exp Med *218*. 10.1084/jem.20201993.
Seow, J., Graham, C., Merrick, B., Acors, S., Pickering, S., Steel, K.J.A., Hemmings, O., O'Byrne, A.,
Kouphou, N., Galao, R.P., et al. (2020). Longitudinal observation and decline of neutralizing antibody
responses in the three months following SARS-CoV-2 infection in humans. Nat Microbiol *5*, 1598-1607.
10.1038/s41564-020-00813-8.

Shinde, V., Bhikha, S., Hossain, Z., Archary, M., Bhorat, Q., Fairlie, L., Lalloo, U., Masilela, M.L.S.,
Moodley, D., Hanley, S., et al. (2021). Preliminary Efficacy of the NVX-CoV2373 Covid-19 Vaccine Against
the B.1.351 Variant. medRxiv, 2021.2002.2025.21252477. 10.1101/2021.02.25.21252477.
Smith, K., Garman, L., Wrammert, J., Zheng, N.Y., Capra, J.D., Ahmed, R., and Wilson, P.C. (2009). Rapid
generation of fully human monoclonal antibodies specific to a vaccinating antigen. Nat Protoc *4*, 372-
384. 10.1038/nprot.2009.3.
Song, G., He, W.T., Callaghan, S., Anzanello, F., Huang, D., Ricketts, J., Torres, J.L., Beutler, N., Peng, L.,
Vargas, S., et al. (2020). Cross-reactive serum and memory B cell responses to spike protein in SARS-CoV-
2 and endemic coronavirus infection. bioRxiv. 10.1101/2020.09.22.308965.
Stadlbauer, D., Amanat, F., Chromikova, V., Jiang, K., Strohmeier, S., Arunkumar, G.A., Tan, J., Bhavsar,
D., Capuano, C., Kirkpatrick, E., et al. (2020). SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol
for a Serological Assay, Antigen Production, and Test Setup. Curr Protoc Microbiol *57*, e100.
10.1002/cpmc.100.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M.,
Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-
1902.e1821. 10.1016/j.cell.2019.05.031.
Sun, W., Leist, S.R., McCroskery, S., Liu, Y., Slamanig, S., Oliva, J., Amanat, F., Schafer, A., Dinnon, K.H.,
3rd, Garcia-Sastre, A., et al. (2020). Newcastle disease virus (NDV) expressing the spike protein of SARS-
CoV-2 as a live virus vaccine candidate. EBioMedicine *62*, 103132. 10.1016/j.ebiom.2020.103132.
Tegally, H., Wilkinson, E., Giovanetti, M., Iranzadeh, A., Fonseca, V., Giandhari, J., Doolabh, D., Pillay, S.,
San, E.J., Msomi, N., et al. (2020). Emergence and rapid spread of a new severe acute respiratory
syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa.
medRxiv, 2020.2012.2021.20248640. 10.1101/2020.12.21.20248640.
Thomson, E.C., Rosen, L.E., Shepherd, J.G., Spreafico, R., da Silva Filipe, A., Wojcechowskyj, J.A., Davis,
C., Piccoli, L., Pascall, D.J., Dillen, J., et al. (2021). Circulating SARS-CoV-2 spike N439K variants maintain
fitness while evading antibody-mediated immunity. Cell. 10.1016/j.cell.2021.01.037.
Turner, J.S., Zhou, J.Q., Han, J., Schmitz, A.J., Rizk, A.A., Alsoussi, W.B., Lei, T., Amor, M., McIntire, K.M.,
Meade, P., et al. (2020). Human germinal centres engage memory and naive B cells after influenza
vaccination. Nature *586*, 127-132. 10.1038/s41586-020-2711-0.
Wajnberg, A., Amanat, F., Firpo, A., Altman, D.R., Bailey, M.J., Mansour, M., McMahon, M., Meade, P.,
Mendu, D.R., Muellers, K., et al. (2020). Robust neutralizing antibodies to SARS-CoV-2 infection persist
for months. Science *370*, 1227-1230. 10.1126/science.abd7728.
Walsh, E.E., Frenck, R., Falsey, A.R., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Neuzil, K., Mulligan,
M.J., Bailey, R., et al. (2020). RNA-Based COVID-19 Vaccine BNT162b2 Selected for a Pivotal Efficacy
Study. medRxiv, 2020.2008.2017.20176651. 10.1101/2020.08.17.20176651.
Wang, Z., Schmidt, F., Weisblum, Y., Muecksch, F., Barnes, C.O., Finkin, S., Schaefer-Babajew, D., Cipolla,
M., Gaebler, C., Lieberman, J.A., et al. (2021). mRNA vaccine-elicited antibodies to SARS-CoV-2 and
circulating variants. Nature. 10.1038/s41586-021-03324-6.
Weisblum, Y., Schmidt, F., Zhang, F., DaSilva, J., Poston, D., Lorenzi, J.C., Muecksch, F., Rutkowska, M.,
Hoffmann, H.H., Michailidis, E., et al. (2020). Escape from neutralizing antibodies by SARS-CoV-2 spike
protein variants. Elife *9*. 10.7554/eLife.61312.
Wibmer, C.K., Ayres, F., Hermanus, T., Madzivhandila, M., Kgagudi, P., Lambson, B.E., Vermeulen, M.,
van den Berg, K., Rossouw, T., Boswell, M., et al. (2021). SARS-CoV-2 501Y.V2 escapes neutralization by
South African COVID-19 donor plasma. bioRxiv, 2021.2001.2018.427166. 10.1101/2021.01.18.427166.
Wilson, P., Stamper, C., Dugan, H., Li, L., Asby, N., Halfmann, P., Guthmiller, J., Zheng, N.Y., Huang, M.,
Stovicek, O., et al. (2020). Distinct B cell subsets give rise to antigen-specific antibody responses against
SARS-CoV-2. Res Sq. 10.21203/rs.3.rs-80476/v1.

Wohlbold, T.J., Chromikova, V., Tan, G.S., Meade, P., Amanat, F., Comella, P., Hirsh, A., and Krammer, F.
(2016). Hemagglutinin Stalk- and Neuraminidase-Specific Monoclonal Antibodies Protect against Lethal
H10N8 Influenza Virus Infection in Mice. J Virol *90*, 851-861. 10.1128/JVI.02275-15.
Wohlbold, T.J., Nachbagauer, R., Xu, H., Tan, G.S., Hirsh, A., Brokstad, K.A., Cox, R.J., Palese, P., and
Krammer, F. (2015). Vaccination with adjuvanted recombinant neuraminidase induces broad
heterologous, but not heterosubtypic, cross-protection against influenza virus infection in mice. mBio *6*,
e02556. 10.1128/mBio.02556-14.
Wrammert, J., Koutsonanos, D., Li, G.M., Edupuganti, S., Sui, J., Morrissey, M., McCausland, M.,
Skountzou, I., Hornig, M., Lipkin, W.I., et al. (2011). Broadly cross-reactive antibodies dominate the
human B cell response against 2009 pandemic H1N1 influenza virus infection. J Exp Med *208*, 181-193.
jem.20101352 [pii]

10.1084/jem.20101352.

Wrammert, J., Smith, K., Miller, J., Langley, W.A., Kokko, K., Larsen, C., Zheng, N.Y., Mays, I., Garman, L.,
Helms, C., et al. (2008). Rapid cloning of high-affinity human monoclonal antibodies against influenza
virus. Nature *453,* 667-671. 10.1038/nature06890.
Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.L., Abiona, O., Graham, B.S., and McLellan,
J.S. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science.
10.1126/science.abb2507.
Wu, K., Werner, A.P., Moliva, J.I., Koch, M., Choi, A., Stewart-Jones, G.B.E., Bennett, H., Boyoglu-Barnum,
S., Shi, W., Graham, B.S., et al. (2021). mRNA-1273 vaccine induces neutralizing antibodies against spike
mutants from global SARS-CoV-2 variants. bioRxiv, 2021.2001.2025.427948.
10.1101/2021.01.25.427948.
Ye, J., Ma, N., Madden, T.L., and Ostell, J.M. (2013). IgBLAST: an immunoglobulin variable domain
sequence analysis tool. Nucleic Acids Res 4*1,* W34-40. 10.1093/nar/gkt382.
822

Figure Legends

Figure 1: Antibody responses in individuals vaccinated with mRNA-based SARS-CoV-2 vaccines.
Antibody responses of convalescent individuals and vaccinees to full length spike protein (**A**) and RBD (**B**)
as measured by ELISA and neutralizing activity of the sera of the same individuals in a microneutralization
assay against authentic SARS-CoV-2 (**C**). Convalescent individuals were grouped based on their initial
antibody response (measured in a CLIA laboratory) to spike protein into +, ++, and +++. **D** shows ratios
between binding and neutralizing antibody levels in vaccinees and convalescent individuals. Higher ratios
indicate a bias towards non-neutralizing antibodies. **E, F, G and H** show antibody responses against α-
coronavirus 229E and NL63 and β-coronavirus OC43 and HKU1 spike proteins over time.

Figure 2. Characterization of mAbs derived from vaccine plasmablasts. Binding of plasmablasts derived
from three vaccinees (V3, V5 and V6) against full length spike (**A**), RBD (**B**), NTD (**C**) and S2 (**D**). **E** shows
the percentages of the respective antibodies per subject. **F** shows neutralizing activity of the mABs against
authentic SARS-CoV-2 and the proportion of neutralizing antibodies per subject is shown in **G**. **H and I**
show reactivity of mAbs to spike protein of human β-coronaviruses OC43 and HKU1. MBC = minimal
binding concentration.

Figure 3. Characterization of bulk sorted plasmablasts via single-cell RNA sequencing. (A) Uniform manifold approximation and projection (UMAP) of scRNAseq from bulk plasmablast with recovered BCR sequences (purple) or unrecovered (grey). **(B)** UMAP overlay of percent of cellular population expressing *MZB1*, *PRDM1*, and *XPB1*. Hexbin equals 80 individual cells. (**C**) UMAP overlay of BCR sequences with confirmed spike binding activity. (**D**) Proportional composition of heavy chains genes in the spike binding sequences broken down by sample. **(E)** Comparison of nucleotide-level mutation frequency in immunoglobulin heavy chain variable (IGHV) genes between plasmablasts clonally related to spike binding mAbs from SARS-CoV-2 vaccinees, plasmablasts sorted from PBMCs one week after seasonal influenza vaccination and found in vaccine-responding B cell clones, and naïve B cells found in blood of an influenza vaccinee (left panel); and between plasmablasts from SARS-CoV-2 vaccinees found to be clonally related to spike-binding mAbs that were, respectively, cross-reactive and non-cross-reactive to human β-coronaviruses spike proteins (right panel).

Figure 4. Mapping of the amino-acid substitutions and deletions onto the structure of the SARS-CoV-2 spike glycoprotein. A lists mutations of the three major variants of concern B.1.17, B.1.315 and P.1. **B** shows these mutations mapped onto the structure of the spike glycoprotein (model generated by superposition of PDB 6M0j and 7C2L (Chi *et al.*, 2020; Lan et al., 2020)). One RBD in the up conformation (red) is bound with ACE2 receptor (pink). The NTD is colored blue and the various amino-acid substitutions are shown as yellow spheres. One spike protomer is shown in bold colors while the other two are colored white. **C** shows competition between ACE2 and neutralizing RBD targeting mAbs PVI.V3-9 and PVI.V6-4 for binding to RBD. **D** BLI- measured binding affinities of the RBD mutants to ACE2, as well as the calculated fold change, are shown in the table on the right.

Figure 5. Binding and neutralization of SARS-CoV-2 variants. Binding of serum samples from convalescent individuals, vaccinees and vaccine derived mAbs to a panel of RBD mutants is shown in **A, B and C** respectively. The red line in A indicates the average reduction. Dotted lines in A and B indicate 100%, the line with smaller dots in C indicated reactivity of the anti-his coating control. For vaccinees late samples (V1=d89, V2=d102, V3=d47, V4=d48, V5=49 and V6=48) were assayed. **D** shows the spike mutations of virus isolate PVI14252 modelled on a co-crystal structure of the SARS-CoV-2 spike protein with ACE2 (model generated by superposition of PDB 6M0j and 7C2L (Chi *et al.*, 2020; Lan *et al.*, 2020)). **E and F** show the inhibitory effect of vaccine serum and vaccine derived neutralizing antibodies on both wild type SARS-CoV-2 and PV14252. **G** shows neutralizing activity of the plasmablast derived neutralizing antibodies aginst wild type, B.1.1.7 and B.1.351 virus isolates. Of note, these comparative assays were always performed side by side but sets are run by different operators and on a different Vero cell clone as the neutralization assays shown in Figure 2.

Supplemental Table 1: Study participant and biospecimen information

Vaccinees	Spike IgG response[1]*	Sex	Age group (yrs)	Specimen tested

V1	Strong positive	F	>60	several longitudinal time points including days 34, 68, 89 and 188 post vaccination
V2	Strong positive	M	30-40	several longitudinal time points including days 19, 47, 74, 102, 129, 157 and 186 post vaccination
V3	Strong positive	F	50-60	several longitudinal time points including days 19, 27, 47, 77 and 102 post vaccination
V4	Strong positive	M	>60	several longitudinal time points including days 27, 48 and 89 post vaccination
V5	Strong positive	F	40-50	several longitudinal time points including 27, 49 and 83 days post vaccination
V6	Strong positive	F	30-40	several longitudinal time points including 27, 48 and 83 days post vaccination
Seronegative, post pandemic		**Sex**	**Age group (yrs)**	**Days from last negative serology test when the sample was taken**
N1	Negative	F	40-50	23
N2	Negative	F	20-29	24
N3	Negative	F	20-29	23
N4	Negative	F	30-35	22
Seropositive, natural infection		**Sex**	**Age group (yrs)**	**Days post onset of COVID-19 symptoms when the sample was taken**
P1	Weak positive	M	20-29	260
P2	Weak positive	M	50-59	no data available
P3	Weak positive	F	30-39	111
P4	Weak positive	F	30-39	221

P5	Weak positive	F	30-39	254
P6	Weak positive	F	20-29	247
P7	Weak positive	M	30-39	220
P8	Weak positive	F	20-29	Asymptomatic
P9	Moderate positive	M	30-39	no data available
P10	Moderate positive	F	30-39	197
P11	Moderate positive	F	50-59	Asymptomatic
P12	Moderate positive	F	30-39	Asymptomatic
P13	Moderate positive	M	30-39	234
P14	Moderate positive	F	20-29	273
P15	Moderate positive	M	30-39	Asymptomatic
P16	Moderate positive	F	20-29	258
P17	Moderate positive	F	20-29	246
P18	Moderate positive	M	20-29	Asymptomatic
P19	Moderate positive	F	50-59	204
P20	Strong positive	F	50-59	no data available
P21	Strong positive	F	30-39	245
P22	Strong positive	M	NA	170
P23	Strong positive	F	>60	Asymptomatic
P24	Strong positive	F	40-49	no data available
P25	Strong positive	F	50-59	191
P26	Strong positive	F	30-39	no data available
P27	Strong positive	F	50-59	113
P28	Strong positive	M	>60	Asymptomatic
P29	Strong positive	M	18-19	218
P30	Strong positive	M	50-59	219

[1]Samples were categorized based on initial titers obtained from Mount Sinai's CLIA laboratory test. Weak positive: 1:80 – 1:160 weak positive; 1:320-1:960 moderate positive; 1:960-1: ≥2880 strong positive

* All six vaccinees were sero-negative for SARS-CoV-2 without clinical evidence of COVID10 prior to SARS-CoV-2 spike mRNA vaccination.

Supplemental Table 2: Immunoglobulin gene usage of the spike-mAbs

Name	Native isotype	Heavy chain		Light chain	
		Gene usage	HCDR3 AA sequence	Gene usage	LCDR3 AA sequence
PVI.V5-1	IgG1	VH3-23 DH5-18 JH4	CAPHRGQLWFDYW	VK3-20 JK4	CQQYGSSPPTF
PVI.V5-2	IgG1	VH3-21 DH2-2 JH4	CARDLKLSPAAIGWDYFDYW	VK3-15 JK2	CQQYNNWPRSF
PVI.V5-3	IgG1	VH3-7 DH6-13 JH4	CAIFGAAGTDYW	VL3-16 JL3	CLSADSSGTYWVF
PVI.V5-4	IgG1	VH3-30 DH3-22 JH4	CARENNYYDSSGYSYYFDYW	VK3-20 JK2	CQQYGSSPMCSF
PVI.V5-5	IgA1	VH1-69 DH5-24 JH4	CARDFGREWLQYFYFDCW	VK3-20 JK4	CQQYGSSPTF

PVI.V5-6	IgG1	VH3-7 DH3-3 JH4	CARDNNDFWSGYLYFDYW	VL3-10 JL2	CYSTDSSGNHRGVF
PVI.V6-1	IgG1	VH3-30 DH6-19 JH4	CARGAVAGQHSFDNW	VK2-30 JK4	CMQGTHWPPTF
PVI.V6-2	IgG1	VH3-33 DH6-13 JH4	CARDKRGSSSWLDQYFDYW	VL3-21 JL2	CQVWDSSSDHVVF
PVI.V6-3	IgG1	VH4-31 DH3-22 JH5	CARDMISGRGLFDPW	VK1-33 JK2	CQQYDNLPTF
PVI.V6-4	IgG1	VH1-69 DH4-17 JH3	CARGNYDYGDYLLKGSAFDIW	VK4-1 JK2	CQQYYSTPPNTF
PVI.V6-5	IgG1	VH3-30 DH3-10 JH4	CAKDGGYYYYGSGSYPSYFDYW	VK2D-29 JK4	CMQSIQLPLTF
PVI.V6-6	IgG1	VH4-31 DH3-10 JH6	CASEKFLWGQGYYYGMDVW	VL2-14 JL2	CSSYTSSSTLVF
PVI.V6-7	IgG1	VH4-39 DH3-22 JH4	CATQSDYDSSGLEFDYW	VL2-14 JL3	CSSYTSSSSWVF
PVI.V6-8	IgG1	VH4-31 DH3-22 JH3	CARGREEPIVVVTDAFDIW	VK3-11 JK2	CQQRSNWPPMYTF
PVI.V6-9	IgG1	VH3-30 DH2-15 JH4	CAKSGYPYCGGGTCYSGWFDYW	VK1-33 JK2	CQRYDNPPYTF
PVI.V6-10	IgG1	VH1-2 DH6-19 JH6	CAREIAVAGNDYSYGLDVW	VK3-20 JK4	CQQYGSSLLTF
PVI.V6-11	IgG1	VH1-46 DH6-19 JH6	CASQSHWQWLGGGDSYGMDVW	VK1-9 JK2	CQQLNSYPYTF
PVI.V6-12	IgG1	VH5-51 DH1-26 JH4	CARRFGSYPPYFDYW	VL3-21 JL1	CQVWDSNSDLYVF
PVI.V6-13	IgG1	VH3-30 DH5-18 JH4	CAKAGYSYAYGDYYFDYW	VK1-33 JK3	CQHYDNLPPAVTF
PVI.V6-14	IgG1	VH4-39 DH3-10 JH4	CARCRPEYYFGSGSYLDFDYW	VK1-12 JK4	CQQANSFPLTF
PVI.V6-15	IgG1	VH3-30 DH5-18 JH4	CAKDWGWIQLWGLDYW	VL2-18 JL3	CSSYTSSSTWVF
PVI.V3-1	IgG1	VH4-4 DH1-26 JH4	CVSRGVGATREKDYW	VK3-15 JK4	CQQYNNWPPDLTF
PVI.V3-2	IgG1	VH4-39 DH3-10 JH4	CASLDYYGSGSGPGYFDYW	VK3-11 JK4	CQQRSNWLTF
PVI.V3-3	IgG3	VH3-33 DH6-19 JH4	CASDSSGWYYFDYW	VL3-9 JL2	CQVWDSSTVVF
PVI.V3-4	IgG1	VH3-21 DH2-21 JH3	CAVTLLPTYCGGEWCAFDIW	VK3-15 JK2	CQQYNNWPPYTF
PVI.V3-5	IgG1	VH1-69 DH3-22 JH4	CARNYYDSSGSQGMDYW	VK3-11 JK4	CQQRSNWPPVLTF
PVI.V3-6	IgA1	VH3-66 DH3-3 JH4	CARHLGVVI[1]	VK4-1 JK1	CQQYYSTLWTF
PVI.V3-7	IgG3	VH4-39 DH1-26 JH4	CAKPSGSYLGFDYW	VK1-39 JK3	CQQSYSTPFTF
PVI.V3-8	IgG1	VH4-38 DH3-3 JH4	CARSDFSVRVGFDCW	VK4-1 JK2	CQQSYTTNTF
PVI.V3-9	IgG1	VH3-53 DH3-16 JH6	CARDLMEGGGMDVW	VK3-20 JK1	CQQYGSSLGTF
PVI.V3-10	IgG1	VH4-34 DH3-22 JH4	CARSQPLLWSSGYCCDYW	VL2-11 JL2	CCSYAGSYTLVF
PVI.V3-11	IgG1	VH3-64D DH4-11 JH4	CVRGPTVTTENDFDSW	VK2-30 JK2	CMQGTHSYTF
PVI.V3-12	IgG2	VH1-46 DH5-24 JH4	CASDPNRDGLALDSW	VK3-20 JK2	CQQYGTSPLYTF
PVI.V3-13	IgG1	VH1-69 DH1-14 JH4	CARDRYHGSPVDYW	VK3-11 JK4	CQQRSNWPPSLTF
PVI.V3-14	IgG1	VH4-31 DH5-12 JH4	CARARYSGSARGPPKQYYFDYW	VK3-20 JK1	CQHLVTF
PVI.V3-15	IgG1	VH3-21 DH3-3 JH3	CARDGGRGYDFWSGYYIGAFDIW	VK3-15 JK4	CQQYNDWPPLTF
PVI.V3-16	IgG1	VH3-33 DH2-15 JH6	CARGLGWDIVVVVSGEMDGMDVW	VL1-40 JL1	CQSYDSSLSGPYVF
PVI.V3-17	IgA1	VH3-20 DH2-2 JH4	CARGEGSSDYW	VK3-20 JK1	CQQYGSSPKTF
PVI.V3-18	IgG1	VH3-30 DH2-15 JH4	CAKASGLYCSGGNCLVADFDYW	VL1-39 JL4	CQQSYSTPLSF
PVI.V3-19	IgG1	VH5-51 DH6-19 JH6	CARRNTSAQYSSGWYVHYYYGMDVW	VK2-28 JK3	CMQALQTPGFTF
PVI.V3-20	IgG1	VH3-30 DH3-3 JH6	CAKDQLGAIFAHYYYYGMDVW	VL1-40 JL1	CQSYDSSLSGYVF
PVI.V3-21	IgG1	VH5-51 DH5-24 JH4	CAKLSRDAYRGPFDYW	VL6-57 JL2	CQSYDSSNPDVVF

[1]Indicates absence of the W118 residue from the junction of the CDR3.

Supplementary Figure 1. **Full length spike to RBD ratios (A) and comparison of binding to neutralizing titer ratios between naturally infected and vaccinated individuals (B).**

Supplementary Figure 2. **Gating strategy for sorting plasmablasts from total PBMCs isolated one week after second immunization.**

Supplementary Figure 3. Representative Biolayer Interferometry binding isotherms from two independent experiments. The raw data are show in pink and the Langmuir 1:1 kinetics fit is show in black.

896 **Supplementary Figure 4. Binding of SARS-CoV-2 variant RBDs to ACE2. A.** ELISA curves of the RBD variants
897 binding to human ACE2. Shown are the binding curves calculated with nonlinear regression to the
898 arithmetic mean values from eight replicates ± SEM. The calculated steady-state K_D values ± SEM from
899 end-point ELISA measurements and the fold-change in comparison to wild type RBD are reported in **B.**

A

Recovered Productive BCR ● Yes ● No

n = 12,556

UMAP-2
UMAP-1

B

MZB1   *PRDM1*   *XBP1*

UMAP-2
UMAP-1

Percent of Cells
0   100

C

V3   V5   V6

n = 4,579   n = 3,521   n = 4,456

UMAP-2
UMAP-1

S-binding antibodies ● Yes ● No

D

NA
IGHA1
IGHG1
IGHG2
IGHG3
IGHM

Proportion of Sequences

V3: n=1, n=92, n=174, n=38, n=30
V5: n=1, n=6
V6: n=5, n=21, n=1307, n=1

V3   V5   V6

E

Nucleotide mutation frequency in IGHV

P>0.9999
P=0.0003
P<0.0001
P<0.0001
P=0.0658
P<0.0001

n=2553 (Naive)
n=1238 (Influenza)
n=360 (V3)
n=14 (V5)
n=1403 (V6)

Naive Influenza V3 V5 V6

P<0.0001

n=1318 (Cross-reactive)
n=411 (Non cross-reactive)

Cross-reactive Non cross-reactive

A

	spike mutation	
variant	RBD	outside RBD
B.1.1.7	N501Y	Δ69-70, Δ144, A570D, P681H, T716I, S982A, D1118H
B.1.351	K417N, E484K, N501Y	Δ242-244 L18F, D80A, D215G, A701V
P.1	K417T, E484K, N501Y	L18F, T20N, P26S, D138Y, R190S, H655Y, T1027I

B

ACE2

E484

K417

N501

RBD "up"

142-145

S982

242-244

P26

69-70

NTD

D614

H655

A701

T716

D1118

C

PVI.V3-9

PVI.V6-4

Rmax (nm)

no ACE2 | 1µM ACE2 | 5µM ACE2 | 25µM ACE2

D

RBD variant	K_D (nM)	Fold change
wild-type	35.4±5.6	1
N501Y	7.0±1.1	0.2
Y453F	29.9±18.1	0.8
N439K	15.7±3.8	0.4
Y453F/N439K	7.2±0.1	0.2
E484K	138.3±30.9	4
K417N/E484K/N501Y	34.0±4.6	1
K417T/E484K/N501Y	4.0±1.7	0.1

A

**Spike binding/neutralizing activity ratio
convalescent individuals**

B

**Ratio spike binding/RBD binding antibodies
convalescent individuals**

**Ratio spike binding/RBD binding antibodies
vaccinated individuals**

RBD variant	$K_D(10^{-11}M)$	Fold change
wild-type	38.8±0.5	1
N501Y	12±0.1	0.3
Y453F	18.9±0.1	0.5
N439K	59.8±9.3	1.5
Y453F/N439K	37.4±0.3	1
E484K	181.9±14.5	4.7
K417N/E484K/N501Y	71.3±3.7	1.8
K417T/E484K/N501Y	32.9±1.3	0.8