Transmission of SARS-CoV-2 by children to contacts in schools and households: a prospective
cohort and environmental sampling study in London

Rebecca Cordery[1], Lucy Reeves[2], Jie Zhou[2], Aileen Rowan[2], Patricia Watber[2], Carolina Rosadas[2], Michael Crone[2], Marko Storch[2], Paul
Freemont[2], Lucy Mosscrop[2], Alice Cowley[1], Gina Zelent[1], Kate Bisset[1], Holly Le Blond[1], Sadie Regmi[1], Christian Buckingham[2], Ramlah
Junaideen[2], Nadia Abdulla[2], Joseph Eliahoo[3], Miranda Mindlin[1], Theresa Lamagni[4], Wendy Barclay[2], Graham
P. Taylor[2], Shiranee Sriskandan[2,5]
7
8 **Affiliations**
9 [1] London Health Protection Teams, United Kingdom Health Security Agency; [2]Department of Infectious Disease, Imperial College London, London,
10 UK; [3]Statistical Advisory Service, Imperial College London, London, UK; [4]COVID-19 EpiCell, United Kingdom Health Security Agency; [5]NIHR Health
11 Protection Research Unit in Healthcare-associated infections and AMR
12
13 **Footnotes to title:** United Kingdom Health Security Agency (UK HSA), formerly Public Health England (PHE).
14 **Correspondence**: Professor Shiranee Sriskandan, Department of Infectious Disease, Imperial College London, Hammersmith Hospital, Du Cane
15 Road, London W12 0NN
16

Abstract

Background Assessing transmission of SARS-CoV-2 by children in schools is of critical importance to inform
public health action. We assessed frequency of acquisition of SARS-CoV-2 by contacts of children with COVID-
19 in schools and households, as well as the amount of virus shed into the air and onto fomites in both settings.
Methods Cases of COVID-19 in children in London schools were identified via notification. Weekly sampling
for 3-4 weeks and PCR testing for SARS-CoV-2 of immediate classroom contacts (the "bubble"), non-bubble
school contacts, and household contacts was undertaken supported by genome sequencing, along with surface
and air sampling in the school and home environment.
Results. Within schools, secondary transmission was not detected in 28 individual bubble contacts,
representing 10 distinct bubble classes. Across 8 non-bubble classes, 3/62 pupils tested positive– all three were
asymptomatic and tested positive in one setting on the same day, unrelated to the original index case. In
contrast, the secondary attack rate in naïve household contacts was 14.3% (5/35) rising to 19.1% (9/47) when
considering all household contacts. Environmental contamination with SARS-CoV-2 was rare in schools,
regardless of school type; fomite SARS-CoV-2 RNA was identified in 4/189 (2.1%) samples in bubble
classrooms, 2/127 (1.6%) samples in non-bubble classrooms, and 5/130 (3.8%) samples in washrooms. This
contrasted with fomites in households, where SARS-CoV-2 RNA was identified in 60/248 (24.2%) bedroom
samples, 66/241 (27.4%) communal room samples, and 21/188 (11.2%) bathroom samples. Air sampling
identified SARS-CoV-2 RNA in just 1/68 (1.5%) of school air samples, compared with 21/85 (24.7%) of air
samples taken in homes.
Summary The low levels of environmental contamination in schools are consistent with low transmission
frequency and adequate levels of cleaning and ventilation in schools during the period of study. Secondary
transmission in schools was rare. The high frequency of secondary transmission in households associated with
evident viral shedding throughout the home suggests a need to improve advice to households with infection in
children in order to prevent onward community spread by sibling and adult contacts. The data highlight that
transmission from children is very likely to occur when precautions are reduced.
42

Introduction.

The potential for schools to amplify outbreaks is well-recognised (1-3). School closures were associated with a reduction in COVID-19 incidence and mortality at the start of the pandemic (4,5) albeit effects remain confounded by other non-pharmaceutical interventions. Importantly, any benefits of closures must be weighed against the unquestionable harms to children and wider society.

COVID-19 poses a much lower risk to children than to adults, both in terms of illness severity (6) and risk of acquisition; children appear half as likely as adults to acquire SARS-CoV-2 (7). The onward transmission risk from SARS-CoV-2-infected children has been subject to less rigorous evaluation, though shedding of virus by children is not markedly different to adults (8). Point prevalence studies indicating a low prevalence of SARS-CoV-2 in schools (9) makes large scale monitoring of transmission impractical and uneconomic. Although outbreaks provide an opportunity to study onward transmission, these are often complicated by uncertainty about timing and transmission direction; one study in schools did identify children as a source of onward transmission in a small number of cases, but such outbreaks comprised just two cases on average (10).

Forward contact tracing offers an opportunity to search actively for secondary infections in a controlled manner. Despite this, transmission of respiratory infection in schools is seldom quantified except in the context of major outbreaks. Clinical attack rates of 20-30% are reported in schools affected by influenza A (11), however the role of silent infection and onward transmission from such cases is not well-studied. In a scarlet fever contact tracing study, we found that outbreak strains spread to over one-quarter of classroom contacts, despite treatment and isolation of index cases. (12) The potential for classrooms and asymptomatic 'shedders' to act as an accelerator for respiratory infection is therefore undeniable.

We adapted our contact-tracing protocol to investigate transmission of SARS-COV-2 by children in schools and households. The TraCK (Transmission of Coronavirus-19 in Kids, ISRCTN 13773960) study aimed to assess the risk posed by a SARS-CoV2-infected child who attends school, via longitudinal sampling of the child, school and household contacts, and associated environments, to evaluate and inform interventions to limit spread of COVID-19.

Methods

Study eligibility   Schools in London reporting new cases of SARS-CoV-2 infection to local Health Protection teams were invited to take part if a child (index case) had been attending school in the 48h prior to a positive PCR test for SARS-CoV-2. Contextual information relating to prevailing regulations are in Appendix p8. Parents/guardians of notified cases were invited to allow their child and wider household to participate in the study. If the school was willing to support the study, parents/guardians

of contacts were also invited to allow their child to participate in the study. The study commenced
October 9th 2020 and recruitment ended July 18th 2021.

Case definition. Children aged 2-14 years (extended to <18 years in November 2020) with a new
nose and/or throat swab PCR test result positive for SARS-CoV-2 from an accredited laboratory.
Findings from cases will be reported elsewhere.

Contact definition. Bubble contacts (BC) were children identified by schools who were required to
isolate at home due to direct contact with a case. Non-bubble school contacts (SC) were children
from a different 'control' class in the same school. Household contacts (HC) were adults and children
of any age normally resident with the case, and required to isolate.

Contact sampling Combined nose and throat samples (single swab of throat followed by nostrils) were
taken by the research team from each participating contact (BC, SC, or HC) as soon as possible (<48
hours) after case identification, and thereafter weekly for a total of 4 visits (3 visits from December
2020).

Environmental sampling. In households, surface and air samples were obtained in each of three
rooms (child's bedroom, communal room, bathroom) at the first visit and thereafter weekly for a
minimum of 4 visits (3 visits from December 2020); in some households sampling was undertaken
more frequently in the first two weeks. In schools, surface and air samples were obtained weekly
from the bubble classroom, school contact classroom, and washrooms. (for details see Appendix p9-
10)

Virological testing. Nasopharyngeal swabs were tested for SARS-CoV-2 E-gene RNA and human
RNAseP RNA by an accredited, quantitative RT-PCR followed by genome sequencing (Appendix p9-
10). (13). Results were reported in real-time to participants and positive results subject to statutory
reporting and associated regulations. Environmental samples were tested by a research laboratory
(14) (Appendix p10).

Gingival crevicular fluid (GCF). GCF samples were collected from contacts on each sampling
occasion (Appendix p11) then tested for total IgG against SARS-CoV2 nucleoprotein by the reference
laboratory (15).

Ethical approval. The study was approved by a research ethics committee (Schools Transmission
Study REC reference 18/LO/0025; IRAS Reference 225006). Written, informed consent was obtained
from all participants or parents/guardians, and assent was obtained from participants aged under 18.

Statistical analysis. Analysis was mostly descriptive due to sample size (Appendix p9); Fisher's exact
test was used to compare proportions of household contacts with positive results (Stata version 15).
Human target RNAs were compared using Mann Whitney U test (GraphPad Prism).

Role of the Funding Source None
113

Results.

Eight schools participated, of which 5 were primary, 2 secondary, and 1 was a special educational needs (SEND) school. In the course of the study, 428 combined nose and throat swabs and GCF samples were obtained from contacts of index cases. Environmental sampling included a total of 1620 surface samples, of which 446 were from schools, and 218 air samples, of which 68 were from schools.

Transmission to Bubble Contacts.

BC were recruited from 10 bubbles in 8 schools. In total 28 bubble contacts who were required to quarantine at home, were followed weekly. Onward transmission of SARS-CoV-2 to the 28 participating BC was not detected over the sampling period (Figure 1A, Table 1). Only 4/28 (14.3%) BC had evidence of prior exposure to SARS-CoV-2 from GCF testing. In one setting, a non-participating BC developed a fever and reported a positive community test. That child was recruited as a 'case' along with their household. Subsequent study sample PCR tests were negative, but GCF seroconversion at 4 weeks was consistent with this child being a co-primary case in the class. Participation rate among BC in each school varied widely (median 8.5%, range 2.4% - 26.9%), being lowest in SEND and secondary schools.

Transmission to and between non-bubble School Contacts.

Sixty-two pupil SC and 3 staff were recruited from the same 8 schools. SC participation rates were higher than BC, median 22.4% (range 5.2-54.5%). Of those tested, 13/65 (20%) had GCF antibodies indicating previous SARS-CoV-2 infection.

In 7/8 participating schools, no SC were found to be infected with SARS-CoV-2. In setting E, a secondary school, all SC tested negative in week 1, but in week 2, unexpectedly, SARS-CoV-2 was detected in swabs of 3/10 SC. (Figure 1B, Table 2). All three were asymptomatic; in one, the viral load increased from 293,240 E gene copies/swab to 5,999,560/swab copies 3 days later and onward transmission to a sibling household contact who shared a bedroom (84,040 E gene copies/swab) was observed. The other two asymptomatic SC had very low viral loads; the first had 280 E gene copies/swab but was tested only once. The second had 560 E gene copies/swab; samples 7 days earlier, and 4 days later were PCR-negative; and anti-SARS-CoV-2 antibodies were already present in GCF in weeks 1 and 2. It was felt possible that these low viral levels did not represent true infections, but transient mucosal contamination while in the company of a fellow pupil with active infection. The original index case in setting E had been identified following a community PCR test; by week 1 of SC testing the index case had a negative PCR test and was still quarantined. It was inferred that the infection in SC was not linked directly to the original index case.

Transmission to Household contacts.

Sixteen households took part, comprising 47 HC and 16 index cases who were each an index or co-
primary case to a bubble class. The number of households exceeded the number of bubbles that
participated because, in four settings, HC agreed to take part, but the relevant schools withdrew. In
one setting, the school agreed to take part, but the HC withdrew; a separate case from the same class
was identified by community testing however, and their HC were recruited. For setting E, HC of two
of the three newly identified SC infections were included. All of the index cases were symptomatic
except these two.
Of the HC, 3 children and 9 adults were already reported to be infected at the start of sampling. Initial
analysis focussed on HC who were considered naïve (n=35) i.e. were not reported to be infected at
the start of sampling, of which 11/35 were children. (Table 3)
Over the sampling period, 9 new infections were detected among naïve HC in 8 adults and 1 child
(Table 3, Figure 1C). In two households, genome sequencing revealed that the index case was
unrelated to the new adult HC infections (2 per household), hence these represented secondary
introduction from the community (Table 3, Appendix p5). In all other households genome sequencing
was consistent with clonal household transmission (Appendix p5). Transmission by children therefore
resulted in infection of 5/35 (14.3%) naïve HC. Only 1/35 (2.9%) GCF samples suggested prior
COVID-19 exposure among naïve HC at the start of sampling though this rose to 6/26 (23.1%) by the
end of sampling (p=0.035). Just 6 HC had been partially or fully-vaccinated; these were 2 adults each
in settings K1, K2, and M.
Twelve HC who were reported to be already-infected prior to study team arrival were also sampled
sequentially, but were not included in the initial analysis, due to uncertainty of transmission direction.
To gain greater insight into the frequency of secondary attack rate, symptom and testing history were
reviewed. Three child HC were reported to be positive prior to research sampling; based on symptom
onset and date of testing, it was deduced that these child HC had been secondarily infected by the
index pupil in the home. Nine adults (from 5 households) were reported to be positive prior to research
sampling. For 5/9 adults, test results and/or symptoms pre-dated that of the index child, suggesting
that the child was not the index case within the household. For 4/9 adults, their infection was believed
to arise from the index child. Taking these additional cases into consideration, the 16 index children
resulted in 9 new cases in 47 household contacts (19.1% secondary attack rate).
179
Environmental samples in schools
Surface sampling identified SARS-CoV-2 in only 4/189 (2.1%) samples from bubble classrooms;
2/127 (1.6%) samples in school contact classrooms, and 5/130 (3.8%) samples from school
bathrooms. (Figure 2A-C). Where detected, viral copy numbers were at the lower limits of detection
except the edge of an index child's chair in a bubble classroom that had $>10^4$ E gene copies per swab
in week 1, prior to deep cleaning. The same items were sampled in each location on a weekly basis
(Appendix p2-3); no item became positive on subsequent sampling. Air sampling was undertaken

weekly in bubble classrooms, control classrooms, and washrooms, as soon as possible after children vacated those rooms, except when availability of equipment components limited this. Only 1/68 (1.5%) air samples was positive: This was at the limit of detection, in week 2 in a school that had experienced a number of staff infections, but in a control SC classroom not known to have any pupil COVID-19 cases.

We considered the possibility that air samples might only be positive when a room is in active use. To provide context, we undertook environmental sampling in a university building (appendix p4). We identified SARS-CoV-2 in 3/10 surface samples from a small office 4 days after use by a confirmed case of COVID-19, but not in any other office or location in the same building, or on follow up (0/96 samples). We also detected low levels of SARS-CoV-2 in an air sample from the same office 4 days after use; all air samples were negative when re-tested two weeks later (Appendix p4).

Environmental samples in households

In contrast to findings in schools, overall 262/1174 (22.3%) surface samples were found to be contaminated with SARS-CoV-2 in 16 households. Focussing on samples taken on the first visit and thereafter weekly, there was a trend to declining virus detection over time (Figure 2 D-F). The most frequent surface contamination was identified in index case bedrooms, where 60/248 (24.2%) samples tested positive, and communal rooms, where 66/241 (27.4%) samples tested positive. In bathrooms, 21/188 (11.2%) surface samples tested positive, consistent with increased bathroom surface cleaning. Personal items relating to the child such as pillows, and digital equipment such as mobile phones, remote controls and digital toys were more persistently positive over the sampling period whereas other sample types became negative within 2-3 weeks, including pet fur (Appendix p6). Surface human RNA levels were higher in households than schools (Appendix p7).

Overall, 42/150 (28%) air samples obtained in households were contaminated with SARS-CoV-2. Focussing on samples taken on the first visit and thereafter weekly, air samples were positive in 4/22 (18.2%) samples taken in the index child's bedroom; 13/42 (30.9%) samples in the communal room; and 4/21 (19%) samples in the bathroom (Figure 2D-F). Virus levels in air were highest in the room with an infected child and infected adults. The index child and household contacts were always in the communal (living) room at the time of sampling except three settings where the index child was only in the bedroom during sampling, and one setting where the child moved between rooms. There was no apparent association between the type of dwelling (apartment or house) and air contamination. Air samples in households and schools did not differ significantly with regard to human RNA (Appendix p7).

Discussion

Conducted during a period of enhanced precautions, transmission from index pupils to bubble contacts, and to other pupils in the school who were not close contacts, was not actively detected. Although the study was small, the findings contrasted with a secondary attack rate of at least 14.3%

in household contacts of the same index cases. When household contacts who had already been tested were included in our analysis, the secondary attack rate in households with a child index case was 19.1%.

One apparent transmission incident in a class that were not isolating involved three asymptomatic pupils, who could not be linked to the original index case in that school. One of these pupils had a high viral load, leading to detection of a secondary case in a household contact and, we believe, accounted for transient low viral loads detected in two other pupils. The low viral loads were similar to environmental samples and may be consistent with transient carriage on mucosae rather than early or late infection.

Environmental surface and air sampling was conducted to understand mechanisms of transmission, where transmission occurred. This showed little or no contamination in schools including surfaces touched frequently by children, providing a high level of reassurance regarding the school environment during a period of enhanced vigilance, underlined by a difference in human RNA detection between surfaces in households and schools. This contrasted with repeated identification of virus on household items frequently touched by children, and in the air around the home, particularly where the child was present. This is perhaps not surprising since the dimensions of domestic rooms are ~4 times smaller than classrooms and provides some insight into the risks of virus acquisition in the two settings. The detailed environmental sampling identifies digital equipment and personal items as potential fomite vectors, or as metrics of infectivity. The high proportion of air samples that were positive in the home compared with school underlines the greater risks associated with smaller rooms and is a reminder that air may remain positive for some time if not well ventilated. We considered the possibility that air sampling in schools was negative because the children were not present in the room, however control human RNA was no different in the air between schools and households. Control sampling in a different educational setting demonstrated low levels of SARS-CoV-2 RNA in the air 4 days after an office was used by a staff member with COVID-19. The low or absent levels of SARS-CoV-2 RNA in the bubble classroom also provides reassurance about the potential for ongoing infection in members of the bubble who returned to school by week 2-3.

Our findings are consistent with studies undertaken in other countries that have examined transmission in the school setting; when actively sought, transmission to bubble contacts was uncommon, with 1-2% co-primary or secondary infections identified where larger numbers have been sampled (16, 17). It is also consistent with the ~1.5% asymptomatic infection rate reported in a recent cluster-randomised trial of daily lateral flow-testing in bubble contacts (18). The infrequency of transmission to other pupils contrasts with transmission frequency of other respiratory infections in schools, including group A streptococcus and influenza (11,12); this may reflect the multifold interventions in place during the pandemic period, or it may reflect the heterogeneity of infection in COVID19 where most transmission is caused by only a minority of infections (19, 20).

Our study prospectively examined transmission from the same children to contacts in both schools
and households; the secondary attack rate in households was higher than expected, and was in stark
contrast to that seen in schools. Our findings are consistent with a recent study that reported a
secondary attack rate of 25% in households even when the index case is a child (21), and a recent
meta-analysis (22). While children may be less likely than adults to become secondary cases, the
risk of generating secondary cases is no different whether the index is a child or adult (21, 23); this
pattern is confirmed in other countries (22). Quarantine for household contacts, in place throughout
our study, may increase exposure of household members to index cases unless mitigated by
protective measures, noting household size has been associated with urban caseloads (24). It was
notable that in all households with no onward transmission to naïve contacts, householders had
ensured that the affected child was isolated from others, without sharing a bedroom, whilst still
affording care and supervision.

For ethical reasons, we used GCF to screen for prior SARS-CoV2 exposure, which may under-
estimate exposure compared with serum (14). Prevalence of seropositivity among school pupils
reported by larger scale testing is similar to levels observed in pupils in our study (25). Due to timing
of our study, just six of the adult contacts had been vaccinated. Though vaccination was reported to
impact household SAR (26), a recent study suggests a lesser impact with more transmissible variants
(27).

Our study adopted a forensic approach to contact tracing, to not miss infections that were cleared
early, or those arising late due to ongoing transmission in the class group. We took combined nasal
and pharyngeal swabs to increase opportunity for virus detection and used human RNAseP as a
control to ensure that negative results could be trusted. Furthermore, almost all swabs were taken
by the study team; a small number of contacts were permitted to take swabs themselves if supervised.
Genome sequencing identified transmission events that were genuine while also refuting others,
highlighting a risk of over- or under-estimating transmission rates when relying on PCR results alone.
285

286 The study was designed to investigate bubble sizes of ~10-15, but interpretation of 'bubble' changed
287 over time, and by autumn 2020 bubble sizes routinely included 30-200 primary- and secondary-aged
288 pupils respectively (28). The study relied upon identification of index cases who had been attending
289 school; as such, index cases in this study were almost all symptomatic, with the expectation that
290 asymptomatic cases would be identified among contacts as a comparison group.
291

There are three key limitations to our study. Firstly, the study was conducted at a time of heightened
and constantly changing interventions, in particular social distancing in schools and reduced class
sizes. Transmission in schools may alter when interventions relax, as indicated by more recent
epidemiological reports (29). Secondly, participation rates in contacts were very low, compared with
participation rates of >40% in a previous contact tracing study (12). Deterrents to participation

reported anecdotally were the legal requirement to notify newly-identified infections; quarantine
impact on participants; study team making home visits; low risk in children; and inclusion of older
pupils. Participation by school contacts was consistently higher than bubble contacts, underlining a
resistance to home visits. Recruiting bubble contacts sent home to isolate was challenging, as schools
use an array of methods to contact parents. The greatest barrier to participation was the recognition
that newly-identified infections would result in quarantine for entire households or classes, such that
participation was actively discouraged by some groups, in contrast to predicted responses at study
inception. Finally, although our study benefitted from the objective starting point of positive index
cases who attend school, there is a risk of bias in all studies that rely on voluntary participation, in
terms of individual schools and participants. Representation from a larger number of participants
would however require expansive recruitment.

Future research of this kind may provide more meaningful data if the results are unlinked to identifiable
data, or any form of notification or requirement to isolate, i.e. without real-time reporting. With reduced
interventions and advent of new variants, it may be prudent to evaluate schools-based transmission
in such a silent study.

Funding
UKRI/DHSC (Grant COV0322); NIHR

Acknowledgements The study team acknowledge the support of the school leaders, support staff,
and the participants who took part in the study as well as the NIHR Clinical Research Networks and
Schools Network for supporting the schools in this study.

The input of teaching and parent advisory groups is also acknowledged, as is the practical support of
key staff from Imperial College Healthcare NHS Trust, Imperial College London, and Public Health
England, and the TraCK study collaborators. We are also grateful to Shamez Ladhani, Samreen Ijaz,
John Poh, Justin Shute and members of the Oral Fluid team at PHE Colindale.

Author contributions. Conceptualisation SS and RC; Study site supervision RC; Laboratory
supervision GPT, WB, PF, SS; Study methodology SS, RC, GPT, WB, TL, MM; Data collection LR,
JZ, AR, PW, CR, MC, MS, LM, AC, GZ, KB, HL, SR, CB, RJ, NA; Data visualisation JZ, LR; Data
validation LR, GPT, JZ; Data analysis JZ, JE; Funding acquisition SS, WB, GPT, RC, TL, MM; Writing,
original draft SS; Review and critical editing RC, GPT, WB. Final draft: all authors.

Declaration of competing interests The authors declare no competing interests

333
334

References

1. Cauchemez S, Valleron AJ, Boëlle PY, Flahault A, Ferguson NM. Estimating the impact of school closure on influenza transmission from Sentinel data. Nature. 2008 Apr 10;452(7188):750-4. doi: 10.1038/nature06732.

2. Earn DJ, He D, Loeb MB, Fonseca K, Lee BE, Dushoff J. Effects of school closure on incidence of pandemic influenza in Alberta, Canada. Ann Intern Med. 2012 Feb 7;156(3):173-81. doi: 10.7326/0003-4819-156-3-201202070-00005

3. Bin Nafisah S, Alamery AH, Al Nafesa A, Aleid B, Brazanji NA. School closure during novel influenza: A systematic review. J Infect Public Health. 2018 Sep-Oct;11(5):657-661. doi: 10.1016/j.jiph.2018.01.003.

4. Auger KA, Shah SS, Richardson T, Hartley D, Hall M, Warniment A, et al. Association between Statewide School Closure and COVID-19 Incidence and Mortality in the US. JAMA - Journal of the American Medical Association. 2020;45229:859-70.

5. Brauner JM, Mindermann S, Sharma M, Johnston D, Salvatier J, Gavenčiak T, Stephenson AB, Leech G, Altman G, Mikulik V, Norman AJ, Monrad JT, Besiroglu T, Ge H, Hartwick MA, Teh YW, Chindelevitch L, Gal Y, Kulveit J. Inferring the effectiveness of government interventions against COVID-19. Science. 2021 Feb 19;371(6531):eabd9338. doi: 10.1126/science.abd9338. Epub 2020 Dec 15.

6. Swann OV, Holden KA, Turtle L, Pollock L, Fairfield CJ, Drake TM, Seth S, Egan C, Hardwick HE, Halpin S, Girvan M, Donohue C, Pritchard M, Patel LB, Ladhani S, Sigfrid L, Sinha IP, Olliaro PL, Nguyen-Van-Tam JS, Horby PW, Merson L, Carson G, Dunning J, Openshaw PJM, Baillie JK, Harrison EM, Docherty AB, Semple MG; ISARIC4C Investigators. Clinical characteristics of children and young people admitted to hospital with covid-19 in United Kingdom: prospective multicentre observational cohort study. BMJ. 2020 Aug 27;370:m3249. doi: 10.1136/bmj.m3249.

7. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, Waddington C, Thomas J, Russell S, van der Klis F, Koirala A, Ladhani S, Panovska-Griffiths J, Davies NG, Booy R, Eggo RM. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. JAMA Pediatr. 2020 Sep 25:e204573. doi: 10.1001/jamapediatrics.2020.4573

8. Jones TC, Biele G, Mühlemann B, Veith T, Schneider J, Beheim-Schwarzbach J, Bleicker T, Tesch J, Schmidt ML, Sander LE, Kurth F, Menzel P, Schwarzer R, Zuchowski M, Hofmann J, Krumbholz A, Stein A, Edelmann A, Corman VM, Drosten C. Estimating infectiousness throughout SARS-CoV-2 infection course. Science. 2021 Jul 9;373(6551):eabi5273. doi: 10.1126/science.abi5273.

9. Office for National Statistics. COVID-19 Schools Infection Survey England, Round 6, June
2021.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddi
seases/bulletins/covid19schoolsinfectionsurveyengland/round6june2021 Accessed December
20th 2021

10. Ismail SA, Saliba V, Lopez Bernal J, Ramsay ME, Ladhani SN. SARS-CoV-2 infection and
transmission in educational settings: a prospective, cross-sectional analysis of infection
clusters and outbreaks in England. Lancet Infect Dis. 2021 Mar;21(3):344-353. doi:
10.1016/S1473-3099(20)30882-3.

11. Health Protection Agency West Midlands H1N1v Investigation Team. Preliminary descriptive
epidemiology of a large school outbreak of influenza A(H1N1)v in the West Midlands, United
Kingdom, May 2009. Euro Surveill. 2009 Jul 9;14(27):19264. doi: 10.2807/ese.14.27.19264-
en.

12. Cordery R, Purba A, Begum L, Mills EA, Mosavie M, Vieira A, Jauneikaite E, Leung
RCY, Ready D, Hoffman P, Lamagni T, Sriskandan S. High frequency transmission,
asymptomatic shedding, and airborne spread of *Streptococcus pyogenes* among
schoolchildren exposed to scarlet fever: a longitudinal multi-cohort moleculo-epidemiological
contact tracing study.
medRxiv 2021.07.04.21259990; doi: https://doi.org/10.1101/2021.07.04.21259990

13. Rowan AG, May P, Badhan A, et al. Optimized protocol for a quantitative SARS-CoV-2 duplex
RT-qPCR assay with internal human sample sufficiency control. *J Virol Methods*.
2021;294:114174. doi:10.1016/j.jviromet.2021.114174

14. Zhou J, Otter JA, Price JR, Cimpeanu C, Garcia DM, Kinross J, Boshier PR, Mason S, Bolt F,
Holmes AH, Barclay WS. Investigating SARS-CoV-2 surface and air contamination in an acute
healthcare setting during the peak of the COVID-19 pandemic in London. Clin Infect Dis. 2020
Jul 8:ciaa905. doi: 10.1093/cid/ciaa905

15. Hoschler K, Ijaz S, Andrews N, Ho S, Dicks S, Jegatheesan K, Poh J, Warrener
L, Kankeyan T, Baawuah F, Beckmann J, Okike IO, Ahmad S, Garstang J, Brent AJ, Brent
B, Aiano F, Brown KE, Ramsay ME, Brown D, Parry JV, Ladhani SN, Zambon M. SARS-
CoV-2 IgG detection in human oral fluids.
medRxiv 2021.07.07.21260121; doi: https://doi.org/10.1101/2021.07.07.21260121

16. Brandal Lin T, Ofitserova Trine S, Meijerink Hinta, Rykkvin Rikard, Lund Hilde M, Hungnes
Olav, Greve-Isdahl Margrethe, Bragstad Karoline, Nygård Karin, Winje Brita A. Minimal
transmission of SARS-CoV-2 from paediatric COVID-19 cases in primary schools, Norway,
August to November 2020. Euro Surveill. 2021;26(1):pii=2002011.
https://doi.org/10.2807/1560-7917.ES.2020.26.1.2002011

17. Dawson P, Worrell MC, Malone S, Tinker SC, Fritz S, Maricque B, Junaidi S, Purnell G, Lai
AM, Neidich JA, Lee JS, Orscheln RC, Charney R, Rebmann T, Mooney J, Yoon N, Petit M,
Schmidt S, Grabeel J, Neill LA, Barrios LC, Vallabhaneni S, Williams RW, Goddard C, Newland
JG, Neatherlin JC, Salzer JS; CDC COVID-19 Surge Laboratory Group. Pilot Investigation of
SARS-CoV-2 Secondary Transmission in Kindergarten Through Grade 12 Schools
Implementing Mitigation Strategies - St. Louis County and City of Springfield, Missouri,
December 2020. MMWR Morb Mortal Wkly Rep. 2021 Mar 26;70(12):449-455. doi:
10.15585/mmwr.mm7012e4.

18. Young BC, Eyre DW, Kendrick S, White C, Smith S, Beveridge G, Nonnenmacher T, Ichofu F,
Hillier J, Oakley S, Diamond I, Rourke E, Dawe F, Day I, Davies L, Staite P, Lacey A, McCrae
J, Jones F, Kelly J, Bankiewicz U, Tunkel S, Ovens R, Chapman D, Bhalla V, Marks P, Hicks
N, Fowler T, Hopkins S, Yardley L, Peto TEA. Daily testing for contacts of individuals with
SARS-CoV-2 infection and attendance and SARS-CoV-2 transmission in English secondary
schools and colleges: an open-label, cluster-randomised trial. Lancet. 2021 Oct
2;398(10307):1217-1229. doi: 10.1016/S0140-6736(21)01908-5.

19. Chen PZ, Bobrovitz N, Premji Z, Koopmans M, Fisman DN, Gu FX. Heterogeneity in
transmissibility and shedding SARS-CoV-2 via droplets and aerosols. Elife. 2021 Apr
16;10:e65774. doi: 10.7554/eLife.65774.

20. Adam DC, Wu P, Wong JY, Lau EHY, Tsang TK, Cauchemez S, Leung GM, Cowling BJ.
Clustering and superspreading potential of SARS-CoV-2 infections in Hong Kong. Nat Med.
2020 Nov;26(11):1714-1719. doi: 10.1038/s41591-020-1092-0.

21. Miller E, Waight PA, Andrews NJ, McOwat K, Brown KE, Katja H, Ijaz S, Letley L, Haskins D,
Sinnathamby M, Cuthbertson H, Hallis B, Parimalanathan V, de Lusignan S, Lopez-Bernal J.
Transmission of SARS-CoV-2 in the household setting: A prospective cohort study in children
and adults in England. J Infect. 2021 Aug 1:S0163-4453(21)00380-7. doi:
10.1016/j.jinf.2021.07.037.

22. Thompson HA, Mousa A, Dighe A, Fu H, Arnedo-Pena A, Barrett P, Bellido-Blasco J, Bi Q,
Caputi A, Chaw L, De Maria L, Hoffmann M, Mahapure K, Ng K, Raghuram J, Singh G, Soman
B, Soriano V, Valent F, Vimercati L, Wee LE, Wong J, Ghani AC, Ferguson NM. Severe Acute
Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Setting-specific Transmission Rates: A
Systematic Review and Meta-analysis. Clin Infect Dis. 2021 Aug 2;73(3):e754-e764. doi:
10.1093/cid/ciab100.

23. Hall JA, Harris RJ, Zaidi A, Woodhall SC, Dabrera G, Dunbar JK. HOSTED-England's
Household Transmission Evaluation Dataset: preliminary findings from a novel passive
surveillance system of COVID-19. Int J Epidemiol. 2021 Jul 9;50(3):743-752. doi:
10.1093/ije/dyab057.

24. Federgruen A, Naha S. Crowding Effects Dominate Demographic Attributes in COVID-19
Cases. Int J Infect Dis. 2021 Jan;102:509-516. doi: 10.1016/j.ijid.2020.10.063.

25. Office for National Statistics. COVID-19 Schools Infection Survey England Round 4, pupil
antibody data, March 2021.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddi
seases/bulletins/covid19schoolsinfectionsurveyengland/round4pupilantibodydatamarch2021#
pupils-testing-positive-for-coronavirus-covid-19-antibodies. Accessed December 20th 2021

26. Harris RJ, Hall JA, Zaidi A, Andrews NJ, Dunbar JK, Dabrera G. Effect of Vaccination on
Household Transmission of SARS-CoV-2 in England. N Engl J Med. 2021 Aug 19;385(8):759-
760. doi: 10.1056/NEJMc2107717.

27. Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, Derqui-Fernandez
N, Barnett JL, Whitfield MG, Varro R, Charlett A, Kundu R, Fenn J, Cutajar J, Quinn V, Conibear
E, Barclay W, Freemont PS, Taylor GP, Ahmad S, Zambon M, Ferguson NM, Lalvani A;
ATACCC Study Investigators. Community transmission and viral load kinetics of the SARS-
CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a
prospective, longitudinal, cohort study. Lancet Infect Dis. 2021 Oct 29:S1473-3099(21)00648-
4. doi: 10.1016/S1473-3099(21)00648-4

28. Department for Education Evidence Summary Coronavirus (COVID-19) - children, young
people and education settings July 2021
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_dat
a/file/1000025/Evidence_Summary_-_July_2021.pdf Accessed December 20th 2021

29. Chudasama DY, Tessier E, Flannagan J, Leeman D, Webster H, Demirjian A, Falconer C,
Thelwall S, Kall M, Saliba V, Ramsay M, Dabrera G, Lamagni T. Surge in SARS-CoV-2
transmission in school-aged children and household contacts, England, August to October
2021. Euro Surveill. 2021 Dec;26(48):2101019. doi: 10.2807/1560-
7917.ES.2021.26.48.2101019.

Tables

Table 1. Transmission to Bubble classroom contacts

School	Bubble size (incl. cases)	Case number Bubble exposed to	Bubble participant number	Number of bubble contacts testing PCR positive[†]				Bubble contact crevicular fluid anti-NP total IgG	
				Week 1	Week 2	Week 3	Week 4	Positive on first sample	Positive on last sample
A	29	1	3	0/2	0/3	0/3	0/3	1/3	1/3
B	26	1	7	0/4	0/4	0/6	0/7	2/7	1/7
D	41	1	1	0/1	0/1	ND	0/1	0/1	0/1
E[‡]	39	1	2	0/2	0/2	0/2	ND	0/2	0/2
F[§]	48	2	5	0/5	0/5	0/5	ND	0/5	0/5
G[¶]	16	4	2	0/2	0/2	0/2	0/2	0/2	0/2
K[†]	150	11	6	0/6	0/6	0/6	ND	1/6	1/6
M	30	1	2	0/1	0/2	0/2	ND	0/2	0/2
TOTAL			28	0/23	0/25	0/26	0/13	4/28	3/28

[‡]swabbing delayed until 7d after case confirmed.
[§]Includes 2 different bubbles exposed to one case each. One non-participant bubble contact tested positive in community test (included in household study).
[¶]Bubble exposed to 2 adult and 2 child cases
[†] Includes 2 different bubbles exposed to 4 cases and 7 cases
ND, not done

Table 2. Transmission to non-bubble (control) class contacts

School	Non-bubble class size	No. of cases at start of study in school	No. of non-bubble participants	No. of non-bubble contacts testing PCR positive/no. swabbed				Non-bubble contact crevicular fluid anti-NP total IgG	
				Week 1	Week 2	Week 3	Week 4	Positive on first sample	Positive on last sample
A	30	1	5	0/1	0/5	0/5	0/5	3/5	1/5
B	22	1	12	0/12	0/12	0/12	0/12	1/12	1/12
D	27	1	2	0/2	0/1	0/2	0/2	0/2	1/2
E[§]	30	1	10	0/10	3/8	1/5	1/1	2/10	1/8
F	11	2	2	0/1	0/1	0/2	ND	0/2	0/1
G[‡]	24	4	7	ND	0/7	0/7	0/7	1/7	2/7
			3	ND	0/3	0/3	0/3	0/3	0/3
K	306	26	16	0/14	0/16	0/16	ND	3/16	4/16
M	30	1	8	0/6	0/7	0/8	ND	3/8	3/7
TOTAL			65	0/46	3/60	1/60	1/30	13/65	13/61

[§]Swabbing of school contacts started one week after initial case
[‡]4 cases in school included 2 children and 2 adults. Contacts include 7 children and 3 adults
ND, not done

Table 3. Transmission events in participating household contacts in each setting

Site	Household size[§]	No. of child cases at start of swabbing[¶]	No. of adult cases at start of swabbing	No. of naïve household contacts	Number of naïve household contacts testing positive				Naïve household contact crevicular fluid anti-NP total IgG	
					Week 1	Week 2	Week 3	Week 4	Positive on first sample	Positive on last sample
A	2	1	0	1	0/1	0/1	0/1	0/1	0/1	0/1
B	3	1	1	1	0/1	0/1	0/1	0/1	0/1	0/1
C[¥]	6	1	0	5	0/5	0/5	0/4	ND	0/5	0/5
D[†▽]	4	1	0	3	2/3	1/3	1/3	0/3	0/3	1/3
E.1[†]	4	1	0	3	0/3	0/3	0/3	ND	0/3	0/3
E.2[‡]	4	1	0	3	1/3	ND	ND	ND	0/3	ND
E.3[†]	7	1	0	6	0/6	ND	ND	ND	0/6	ND
F	4	2	0	2	1/2	2/2	ND	ND	0/2	0/2
G.1[†]	4	1	0	3	0/3	1/3	1/3	0/3	0/3	2/3
G.2[†]	3	1	0	2	1/2	1/2	1/1	0/1	1/2	2/2
H	3	1	2	0	0/0	0/0	ND	ND	0/0	0/0
I	3	1	2	0	0/0	0/0	0/0	ND	0/0	0/0
J	4	2	2	0	0/0	0/0	0/0	ND	0/0	0/0
K.1[†▽]	5	1	0	4	1/4	2/4	0/3	0/2	0/4	1/4
K.2	4	2	2	0	0/0	0/0	0/0	0/0	0/0	0/0
M	3	1	0	2	0/2	0/2	0/2	ND	0/2	0/2
TOTAL				35	6/35	7/26	3/21	0/11	**1/35**	**6/26**

[§] includes index child case: each household had 100% participation rate at time of consent
[¶] includes index child case plus any other child already identified as infected. ND, not done due to intervening holiday or withdrawal from study.
Naïve household contacts include 1 child[†]; 2 children[‡]; 3 children[¥]. Household contacts with different genomic sequences to index case are indicated[▽]. Proportions in **bold** are different (p=0.035).

Figure 1. Pictograms of individual contacts in each week of sampling. A. Bubble contacts (n=28). B. School contacts (n=62 pupils, 3 staff) and C. Household
cases and contacts. (n=63). For panel C, the 26 participants reported to the study team as having tested positive prior to research swabbing are indicated by*
(child index cases, adult and child household contacts). Colour of icons indicates research swab test result in each week of study: Blue icons,negative swab
result; pink icons, SARS-CoV2 detected; grey icons, subject not swabbed in that week or not recruited yet. Two of three pupils identified incidentally are included
in both panels B and C (i.e. school and household contact pictograms); although pupils were swabbed weekly, the associated households were recruited only
after week 2. Within each panel, the figure position is consistent in each week and represents individual participants so can be compared between weeks 1-4.
Individual settings are separated by gaps between groups of figures. For presentation purposes, the ordering of settings between panels A, B and C is not the
same. Longitudinal sampling was limited to three weeks rather than four weeks for part of the study hence some subjects were not swabbed in week 4.

Figure 2.

Figure 2 Environmental contamination with SARS-CoV-2 in schools and households. A-C, schools; D-F, households. Samples obtained at start of sampling and thereafter weekly are shown. Red indicates surface samples; blue indicates air samples. Surface and air samples were obtained from the same items and locations weekly in each school and households. Data shown as absolute E gene copy number and represent samples from 8 schools (1 SEND; 2 secondary; 5 primary) and 16 households; note y axis range differs between schools and households A. Bubble contact classroom . B. School contact classroom C. School bathroom used by bubble. D. Child's bedroom. E. Communal room F. Bathroom used by child.