

1 **Factors associated with maternal tetanus vaccination in Myanmar: An analysis of Demographic**
2 **and Health Survey data**

3 Zaw Myo Tun^{*a}, Zau Ring^{*b}, Clarence C Tam^{a, c}

4 ^a Saw Swee Hock School of Public Health, National University of Singapore

5 12 Science Drive 2, #10-01, Singapore 117549

6 ^b Vector Borne Diseases Control Program, Kachin State Public Health Department, Ministry of Health and

7 Sports, Myanmar

8 Kachin State Public Health Department Office, Yone Gyi Street, Ayar Quarter, Myitkyina, Myanmar

9 ^c London School of Hygiene & Tropical Medicine

10 Keppel Street, London WC1E 7HT

11 * ZMT and ZR equally contributed to the manuscript

12

13 Corresponding author:

14 Zaw Myo Tun (Email: zawmyotun@nus.edu.sg)

15 Saw Swee Hock School of Public Health, National University of Singapore

16 12 Science Drive 2, #10-01, Singapore 117549

17

18 Keywords:

19 **Tetanus toxoid vaccination; Myanmar; Universal health coverage; Epidemiology; Demographic Health**
20 **Survey**

21

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1 **Introduction**

2 Maternal and neonatal tetanus (MNT) is a devastating disease with a high case fatality, especially in low-income
3 settings [1]. The World Health Organization (WHO) estimated 25,000 new-borns died from neonatal tetanus in
4 2018 [1]. MNT remains a significant public health problem in 12 countries, most of them are in Asia and Africa
5 [1]. MNT death is preventable by hygienic delivery, cord care practices, and by immunizing children and pregnant
6 women with inexpensive and efficacious tetanus toxoid vaccine [1]. In the last 30 years, vaccinating pregnant
7 women in their second or third trimester has reduced tetanus-related neonatal mortality by 94% [2]. WHO
8 recommends that “all women giving birth and their newborn babies should be protected against tetanus” and that
9 countries should ensure that a national policy and strategies to achieve high vaccination coverage among pregnant
10 women are available and are correctly implemented [3].

11 In Myanmar, a population of 51.5 million [4], tetanus vaccination for pregnant women has been part of the routine
12 Expanded Program on Immunization since 1978 [5]. Although MNT was eliminated (defined as <1 MNT per
13 1000 live births in every district) in Myanmar in 2010 [6], the MNT burden and gaps in MNT preventive services
14 remains: 231 neonatal tetanus cases were reported from 2011 to 2018, and 64% of mothers of neonatal tetanus
15 cases in 2018 were unimmunized [7]; only 72% of women giving birth in the previous five years are protected
16 against neonatal tetanus [5].

17 Factors associated with maternal tetanus vaccination are not well understood in Myanmar. The Myanmar Ministry
18 of Health and Sports launched the five-year National Health Plan in December 2016 to achieve universal health
19 coverage by 2030, aiming to ensure access to a basic essential package of health services by the entire population
20 [8]. Maternal tetanus vaccination is listed as a component of the essential package [9]. Information on factors
21 associated with maternal tetanus under-vaccination would be useful to inform the implementation strategy. We
22 estimated maternal tetanus vaccination coverage at both national and subnational levels using 2015-16 Myanmar
23 Demographic and Health Survey (MDHS) data. We also investigated factors associated with being unvaccinated
24 among women reporting pregnancy in the previous five years.

25 **Methods**

26 We used maternal health data from the 2015-16 MDHS which was implemented by the Ministry of Health and
27 Sports with support from the Inner City Fund (ICF) (Rockville, Maryland, USA). The DHS collected data on
28 demographic and health indicators of women and their household members using a nationally representative
29 sample. The survey methodology and data collection procedures have been published in the main report [5].

1 Briefly, the survey design employed a two-stage sampling (441 clusters, 30 households per cluster) stratified by
2 urban and rural areas in 15 subnational administrative divisions (seven states, seven regions, and Nay Pyi Taw
3 Union Territory). In this analysis, Nay Pyi Taw Union Territory was categorized as a region. The states,
4 representing the mountainous areas, are the areas of residence for seven ethnic minority groups while the majority
5 Burmese ethnicity primarily live in the eight regions which represent the plain areas. The states are generally less
6 developed than the regions [4]. In this cross-sectional analysis, we included women aged 15-49 years who reported
7 giving birth in the last five years.

8 *Outcome variable*

9 Outcome was being unvaccinated with tetanus toxoid containing vaccine (TTCV) self-reported by mothers.
10 Maternal tetanus vaccination was defined as newborns protected against neonatal tetanus (See Appendix 1 for the
11 full definition) according to the Myanmar National Guidelines for Antenatal Care [10]. The following variables
12 were used to construct the outcome variable: (1) number of tetanus injection before birth (during the last
13 pregnancy); (2) number of tetanus injections before the last pregnancy; (3) number of years ago received last
14 tetanus injection before the last pregnancy. We estimated the percentage of unvaccinated mothers with
15 corresponding 95% confidence interval (CI), at national and subnational levels, accounting for the DHS sampling
16 weights. We also compared vaccine coverage with the percentage of mothers who received at least one dose of
17 TTCV during their last pregnancy in each state and region. Their differences indicate mothers who missed a
18 second dose of TTCV as the vast majority of vaccinated mothers received two TTCV injections (See results
19 section below for more details).

20 *Explanatory variables*

21 We selected 13 variables and grouped them into three domains: (1) Socio-demographic: residence in a state or a
22 region, urban or rural residence, age (categorized as 15-24, 25-29, 30-34, 35-49), birth order of the last pregnancy
23 (first, second, third, fourth or higher), educational attainment (no education, primary, secondary, post-secondary
24 or higher), occupation type (manual, sales, agricultural, other, not working), wealth index quintiles (specific to
25 urban or rural areas); (2) Women's agency: whether getting permission and going alone for getting medical help
26 were big problems, person who usually decide the respondent's healthcare; (3) Healthcare access: whether getting
27 money needed for treatment and the distance to the nearest health facility were big problems, and the type of
28 health facility where the respondent received antenatal care (ANC) (at a public health facility, a private health
29 facility or a non-governmental organization clinic, at home, or no ANC).

1 *Conceptual framework*

2 Among factors influencing maternal and child health, socio-demographic factors unlikely influence their health
3 outcomes directly. Rather, their influence is most likely mediated through multiple inter-related proximal factors.
4 For example, mothers with higher educational attainment and greater household wealth may be empowered to
5 exercise their agency and actively seek the necessary health services. In this example, adjusting for proximal
6 factors such as women's agency and healthcare access may reduce the overall effects of the distal factors (socio-
7 demographic).

8 Following this reasoning, we adapted a conceptual framework recommended by Victora and colleagues [11],
9 based on hypotheses derived from the literature and plausibility reasoning. We hypothesized that TTCV
10 vaccination was influenced by proximal factors related to healthcare access, which were themselves influenced
11 by factors related to women's agency in seeking care. These, in turn, were influenced by socio-demographic
12 factors. The aim of the conceptual framework is to understand the extent of the role of more distal factors (e.g.,
13 socio-demographic variables) on vaccination that is explained by more proximal factors (e.g., healthcare access).
14 Appendix 2 shows a simple schema of the conceptual framework.

15 A review on maternal immunization in developing countries summarized various socio-economic factors
16 associated with tetanus vaccine uptake by pregnant women in Bangladesh, India, Kenya, and Ethiopia [12]. These
17 factors included mother's age, birth order, urban or rural residence, educational level, socioeconomic status,
18 woman's occupation, among the others.

19 As ANC is the most important strategy to provide maternal immunisation [13], including TTCV, it seems
20 reasonable to assume factors related to healthcare and ANC access as the most proximal to being unvaccinated.
21 Studies have shown that ANC uptake is influenced by many socio-demographic factors. In Ethiopia, an analysis
22 of DHS 2016 data found that higher maternal education status and mothers with greater household wealth were
23 more likely to access ANC while the uptake was low among mothers who had traditional belief and those with
24 five children and more [14]. In addition, a systematic review summarizing the factors influencing ANC uptake
25 showed that most factors were related to socio-demographic domain: maternal education, husband's education,
26 marital status, household income, mother's employment, media exposure, having a history of obstetric
27 complications, and parity [15].

28 Women's agency is the extent of belief in herself, the awareness of her rights, and make choices to control
29 acquired resources. A 2016 systematic review showed that, in developing countries, women's empowerment was

1 associated with improved maternal and child health outcomes such as access to ANC and maternal vaccination
2 [16]. A previous analysis of MDHS 2015-16 dataset also demonstrated the association between women
3 empowerment and fewer barriers in accessing healthcare, especially among women with a higher educational
4 attainment, fewer children, from rich households, and those residing in urban areas [17].

5 *Statistical analysis*

6 We described continuous variables using median and interquartile range; categorical variables using frequency
7 and percentage. Logistic regression models were used in univariable and multivariable analyses accounting for
8 the MDHS sampling weight to investigate factors associated with being unvaccinated. Odds ratios (OR) with their
9 corresponding confidence intervals (CI) were reported. Independent variables were assessed for possible
10 collinearity using Phi and Cramer's V correlation coefficients (further details can be found in Appendix 4).
11 Multivariable analysis steps and model interpretation under the conceptual framework are summarized in
12 Appendix 3. All analyses were performed using R software (version 4.1.0) [18] and survey analysis was performed
13 using survey package [19]. Correlation coefficients of independent variables were computed using vcd package
14 [20].

15 *Ethics statement*

16 The 2015-16 MDHS has obtained ethics approval from both ICF Institutional Review Board and Ethics Review
17 Committee of the Department of Medical Research, Ministry of Health and Sports, Myanmar. We obtained the
18 permission from ICF to access the MDHS datasets.

19 *Role of the funding source*

20 No funding was received for this work.

21 **Results**

22 Our results showed that 69% resided in regions and 77% in rural areas. The median age was 30 years (interquartile
23 range: 26, 36). A third of mothers had their first childbirth in the previous five years. Most mothers (84%) had at
24 least primary school education. Sixteen percent of mothers reported that their decisions of healthcare were made
25 by husband, partner, or others. One in eight mothers did not receive ANC during their last pregnancy (Table 1).

1 *Unvaccinated mothers*

2 Overall, 28% (95% CI: 69%, 74%) of mothers reported that they were unvaccinated. The vast majority of
3 vaccinated mothers received two TTCV injections during the last pregnancy (the first definition in Appendix 1)
4 while fewer than 30 mothers fulfilled the remaining definitions each. At subnational level, the percent
5 unvaccinated was the highest in Shan State (42%), followed by Magway Region (33%), Kayin State (33%), and
6 Sagaing Region (32%) (Figure 1). The percentage of being unvaccinated in each category of explanatory variables
7 is shown in Table 1.

8 We also estimated that mothers who received at least one dose of TTCV during their last pregnancy was 84%
9 (95%CI: 82%, 87%) with lowest estimates in Shan (69%), Kayin (75%) and Chin (79%) States. Figure 2 shows
10 that the percentage of mothers who missed a second dose of TTCV varied between 5% and 17% at subnational
11 level. Notably, the subnational administrative areas with the highest percentage were regions: Magway (17%),
12 Sagaing (17%), Nay Pyi Taw Union Territory (16%), Bago (16%), and Tanintharyi (14%) (Figure 2).


13 *Factors associated with being unvaccinated*

14 Table 2 shows the results of multivariable analysis under the conceptual framework. Model 1 results represent
15 overall effects of socio-demographic variables. Model 2 provided the effect estimates of women's agency
16 variables after adjusting for the confounding effects of socio-demographic factors. Model 3 results show the effect
17 estimates of healthcare access factors after adjusting for socio-demographic and women's agency variables. We
18 found that older age and higher educational attainment of mothers were associated with lower odds of being
19 unvaccinated. On the other hand, not receiving ANC services or receiving ANC in non-public health facilities
20 (private health facilities or non-governmental organization clinic, at home, or mixed, compared to public health
21 facilities) was associated with higher odds of being unvaccinated. Maternal educational attainment remained
22 associated with being unvaccinated after adjusting for the confounding effects of variables in women's agency
23 and healthcare access domains. Additionally, there was little evidence that maternal age was associated with being
24 unvaccinated in Model 3, suggesting that the effects of maternal age is mediated primarily through mothers'
25 healthcare access.

26

27


1 **Figure 1. Percentage of unvaccinated mothers with tetanus toxoid containing vaccine in each state or**
2 **region**



3
4

1 **Figure 2. Mothers who missed a second dose of maternal tetanus vaccine in each state and region**

2 Mothers who missed a second dose of maternal tetanus vaccine is the percentage difference between the vaccine
3 coverage and the percentage of mothers who received at least one dose of tetanus toxoid containing vaccine
4 during the last pregnancy



5
6
7
8

9 **Table 1. Characteristics of mothers and the percentage of being unvaccinated with tetanus toxoid**
10 **containing vaccine**

	Number of mothers	%	Unvaccinated*		
			n	%	(95% CI)
Overall	3,582		990	27.6	(25.1, 30.0)
Socio-demographic					
State or region of residence					
Region	2,452	68.5	639	26.1	(24.3, 27.9)
State	1,130	31.5	351	31.1	(28.4, 33.9)
Urban or rural residence					
Urban	838	23.4	163	19.5	(16.9, 22.3)
Rural	2,744	76.6	827	30.1	(28.4, 31.9)
Age of women					
15-24	660	18.4	211	32.0	(28.5, 35.7)
25-29	924	25.8	248	26.8	(24.0, 29.8)
30-34	912	25.5	226	24.8	(22.0, 27.7)
35-49	1,087	30.3	306	28.2	(25.5, 30.9)
Birth order of last pregnancy					
1st	1,222	34.1	302	24.7	(22.3, 27.3)

2nd	920	25.7	220	23.9	(21.2, 26.8)
3rd	617	17.2	178	28.8	(25.3, 32.6)
4th or higher	824	23.0	290	35.2	(31.9, 38.6)
Highest education level of mothers					
No education	587	16.4	261	44.5	(40.4, 48.6)
Primary	1,629	45.5	480	29.5	(27.3, 31.8)
Secondary	1,069	29.8	205	19.2	(16.9, 21.7)
More than secondary	298	8.3	45	15.1	(11.3, 19.8)
Mother's occupation					
Manual	1,060	29.6	323	30.5	(27.7, 33.4)
Sales	541	15.1	129	23.8	(20.4, 27.7)
Agricultural	512	14.3	174	34.0	(29.9, 38.3)
Other	210	5.9	46	21.9	(16.6, 28.2)
Not working	1,249	34.9	313	25.1	(22.7, 27.6)
(Missing)	10	0.3			
Quintiles of woman's wealth index (specific to urban/rural areas)					
Lowest	949	26.5	341	35.9	(32.9, 39.1)
Second	812	22.7	229	28.2	(25.2, 31.5)
Third	669	18.7	171	25.6	(22.3, 29.1)
Fourth	585	16.3	131	22.4	(19.1, 26.0)
Highest	567	15.8	119	21.0	(17.8, 24.6)
Women's agency					
Getting medical help for self: getting permission to go					
Not a big problem	3,407	95.1	928	27.2	(25.8, 28.8)
Big problem	175	4.9	62	35.4	(28.5, 43.1)
Person who usually decides on woman's healthcare					
Self	1,314	36.7	355	27.0	(24.6, 29.5)
With husband or partner	1,534	42.8	424	27.6	(25.4, 30.0)
By husband, partner, or others	575	16.1	157	27.3	(23.7, 31.2)
(Missing)	159	4.4			
Getting medical help for self: not wanting to go alone					
Not a big problem	2,408	67.2	602	25.0	(23.3, 26.8)
Big problem	1,174	32.8	388	33.0	(30.4, 35.8)
Healthcare access					
Getting medical help for self: getting money needed for treatment					
Not a big problem	2,192	61.2	523	23.9	(22.1, 25.7)
Big problem	1,390	38.8	467	33.6	(31.1, 36.2)
Getting medical help for self: distance to health facility					
Not a big problem	2,601	72.6	624	24.0	(22.4, 25.7)
Big problem	981	27.4	367	37.4	(34.4, 40.5)
Place of antenatal care					
Public health facilities	2,133	59.5	412	19.3	(17.7, 21.1)
Private health facilities or NGO clinics	289	8.1	68	23.5	(18.8, 28.9)
Home	565	15.8	163	28.8	(25.2, 32.8)
Mixed	136	3.8	32	23.5	(16.9, 31.7)
None	460	12.8	317	68.9	(64.4, 73.1)

1 CI, Confidence Interval

2 * Vaccinated mothers include those who has received two injections of tetanus toxoid containing vaccine during
3 the last pregnancy, two or more injections with the last one within three years of the last pregnancy, three or
4 more injections with the last one within five years of the last pregnancy, four or more injections with the last
5 one within 10 years of the last pregnancy, and five or more injections at any time prior to the last pregnancy

1 **Table 2. Factors associated with being unvaccinated with tetanus toxoid containing vaccine**

	Univariable OR (95% CI)	Adjusted OR (95% CI)		
		Model 1	Model 2	Model 3
Socio-demographic				
State or region of residence				
Region	1	1	1	1
State	1.28 (0.95, 1.72)	0.97 (0.75, 1.25)	0.91 (0.70, 1.18)	0.78 (0.60, 1.01)
Urban or rural residence				
Urban	1***	1	1	1
Rural	1.78 (1.35, 2.35)	1.33 (1.00, 1.77)	1.23 (0.92, 1.65)	1.14 (0.85, 1.52)
Age of mothers				
15-24	1*	1**	1*	1
25-29	0.78 (0.58, 1.04)	0.78 (0.57, 1.04)	0.75 (0.55, 1.02)	0.78 (0.56, 1.07)
30-34	0.70 (0.55, 0.90)	0.61 (0.45, 0.82)	0.61 (0.44, 0.83)	0.66 (0.48, 0.92)
35-49	0.84 (0.67, 1.05)	0.62 (0.45, 0.85)	0.63 (0.45, 0.88)	0.72 (0.51, 1.01)
Birth order of last pregnancy				
1st	1***	1	1	1
2nd	0.96 (0.75, 1.23)	0.98 (0.74, 1.28)	0.94 (0.71, 1.26)	0.88 (0.65, 1.20)
3rd	1.24 (0.95, 1.62)	1.25 (0.92, 1.70)	1.20 (0.87, 1.66)	1.03 (0.73, 1.44)
4th and higher	1.66 (1.27, 2.16)	1.40 (1.00, 1.98)	1.37 (0.96, 1.95)	1.02 (0.73, 1.44)
Mother's highest education level				
No education	1***	1***	1***	1**
Primary	0.52 (0.36, 0.75)	0.58 (0.41, 0.81)	0.58 (0.41, 0.81)	0.79 (0.56, 1.11)
Secondary	0.30 (0.20, 0.43)	0.38 (0.26, 0.56)	0.36 (0.25, 0.54)	0.53 (0.35, 0.81)
More than secondary	0.22 (0.13, 0.37)	0.38 (0.21, 0.70)	0.34 (0.18, 0.65)	0.44 (0.22, 0.85)
Mother's occupation				
Manual	1*	1	1	1
Sales	0.72 (0.52, 1.00)	1.00 (0.72, 1.38)	1.02 (0.73, 1.43)	1.19 (0.85, 1.66)
Agricultural	1.18 (0.81, 1.72)	1.14 (0.80, 1.61)	1.19 (0.85, 1.67)	1.15 (0.84, 1.56)
Other	0.64 (0.41, 0.99)	1.00 (0.60, 1.64)	1.03 (0.62, 1.73)	1.17 (0.70, 1.98)
Not working	0.76 (0.59, 0.99)	0.87 (0.67, 1.13)	0.88 (0.68, 1.14)	0.91 (0.70, 1.18)
Quintiles of mother's wealth index (specific to urban or rural areas)				
Lowest	1***	1	1	1
Second	0.70 (0.53, 0.93)	0.80 (0.59, 1.08)	0.77 (0.57, 1.05)	0.84 (0.61, 1.14)
Third	0.61 (0.47, 0.80)	0.75 (0.56, 1.01)	0.80 (0.59, 1.09)	0.89 (0.65, 1.24)
Fourth	0.51 (0.38, 0.69)	0.71 (0.51, 0.99)	0.76 (0.54, 1.06)	0.82 (0.57, 1.19)
Highest	0.47 (0.31, 0.72)	0.70 (0.45, 1.11)	0.74 (0.47, 1.18)	0.94 (0.57, 1.55)
Women's agency				
Getting medical help for self: getting permission to go				
Not a big problem	1*		1	1
Big problem	1.49 (1.01, 2.18)		1.02 (0.69, 1.49)	0.82 (0.52, 1.29)
Person who usually decides on mother's healthcare				
Self	1		1	1
With husband or partner	1.03 (0.85, 1.26)		1.02 (0.84, 1.25)	1.01 (0.82, 1.25)
By husband, partner or others	1.02 (0.78, 1.33)		0.89 (0.67, 1.18)	0.80 (0.59, 1.08)
Getting medical help for self: not wanting to go alone				
Not a big problem	1**		1	1
Big problem	1.48 (1.14, 1.93)		1.19 (0.93, 1.51)	1.02 (0.77, 1.35)
Healthcare Access				
Getting medical help for self: getting money needed for treatment				
Not a big problem	1***			1
Big problem	1.61 (1.29, 2.02)			1.19 (0.92, 1.54)
Getting medical help for self: distance to health facility				
Not a big problem	1***			1
Big problem	1.89 (1.46, 2.46)			1.15 (0.85, 1.55)
Place of antenatal care				

Public health facilities	1***	1***
Private health facilities or NGO clinics	1.28 (0.89, 1.83)	1.68 (1.16, 2.43)
Home	1.69 (1.31, 2.18)	1.49 (1.13, 1.96)
Mixed	1.28 (0.79, 2.07)	1.44 (0.86, 2.42)
None	9.23 (6.43, 13.26)	7.65 (5.29, 11.07)

CI, Confidence interval; OR, Odds ratio

Overall p value using the Wald-type test: *** = <0.001, ** = <0.01, * = <0.05

1
2

3 **Discussion**

4 We found important regional variations in the percentage of unvaccinated mothers (ranging from 42% in Shan
5 State to 15% in Mon State), and the percentage of mothers who missed a second dose of TTCV. Notably, we
6 found that the percentage of unvaccinated mothers was higher among younger mothers, and mothers with lower
7 educational attainment. Among these factors, differences in the percentage of unvaccinated mothers maternal age
8 were largely mediated through their access to healthcare and ANC services. Maternal educational attainment
9 remained associated with vaccination status after controlling for healthcare access variables. A lack of ANC or
10 receiving ANC in non-public healthcare facilities were strongly associated with being unvaccinated.

11 Our analysis showed that the overall percentage of being unvaccinated with TTCV was 28%, translating to the
12 vaccine coverage of 72%. This was lower than the vaccine coverage (88%) officially reported to the WHO by the
13 Myanmar health ministry in 2016 [21]. Although this gap may be explained by methodological differences, it
14 remains a possibility that the official vaccine coverage may underestimate the gap in maternal tetanus vaccination
15 services in the country. In the current analysis, the national vaccine coverage was lower than most countries in
16 the WHO Southeast Asia region in 2016, except Nepal (67%), Indonesia (65%), and Timor-Leste (37%) [21].

17 We also analysed the percentage of mothers who missed a second dose of TTCV (Figure 2). This information is
18 useful because reasons of not receiving even the first dose of TTCV may differ from reasons of not receiving a
19 second dose. The former is likely due to the lack of access to vaccination services, to information related to these
20 services, or vaccine hesitancy. However, mothers who missed a second dose of TTCV may be due to different
21 reasons, for example, supply side problems: inadequate coverage of vaccination services, poor service quality,
22 frequent stock out, inadequate competent healthcare workers. Our analysis identified regional differences in the
23 percentage of mothers who missed a second dose of TTCV. These percentages are greater in the regions compared
24 to the states. These findings point to an opportunity to significantly improve the vaccine coverage in Myanmar
25 and warrant further investigation.

1 At subnational level, the percentage of mothers who received at least one dose of TTCV was lowest in Shan,
2 Kayin and Chin States. In addition, the percentage of unvaccinated mothers in these states was among the highest
3 of states and regions (Figure 1). This may be due to limited access to tetanus vaccine. Kayin State and certain
4 areas in Shan State have long been known as chronic conflict-affected areas. Although Myanmar has transitioned
5 to democracy since 2011, conflicts in these areas continued while a multilateral ceasefire agreement involving the
6 military, the newly elected civilian government, and several ethnic armed organizations were being negotiated in
7 parallel [22, 23]. As such, pregnant women residing in these areas faced tremendous challenges in accessing
8 essential maternal health services. Studies aiming to understand the maternal health in these areas are limited. A
9 survey was conducted in 2006 in several accessible conflict-affected areas in Eastern Myanmar (including Shan
10 and Kayin States) near the border with Thailand. The survey found that only 18% and 4% of mothers in Shan and
11 Kayin States, respectively, reported that they have received two doses of tetanus toxoid. The corresponding
12 percentages of mothers attending ANC at least four visits were 45% and 7%, respectively. In addition, women
13 also reported their experience of human rights violations such as destruction or stealing of field and livestock,
14 forced labor, forced relocation, direct physical attacks by soldiers or authorities, and landmines [24]. Separately,
15 Chin State is highly mountainous with few transportation links, which poses challenges for access to maternal
16 health services [25]. A study analyzing MDHS data observed regional differences in ANC coverage (at least four
17 antenatal visits): the coverage in Shan, Kayin and Chin States ranged from 37 to 50%, compared with the coverage
18 in Yangon Region of 82% [26].

19 Of particular concern is the high percentage of unvaccinated mothers among those who did not access or receive
20 ANC. Approximately 1 in 8 mothers reported not receiving any ANC and of these, more than half were
21 unvaccinated. In Myanmar, ANC services including maternal tetanus vaccination are primarily provided through
22 networks of public health facilities/centres at the village level [10]. Our results show that this system can
23 effectively provide immunization services to most mothers, although the lower coverage among mothers receiving
24 ANC outside the public health sector – private or NGO facilities or at home – points to an opportunity to increase
25 vaccination coverage in these sectors. ANC at home is usually provided by trained midwives and lady health
26 visitors, but in some rural areas ANC is provided by traditional birth attendants who are not trained on vaccination.
27 A study in Krabi Province, Thailand, found that although 91% of traditional birth attendants were aware of
28 injections during ANC at the health center, only half knew that it was for tetanus prevention [27]. A similar
29 knowledge gap may exist among traditional birth attendants in Myanmar. These gaps in ANC provision need to
30 be addressed as Myanmar works towards achieving universal health coverage.

1 Lower maternal educational attainment was strongly associated with being unvaccinated. The association
2 remained after controlling for healthcare access variables. This suggests a synergistic opportunity for promoting
3 access to both education and healthcare in generating demand for and acceptance of vaccination and improving
4 maternal health outcomes. Lower maternal education is related to lower use of antenatal care services [28], while
5 higher maternal education is also associated with higher uptake of maternal immunisation in several low-income
6 countries, including India [29], Bangladesh [30], Kenya [31], and Malawi [12].

7 Our analysis has several limitations. In common with other DHS surveys, maternal tetanus vaccination status is
8 self-reported and subject to inaccurate recall. However, a previous study from Brazil showed that self-reported
9 uptake of maternal tetanus vaccination can be higher compared with medical records [32]. In addition, ethnicity
10 and religion could be important determinants of maternal vaccination [33]. However, this information was not
11 available in the survey. Nonetheless, the use of a large, nationally representative, and geographically stratified
12 survey dataset is a major strength.

13 In our analysis, we were able to identify regional, structural, and individual differences in the percentage of
14 mothers unvaccinated with TTCV. Our findings indicate that further research is necessary to investigate the
15 reasons of mothers not receiving a second dose of TTCV. Myanmar's progress toward achieving WHO's
16 recommended target of protecting all pregnant women and babies against tetanus will largely depend on the ability
17 to provide accessible ANC to the substantial fraction of mothers who currently do not currently receive it.

18 **Data sharing**

19 The 2015-16 DHS Myanmar dataset is accessible upon request from: <https://dhsprogram.com/>

20 **Conflict of interest**

21 We declare no competing interest.

22 **Contributors**

23 ZMT, ZR, and CCT conceived the study. ZR and ZMT obtained the data. ZR and ZMT analysed and interpreted
24 the data. ZMT and ZR wrote the first draft of the report. ZMT, ZR, and CCT contributed to the writing of the
25 report. All authors agreed with report results and conclusions. ZMT and CCT accessed and verified the underlying
26 data.

1 **Acknowledgements**

- 2 We thank the Demographic and Health Survey Program for providing data for this analysis.

1 **References**

- 2 [1] World Health Organization. Maternal and Neonatal Tetanus Elimination (MNTE),
3 https://www.who.int/immunization/diseases/MNTE_initiative/en/; 2020 [Accessed 23 December 2020]
- 4 [2] World Health Organization. Maternal immunization and antenatal care service delivery situation analysis
5 project, [https://www.who.int/maternal_child_adolescent/topics/maternal/immunization-antenatal-care-](https://www.who.int/maternal_child_adolescent/topics/maternal/immunization-antenatal-care-delivery/en/)
6 [delivery/en/](https://www.who.int/maternal_child_adolescent/topics/maternal/immunization-antenatal-care-delivery/en/); 2020 [Accessed 23 May 2020]
- 7 [3] World Health Organization. Maternal Immunization Against Tetanus. Geneva, Switzerland; 2006.
- 8 [4] Ministry of Immigration and Population/Myanmar. The 2014 Population and Housing Census. Nay Pyi Taw,
9 Myanmar; 2014.
- 10 [5] Ministry of Health Sports - MoHS/Myanmar, ICF. Myanmar Demographic and Health Survey 2015-16. Nay
11 Pyi Taw, Myanmar; 2017.
- 12 [6] United Nations. UN declares Myanmar free of maternal and neonatal tetanus,
13 <https://news.un.org/en/story/2010/06/340762-un-declares-myanmar-free-maternal-and-neonatal-tetanus>;
14 [Accessed 29 December 2020]
- 15 [7] Ministry of Health Sports - MoHS/Myanmar. Neonatal Tetanus. Monthly Epidemiology Bulletin. 2019;3, 7.
- 16 [8] Ministry of Health and Sports/Myanmar. National Health Plan 2017-2021. Nay Pyi Taw, Myanmar; 2016.
- 17 [9] Ministry of Health and Sports/Myanmar. National Health Plan: Annual Operational Plan (2018-2019). Nay
18 Pyi Taw, Myanmar; 2018.
- 19 [10] Ministry of Health Sports - MoHS/Myanmar. National Guidelines for Antenatal Care For Service Providers.
20 Nay Pyi Taw, Myanmar; 2018.
- 21 [11] Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in epidemiological
22 analysis: a hierarchical approach. Int J Epidemiol. 1997;26:224-7.
- 23 [12] Pathirana J, Nkambule J, Black S. Determinants of maternal immunization in developing countries. Vaccine.
24 2015;33:2971-7.
- 25 [13] World Health Organization. Protecting all against tetanus: guide to sustaining maternal and neonatal tetanus
26 elimination (MNTE) and broadening tetanus protection for all populations. Geneva; 2019.
- 27 [14] Tsegaye B, Ayalew M. Prevalence and factors associated with antenatal care utilization in Ethiopia: an
28 evidence from demographic health survey 2016. BMC Pregnancy and Childbirth. 2020;20:528.
- 29 [15] Simkhada B, Teijlingen ERv, Porter M, Simkhada P. Factors affecting the utilization of antenatal care in
30 developing countries: systematic review of the literature. Journal of Advanced Nursing. 2008;61:244-60.

- 1 [16] Pratley P. Associations between quantitative measures of women's empowerment and access to care and
2 health status for mothers and their children: A systematic review of evidence from the developing world. *Social
3 Science & Medicine*. 2016;169:119-31.
- 4 [17] Htun NMM, Hnin ZL, Khaing W. Empowerment and health care access barriers among currently married
5 women in Myanmar. *BMC Public Health*. 2021;21:139.
- 6 [18] R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for
7 Statistical Computing; 2019.
- 8 [19] Lumley T. survey: analysis of complex survey samples. 2020.
- 9 [20] Zeileis A, Meyer D, Hornik K. vcd: Visualizing Categorical Data. 2020.
- 10 [21] World Health Organization. Second and subsequent doses of tetanus toxoid,
11 https://apps.who.int/immunization_monitoring/globalsummary/timeseries/tscoveragett2plus.html; 2021
12 [Accessed 17 October 2021]
- 13 [22] Institute for Security & Development Policy. A Return to War: Militarized Conflicts in Northern Shan State.
14 Stockholm-Nacka, Sweden; 2018.
- 15 [23] Sandford S. Conflict Resumes in Karen State After Myanmar Army Returns, [https://www.voanews.com/east-
16 asia-pacific/conflict-resumes-karen-state-after-myanmar-army-returns](https://www.voanews.com/east-
16 asia-pacific/conflict-resumes-karen-state-after-myanmar-army-returns); 2018 [Accessed 30 December 2020]
- 17 [24] Mullany LC, Lee CI, Yone L, Paw P, Oo EKS, Maung C, et al. Access To Essential Maternal Health
18 Interventions and Human Rights Violations among Vulnerable Communities in Eastern Burma. *PLOS Medicine*.
19 2008;5:e242.
- 20 [25] Soe KH. Transport infrastructure upgrades take shape in Chin State, [https://www.gnlm.com.mm/transport-
21 infrastructure-upgrades-take-shape-in-chin-state/](https://www.gnlm.com.mm/transport-
21 infrastructure-upgrades-take-shape-in-chin-state/); 2020 [Accessed 29 December 2020]
- 22 [26] Han SM, Rahman MM, Rahman MS, Swe KT, Palmer M, Sakamoto H, et al. Progress towards universal
23 health coverage in Myanmar: a national and subnational assessment. *The Lancet Global Health*. 2018;6:e989-e97.
- 24 [27] Chongsuvivatwong V, Bucharkorn L, Treerong R. Traditional birth attendants in an endemic area of tetanus
25 neonatorum in Thailand: pitfalls in the control program. *J Community Health*. 1991;16:325-31.
- 26 [28] Titaley CR, Dibley MJ, Roberts CL. Factors associated with underutilization of antenatal care services in
27 Indonesia: results of Indonesia Demographic and Health Survey 2002/2003 and 2007. *BMC Public Health*.
28 2010;10:485.
- 29 [29] Khan REA, Raza MA. Maternal health-care in India: The case of tetanus toxoid vaccination. *Asian
30 Development Policy Review*. 2013;1:1-14.

- 1 [30] Rahman M. Tetanus toxoid vaccination coverage and differential between urban and rural areas of
2 Bangladesh. *East Afr J Public Health*. 2009;6:26-31.
- 3 [31] Haile ZT, Chertok IR, Teweldeberhan AK. Determinants of utilization of sufficient tetanus toxoid
4 immunization during pregnancy: evidence from the Kenya Demographic and Health Survey, 2008-2009. *J*
5 *Community Health*. 2013;38:492-9.
- 6 [32] Morón-Duarte LS, Ramirez Varela A, Bassani DG, Bertoldi AD, Domingues MR, Wehrmeister FC, et al.
7 Agreement of antenatal care indicators from self-reported questionnaire and the antenatal care card of women in
8 the 2015 Pelotas birth cohort, Rio Grande do Sul, Brazil. *BMC Pregnancy and Childbirth*. 2019;19:410.
- 9 [33] McGlynn N, Wilk P, Luginaah I, Ryan BL, Thind A. Increased use of recommended maternal health care as
10 a determinant of immunization and appropriate care for fever and diarrhoea in Ghana: an analysis pooling three
11 demographic and health surveys. *Health Policy and Planning*. 2014;30:895-905.
- 12