

*Infections in childhood and subsequent diagnosis of autism spectrum disorders and intellectual disability: a register-based cohort study*

Håkan Karlsson<sup>a</sup> PhD, Hugo Sjöqvist<sup>b</sup> MSc, Martin Brynge<sup>b</sup> MD, Renee Gardner<sup>b</sup> PhD,  
Christina Dalman<sup>b</sup> MD, PhD

*Previous presentations:* None

*Location of work:* The Departments of Neuroscience<sup>a</sup> and Global Public Health<sup>b</sup>, Karolinska Institutet, 171 77 Stockholm, Sweden.

Correspondence to: Håkan Karlsson, Karolinska Institutet, Neuroscience, Biomedicum B8, 171 77 Stockholm, Sweden, ([hakan.karlsson.2@ki.se](mailto:hakan.karlsson.2@ki.se))

*E-mail addresses:* Hugo Sjöqvist ([hugo.sjoqvist@ki.se](mailto:hugo.sjoqvist@ki.se)), Martin Brynge ([m.brynge@ki.se](mailto:m.brynge@ki.se)), Reneé Gardner ([renee.gardner@ki.se](mailto:renee.gardner@ki.se)), Christina Dalman ([christina.dalman@ki.se](mailto:christina.dalman@ki.se))

*Disclosure:* None of the authors have any financial relationships with commercial interests to disclose.

*Acknowledgments:* This work was generously supported by the Stanley Medical Research Institute (V-002, HK) and the Swedish Research Council, grant numbers 2018-02907 (HK), 2016-01477, 2012-2264 and 523-2010-1052 (CD)

3,521 words in main text, 1 table, 4 figures, 2 supplementary tables, 5 supplementary figures

**NOTE:** This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

## **Abstract**

*Objective.* To explore the associations between childhood infections and subsequent diagnoses of autism spectrum disorder (ASD), intellectual disability (ID) and their co-occurrence.

*Methods.* The association between specialized care for any infection, defined by ICD-codes and later ASD or ID was investigated in a register-based cohort of 556,732 individuals born 1987-2010, resident in Stockholm County, followed from birth to their 18th birthday or December 31, 2016. We considered as potential confounders children's characteristics, family socioeconomic factors, obstetric complications, and parental histories of both treatment for infection and psychiatric disorders in survival analyses with extended Cox regression models. Residual confounding by shared familial factors was addressed in sibling analyses using within-strata estimation in Cox regression models. Sensitivity analyses with exclusion of congenital causes of ASD/ID and documented risk for infections were also performed.

*Results.* Crude estimates indicated that infections during childhood were associated with later ASD and ID with largest risks observed for diagnoses involving ID. Inclusion of covariates, exclusion of congenital causes of ASD/ID from the population and sibling comparisons highlighted the potential for confounding by both heritable and non-heritable factors, though risks remained in all adjusted models. In adjusted sibling comparisons, excluding congenital causes, infections were associated with later 'ASD without ID' (HR 1.24, 95%CI 1.15-1.33), 'ASD with ID' (1.57, 1.35-1.82) and 'ID without ASD' (2.01, 1.76-2.28). Risks associated with infections varied by age at exposure and by age at diagnosis of ASD/ID.

*Conclusions.* Infections during childhood cannot be excluded in the etiology of ASD, particularly ASD with co-occurring ID.

## **1. Introduction**

Autism spectrum disorder (ASD) is diagnosed throughout childhood and age at diagnosis appears to be determined by both internal (severity, sex) and external factors (parental educational level, access to care) <sup>1</sup>. While parents often report deviations from normal developmental trajectories at an early age, the pattern of onset varies substantially. It remains to be established whether children later diagnosed are ‘born’ with ASD <sup>2,3</sup> or if the early childhood environment may contribute to a trajectory towards ASD in vulnerable children.

The etiology of ASD remains elusive but appears to involve both heritable and non-heritable genetic variants and environmental exposures <sup>4</sup>. The relative contributions of these plausibly causal factors vary across diagnostic sub-groups <sup>5-7</sup>. Intellectual disability (ID, IQ<70) is diagnosed in around 1/3 of those diagnosed with ASD <sup>8</sup>. Individuals diagnosed with ASD without co-occurring ID are more likely to have a family history of ASD and other psychiatric diagnoses than individuals diagnosed with ASD with ID <sup>9,10</sup>. Moreover, ID in individuals with ASD appears less familial compared to ID in the absence of ASD and may therefore have non-heritable causes <sup>11</sup>.

Infections that can invade the fetal brain are established causes of life-long behavioral and intellectual disabilities <sup>12,13</sup>. Infections targeting the brain during childhood have also been associated with cognitive impairments and psychiatric illness which become evident only later in life <sup>14-18</sup>. A growing literature associates childhood infections with no apparent CNS

involvement with schizophrenia risk in adulthood<sup>19-22</sup>, with a significant portion of risk mediated by reduced cognitive abilities<sup>21</sup>. Observations of deficits in innate immunity among neonates who later develop schizophrenia<sup>23</sup> or ASD<sup>24</sup> further highlight the potential contribution of postnatal infections to these outcomes<sup>25-27</sup>. However, only a handful of studies, with equivocal results, have investigated the relationship between childhood infections and the risk of ASD<sup>18, 28-31</sup>.

We aimed to explore the associations between infections and subsequent diagnoses of ASD, ID and their co-occurrence, and to understand whether any observed associations varied by age at infection.

## **2. Methods**

### *2.1 Study population and general design*

We defined a register-based cohort of 556,732 individuals born 1987-2010, resident in Stockholm County for  $\geq 3$  years<sup>32</sup>. Individuals not born in Sweden, adopted, not registered in the Medical Birth Register (MBR), or missing maternal data were excluded from the study (Figure S1).

Exposure, outcome and covariate data were extracted from national and regional registers containing routinely collected health and sociodemographic data cross-linked via each resident's unique national identification number<sup>33</sup>. The study was approved by the Stockholm regional ethics review board (DNR 2010/1185-31/5). Informed consent was not required to use anonymized register data.

### *2.2 Diagnoses of ASD and ID*

Diagnostic outcomes as of December 31, 2016 were defined by validated procedures covering all inpatient and outpatient pathways to care and diagnosis in Stockholm County <sup>11, 32</sup> (ICD-9: ASD 299, ID 317-19, ICD-10: ASD F84, ID F70-79). We considered two overlapping diagnostic groups (individuals who received any diagnosis of ASD or any diagnosis of ID), as well as the three mutually exclusive diagnostic outcomes: ASD with co-occurring ID ('ASD with ID'), ASD without co-occurring ID ('ASD without ID') and ID without co-occurring ASD ('ID without ASD'), Figure 1A.

### *2.3 Assessment of childhood infections*

All in- or outpatient specialized care diagnoses of infection, defined by ICD-9/10 codes as detailed previously <sup>19</sup>, were identified in the National Patient Register (NPR). We omitted all codes "sequel" and "post."

### *2.4 Covariates*

Covariates whose distribution varied by exposure to infection between birth and age 18 (Table 1) and that were associated with ASD or ID (Tables S1 & S2) were considered as potential confounders. Parental region of origin was identified from the Multigenerational Register. The Longitudinal Integrated Database for Health Insurance and Labor Markets Studies (LISA) supplied quintiles of family income adjusted by inflation and size of family (recorded at the birth of the index person) and highest parental education. MBR provided information on maternal body mass index (BMI, categorized according to WHO standards <sup>34</sup>), birth order, parental age at birth, birth season, maternal pre-eclampsia during pregnancy, size for gestational age (GA) at birth, Apgar score at 5 min after birth, birth by cesarean section, and GA at birth. NPR provided information on parental history of infections before birth of the index person. NPR and regional registers provided information on history of

parental psychiatric illness (defined as ICD10 F-chapter diagnosis, ICD9 290-315, or ICD8 290-315).

## *2.5 Statistical analyses*

We employed survival analysis with the extended Cox regression model using within-model stratification to account for differences by the children's year of birth. Age of the child was used as the underlying timescale, beginning at birth. Individuals were followed until death, emigration from Stockholm, ASD/ID diagnosis, or end of study. The infection variables were treated as time-dependent exposures, in which the individual could have had an infection from birth up to 18 years of follow-up, given that the child was not diagnosed with an outcome of interest or otherwise censored. To account for the possible non-proportional risk of the outcomes over follow-up time, we considered both cumulative time wherein the exposure to infection could occur from birth up to the 18<sup>th</sup> birthday, and stratified according to age at infection; 0-<1 year, 1-<3, 3-<6, 6-<12, and 12-<18. To test for differences in risk related to age at infection, we used the Wald test to evaluate the beta coefficients for each strata of age at exposure in the cumulative models. To assess if the associations varied by age at outcome, we also stratified the outcome according to age at first diagnosis of ASD and/or ID: 0-<6, 6-<12 and 12-<18 years.

To explore potential confounding by familial factors shared between siblings not accounted for by adjustment for covariates, we performed sibling analyses using within-strata estimation in Cox regression models, adjusted for sex, parity, birth season, GA at birth and cesarean section in which we discarded all only-child observations (n=131,335). Data management was done in SAS 9.4, data analyses in Stata/MP 15.1.

## *2.6 Sensitivity analyses*

Individuals with genetic disorders are at increased risk for both ASD/ID and repeated specialized care for infections during infancy and childhood<sup>35, 36</sup>. To address if such individuals were driving potential associations observed, we repeated the population and sibling comparison analyses after excluding individuals affected by a study outcome who were also diagnosed with a congenital disorder, as detailed in<sup>11</sup>.

### **3. Results**

#### *3.1 Description of the study population*

Of the 556,732 children included in our study (Figure S1), 282,595 (50.8%) had at least one registered infection before age 18. All investigated covariates differed significantly ( $p < 0.001$ ) between those who received specialized care for infection and those who did not (Table 1). Among those who were not diagnosed with an infection during childhood, 5,803 (2.1%) were diagnosed with ASD and 2,051 (0.7%) with ID. Among those diagnosed with one or more infection, 8,802 (3.1%) were diagnosed with ASD and 4,465 (1.6%) with ID (Table 1). As compared to 'ASD without ID' or 'ID without ASD', 'ASD with ID' was diagnosed at an earlier age (Figure 1B). CNS infections were rare as compared to non-CNS infections. Both types of infections were overrepresented in individuals with the investigated outcomes, particularly those involving ID (Figure 1C-D). Infections were more common during the first few years of childhood in all diagnostic groups (Figure 1E-G), particularly among those with an early diagnosis of 'ID without ASD' or 'ASD with ID', as compared to those without a diagnosis of ASD/ID. Infections were more common throughout childhood among children diagnosed with ID before age 12 as compared to those unaffected by ASD/ID or those diagnosed with 'ASD without ID'.

All included covariates were associated with ASD and age at ASD diagnosis. For example, ASD was more common among males and first-born children, particularly among those diagnosed before age 6 (Table S1). Similarly, obstetric complications, such as pre-term birth and cesarean section were associated with early diagnosis whereas parental psychiatric history was more common among cases diagnosed after age 12. Similar observations were made for ID (Table S2).

### *3.2 Main analyses*

#### *3.2.1 Overlapping diagnostic outcomes*

Inclusion of covariates in our fully adjusted models (Figure 2) attenuated risk estimates compared to crude models (Figure S2) but did not change the overall findings. In fully adjusted models, infections during childhood (0-<18 years) were associated with later ASD (HR 1.23, 95%CI 1.19-1.28) and ID diagnoses (HR 2.16, 95%CI 2.05-2.28). Infections at all age intervals except 6-<12 were associated with later ASD, with the strongest association observed for infections in the age interval 1-<3 (HR 1.26, 95%CI 1.21-1.31) Figure 2A.

Infections at all ages were associated with a later ID diagnosis, with the strongest association observed for infections during age 1-<3 (HR 2.12, 95%CI 2.00-2.24), Figure 2E.

Infections were associated with ASD diagnosed at age 0-<6 (HR 1.50, 95%CI 1.40-1.61, Figure 2B) and 6-<12 (HR 1.21, 95%CI 1.14-1.28, Figure 2C), but not 12-<18 (Figure 2D). Infections were associated with ID diagnosed throughout the follow-up, Figures 2F-H, with the strongest association during ages 0-<6 (HR 2.55, 95%CI 2.33-2.80, Figure 2F).

Sibling comparison models resulted in generally comparable associations to the total population analysis, see Figure 2I-P. Notably, infections during age 1-<3 exhibited a stronger

association with ASD diagnosed before age 12 (Figure 2J-K). Only infections 6-<12 were associated with ASD diagnosed at 12-<18 (Figure 2L). Infections at 12-<18 were not associated with ASD or ID.

### 3.2.2 Mutually exclusive outcomes

Similar results regarding the mutually exclusive outcome groups were observed in crude (Figure S3) and adjusted models (Figure 3). Infections during childhood were associated with the mutually exclusive diagnoses with the largest risk estimates observed for 'ID without ASD' (HR 2.56, 95%CI 2.39-2.75, Figure 3I) and the smallest for 'ASD without ID' (HR 1.15, 95%CI 1.10-1.19, Figure 3A). Whereas infections at all ages were associated with 'ID without ASD' and 'ASD with ID', only those at 1-<3 and 3-<6 were significantly associated with 'ASD without ID'. Larger point-estimates were observed among those diagnosed early. Infections were not associated with 'ASD without ID' diagnosed after age 12 (Figure 3B-D, F-H & J-L).

Sibling analyses resulted in slightly modified associations as compared to those observed in the general population, with enhanced risk for 'ASD without ID' (Figure 3M-P) and mostly attenuated risks for 'ASD with ID' (Figure 3Q-T) or 'ID without ASD' (Figure 3U-Y).

Associations with infections at 12-<18 were attenuated towards unity for all outcomes.

Higher risk estimates for 'ASD without ID' were largely driven by infections in the age intervals 1-<3 and 3-<6 (Figure 3M) among those diagnosed before age 6 (Figure 3N) or at 6-<12 (Figure 3O). Infections during the first year of life were associated with later 'ASD with ID' (Figure 3Q) with the largest risk estimate observed among those diagnosed before age 6 (Figure 3R). Infections before age 3 were no longer associated with 'ASD with ID' diagnosed at 12-<18, though elevated risks were observed for infections at 3-<6 and 6-<12 (HR 3.44, 95%CI 1.87-6.33, and HR 2.21, 95%CI 1.39-3.51), respectively, Figure 3T). Infections during

early childhood remained strongly associated with 'ID without ASD' (Figure 3U), particularly among those diagnosed before age 6 (Figure 3V) or at 6-<12 (Figure 3X), but also among those diagnosed at 12-<18 (Figure 3Y).

### *3.3 Sensitivity analyses*

Exclusion of children diagnosed with congenital disorders from the total population (Figure S4) and sibling (Figure S5) comparisons attenuated the associations observed between infections and the mutually exclusive diagnoses involving ID. In the sibling comparison, infections at 1-<3 and 3-<6 remained associated with 'ASD without ID'. Infections before age 12 remained associated with both 'ASD with ID' and 'ID without ASD' throughout the follow-up. Among those diagnosed at 12-<18, only infections before age 3 were associated with 'ID without ASD', whereas infections at 3-<6 and 6-<12 were associated with 'ASD with ID' (Figure S5).

### *3.4 CNS and non-CNS infections*

Larger risk estimates for all outcomes were generally observed for CNS infections than for non-CNS infections (Figure 4), e.g. risk for 'ID without ASD' was 3.66 (95% CI: 2.91-4.60, figure 4E) for CNS infections and 1.42 (95% CI: 1.27-1.58, figure 4F) for non-CNS infections.

## **4. Discussion**

In the present study, infections requiring specialized care between ages 1 and 6 were associated with an excess risk of 'ASD without ID' before age 12. Infections before age 12 were associated with 'ASD with ID' diagnosed throughout childhood. Similar observations were made for 'ID without ASD' with even larger point estimates, particularly for infections during the first few years of childhood. We noted larger risks for all outcomes associated

with CNS infections than for non-CNS infections, particularly for diagnoses involving ID.

While our analyses highlighted the potential for confounding by both familial factors and factors not shared between family members, they also indicate that childhood infections cannot be excluded in the etiologies of ASD/ID, including those diagnosed years later.

#### *4.1 Agreement with previous studies*

##### *4.1.1 Infections and ASD*

Long-term follow-up studies of individuals with congenital infections have reported autism-like behaviors<sup>37-39</sup>. There are, however, few published studies of the potential association between infections overall and ASD in the general population, none of which considered the potential differential effect on ASD with or without co-occurring ID or the potential for confounding by factors shared between family members.

Rosen et al. reported no association between inpatient, outpatient, emergency and referral visits for infections registered within the Kaiser-Permanente health plan during the first two years of life and later ASD in a sample consisting of 500 cases and 2000 matched controls<sup>30</sup>.

Atladottir et al., on the other hand, reported associations between hospitalization for infections during childhood and later ASD (HR 1.38) in a Danish population in 2010<sup>29</sup>, similar to those observed here (HR 1.23). More recently, Sabourin and co-workers reported that individuals with ASD were more likely to have had an infection during the first three years of life than comparison individuals using a combination of neonatal records and care-giver interviews prone to bias<sup>31</sup>. Our observations regarding CNS-infections agree with those reported by Pedersen et al.<sup>18</sup>, who found a weak association between CNS-infections and ASD that disappeared after exclusion of individuals with co-occurring ID, although we lacked power to distinguish between effects of CNS and non-CNS infections.

#### 4.1.2 Infections and ID

With regard to the observed risk for ID, our results are in agreement with a number of clinical follow-up studies of neonates with congenital infections<sup>13</sup> and CNS-infections<sup>40</sup> of both viral and bacterial origin during both infancy and childhood<sup>14-16, 41-43</sup>. In a study employing Danish population registers<sup>18</sup>, the authors explored a wide range of mental disorders in relation to previous CNS infections and reported a significant association (HR 3.29) between CNS-infections and later ID, similar to those observed here for 'ID without ASD' (HR 3.66) and for 'ASD with ID' (HR 3.21).

Fewer studies have focused on cognitive function after childhood infections not involving the CNS. Kariuki *et al.*, reported impairments in neurocognitive testing at age 7 in children who had been hospital treated for infections before age four<sup>44</sup>. Moreover, the Danish register study by Atladottir *et al.* reported an association between infections during childhood and later ID<sup>29</sup> of a magnitude similar to that observed here. Our current findings are in line with previous studies where both CNS and non-CNS infections during early childhood were associated with lower scores on a mandatory cognitive test administered to conscripts<sup>21, 45</sup>. Taken together with these previous studies, our current observations, suggest that infections without a recognized CNS involvement during childhood may contribute to impaired cognitive function.

#### 4.2 Strengths and weaknesses

Strengths of the study include the large, population-based cohort with prospectively collected data in a setting with universal access to health care with minimal risk for selection or recall bias. Though all clinical pathways to ASD/ID diagnosis are covered, we do not have access to individual assessments of cognitive function. Thus, children diagnosed as 'ASD

without ID' could experience impairments that do not meet criteria for ID. Despite the large study population, we had limited power to analyze specific infections.

Potential confounding was addressed in multiple ways. We adjusted our total-population estimates for wide range of potentially confounding factors. We used sibling comparison models to account for residual confounding by factors shared between siblings and adjusted for factors not necessarily shared between siblings (e.g. pre-term birth). Sensitivity analyses indicated that the associations observed were not fully explained by higher risk of infection among children with congenital diagnoses, although confounding by undiagnosed genetic abnormalities cannot be excluded.

Age at diagnosis depends on a wide range of factors such as clinical practice, severity and parental vigilance <sup>1</sup>. Results from studies using register-based data to investigate timing of an exposure in relation to the onset of a childhood disorder should therefore be interpreted cautiously. While infections during early childhood were more common in children diagnosed with ASD or ID than in children without such diagnoses, we also note that the group diagnosed with ID at a young age had an excess of infections throughout childhood. Such an excess of infections in the group of individuals diagnosed with ASD without ID, however, seemed to be limited to the first decade of life. While these groups contain individuals with underlying hereditary and non-hereditary causes of ASD/ID that also receive more specialized care for infections during childhood (as addressed by our total population, sibling and sensitivity analyses), this does not preclude infections from being casually linked to some cases of ID and ASD. A higher liability to infections would increase the risk of contracting an infection requiring specialized care during early childhood, arguably a period of development more sensitive to external insults than later parts of childhood. It should

also be kept in mind that our exposure definition is broad and encompasses a range of infectious agents, each with a prevalence and consequences that are likely to vary by age<sup>46</sup>, which complicates comparisons of effects of exposures at different ages.

#### *4.3 Potential mechanisms underlying the observed associations.*

A number of different, but not mutually exclusive, mechanisms can potentially explain our current observations. Targeting of the nervous system by infectious agents during periods of rapid growth and plasticity<sup>47</sup> is one plausible mechanism. Complications involving the CNS are, albeit rarely, observed for many different infectious agents, including those not normally associated with neuro-invasion, particularly in young individuals<sup>46, 48, 49</sup>. For example, respiratory viruses (e.g. influenza A virus, respiratory syncytial virus) are important causes of encephalitis in young children<sup>50</sup>. It is possible that for some registered infections, mild CNS-involvement may have gone unnoticed or undocumented, as has been reported for uncomplicated measles virus infections<sup>51</sup>. Moreover, low levels of several molecules involved in the innate immune response at birth were recently associated with ASD risk<sup>24</sup>, supporting the notion of an increased vulnerability to infections in ASD that was proposed based on a genetic correlation between hospitalization for infection and ASD<sup>52</sup>. Further studies are needed to investigate if deficits in immunity associated with ASD risk could contribute to more severe outcomes of childhood infections, such as ID, and potentially contribute to the high co-morbidity between these disorders<sup>53</sup>.

Direct support for immune mediated influences on the developing CNS resulting in cognitive and behavioral abnormalities is provided by experimental studies. Various protocols mimicking viral or bacterial infections during certain stages of development produce persistent cognitive deficits and behavioral changes in adult animals reminiscent of those

observed in individuals with ASD <sup>54</sup>. Moreover, transient treatment with interferon- $\gamma$  during neuronal differentiation of human stem cells was reported to cause induction of ASD-associated genes <sup>55</sup>. While studies of individuals diagnosed with ASD support activation of the immune system in ASD, the pathogenetic role of such immune activity remains unclear <sup>56-59</sup>. Nevertheless, many of the pathways activated by infections have pleiotropic effects during brain development <sup>60</sup>. Thus, the association between infections during periods of rapid growth and plasticity of the brain and later diagnosis of ASD/ID may be explained by components of the immune system interfering with the normal trajectories of brain development.

Infections cause changes in the distribution of both macro- and micronutrients in the host, with the purpose of promoting the function of the immune system while limiting pathogen proliferation <sup>61, 62</sup>. Indeed, febrile episodes during childhood have been associated with reduced adult height, independent of genetic background and nutritional intake <sup>63</sup>. Brain development during the first years of life is characterized by rapid, sequential patterns of cellular differentiation, migration and synapse formation during which the demand for macro- and micronutrients is high <sup>64</sup>. Diversion of macro- and micronutrients necessary to mount an adequate immune response and clear an infection may explain part of our observed association between infections during childhood and later ID/ASD <sup>65, 66</sup>.

#### *4.4 Conclusions*

Infections at an early age, appear to confer risk for a later diagnosis of ASD involving ID. Our observation that infections during early childhood exhibit weak but significant associations with 'ASD without ID' requires replication in future studies. Pending independent replication, future studies should focus on identifying specific infectious agents, their mechanisms of

action, and individuals at risk. Such information will be critically important for guiding the development of preventive strategies.

## 5. References

1. Shulman C, Esler A, Morrier MJ, Rice CE. Diagnosis of Autism Spectrum Disorder Across the Lifespan. *Child Adolesc Psychiatr Clin N Am*. 2020;29(2):253-73.  
10.1016/j.chc.2020.01.001
2. Ozonoff S, Iosif AM. Changing conceptualizations of regression: What prospective studies reveal about the onset of autism spectrum disorder. *Neurosci Biobehav Rev*. 2019;100:296-304. 10.1016/j.neubiorev.2019.03.012
3. Stoner R, Chow ML, Boyle MP, Sunkin SM, Mouton PR, Roy S, et al. Patches of disorganization in the neocortex of children with autism. *N Engl J Med*. 2014;370(13):1209-19. 10.1056/NEJMoa1307491
4. Gaugler T, Klei L, Sanders SJ, Bodea CA, Goldberg AP, Lee AB, et al. Most genetic risk for autism resides with common variation. *Nat Genet*. 2014;46(8):881-5.  
10.1038/ng.3039
5. Grove J, Ripke S, Als TD, Mattheisen M, Walters RK, Won H, et al. Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet*. 2019;51(3):431-44. 10.1038/s41588-019-0344-8
6. Iossifov I, O'Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, et al. The contribution of de novo coding mutations to autism spectrum disorder. *Nature*. 2014;515(7526):216-21. 10.1038/nature13908
7. Lee BK, Magnusson C, Gardner RM, Blomstrom A, Newschaffer CJ, Burstyn I, et al. Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun*. 2015;44:100-5. 10.1016/j.bbi.2014.09.001
8. Maenner MJ, Shaw KA, Baio J, EdS, Washington A, Patrick M, et al. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years - Autism and Developmental

Disabilities Monitoring Network, 11 Sites, United States, 2016. MMWR Surveill Summ.

2020;69(4):1-12. 10.15585/mmwr.ss6904a1

9. Robinson EB, Samocha KE, Kosmicki JA, McGrath L, Neale BM, Perlis RH, et al.

Autism spectrum disorder severity reflects the average contribution of de novo and familial influences. Proc Natl Acad Sci U S A. 2014;111(42):15161-5. 10.1073/pnas.1409204111

10. Xie S, Karlsson H, Dalman C, Widman L, Rai D, Gardner RM, et al. Family History of Mental and Neurological Disorders and Risk of Autism. JAMA Netw Open.

2019;2(3):e190154. 10.1001/jamanetworkopen.2019.0154

11. Gardner RM, Dalman C, Rai D, Lee BK, Karlsson H. The Association of Paternal IQ With Autism Spectrum Disorders and Its Comorbidities: A Population-Based Cohort Study.

J Am Acad Child Adolesc Psychiatry. 2020;59(3):410-21. 10.1016/j.jaac.2019.04.004

12. Johnson RT. Effects of viral infection on the developing nervous system. N Engl J Med. 1972;287(12):599-604. 10.1056/NEJM197209212871208

13. Ostrander B, Bale JF. Congenital and perinatal infections. Handb Clin Neurol. 2019;162:133-53. 10.1016/B978-0-444-64029-1.00006-0

14. Chamberlain RN, Christie PN, Holt KS, Huntley RM, Pollard R, Roche MC. A study of school children who had identified virus infections of the central nervous system during infancy. Child: care, health and development. 1983;9(1):29-47.

15. Chang LY, Huang LM, Gau SS, Wu YY, Hsia SH, Fan TY, et al. Neurodevelopment and cognition in children after enterovirus 71 infection. N Engl J Med. 2007;356(12):1226-34. 10.1056/NEJMoa065954

16. Michaeli O, Kassis I, Shachor-Meyouhas Y, Shahar E, Ravid S. Long-term motor and cognitive outcome of acute encephalitis. Pediatrics. 2014;133(3):e546-52.

10.1542/peds.2013-3010

17. Dalman C, Allebeck P, Gunnell D, Harrison G, Kristensson K, Lewis G, et al. Infections in the CNS during childhood and the risk of subsequent psychotic illness: a cohort study of more than one million Swedish subjects. *Am J Psychiatry*. 2008;165(1):59-65.
18. Pedersen EMJ, Kohler-Forsberg O, Nordentoft M, Christensen RHB, Mortensen PB, Petersen L, et al. Infections of the central nervous system as a risk factor for mental disorders and cognitive impairment: a nationwide register-based study. *Brain Behav Immun*. 2020. [10.1016/j.bbi.2020.04.072](https://doi.org/10.1016/j.bbi.2020.04.072)
19. Blomstrom A, Karlsson H, Svensson A, Frisell T, Lee BK, Dal H, et al. Hospital Admission With Infection During Childhood and Risk for Psychotic Illness--A Population-based Cohort Study. *Schizophr Bull*. 2013. sbt195 [pii]  
[10.1093/schbul/sbt195](https://doi.org/10.1093/schbul/sbt195)
20. Nielsen PR, Benros ME, Mortensen PB. Hospital contacts with infection and risk of schizophrenia: a population-based cohort study with linkage of Danish national registers. *Schizophr Bull*. 2014;40(6):1526-32. [10.1093/schbul/sbt200](https://doi.org/10.1093/schbul/sbt200)
21. Khandaker GM, Dalman C, Kappelmann N, Stochl J, Dal H, Kosidou K, et al. Association of Childhood Infection With IQ and Adult Nonaffective Psychosis in Swedish Men: A Population-Based Longitudinal Cohort and Co-relative Study. *JAMA Psychiatry*. 2018. [10.1001/jamapsychiatry.2017.4491](https://doi.org/10.1001/jamapsychiatry.2017.4491)
22. Benros ME, Nielsen PR, Nordentoft M, Eaton WW, Dalton SO, Mortensen PB. Autoimmune diseases and severe infections as risk factors for schizophrenia: a 30-year population-based register study. *Am J Psychiatry*. 2011;168(12):1303-10. [10.1176/appi.ajp.2011.11030516](https://doi.org/10.1176/appi.ajp.2011.11030516)

23. Gardner RM, Dalman C, Wicks S, Lee BK, Karlsson H. Neonatal levels of acute phase proteins and later risk of non-affective psychosis. *Transl Psychiatry*. 2013;3:e228. 10.1038/tp.2013.5
24. Gardner RM, Lee BK, Brynne M, Sjoqvist H, Dalman C, Karlsson H. Neonatal Levels of Acute Phase Proteins and Risk of Autism Spectrum Disorder. *Biol Psychiatry*. 2021;89(5):463-75. 10.1016/j.biopsych.2020.09.005
25. Blomstrom A, Gardner RM, Dalman C, Yolken RH, Karlsson H. Influence of maternal infections on neonatal acute phase proteins and their interaction in the development of non-affective psychosis. *Transl Psychiatry*. 2015;5:e502. 10.1038/tp.2014.142
26. Hartwig FP, Borges MC, Horta BL, Bowden J, Davey Smith G. Inflammatory Biomarkers and Risk of Schizophrenia: A 2-Sample Mendelian Randomization Study. *JAMA Psychiatry*. 2017;74(12):1226-33. 10.1001/jamapsychiatry.2017.3191
27. Ligthart S, Vaez A, Vosa U, Stathopoulou MG, de Vries PS, Prins BP, et al. Genome Analyses of >200,000 Individuals Identify 58 Loci for Chronic Inflammation and Highlight Pathways that Link Inflammation and Complex Disorders. *Am J Hum Genet*. 2018;103(5):691-706. 10.1016/j.ajhg.2018.09.009
28. Atladottir HO, Henriksen TB, Schendel DE, Parner ET. Autism after infection, febrile episodes, and antibiotic use during pregnancy: an exploratory study. *Pediatrics*. 2012;130(6):e1447-54. 10.1542/peds.2012-1107
29. Atladottir HO, Thorsen P, Schendel DE, Ostergaard L, Lemcke S, Parner ET. Association of hospitalization for infection in childhood with diagnosis of autism spectrum disorders: a Danish cohort study. *Arch Pediatr Adolesc Med*. 2010;164(5):470-7. 10.1001/archpediatrics.2010.9

30. Rosen NJ, Yoshida CK, Croen LA. Infection in the first 2 years of life and autism spectrum disorders. *Pediatrics*. 2007;119(1):e61-9. 10.1542/peds.2006-1788
31. Sabourin KR, Reynolds A, Schendel D, Rosenberg S, Croen LA, Pinto-Martin JA, et al. Infections in children with autism spectrum disorder: Study to Explore Early Development (SEED). *Autism Res*. 2019;12(1):136-46. 10.1002/aur.2012
32. Idring S, Rai D, Dal H, Dalman C, Sturm H, Zander E, et al. Autism spectrum disorders in the Stockholm Youth Cohort: design, prevalence and validity. *PLoS One*. 2012;7(7):e41280. 10.1371/journal.pone.0041280
33. Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, Ekblom A. The Swedish personal identity number: possibilities and pitfalls in healthcare and medical research. *Eur J Epidemiol*. 2009;24(11):659-67. 10.1007/s10654-009-9350-y
34. Mackay E, Dalman C, Karlsson H, Gardner RM. Association of Gestational Weight Gain and Maternal Body Mass Index in Early Pregnancy With Risk for Nonaffective Psychosis in Offspring. *JAMA Psychiatry*. 2017. 10.1001/jamapsychiatry.2016.4257
35. Sanchez-Luna M, Medrano C, Lirio J, Group R-S. Down syndrome as risk factor for respiratory syncytial virus hospitalization: A prospective multicenter epidemiological study. *Influenza Other Respir Viruses*. 2017;11(2):157-64. 10.1111/irv.12431
36. Soorya L, Kolevzon A, Zweifach J, Lim T, Dobry Y, Schwartz L, et al. Prospective investigation of autism and genotype-phenotype correlations in 22q13 deletion syndrome and SHANK3 deficiency. *Mol Autism*. 2013;4(1):18. 10.1186/2040-2392-4-18
37. Ivarsson SA, Bjerre I, Vegfors P, Ahlfors K. Autism as one of several disabilities in two children with congenital cytomegalovirus infection. *Neuropediatrics*. 1990;21(2):102-3. 10.1055/s-2008-1071471


38. Chess S, Fernandez P, Korn S. Behavioral consequences of congenital rubella. *The Journal of pediatrics*. 1978;93(4):699-703. 10.1016/s0022-3476(78)80921-4
39. Deykin EY, MacMahon B. Viral exposure and autism. *Am J Epidemiol*. 1979;109(6):628-38. 10.1093/oxfordjournals.aje.a112726
40. Christie D, Rashid H, El-Bashir H, Sweeney F, Shore T, Booy R, et al. Impact of meningitis on intelligence and development: A systematic review and meta-analysis. *PLoS One*. 2017;12(8):e0175024. 10.1371/journal.pone.0175024
41. Engman ML, Adolfsson I, Lewensohn-Fuchs I, Forsgren M, Mosskin M, Malm G. Neuropsychologic outcomes in children with neonatal herpes encephalitis. *Pediatr Neurol*. 2008;38(6):398-405. 10.1016/j.pediatrneurol.2008.02.005
42. Pace D, Pollard AJ. Meningococcal disease: clinical presentation and sequelae. *Vaccine*. 2012;30 Suppl 2:B3-9. 10.1016/j.vaccine.2011.12.062
43. Grimwood K, Anderson P, Anderson V, Tan L, Nolan T. Twelve year outcomes following bacterial meningitis: further evidence for persisting effects. *Arch Dis Child*. 2000;83(2):111-6. 10.1136/adc.83.2.111
44. Kariuki M, Raudino A, Green MJ, Laurens KR, Dean K, Brinkman SA, et al. Hospital admission for infection during early childhood influences developmental vulnerabilities at age 5 years. *J Paediatr Child Health*. 2016;52(9):882-8. 10.1111/jpc.13239
45. Benros ME, Sorensen HJ, Nielsen PR, Nordentoft M, Mortensen PB, Petersen L. The Association between Infections and General Cognitive Ability in Young Men - A Nationwide Study. *PLoS One*. 2015;10(5):e0124005. 10.1371/journal.pone.0124005
46. Fowler A, Stodberg T, Eriksson M, Wickstrom R. Childhood encephalitis in Sweden: etiology, clinical presentation and outcome. *Eur J Paediatr Neurol*. 2008;12(6):484-90. 10.1016/j.ejpn.2007.12.009

47. Semple BD, Blomgren K, Gimlin K, Ferriero DM, Noble-Haeusslein LJ. Brain development in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across species. *Prog Neurobiol.* 2013;106-107:1-16.  
10.1016/j.pneurobio.2013.04.001
48. Boucher A, Herrmann JL, Morand P, Buzele R, Crabol Y, Stahl JP, et al. Epidemiology of infectious encephalitis causes in 2016. *Med Mal Infect.* 2017;47(3):221-35.  
10.1016/j.medmal.2017.02.003
49. Venkatesan A, Jagdish B. Imaging in Encephalitis. *Semin Neurol.* 2019;39(3):312-21. 10.1055/s-0039-1687838
50. Beattie GC, Glaser CA, Sheriff H, Messenger S, Preas CP, Shahkarami M, et al. Encephalitis with thalamic and basal ganglia abnormalities: etiologies, neuroimaging, and potential role of respiratory viruses. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America.* 2013;56(6):825-32. 10.1093/cid/cis990
51. Hanninen P, Arstila P, Lang H, Salmi A, Panelius M. Involvement of the central nervous system in acute, uncomplicated measles virus infection. *J Clin Microbiol.* 1980;11(6):610-3.
52. Nudel R, Wang Y, Appadurai V, Schork AJ, Buil A, Agerbo E, et al. A large-scale genomic investigation of susceptibility to infection and its association with mental disorders in the Danish population. *Transl Psychiatry.* 2019;9(1):283. 10.1038/s41398-019-0622-3
53. Thurm A, Farmer C, Salzman E, Lord C, Bishop S. State of the Field: Differentiating Intellectual Disability From Autism Spectrum Disorder. *Front Psychiatry.* 2019;10:526. 10.3389/fpsy.2019.00526

54. Mora S, Martin-Gonzalez E, Flores P, Moreno M. Neuropsychiatric consequences of childhood group A streptococcal infection: A systematic review of preclinical models. *Brain Behav Immun.* 2020;86:53-62. 10.1016/j.bbi.2019.02.027
55. Warre-Cornish K, Perfect L, Nagy R, Duarte RRR, Reid MJ, Raval P, et al. Interferon-gamma signaling in human iPSC-derived neurons recapitulates neurodevelopmental disorder phenotypes. *Sci Adv.* 2020;6(34):eaay9506. 10.1126/sciadv.aay9506
56. Gandal MJ, Haney JR, Parikshak NN, Leppa V, Ramaswami G, Hartl C, et al. Shared molecular neuropathology across major psychiatric disorders parallels polygenic overlap. *Science.* 2018;359(6376):693-7. 10.1126/science.aad6469
57. Nazeen S, Palmer NP, Berger B, Kohane IS. Integrative analysis of genetic data sets reveals a shared innate immune component in autism spectrum disorder and its comorbidities. *Genome biology.* 2016;17(1):228. 10.1186/s13059-016-1084-z
58. Voineagu I, Wang X, Johnston P, Lowe JK, Tian Y, Horvath S, et al. Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature.* 2011;474(7351):380-4. <http://www.nature.com/nature/journal/v474/n7351/abs/10.1038-nature10110-unlocked.html#supplementary-information>
59. Ashwood P, Wills S, Van de Water J. The immune response in autism: a new frontier for autism research. *J Leukoc Biol.* 2006;80(1):1-15. 10.1189/jlb.1205707
60. Filiano AJ, Xu Y, Tustison NJ, Marsh RL, Baker W, Smirnov I, et al. Unexpected role of interferon-gamma in regulating neuronal connectivity and social behaviour. *Nature.* 2016;535(7612):425-9. 10.1038/nature18626
61. Calder PC. Feeding the immune system. *Proc Nutr Soc.* 2013;72(3):299-309. 10.1017/S0029665113001286


62. Powanda MC, Beisel WR. Metabolic effects of infection on protein and energy status. *J Nutr*. 2003;133(1):322S-7S. [10.1093/jn/133.1.322S](https://doi.org/10.1093/jn/133.1.322S)
63. Hwang AE, Mack TM, Hamilton AS, Gauderman WJ, Bernstein L, Cockburn MG, et al. Childhood infections and adult height in monozygotic twin pairs. *Am J Epidemiol*. 2013;178(4):551-8. [10.1093/aje/kwt012](https://doi.org/10.1093/aje/kwt012)
64. Cusick SE, Georgieff MK. The Role of Nutrition in Brain Development: The Golden Opportunity of the "First 1000 Days". *The Journal of pediatrics*. 2016;175:16-21. [10.1016/j.jpeds.2016.05.013](https://doi.org/10.1016/j.jpeds.2016.05.013)
65. Lee BK, Eyles DW, Magnusson C, Newschaffer CJ, McGrath JJ, Kvaskoff D, et al. Developmental vitamin D and autism spectrum disorders: findings from the Stockholm Youth Cohort. *Mol Psychiatry*. 2019. [10.1038/s41380-019-0578-y](https://doi.org/10.1038/s41380-019-0578-y)
66. Wieggersma AM, Dalman C, Lee BK, Karlsson H, Gardner RM. Association of Prenatal Maternal Anemia With Neurodevelopmental Disorders. *JAMA Psychiatry*. 2019:1-12. [10.1001/jamapsychiatry.2019.2309](https://doi.org/10.1001/jamapsychiatry.2019.2309)

## Figures and legends




**Figure 1. Description of outcomes and exposures in the study population.** Individuals diagnosed with ASD and/or ID identified in the final study population (n=556,732) as of

December 31, 2016 (A). Age at first diagnosis of ASD/ID for the mutually exclusive diagnostic groups (B). Incidence of CNS (C) and non-CNS infections (D) according to age among children unaffected by ASD/ID and among those diagnosed with 'ASD without ID', 'ASD with ID', 'ID without ASD'. Note the different scales of the y-axes in C and D. Incidence of one or more infections according to age among children unaffected by ASD/ID and those diagnosed with 'ASD without ID' (E), 'ASD with ID' (F) and 'ID without ASD' (G), throughout childhood and stratified by age at diagnosis of ASD/ID.




**Figure 2. Infections during childhood and later diagnosis of ASD or ID.** Associations between specialized care for infections during childhood and later, non-mutually exclusive, diagnosis of ASD and ID. Associations between exposure at different age intervals and diagnoses at different age intervals are also shown. Comparison between unrelated individuals in the general population are shown (A-H) and comparisons between full biological siblings (I-P). Hazard ratios presented here are from fully adjusted models. Population-based estimates (A-H) are adjusted for sex, parity, maternal body mass index, pre-eclampsia, parental age, education, income, region of origin, histories of psychiatric

illness and infections, season of birth, gestational age at birth, size for gestational age, cesarean section, Apgar score. Estimates from the sibling analyses (I-P) are adjusted for sex, parity, gestational age at birth and cesarean section.



**Figure 3. Infections during childhood and mutually exclusive diagnoses.** Associations between specialized care for infections during childhood and later diagnosis of ‘ASD without ID’, ‘ASD with ID’ or ‘ID without ASD’. Associations between exposure at different age intervals and diagnoses at different age intervals are also shown. Comparisons between unrelated individuals in the general population (A-L) and between full biological siblings (M-X) are shown. Hazard ratios presented here are from fully adjusted models. Population-

based estimates (A-L) are adjusted for sex, parity, maternal body mass index, pre-eclampsia, parental age, education, income, region of origin, histories of psychiatric illness and infections, season of birth, gestational age at birth, size for gestational age, cesarean section, Apgar score. Estimates from the sibling analyses (M-X) are adjusted for sex, parity, gestational age at birth and cesarean section.




**Figure 4. CNS/non-CNS infections and mutually exclusive diagnostic groups.** Associations between specialized care for CNS infections (left) or non-CNS infections (right) during childhood and later diagnosis of ‘ASD without ID’ (A-B), ‘ASD with ID’ (C-D) or ‘ID without ASD’ (E-F). Associations between exposure at different age intervals and later diagnoses are also shown. Hazard ratios are adjusted for sex, parity, maternal body mass index, pre-eclampsia, parental age, education, income, region of origin, histories of psychiatric illness


and infections, season of birth, gestational age at birth, size for gestational age, cesarean section and Apgar score.

## Supplementary material

*Infections in childhood and subsequent diagnosis of autism spectrum disorders and intellectual disability: a nation-wide register-based cohort study*




**Supplementary figure S1.** Flow chart delineating the derivation of the populations included in the present study.




### Supplementary figure S2. Infections during childhood and later diagnosis of ASD or ID.

Crude associations between specialized care for infections and later, non-mutually exclusive, diagnosis of ASD and ID. Associations between exposure and diagnoses at different age interval are also shown. Comparison between unrelated individuals in the general population are shown (A-H) and comparisons between full biological siblings (I-P).




### Supplementary figure S3. Infections during childhood and mutually exclusive diagnoses.

Crude associations between infections between birth and age 18 and later diagnosis of ‘ASD without ID’, ‘ASD with ID’ or ‘ID without ASD’. Associations between exposures and diagnoses at different ages are also shown. Comparisons between unrelated individuals in the general population (A-L) and between full biological siblings (M-X) are shown.



**Supplementary figure S4. Infections during childhood and mutually exclusive diagnoses-sensitivity analyses.** Associations between infections between birth and age 18 and later diagnosis of ‘ASD without ID’, ‘ASD with ID’ or ‘ID without ASD’ in the total population following exclusion of individuals with congenital diagnoses and ASD/ID. Associations between exposures and diagnoses at different ages are also shown. Estimates are adjusted for sex, parity, maternal body mass index, pre-eclampsia, parental age, education, income, region of origin, histories of psychiatric illness and infections, season of birth, gestational age at birth, size for gestational age, cesarean section and Apgar score.



**Supplementary figure S5. Infections during childhood and mutually exclusive diagnoses-sensitivity analyses.** Associations between infections between birth and age 18 and later diagnosis of 'ASD without ID', 'ASD with ID' or 'ID without ASD' among biological siblings following exclusion of individuals with congenital diagnoses and ASD/ID. Associations between exposures and diagnoses at different ages are also shown. Estimates from are adjusted for sex, parity, gestational age at birth and cesarean section.

**Supplementary Table S1. Association between covariates and ASD and by age at first diagnosis.**

		<b>No ASD</b>	<b>ASD</b>	<b>p-value</b>	<b>0-&lt;6 years</b>	<b>6-&lt;12 years</b>	<b>12-&lt;18 years</b>	<b>p-value</b>
<b>N</b>		542127	14605		3704	5845	5056	
<b>Sex</b>	Male	275164 (50.8%)	10290 (70.5%)	<0.001	2870 (77.5%)	4446 (76.1%)	2974 (58.8%)	<0.001
	Female	266963 (49.2%)	4315 (29.5%)		834 (22.5%)	1399 (23.9%)	2082 (41.2%)	
<b>Maternal BMI</b>	Underweight	14729 (2.7%)	371 (2.5%)	<0.001	100 (2.7%)	144 (2.5%)	127 (2.5%)	<0.001
	Normal	274989 (50.7%)	6885 (47.1%)		1900 (51.3%)	2672 (45.7%)	2313 (45.7%)	
	Overweight	78155 (14.4%)	2600 (17.8%)		705 (19.0%)	1063 (18.2%)	832 (16.5%)	
	Obese	26353 (4.9%)	1217 (8.3%)		338 (9.1%)	531 (9.1%)	348 (6.9%)	
	Missing	147901 (27.3%)	3532 (24.2%)		661 (17.8%)	1435 (24.6%)	1436 (28.4%)	
<b>Birth order</b>	First	244369 (45.1%)	7051 (48.3%)	<0.001	1868 (50.4%)	2876 (49.2%)	2307 (45.6%)	<0.001
	Second	196328 (36.2%)	4832 (33.1%)		1196 (32.3%)	1906 (32.6%)	1730 (34.2%)	
	≥Third	101430 (18.7%)	2722 (18.6%)		640 (17.3%)	1063 (18.2%)	1019 (20.2%)	
<b>Family income</b>	Low (0<20%)	77873 (14.4%)	2024 (13.9%)	<0.001	646 (17.4%)	731 (12.5%)	647 (12.8%)	<0.001
	20<40%	113692 (21.0%)	3720 (25.5%)		933 (25.2%)	1503 (25.7%)	1284 (25.4%)	
	40<60%	117331 (21.6%)	3484 (23.9%)		825 (22.3%)	1375 (23.5%)	1284 (25.4%)	
	60<80%	116754 (21.5%)	2966 (20.3%)		674 (18.2%)	1247 (21.3%)	1045 (20.7%)	
	High (80-100%)	115761 (21.4%)	2392 (16.4%)		618 (16.7%)	982 (16.8%)	792 (15.7%)	
	Missing	716 (0.1%)	19 (0.1%)		8 (0.2%)	7 (0.1%)	4 (0.1%)	
<b>Maternal age, years</b>	<25	80391 (14.8%)	2301 (15.8%)	<0.001	504 (13.6%)	903 (15.4%)	894 (17.7%)	<0.001
	25-29	159722 (29.5%)	4199 (28.8%)		994 (26.8%)	1695 (29.0%)	1510 (29.9%)	
	30-34	188569 (34.8%)	4781 (32.7%)		1211 (32.7%)	1955 (33.4%)	1615 (31.9%)	
	35-39	93935 (17.3%)	2662 (18.2%)		783 (21.1%)	1040 (17.8%)	839 (16.6%)	
	>39	19510 (3.6%)	662 (4.5%)		212 (5.7%)	252 (4.3%)	198 (3.9%)	
<b>Paternal age, years</b>	<25	36915 (6.9%)	1147 (7.9%)	<0.001	205 (5.6%)	462 (8.0%)	480 (9.6%)	<0.001
	25-29	117557 (21.9%)	3112 (21.6%)		676 (18.6%)	1263 (21.8%)	1173 (23.4%)	
	30-34	180109 (33.6%)	4601 (31.9%)		1127 (31.0%)	1904 (32.9%)	1570 (31.4%)	
	35-39	124697 (23.2%)	3227 (22.4%)		904 (24.9%)	1275 (22.0%)	1048 (20.9%)	
	>39	77233 (14.4%)	2343 (16.2%)		718 (19.8%)	891 (15.4%)	734 (14.7%)	
<b>Birth season</b>	Dec-Feb	127267 (23.5%)	3370 (23.1%)	<0.001	821 (22.2%)	1402 (24.0%)	1147 (22.7%)	<0.001

<b>Pre-eclampsia</b>	Mar-May	146735 (27.1%)	3690 (25.3%)		912 (24.6%)	1423 (24.3%)	1355 (26.8%)	
	Jun-Aug	140652 (25.9%)	3809 (26.1%)		996 (26.9%)	1483 (25.4%)	1330 (26.3%)	
	Sep-Nov	127473 (23.5%)	3736 (25.6%)		975 (26.3%)	1537 (26.3%)	1224 (24.2%)	
	No	527700 (97.3%)	14057 (96.2%)	<0.001	3542 (95.6%)	5633 (96.4%)	4882 (96.6%)	<0.001
	Yes	14427 (2.7%)	548 (3.8%)		162 (4.4%)	212 (3.6%)	174 (3.4%)	
<b>Parental education</b>	<10 years	33479 (6.2%)	1014 (6.9%)	<0.001	234 (6.3%)	369 (6.3%)	411 (8.1%)	<0.001
	10-12	206592 (38.1%)	6208 (42.5%)		1292 (34.9%)	2537 (43.4%)	2379 (47.1%)	
	>12	300631 (55.5%)	7357 (50.4%)		2165 (58.5%)	2931 (50.1%)	2261 (44.7%)	
	Missing	1425 (0.3%)	26 (0.2%)		13 (0.4%)	8 (0.1%)	5 (0.1%)	
<b>Size for GA</b>	Small	12569 (2.3%)	540 (3.7%)	<0.001	177 (4.8%)	203 (3.5%)	160 (3.2%)	<0.001
	Normal	495982 (91.5%)	12942 (88.6%)		3237 (87.4%)	5186 (88.7%)	4519 (89.4%)	
	Large	16030 (3.0%)	592 (4.1%)		153 (4.1%)	240 (4.1%)	199 (3.9%)	
	Missing	17546 (3.2%)	531 (3.6%)		137 (3.7%)	216 (3.7%)	178 (3.5%)	
<b>Low Apgar score (&lt;7)</b>	No	530639 (97.9%)	14194 (97.2%)	<0.001	3585 (96.8%)	5693 (97.4%)	4916 (97.2%)	<0.001
	Yes	4856 (0.9%)	223 (1.5%)		85 (2.3%)	78 (1.3%)	60 (1.2%)	
	Missing	6632 (1.2%)	188 (1.3%)		34 (0.9%)	74 (1.3%)	80 (1.6%)	
<b>Cesarean delivery</b>	No	488835 (90.2%)	12764 (87.4%)	<0.001	3126 (84.4%)	5091 (87.1%)	4547 (89.9%)	<0.001
	Yes	35746 (6.6%)	1310 (9.0%)		441 (11.9%)	538 (9.2%)	331 (6.5%)	
	Missing	17546 (3.2%)	531 (3.6%)		137 (3.7%)	216 (3.7%)	178 (3.5%)	
<b>GA</b>	Preterm	30044 (5.5%)	1141 (7.8%)	<0.001	343 (9.3%)	459 (7.9%)	339 (6.7%)	<0.001
	Term	507893 (93.7%)	13338 (91.3%)		3339 (90.1%)	5343 (91.4%)	4656 (92.1%)	
	Post term	3279 (0.6%)	96 (0.7%)		20 (0.5%)	32 (0.5%)	44 (0.9%)	
	Missing	911 (0.2%)	30 (0.2%)		2 (0.1%)	11 (0.2%)	17 (0.3%)	
<b>Paternal psychiatric diagnosis</b>	No	395044 (72.9%)	8982 (61.5%)	<0.001	2405 (64.9%)	3557 (60.9%)	3020 (59.7%)	<0.001
	Yes	147083 (27.1%)	5623 (38.5%)		1299 (35.1%)	2288 (39.1%)	2036 (40.3%)	
<b>Maternal psychiatric diagnosis</b>	No	309103 (57.0%)	5492 (37.6%)	<0.001	1643 (44.4%)	2067 (35.4%)	1782 (35.2%)	<0.001

<b>Paternal infection</b>	Yes	233024 (43.0%)	9113 (62.4%)		2061 (55.6%)	3778 (64.6%)	3274 (64.8%)	
	No	313856 (57.9%)	7824 (53.6%)	<0.001	1899 (51.3%)	3078 (52.7%)	2847 (56.3%)	<0.001
<b>Maternal infection</b>	Yes	228271 (42.1%)	6781 (46.4%)		1805 (48.7%)	2767 (47.3%)	2209 (43.7%)	
	No	216953 (40.0%)	4724 (32.3%)	<0.001	1052 (28.4%)	1806 (30.9%)	1866 (36.9%)	<0.001
<b>Mother's region of origin</b>	Yes	325174 (60.0%)	9881 (67.7%)		2652 (71.6%)	4039 (69.1%)	3190 (63.1%)	
	Sweden	397154 (73.3%)	10919 (74.8%)	<0.001	2359 (63.7%)	4546 (77.8%)	4014 (79.4%)	<0.001
	Nordics	14984 (2.8%)	451 (3.1%)		58 (1.6%)	186 (3.2%)	207 (4.1%)	
	West Eu and NA	23954 (4.4%)	552 (3.8%)		178 (4.8%)	215 (3.7%)	159 (3.1%)	
	Africa	24973 (4.6%)	642 (4.4%)		332 (9.0%)	181 (3.1%)	129 (2.6%)	
	Asia and Oceania	11384 (2.1%)	278 (1.9%)		146 (3.9%)	77 (1.3%)	55 (1.1%)	
	Middle-East	45453 (8.4%)	969 (6.6%)		380 (10.3%)	352 (6.0%)	237 (4.7%)	
	East Eu and Russia	7262 (1.3%)	200 (1.4%)		65 (1.8%)	75 (1.3%)	60 (1.2%)	
	South America	11315 (2.1%)	417 (2.9%)		112 (3.0%)	161 (2.8%)	144 (2.8%)	
	Unknown/Missing	5648 (1.0%)	177 (1.2%)		74 (2.0%)	52 (0.9%)	51 (1.0%)	
<b>Father's region of origin</b>	Sweden	406563 (75.0%)	11253 (77.0%)	<0.001	2346 (63.3%)	4729 (80.9%)	4178 (82.6%)	<0.001
	Nordics	19321 (3.6%)	495 (3.4%)		84 (2.3%)	179 (3.1%)	232 (4.6%)	
	West Eu and NA	15974 (2.9%)	366 (2.5%)		124 (3.3%)	140 (2.4%)	102 (2.0%)	
	Africa	21694 (4.0%)	525 (3.6%)		323 (8.7%)	136 (2.3%)	66 (1.3%)	
	Asia and Oceania	17133 (3.2%)	427 (2.9%)		206 (5.6%)	113 (1.9%)	108 (2.1%)	
	Middle-East	39176 (7.2%)	787 (5.4%)		338 (9.1%)	289 (4.9%)	160 (3.2%)	
	East Eu and Russia	11764 (2.2%)	364 (2.5%)		146 (3.9%)	137 (2.3%)	81 (1.6%)	
	South America	10485 (1.9%)	388 (2.7%)		137 (3.7%)	122 (2.1%)	129 (2.6%)	
	Unknown/Missing	17 (<1%)	0 (0.0%)		0 (0.0%)	0 (0.0%)	0 (0.0%)	

Autism spectrum disorder (ASD), body mass index (BMI), Gestational age (GA), North America (NA), Europe (Eu)

**Supplementary Table S2. Association between covariates and ID and by age at first diagnosis.**

		<b>No ID</b>	<b>ID</b>	<b>p-value</b>	<b>0-&lt;6 years</b>	<b>6-&lt;12 years</b>	<b>12-&lt;18 years</b>	<b>p-value</b>
<b>N</b>		550216	6516		2410	2710	1396	
<b>Sex</b>	Male	281322 (51.1%)	4132 (63.4%)	<0.001	1556 (64.6%)	1754 (64.7%)	822 (58.9%)	<0.001
	Female	268894 (48.9%)	2384 (36.6%)		854 (35.4%)	956 (35.3%)	574 (41.1%)	
<b>Maternal BMI</b>	Underweight	14911 (2.7%)	189 (2.9%)	<0.001	63 (2.6%)	77 (2.8%)	49 (3.5%)	<0.001
	Normalweight	279127 (50.7%)	2747 (42.2%)		1163 (48.3%)	1103 (40.7%)	481 (34.5%)	
	Overweight	79730 (14.5%)	1025 (15.7%)		447 (18.5%)	391 (14.4%)	187 (13.4%)	
	Obese	27037 (4.9%)	533 (8.2%)		211 (8.8%)	230 (8.5%)	92 (6.6%)	
	Missing	149411 (27.2%)	2022 (31.0%)		526 (21.8%)	909 (33.5%)	587 (42.0%)	
<b>Birth order</b>	First born	248771 (45.2%)	2649 (40.7%)	<0.001	1003 (41.6%)	1103 (40.7%)	543 (38.9%)	<0.001
	Second born	198901 (36.1%)	2259 (34.7%)		843 (35.0%)	932 (34.4%)	484 (34.7%)	
	Third born or later	102544 (18.6%)	1608 (24.7%)		564 (23.4%)	675 (24.9%)	369 (26.4%)	
<b>Family income at birth</b>	Low (0-20%)	78377 (14.2%)	1520 (23.3%)	<0.001	511 (21.2%)	629 (23.2%)	380 (27.2%)	<0.001
	Medium-low (20-40%)	115721 (21.0%)	1691 (26.0%)		606 (25.1%)	727 (26.8%)	358 (25.6%)	
	Medium (40-60%)	119491 (21.7%)	1324 (20.3%)		495 (20.5%)	539 (19.9%)	290 (20.8%)	
	Medium-high (60-80%)	118690 (21.6%)	1030 (15.8%)		379 (15.7%)	436 (16.1%)	215 (15.4%)	
	High (80-100%)	117294 (21.3%)	859 (13.2%)		355 (14.7%)	358 (13.2%)	146 (10.5%)	
	Missing	643 (0.1%)	92 (1.4%)		64 (2.7%)	21 (0.8%)	7 (0.5%)	
<b>Maternal age, years</b>	<25	81411 (14.8%)	1281 (19.7%)	<0.001	363 (15.1%)	534 (19.7%)	384 (27.5%)	<0.001
	25-29	161994 (29.4%)	1927 (29.6%)		663 (27.5%)	833 (30.7%)	431 (30.9%)	
	30-34	191406 (34.8%)	1944 (29.8%)		780 (32.4%)	816 (30.1%)	348 (24.9%)	
	35-39	95542 (17.4%)	1055 (16.2%)		457 (19.0%)	414 (15.3%)	184 (13.2%)	
	>39	19863 (3.6%)	309 (4.7%)		147 (6.1%)	113 (4.2%)	49 (3.5%)	
<b>Paternal age, years</b>	<25	37471 (6.9%)	591 (9.2%)	<0.001	137 (5.8%)	254 (9.5%)	200 (14.4%)	<0.001
	25-29	119222 (21.9%)	1447 (22.5%)		469 (19.8%)	615 (23.0%)	363 (26.2%)	
	30-34	182782 (33.6%)	1928 (30.0%)		715 (30.2%)	825 (30.8%)	388 (28.0%)	

<b>Birth season</b>	35-39	126569 (23.2%)	1355 (21.1%)	0.21	561 (23.7%)	555 (20.7%)	239 (17.2%)	0.014
	>39	78467 (14.4%)	1109 (17.2%)		487 (20.6%)	426 (15.9%)	196 (14.1%)	
	Dec-Feb	129074 (23.5%)	1563 (24.0%)		569 (23.6%)	664 (24.5%)	330 (23.6%)	
	Mar-May	148692 (27.0%)	1733 (26.6%)		597 (24.8%)	724 (26.7%)	412 (29.5%)	
	Jun-Aug	142825 (26.0%)	1636 (25.1%)		654 (27.1%)	670 (24.7%)	312 (22.3%)	
	Sep-Nov	129625 (23.6%)	1584 (24.3%)		590 (24.5%)	652 (24.1%)	342 (24.5%)	
<b>Pre-eclampsia</b>	No	535497 (97.3%)	6260 (96.1%)	<0.001	2306 (95.7%)	2599 (95.9%)	1355 (97.1%)	<0.001
	Yes	14719 (2.7%)	256 (3.9%)		104 (4.3%)	111 (4.1%)	41 (2.9%)	
<b>Parental education</b>	<10 years	33669 (6.1%)	824 (12.6%)	<0.001	222 (9.2%)	335 (12.4%)	267 (19.1%)	<0.001
	10-12	209705 (38.1%)	3095 (47.5%)		960 (39.8%)	1364 (50.3%)	771 (55.2%)	
	>12	305487 (55.5%)	2501 (38.4%)		1162 (48.2%)	989 (36.5%)	350 (25.1%)	
	Missing	1355 (0.2%)	96 (1.5%)		66 (2.7%)	22 (0.8%)	8 (0.6%)	
<b>Size for GA</b>	Small	12603 (2.3%)	506 (7.8%)	<0.001	208 (8.6%)	210 (7.7%)	88 (6.3%)	<0.001
	Normal	503478 (91.5%)	5446 (83.6%)		1981 (82.2%)	2270 (83.8%)	1195 (85.6%)	
	Large	16393 (3.0%)	229 (3.5%)		85 (3.5%)	88 (3.2%)	56 (4.0%)	
	Missing	17742 (3.2%)	335 (5.1%)		136 (5.6%)	142 (5.2%)	57 (4.1%)	
<b>Low Apgar score (&lt;7)</b>	No	538709 (97.9%)	6124 (94.0%)	<0.001	2249 (93.3%)	2558 (94.4%)	1317 (94.3%)	<0.001
	Yes	4824 (0.9%)	255 (3.9%)		118 (4.9%)	92 (3.4%)	45 (3.2%)	
	Missing	6683 (1.2%)	137 (2.1%)		43 (1.8%)	60 (2.2%)	34 (2.4%)	
<b>Cesarean delivery</b>	No	496004 (90.1%)	5595 (85.9%)	<0.001	1991 (82.6%)	2327 (85.9%)	1277 (91.5%)	<0.001
	Yes	36470 (6.6%)	586 (9.0%)		283 (11.7%)	241 (8.9%)	62 (4.4%)	
	Missing	17742 (3.2%)	335 (5.1%)		136 (5.6%)	142 (5.2%)	57 (4.1%)	
<b>GA at birth</b>	Preterm	30345 (5.5%)	840 (12.9%)	<0.001	343 (14.2%)	344 (12.7%)	153 (11.0%)	<0.001
	Term	515635 (93.7%)	5596 (85.9%)		2041 (84.7%)	2331 (86.0%)	1224 (87.7%)	
	Post term	3321 (0.6%)	54 (0.8%)		19 (0.8%)	21 (0.8%)	14 (1.0%)	
	Missing	915 (0.2%)	26 (0.4%)		7 (0.3%)	14 (0.5%)	5 (0.4%)	
<b>Paternal psychiatric diagnosis</b>	Not present	399806 (72.7%)	4220 (64.8%)	<0.001	1628 (67.6%)	1751 (64.6%)	841 (60.2%)	<0.001
	Present	150410 (27.3%)	2296 (35.2%)		782 (32.4%)	959 (35.4%)	555 (39.8%)	

<b>Maternal psychiatric diagnosis</b>	Not present	311676 (56.6%)	2919 (44.8%)	<0.001	1147 (47.6%)	1228 (45.3%)	544 (39.0%)	<0.001
	Present	238540 (43.4%)	3597 (55.2%)		1263 (52.4%)	1482 (54.7%)	852 (61.0%)	
<b>Paternal infection</b>	No	317992 (57.8%)	3688 (56.6%)	0.052	1306 (54.2%)	1545 (57.0%)	837 (60.0%)	0.001
	Yes	232224 (42.2%)	2828 (43.4%)		1104 (45.8%)	1165 (43.0%)	559 (40.0%)	
<b>Maternal infection</b>	No	219388 (39.9%)	2289 (35.1%)	<0.001	725 (30.1%)	977 (36.1%)	587 (42.0%)	<0.001
	Yes	330828 (60.1%)	4227 (64.9%)		1685 (69.9%)	1733 (63.9%)	809 (58.0%)	
<b>Mother's region of origin</b>	Sweden	404103 (73.4%)	3970 (60.9%)	<0.001	1362 (56.5%)	1687 (62.3%)	921 (66.0%)	<0.001
	Nordics	15221 (2.8%)	214 (3.3%)		43 (1.8%)	98 (3.6%)	73 (5.2%)	
	West Eu and NA	24207 (4.4%)	299 (4.6%)		136 (5.6%)	103 (3.8%)	60 (4.3%)	
	Africa	25044 (4.6%)	571 (8.8%)		260 (10.8%)	233 (8.6%)	78 (5.6%)	
	Asia and Ocean	11481 (2.1%)	181 (2.8%)		96 (4.0%)	65 (2.4%)	20 (1.4%)	
	Middle east	45507 (8.3%)	915 (14.0%)		377 (15.6%)	373 (13.8%)	165 (11.8%)	
	East Eu and							
	Russia	7394 (1.3%)	68 (1.0%)		26 (1.1%)	24 (0.9%)	18 (1.3%)	
	South America	11520 (2.1%)	212 (3.3%)		69 (2.9%)	92 (3.4%)	51 (3.7%)	
	Unknown/Missing	5739 (1.0%)	86 (1.3%)		41 (1.7%)	35 (1.3%)	10 (0.7%)	
<b>Father's region of origin</b>	Sweden	413612 (75.2%)	4204 (64.5%)	<0.001	1439 (59.7%)	1774 (65.5%)	991 (71.0%)	<0.001
	Nordics	19580 (3.6%)	236 (3.6%)		62 (2.6%)	89 (3.3%)	85 (6.1%)	
	West Eu and NA	16143 (2.9%)	197 (3.0%)		87 (3.6%)	63 (2.3%)	47 (3.4%)	
	Africa	21701 (3.9%)	518 (7.9%)		239 (9.9%)	222 (8.2%)	57 (4.1%)	
	Asia and Ocean	17317 (3.1%)	243 (3.7%)		129 (5.4%)	89 (3.3%)	25 (1.8%)	
	Middle east	39192 (7.1%)	771 (11.8%)		328 (13.6%)	314 (11.6%)	129 (9.2%)	
	East Eu and							
	Russia	11985 (2.2%)	143 (2.2%)		59 (2.4%)	60 (2.2%)	24 (1.7%)	
	South America	10669 (1.9%)	204 (3.1%)		67 (2.8%)	99 (3.7%)	38 (2.7%)	
	Unknown/Missing	17 (<1%)	0 (0.0%)		0 (0.0%)	0 (0.0%)	0 (0.0%)	

Intellectual disability (ID), body mass index (BMI), Gestational age (GA), North America (NA), Europe (Eu)