Oxygen provision to severely ill COVID-19 patients at the peak of the 2020 pandemic in a Swedish district hospital

Anna Hvarfner[1, 2], Ahmed Al-Djaber[3,4], Hampus Ekström[3, 4], Malin Enarsson[3,5], Markus Castegren[3,6], Tim Baker[7,8]*, Carl Otto Schell[3,7]*

[1] Center for Clinical Research Dalarna, Uppsala University, Falun, Sweden

[2] Malung health center, Dalarna Region, Malung, Sweden

[3] Center for Clinical Research Sörmland, Uppsala University, Eskilstuna, Sweden

[4] Department of Medicine Nyköping Hospital, Nyköping, Sweden

[5] Department of Infectious diseases, Eskilstuna Hospital , Eskilstuna, Sweden

[6] CLINTEC and FyFa, Karolinska Institute, Stockholm, Sweden

[7] Department of Global Public Health, Karolinska Institute, Stockholm, Sweden

[8] Department of Clinical Research, London School of Hygiene and Tropical Medicine, London, UK

* joint last authors

Correspondence should be addressed to Anna Hvarfner, anna.hvarfner@gmail.com.

Abstract

Oxygen is a low-cost and life-saving therapy for patients with COVID-19. Yet, it is a limited resource in many hospitals in low income countries and in the 2020 pandemic even hospitals in richer countries reported oxygen shortages. An accurate understanding of oxygen requirements is needed for capacity planning. The World Health Organization estimates the average flow-rate of oxygen to severe COVID-19-patients to be 10 l/min. However, there is a lack of empirical data about the oxygen provision to patients. This study aimed to estimate the oxygen provision to COVID-19 patients with severe disease in a Swedish district hospital.

A retrospective, medical records-based cohort study was conducted in March to May 2020 in a Swedish district hospital. All adult patients with severe COVID-19 – those who received oxygen in the ward and had no ICU-admission during their hospital stay – were included. Data were collected

on the oxygen flow-rates provided to the patients throughout their hospital stay, and summary

measures of oxygen provision calculated.

One-hundred and twenty six patients were included, median age was 70 years and 43% were

female. On admission, 27% had a peripheral oxygen saturation of ≤91% and 54% had a respiratory

rate of ≥25/min. The mean oxygen flow-rate to patients while receiving oxygen therapy was 3.0

l/min (SD 2.9) and the mean total volume of oxygen provided per patient admission was 16,000 l (SD

23,000).

In conclusion, the provision of oxygen to severely ill COVID-19-patients was lower than

previously estimated. Further research is required before global estimates are adjusted.

Introduction

The world has been tackling the COVID-19 pandemic since early 2020 (1). The pandemic has led to

high demands on health systems and many lives have been lost because of the disease (2). The

severe acute respiratory syndrome coronvarius-2 (SARS-CoV-2) causing COVID-19 can affect most

organs but is primarily a respiratory virus (3). Viral pneumonia is the most common serious

manifestation and can result in hypoxemia and acute respiratory distress syndrome (ARDS) (3,4).

To treat severe and critical COVID-19, much global focus has been on expanding intensive

care capacity including the use of mechanical ventilation (5). Whilst many critically ill patients can

benefit from mechanical ventilation if it is administrated by experienced personnel, it is a staff

intensive measure and requires training to be effective and avoid harm (6).

Oxygen therapy is a low cost treatment, less complex than mechanical ventilation and saves

lives in COVID-19 (4,7–9). It is the first-line treatment of hypoxemia and has been listed as a World

Health Organization (WHO) essential medicine (10). Oxygen is a limited resource in many hospitals in

low income countries (11–14), and during the peaks of the pandemic waves there have been reports

of hospitals running out of oxygen in high and middle income countries such as the UK, USA, South

Africa, Portugal, Egypt and Brazil (15–21). In addition, sudden failures of the oxygen systems

requiring emergency transport from other sites may happen to hospitals anywhere in the world.

Capacity planning is needed to optimise the distribution of oxygen and reach highest positive

impact on patient outcomes. This requires an accurate understanding of oxygen requirements for

patients with COVID-19. The WHO estimates the average flow-rate of oxygen to severely ill COVID-

19-patients (referring to those requiring oxygen but not intensive care unit treatment) to be 10 l/min

(22). This estimate is not based on empirical findings and there is a lack of quantitative data on

oxygen provision to patients. This study aims to estimate oxygen provision to severely ill COVID-19

patients in a Swedish district hospital.

Method

65 A retrospective, electronic medical records-based cohort study in the medical department in

66 Nyköping Hospital, Sweden.

Setting

Nyköping Hospital is a first-line district hospital in Sörmland Region in Sweden with a catchment area

of 90,000 people. The medical department in the hospital has a usual capacity for managing 35

inpatients at a time. Sörmland was one of the first Swedish regions to be significantly affected by

COVID-19 in 2020 (23) – the first COVID-19 patient was admitted to the department in early March.

Less than a month later, the number of in-patients with COVID-19 was 43, with an average of seven

new COVID-19 admissions per day.

In the medical department, 40 beds on two new wards were opened within two weeks of the

first COVID-19-admission and extra staff were drawn from other areas of the hospital. The patients'

respiratory rate and peripheral oxygen saturation (SpO2) were assessed at least every two hours and

oxygen therapy adjusted to maintain SpO2 within an individualized target range (standard 92-96%,

lower to patients with assumed hypercapnic respiratory failure). The hospital has an intensive care

unit (ICU), which was expanded from the usual 5-beds to 14-beds during the pandemic. There was no

shortage of oxygen during the study period.

Before rapid polymerase chain reaction (PCR) testing capacity was widely available, an initial

diagnosis of COVID-19 was based on the clinical picture together with typical findings on a thoracic

computed tomography (CT) scan (24). Often the initial diagnosis was confirmed by later PCR-testing.

This early diagnostic reliance on CT scans shifted towards PCR-testing during the time of the study.

Standard treatments protocols evolved during the study period and included at different

times chloroquine, antibiotics and anticoagulation prophylaxis. Corticosteroids, Remdesivir and other

COVID-therapeutics were not included as standard treatment during the time of the study. High-flow

nasal oxygen and non-invasive ventilation were neither recommended for ARDS in COVID-19, nor

available in the medical wards at this time. Patients that needed more respiratory support than low-

flow oxygen were transferred to the ICU where most received invasive mechanical ventilation. As

many of the admitted patients were elderly and frail, in which ICU care may cause harm or would not

be in the patient´s best interest, a policy was introduced to make patient-centred decisions for every

admitted patient about the appropriateness of care-escalation to ICU in the event of clinical

deterioration. The decision of no escalation of care to ICU (no-ICU) was documented in the patient´s

notes and, importantly, was not regarded as synonymous with end-of-life or palliative care. Patients

for whom a no-ICU decision had been made received all other therapies, including oxygen therapy

when indicated, unless an additional clinical decision was made to provide end-of-life palliative care.

Study cohort

The Sörmland Region database of COVID-19 patients was used to identify participants. All adult

patients (age ≥18) admitted to the department of medicine in Nyköping Hospital from March 13 to

May 10 who had been diagnosed with COVID-19 during their admission and who fulfilled the criteria

for "severe" disease were included. Severe patients were those who received oxygen at some point

during their care in the ward and had no ICU-admission during their hospital-stay, in-line with the

WHO's classification (22). Non-severe COVID-19 patients – patients with either moderate disease

(never received oxygen) or critical disease (admitted to the ICU, either from the wards or directly

from the emergency department) (22) – were not included in the study cohort. However, to describe

the study cohort in context, data about admission findings, the most advanced mode of respiratory

support and outcomes were collected for all COVID-19 patients admitted to the department during

the study period. For patients with more than one admission, only data from the first admission were

included.

Sub-groups

115 Two a-priori defined sub-groups were analysed, as it was hypothesised that their oxygen provision

116 may differ substantially from other patients. The first group were those patients for whom a no-ICU-

117 decision had been made. The second group consisted of patients younger than 70 years old.

Data extraction

Data were extracted in two ways. Data on vital signs and oxygen treatment on admission, patient

characteristics including Charlson´s Age Adjusted Comorbidity score (CACI) (25,26), clinical and

laboratory findings, pharmaceutical treatments and the presence of a no-ICU decision were extracted

manually from the patients' electronic medical records. Other data were collected through a

computerised search in the electronic medical records system, including for dates of admission,

discharge, outcomes, and all registered SpO2 values and oxygen flow-rates throughout the patients'

care. The data extracted with a computerised search were validated by manual cross-checking a 10%

sample of the data with the medical records. Data were used from the patients' entire stay in the

medical department. The patients' outcome in the medical department were noted as either transfer

to another department, discharge home or died, with an additional outcome of died within 60 days.

Analysis

All data were anonymised before analysis. Oxygen provision to each patient was estimated by

multiplying the oxygen flow-rate during each assessment of the patient (l/min) and the time since

the previous assessment (minutes). Thereby, the volumes of oxygen provided to the patient over

time were generated, and the sum of these made up the total volume of oxygen provided to the

patient (formula 1).

Formula 1,   Total volume of oxygen provided to a patient:

$$\sum_n X_n \left(T_{(n+1)} - T_n \right)$$

n: one time point in a series of consecutive time points, **T$_n$**: time at the time point n, **X$_n$**: oxygen flow (l/min) at

the time point n

The mean oxygen flow-rate for each patient while receiving oxygen was calculated as the total

volume of oxygen provided divided by time spent receiving oxygen treatment. An additional analysis

of the mean oxygen flow-rate for each patient during their care in the ward used the total oxygen

volume provided divided by total time in the ward. Means were used to provide an estimate of the

total amount of oxygen that needs to be supplied to such a patient cohort. Medians were also

calculated for both analyses to provide an estimate of a "typical" patient, given the non-normally

distributed data. For other variables, median and interquartile range (IQR) were used for non-

normally distributed values and mean and standard deviation (SD) for normally distributed values.

For all admitted patients, the clinical progression score was determined as has been recently defined

(27). Admission vital signs were analysed as the proportion of vital signs corresponding to a red

NEWS-2 parameter (28). Missing data was omitted from analysis. STATA IC/15.1(StataCorpLLC) was

used for the analysis.

Ethical considerations

155   The study was approved by the Ethical Review Board in Lund, reference number 2020-04012. As the

156   study did not alter patient care and data were anonymised before analysis, individual patient consent

157   was waived.

Results

Patient characteristics

In total, 206 COVID-19 patients were admitted to the department during the study period (Table 1,

Supplementary table 1). Of these, 126 had severe disease and were included as the study cohort. The

median age was 70 years (IQR 57-82), 54 (43%) were female and 42% had a body mass index (BMI) of

30 or above. CACI scores were 4 or above (indicating <50% estimated 10-years survival) in 50% of the

patients (26,27). A no-ICU decision was documented for 48% of the patients. On admission, 34 (27%)

had an SpO2 of 91% or below and 68 (54%) had a respiratory rate of 25 or above. The length of stay

was a median of 4.9 days (IQR 2.8-7.8). Eight (6.4%) patients were transferred to another

department. The in-hospital mortality was 26% and the 60-day mortality was 32%. For the subset of

patients aged <70, in-hospital and 60-day mortalities were 6.5% and 8.1% respectively and for

patients with a documented no-ICU-decision 55% and 65% respectively. The clinical progression

scores for all admitted patients are presented in supplementary table 2.

Table 1 Patient characteristics and outcomes

	Study cohort % (n/N), unless otherwise stated	All patients admitted to the department % (n/N), unless otherwise stated
Age (years), median (IQR)	70 (57-82)	65 (54-78)
Female	43% (54/126)	42% (87/206)
Diagnosis of COVID-19 confirmed by PCR	90% (114/126)	89% (184/206)
BMI ≥30	42% (38/91)	40% (56/141)
CACI ≥4	50% (63/126)	43% (88/206)
No-ICU-decision documented	48% (60/126)	37% (77/206)
Red NEWS-2 (28) parameter on first measurements of vital signs		
SpO2 (≤91%)	27% (34/126)	28% (57/206)
Respiratory rate (≤8 or ≥25 breaths/min)	54% (68/126)	49% (100/203)
Heart rate	3.2% (4/125)	2.4% (5/205)

(≤40 or ≥131 beats/min)		
Systolic blood pressure (≤90 or ≥220 mmHg)	2.4% (3/125)	2.0% (4/205)
Consciousness (Non-alert)	13% (17/126)	10% (20/204)
Temperature (≤35.0 or ≥39.1°C)	14% (18/125)	16% (32/205)
Treatments during hospital-stay		
Antibiotics	79% (99/126)	77% (158/206)
Chloroquine	12% (15/126)	14% (28/206)
Anticoagulants	52% (65/126)	55% (114/206)
Patient outcomes		
Length of stay (days), median (IQR)	4.9 (2.8-7.8)	4.3 (2.2-9.0)
Transfer to another department	6.4% (8/126)	13% (27/206)
Dead in-hospital	26% (33/126)	19% (40/206)
Dead at 60 days	32% (40/126)	23% (48/206)

175 *Abbreviations: PCR: polymerase chain reaction, BMI: body mass index, CACI: Charlson´s age

176 adjusted comorbidity score, ICU: intensive care unit, SpO2: peripheral oxygen saturation

Oxygen provision

The mean oxygen flow-rate to the patients while receiving oxygen therapy was 3.0 l/min (SD 2.9) and

the median was 2.0 l/min (IQR 1.3-3.5). Results for oxygen provision to the study cohort are shown in

Table 2 and for patients that were initially cared for in the wards but later transferred to ICU

(therefore not part of the study cohort) in supplementary table 3. The highest oxygen flow-rate

provided to the patients was median 4.0 l/min (IQR 2.0-8.0) and mean 5.4 l/min (SD 4.1) (figure 1).

Table 2 Oxygen provision

	Study cohort (n= 126)	Subgroups	
		Patients aged <70 (n=62)	Patients with a no-ICU-decision (n=60)
Age, median (IQR)	70 (57-82)	57 (48-61)	83 (75-88)
Days on oxygen treatment, median (IQR)	2.3 (0.68-4.2)	1.8 (0.68-3.9)	2.5 (0.77-4.8)
Oxygen flow to patients during oxygen therapy (l/min)			
- Mean (SD)	3.0 (2.9)	2.6 (2.3)	4.0 (3.7)
- Median (IQR)	2.0 (1.3-3.5)	1.9 (1.3-2.9)	2.9 (1.6-5.6)
Oxygen flow to patients during time in the ward (l/min)			
- Mean (SD)	2.2 (2.9)	1.8 (2.3)	3.1 (3.7)
- Median (IQR)	1.2 (0.32-2.6)	1.1 (0.43-2.0)	1.6 (0.28-4.6)
Total volume of oxygen provided per patient admission (l)			
- Mean (SD)	16,000 (23,000)	12,000 (16,000)	22,000 (26,000)

- Median (IQR)	7,400 (1,200-21,000)	4,8000 (1,800-17,000)	12,000 (2,000-32,000)

Figure 1 Highest oxygen flow-rate received by patients with severe COVID-19 during their care in the medical wards

Discussion

We have found that a mean of 3.0 l/min (median 2.0 l/min) oxygen were provided to severely ill COVID-19 patients while receiving oxygen in a Swedish district hospital during the first peak of the pandemic in 2020. This calculated oxygen flow-rate is lower than the 10 l/min estimated by the WHO for COVID-19 patients with severe disease (22).

A strength of this study is the use of a method for calculating oxygen provision to hospitalized patients by summing up documented flow-rates in the medical records. Previously, findings of the proportions of patients receiving oxygen (8,29–32) and assumptions on required oxygen flow (22) have been presented. To the best of our knowledge, this study is the first to use all the necessary data to be able to calculate total oxygen provision to patients.

The 206 patients admitted to the medical department in Nyköping with COVID-19 during this study period had similar age, gender proportions and in-hospital mortality as other described cohorts of hospitalised COVID-19 patients (8,29,31,33). The median length of stay of 4.3 days was shorter in the Nyköping patient group than in two cohorts from New York and Madrid described during the early pandemic, as well as in the multinational ISARIC survey, with stays of seven to nine days (31,33,34). This could be explained by Sweden´s early discharge policies (35,36) and a considerable

amount of patient transfers in this study (13% of all admitted patients and 6.4% of the study cohort)

due to overcrowding.

The lower flow-rates of oxygen to patients with severe COVID-19 in this study compared to

previous estimates are interesting but, for several reasons, should be interpreted with caution when

planning health care services. Firstly, it is treacherous for health care systems to make plans based on

the COVID-19 patient categories "severe" and "critical" as there are currently several definitions in

use and many of them categorise patients based on the care they receive rather than the care they

require (22,27,37). Since resources and practices for ICU-admission vary greatly around the world, for

example ICU beds per 100,000 population vary from 29 in Germany to 6 in Sweden and <1 in Uganda

(38), very few COVID-19 patients would be classified as critical in Uganda according to the definition

used by WHO and in this study. Indeed, in settings with less availability of ICU beds, sicker patients

will be classified as "severe" and require higher flows of oxygen than the cohort in this study. In

addition, suboptimal respiratory support might cause shorter survival or survival with prolonged time

to recovery – hence it is difficult to make assumptions on the quantity of oxygen they need without

available data. This study aimed to study a patient group that was truly severe according to the WHO

classification, and due to the well-resourced hospital system in Sweden succeeded with this aim, but

a similar group may not be easy to delineate in other settings.

Secondly, decisions on treatment limitations were important to the assignment of patients to

the severe group in this study and such practices vary largely across countries. Of the study patients,

48% had a no-ICU-decision meaning that even if they deteriorated, they were kept in the general

wards and received targeted oxygen treatment. Although oxygen needs for this subgroup was also

considerably lower than the WHO estimate (4.0 l/min), it is possible that in settings with other norms

around treatment limitations, these patients may, at some point, have received ICU-treatments such

as mechanical ventilation with substantially higher oxygen flow-rates.

Thirdly, the flow-rates of oxygen provided to patients depend on target saturation and duration

between treatment modifications. The patients in this study were cared for with defined targets and

frequent saturation controls. In wards where oxygen flow is not − or cannot be − adjusted as

frequently, oxygen flow-rates and target saturations may be higher to provide patients with a safe

margin for avoiding hypoxia (39). While the optimal target saturation for hospitalised patients

receiving oxygen is debated (40,41), the target range in Nyköping was set following the Surviving

Sepsis Campaign guidelines (42).

Our findings suggest that the oxygen need for severely ill COVID-19 patients may be lower

than previously estimated. Future research using the methods described in this study in larger

cohorts and from other settings would be useful to inform capacity planning with updated estimates

of oxygen need. Additionally, as oxygen is essential in many other conditions such as sepsis, trauma

(21,43–45) and notably child pneumonia, a disease killing 800,000 children under 5 each year (14,46),

estimates for the oxygen needs for treating these conditions would be beneficial.

Conclusion

247   The provision of oxygen to severely ill COVID-19-patients was lower than previously estimated.

248   Further research is required before global estimates are adjusted.

Acknowledgements

251   The authors would like to thank Johan Printz and Josefin Back for conducting the computerised

252   search in the electronic medical records system, Karin Näslund for her general IT-support throughout

253   the study and the heads of the medical department in Nyköping, Björn Alpe and Karin Marminge, for

254   their support of this study.

References

1. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020

[Internet]. [cited 2021 Jan 6]. Available from: https://www.who.int

2. COVID-19 Map [Internet]. Johns Hopkins Coronavirus Resource Center. [cited 2021 Feb 18].

Available from: https://coronavirus.jhu.edu

3. Gupta A, Madhavan MV, Sehgal K, Nair N, Mahajan S, Sehrawat TS, et al. Extrapulmonary

manifestations of COVID-19. Nat Med. 2020 Jul;26(7):1017–32.

4. Brosnahan SB, Jonkman AH, Kugler MC, Munger JS, Kaufman DA. COVID-19 and Respiratory

System Disorders. Arterioscler Thromb Vasc Biol. 2020 Nov;40(11):2586–97.

5. Baker T, Schell CO, Petersen DB, Sawe H, Khalid K, Mndolo S, et al. Essential care of critical illness

must not be forgotten in the COVID-19 pandemic. Lancet Lond Engl. 2020 Apr

18;395(10232):1253–4.

6. Katira BH. Ventilator-Induced Lung Injury: Classic and Novel Concepts. Respir Care. 2019

Jun;64(6):629–37.

7. Duke T, Wandi F, Jonathan M, Matai S, Kaupa M, Saavu M, et al. Improved oxygen systems for

childhood pneumonia: a multihospital effectiveness study in Papua New Guinea. Lancet Lond

Engl. 2008 Oct 11;372(9646):1328–33.

8. Xie J, Covassin N, Fan Z, Singh P, Gao W, Li G, et al. Association Between Hypoxemia and Mortality

in Patients With COVID-19. Mayo Clin Proc. 2020;95(6):1138–47.

9. Dondorp AM, Hayat M, Aryal D, Beane A, Schultz MJ. Respiratory Support in Novel Coronavirus

Disease (COVID-19) Patients, with a Focus on Resource-Limited Settings. Am J Trop Med Hyg

[Internet]. 2020 Apr 21 [cited 2021 Feb 18]; Available from: https://www.ncbi.nlm.nih.gov

10. WHO model list of essential medicines [Internet]. [cited 2021 Jan 22]. Available from:

https://www.who.int

11. Dat VQ, Long NT, Giang KB, Diep PB, Giang TH, Diaz JV. Healthcare infrastructure capacity to

respond to severe acute respiratory infection (SARI) and sepsis in Vietnam: A low-middle

income country. J Crit Care. 2017;42:109–15.

12. Bradley BD, Light JD, Ebonyi AO, N'Jai PC, Ideh RC, Ebruke BE, et al. Implementation and 8-year

follow-up of an uninterrupted oxygen supply system in a hospital in The Gambia. Int J Tuberc

Lung Dis Off J Int Union Tuberc Lung Dis. 2016;20(8):1130–4.

13. Belle J, Cohen H, Shindo N, Lim M, Velazquez-Berumen A, Ndihokubwayo J-B, et al. Influenza

preparedness in low-resource settings: a look at oxygen delivery in 12 African countries. J Infect

Dev Ctries. 2010 Aug 4;4(7):419–24.

14. Fore HH, Ghebreyesus TA, Watkins K, Greenslade L, Berkley S, Bassat Q, et al. Leveraging the

COVID-19 response to end preventable child deaths from pneumonia. Lancet Lond Engl. 2020

Nov 28;396(10264):1709–11.

15. Denis Campbell H policy editor. Southend hospital's oxygen supply reaches 'critical situation'. The

Guardian [Internet]. 2021 Jan 11 [cited 2021 Jan 29]; Available from:

https://www.theguardian.com

16. Tompkins L, Ives M, Shaver JW. Running low on oxygen, emergency workers in Los Angeles

County are told to administer the minimum necessary. The New York Times [Internet]. 2021 Jan

4 [cited 2021 Jan 29]; Available from: https://www.nytimes.com/

17. Naidoo Y, Singh O. Staff and oxygen shortages in second wave tremor, front-line workers fear

Covid-19 tsunami. TimesLIVE [Internet]. 2021 Jan 3 [cited 2021 Jan 29]; Available from:

https://www.timeslive.co.za

18. Covid-19: Brazil hospitals 'run out of oxygen' for virus patients. BBC News [Internet]. 2021 Jan 15

[cited 2021 Jan 29]; Available from: https://www.bbc.com/

19. Michaelson R. Egypt denies 'oxygen crisis' as Covid-19 ward videos allege shortage. The Guardian

[Internet]. 2021 Jan 12 [cited 2021 Jan 29]; Available from: https://www.theguardian.com

20. Demony VW Catarina. Germany sends medical experts to Portugal as hospitals face COVID-19

crisis. Reuters [Internet]. 2021 Jan 28 [cited 2021 Feb 19]; Available from:

https://www.reuters.com

21. Usher AD. Medical oxygen crisis: a belated COVID-19 response. The Lancet. 2021 Mar

6;397(10277):868–9.

22. World Health Organization. Oxygen sources and distribution for COVID-19 treatment centres:

interim guidance [Internet]. [cited 2021 Feb 18]. Report No.: WHO/2019-

nCoV/Oxygen_sources/2020.1. Available from: https://www.who.int

23. The Public Health Agency of Sweden. Veckorapport om covid-19, vecka 21 [Internet]. 2020 May

[cited 2021 Feb 18]. Available from: https://www.folkhalsomyndigheten.se

24. Kwee TC, Kwee RM. Chest CT in COVID-19: What the Radiologist Needs to Know. RadioGraphics.

2020 Oct 23;40(7):1848–65.

25. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic

comorbidity in longitudinal studies: development and validation. J Chronic Dis. 1987;40(5):373–

83.

26. Charlson Comorbidity Index (CCI) [Internet]. MDCalc. [cited 2020 Aug 28]. Available from:

https://www.mdcalc.com

27. Marshall JC, Murthy S, Diaz J, Adhikari NK, Angus DC, Arabi YM, et al. A minimal common

outcome measure set for COVID-19 clinical research. Lancet Infect Dis. 2020 Aug 1;20(8):e192–

7.

28. Royal College of Physicians. National Early Warning Score (NEWS) 2. Standardising the

assessment of acute-illness severity in the NHS. Updated report of a working party. [Internet].

London: RCP; 2017 [cited 2021 Jan 6]. Available from: https://www.rcplondon.ac.uk

29. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory

Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan,

China. JAMA Intern Med. 2020 Jul 1;180(7):934.

30. Hong L, Ye E, Sun G, Wang X, Zhang S, Wu Y, et al. Clinical and radiographic characteristics,

management and short-term outcomes of patients with COVID-19 in Wenzhou, China. BMC

Infect Dis. 2020 Nov 13;20(1):841.

31. Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, et al. Factors associated

with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in

New York City: prospective cohort study. BMJ. 2020 May 22;369:m1966.

32. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult

inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet Lond Engl.

2020 Mar 28;395(10229):1054–62.

33. Jiménez E, Fontán-Vela M, Valencia J, Fernandez-Jimenez I, Álvaro-Alonso EA, Izquierdo-García E,

et al. Characteristics, complications and outcomes among 1549 patients hospitalised with

COVID-19 in a secondary hospital in Madrid, Spain: a retrospective case series study. BMJ Open.

2020 Nov 1;10(11):e042398.

34. Consortium ISAR and emerging I, Hall M, Pritchard M, Dankwa EA, Baillie JK, Carson G, et al.

ISARIC Clinical Data Report 20 November 2020. medRxiv. 2020 Nov 23;2020.07.17.20155218.

35. Average length of stay in hospitals | Health at a Glance 2019: OECD Indicators | OECD iLibrary

[Internet]. [cited 2021 Mar 10]. Available from: https://www.oecd-ilibrary.org

36. Average length of stay in hospital | Health at a Glance: Europe 2020: State of Health in the EU

Cycle | OECD iLibrary [Internet]. [cited 2021 Mar 10]. Available from: https://www.oecd-

ilibrary.org

37. WHO Global. Clinical management of COVID-19 interim guidance [Internet]. 2020 May p. 62.

Report No.: WHO/2019-nCoV/clinical/2020.5. Available from: https://www.who.int

38. Ma X, Vervoort D. Critical care capacity during the COVID-19 pandemic: Global availability of

intensive care beds. J Crit Care. 2020 Aug;58:96–7.

39. Neto AS, Checkley W, Sivakorn C, Hashmi M, Papali A, Schultz MJ. Pragmatic Recommendations

for the Management of Acute Respiratory Failure and Mechanical Ventilation in Patients with

COVID-19 in Low- and Middle-Income Countries. Am J Trop Med Hyg [Internet]. 2021 Jan 13

[cited 2021 Mar 10];1(aop). Available from: https://www.ajtmh.org

40. World Health Organization. Clinical care for severe acute respiratory infection: toolkit: COVID-19

adaptation [Internet]. World Health Organization; 2020. Available from: https://apps.who.int

41. Chu DK, Kim LH-Y, Young PJ, Zamiri N, Almenawer SA, Jaeschke R, et al. Mortality and morbidity in

acutely ill adults treated with liberal versus conservative oxygen therapy (IOTA): a systematic

review and meta-analysis. The Lancet. 2018 Apr 28;391(10131):1693–705.

42. Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving Sepsis Campaign:

Guidelines on the Management of Critically Ill Adults with Coronavirus Disease 2019 (COVID-

19). Crit Care Med. 2020 Jun;48(6):e440.

43. Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, et al. Surviving Sepsis

Campaign: international guidelines for management of severe sepsis and septic shock, 2012.

Intensive Care Med. 2013 Feb;39(2):165–228.

44. Mock C, Essential Trauma Care Project (World Health Organization), World Health Organization,

International Society of Surgery, International Association for the Surgery of Trauma and

Surgical Intensive Care, editors. Guidelines for essential trauma care. Geneva: World Health

Organization; 2004. 93 p.

45. Schell CO, Gerdin Wärnberg M, Hvarfner A, Höög A, Baker U, Castegren M, et al. The global need

for essential emergency and critical care. Crit Care Lond Engl. 2018 Oct 29;22(1):284.

46. Pneumonia in Children Statistics [Internet]. UNICEF DATA. [cited 2021 Feb 1]. Available from:

https://data.unicef.org