ORIGINAL RESEARCH

Sleep quality mediates the relationship between risk of obstructive sleep apnea and acute stress in young adults

Author name: Kunal Aggarwal[1]

Author name: Nasreen Akhtar[2][*]

Author name: Hrudananda Mallick[3]

[1]Author affiliations: MBBS student, All India institute of medical Sciences, New Delhi-110029, India

[2]Author affiliations: Associate Professor, Department of Physiology, All India institute of medical Sciences, New Delhi-110029, India

[3]Author affiliations: Professor, Department of Physiology, SGT University, Gurgaon, Haryana-122505, India

Contact details: Email: drnasreenakhtar@gmail.com

[*]Correspondence: Nasreen Akhtar, Room no. 2024, Department of Physiology, AIIMS, New Delhi-110029, India

Tel +911126594053

Email : drnasreenakhtar@gmail.com

Abstract:

Purpose:

Intermittent hypoxia and transient arousals in obstructive sleep apnea (OSA) can lead to poor

sleep quality and acute stress. Rising levels of obesity and increased incidence of OSA in young

adults predisposes them to acute stress. We propose a mediation model to assess if risk of OSA

is associated with acute stress and if the relationship between risk for OSA and acute stress is

mediated by sleep quality.

Methods:

493 healthy individuals (F=237, M=256) from 18-25 years of age (mean age=20.3±1.53 years)

were screened for OSA, sleep quality and acute stress using STOP-BANG questionnaire,

Pittsburg Sleep Quality Index and American Psychiatry Association's National Stressful Events

Survey Acute Stress Disorder Short Scale (NSESS-S) respectively. Binary and logistic regression

were used establish the relationships between the variables. Sobel test for mediation analysis

was conducted.

Results:

73 participants (17.3%) were found at an intermediate and high risk of OSA by STOP BANG

questionnaire. 79 (16%) participants reported level of stress as 'None'. Mild, moderate and

severe stress was present in 248 (50.3%), 109 (22.1%), 51 (10.3%) and 16 (3.2%) participants

respectively. The odds of having severe and extreme stress among those at risk of sleep apnea

is 2.18 times higher than that among those not at risk of sleep apnea (OR: 2.18, 95% Confidence

Interval: 1.37-3.51). Sobel test established that the relationship between OSA and acute stress is

mediated by sleep quality.

Conclusion

Sleep quality mediates the relationship between risk for sleep apnea and acute stress. This

highlights the importance of screening for OSA in young adults, particularly young men with high

BMI, presenting with high stress levels.

Keywords: Emotional stress, sleep apnea, sleep quality, logistic regression, mediation analysis

Introduction

Obstructive sleep apnea (OSA) is characterized by repeated upper airway obstruction causing oxygen desaturation and frequent arousals from sleep[1]. The prevalence of OSA is increasing worldwide. A study from India reported a prevalence of 13.74% of OSA in general population[2]. The frequent desaturations, electroencephographic microarousals and sleep fragmentation in OSA lead to cardiovascular and neuropsychiatric sequalae [3]. OSA has an effect on intellectual abilities related mainly to verbal ability, attention and learning skills including planning and object categorisation [3]. These represent an organic damage to the central nervous system. The awareness of these intellectual impairments, sleep fragmentation and reduced daytime alertness makes the patients feel restless, anxious or depressed [4,5]. This can manifest emotionally as stress. Frequent arousals also cause an increase in catecholamines and cortisol levels, exacerbating the stress. Emotional stress has been found to be higher in patients of OSA and reduces after treatment [6]. OSA has been linked to quality of life [7], [7] depression, [8] and anxiety, [9] but the association with acute stress has been reported by very few studies. [6,10] Most of the studies on OSA have collected data from moderate to severe sleep apnea in older patients and in clinical settings. Occurrence of OSA in general population, particularly among young adults, has been neglected. Young adults are a vulnerable population for higher emotional stress. The current study was conducted in young adults in general population with the objectives to investigate the association of between risk of OSA and acute stress levels and to assess if the association between risk of OSA and acute stress level is mediated by sleep quality.

Material and methods

Sample and data Collection

This data is obtained from a study conducted in undergraduate college students from various courses in New Delhi from July to September 2019. [11] Ethical clearance was obtained from Institutional Ethical Committee of All India Institute of Medical Sciences, New Delhi (No IECPG-343/29.05.2019 dated 03.07.2019). The primary objective of the study was to investigate the association between acute stress and risk for obstructive sleep apnea in young adults.

This was a cross-sectional and observational study. There was no control group or blinding of assessors. Participants were approached by advertising in college social media groups and announcements in classrooms. The complete information regarding the study and consent form were provided online. All participants gave informed consent for the study and filled the online questionnaire. The questionnaire collected information about the student's age, gender, height, weight, acute stress, risk for sleep apnea and sleep quality. 503 participants between the ages of 18 and 25 years consented for the study. Those who were previously diagnosed for or on treatment for sleep disorders, psychiatric disorders or any other chronic disease were excluded.

Measures

Risk of sleep apnea was assessed by using STOP-BANG questionnaire (SBQ). [12] Even though Berlin questionnaire is the one most used in assessments in the clinic, SBQ is the more accurate tool for assessing risk of OSA in the general population. [13] There were 8-items in the questionnaire. Each positive item was scored as one. The risk was marked low, intermediate and high for a score of 0-2, 3-4 and 5-8 respectively. Intermediate and high risk categories were clubbed into one category for the purpose of analysis in this study as the number of participants in high risk category were very few and would skew the analysis.

Sleep quality was assessed by using Pittsburg Sleep Quality Index. [14] It is a 9-item questionnaire

widely used for assessing sleep quality. A score of less than or equal to five was considered good

quality sleep, and a score of more than five was considered as poor-quality sleep. Sleep time was

distinguished from other activities after going to bed like listening to music or reading books.

Acute stress was assessed by American Psychiatry Association Severity of Acute Stress

Symptoms—Adult (National Stressful Events Survey Acute Stress Disorder Short Scale

[NSESSS]) questionnaire. [15] It was a 7-item questionnaire. Each item on the measure was rated

by the subject on a 5-point scale (0=Not at all; 1=A little bit; 2=Moderately; 3=quite a bit, and

4=Extremely). The total raw score ranged from 0 to 28 and a higher score indicated a higher

degree of acute stress. This score was divided by 7 (the number of items) to provide a total

average score which was allocated a 5-point scale. It was used to assess severity of acute stress

as none (0), mild (1), moderate (2), severe (3) and extreme (4). [15]

Analysis

Acute stress, sleep quality and risk for obstructive sleep apnea were correlated to each other

using Pearson's bivariate correlation analysis. The Statistical Package for the Social Sciences

(SPSS®) software version 25.0 (IBM, Armonk NY, USA) for Windows® was used for the

correlation analyses. To investigate if risk for sleep apnea is associated with acute stress and if

sleep quality mediates the relationship between sleep apnea and acute stress, we created a

mediation model (Figure 1). A simple mediation analysis was performed based on the method

described by Baron and Kenny. [16] The dependent (outcome) variable for analysis was acute

stress. The independent variable was risk for obstructive sleep apnea. Mediator variable was

sleep quality. In our model, we assume there are two causal paths to the dependent variable

(acute stress), the direct impact of risk of OSA (path c) and the impact of sleep quality as the

mediator (path a and b). The mediated path from independent variable (Risk of OSA) to the

dependant variable (acute stress) is path c'. Binary and ordinal logistic regression were

performed using R software (R Core Team 2014 v3.4.4). Mediation analysis was performed using

Sobel test. [17] Sobel test is an accepted test of indirect effect and mediation. [18]

Results

503 volunteers filled up the questionnaire. After applying inclusion and exclusion criteria, 493 were included for data analysis. Female: male ratio was 0.91 (Females= 237, Males = 256). Mean age of participants was 20.3±1.53 years, ranging from 18 to 25 years. The basic descriptives of the study participants are depicted in Table 1. 73 participants (17.3%) were found at an intermediate and high risk of OSA. 79 (16%) participants reported level of stress as 'None'. The prevalence of mild, moderate and severe stress was 248 (50.3%), 109 (22.1%), 51 (10.3%) and 16 (3.2%) respectively. Mild stress was found higher in males (males- 27.2%, females- 23.1%), whereas moderate, severe and extreme stress was higher in females (Table 1).

Correlation analysis: Acute stress was significantly correlated with sleep quality (r=0.328, p<0.001), and risk for OSA (r=0.160, p<0.001) (Table 2). Sleep quality also correlates with risk of OSA (r=0.150, p<0.001).

Logistic Regression and mediation analysis: The assumptions of regression were checked and found satisfactory. First, using steps described by Baron and Kenny, [16] using acute stress as the dependant variable and risk for obstructive apnea as the independent variable, ordinal logistic regression was applied and risk for obstructive apnea was shown to be a significant predictor (p<0.001) of acute stress for all levels of stress (none, mild, moderate, severe and extreme) (c pathway). The odds of having severe and extreme stress among those among those at risk of sleep apnea is 2.18 times higher than that among those not at risk of sleep apnea (OR: 2.18, 95% Confidence Interval: 1.37-3.51). Second, risk for sleep apnea was used to predict the mediator variable of sleep quality (a pathway) using binary logistic regression. The odds ratio of poor sleep quality was 2.3 times in a person at high risk of OSA compared to a participant at low risk of OSA (OR: 2.3, 95% Confidence interval: 1.40-3.83). Third, the relationship between the mediator sleep quality and acute stress was established using ordinal logistic regression (b pathway). On change in sleep quality from good to poor, the odds of mild stress vs moderate or

severe combined are 3.5 times greater (OR: 3.55, 95% Confidence interval: 2.48-5.11). Last, the

mediated relationship between SBQ and acute stress was examined for a reduction in prediction

when the mediator was added to the model (c' pathway). Based on a previous study, [19] the three

regression equations were used to test the mediation effects of SQ on relation between OSA and

acute stress. Utilizing the coefficients of pathways 'a' and 'b', and Standard Errors (S_a, S_b) of

these coefficients, mediation was found (Test statistic: 0.7255; p-value 0.049, using Sobel test),

[8][7] showing that the relationship between SB and acute stress was no longer significant after

controlling for sleep quality. The Sobel test was used to determine that the 'ab' effect was

significantly greater than zero. This provides evidence that the relationship between OSA and

acute stress is mediated by sleep quality.

Discussion

Recurring upper-airway obstruction, intermittent hypoxemia and transient arousals leading to

sleep fragmentation and poor sleep quality, are the hall marks of OSA. The risk factors for OSA

are male sex, obesity, and advancing age, among others. [6]

Rate of weight gain is greatest in young adults, averaging 1 to 2 pounds per year. [20] Up to 19% of

teenagers and young adults are reportedly obese in India. [21] Rising levels of obesity are

increasing the prevalence of OSA in young adults. The role of OSA in development of stress in

young adults has not been studies much. When presenting with acute stress, young adults are

seldom screened for OSA. In this study, our aim is to highlight the importance of screening for

OSA in young adults, particularly young men with high BMI, presenting with high stress levels.

SBQ has been recommended as the screening tool for OSA in community settings and where

PSG is not available. [13] It is more accurate than Berlin and STOP questionnaire for screening of

OSA in general population. [13,22] SBQ has a reported sensitivity of is 88,90 and 93%, specificity of

42,36 and 35%, and a diagnostic odds ratio of 5.13, 5.05 and 6.51 for mild, moderate and severe

OSA, respectively. [13] We found a prevalence of intermediate and high risk of sleep apnea to be

14.8% in our study. It is similar to the prevalence rate of 13.7% reported by a previous study

conducted in Delhi. [2] We have found a significant association of risk of OSA with acute stress. We

presented the result in World Sleep Congress 2019. [23] The risk of being under severe acute

stress was 2.18 times higher in those at risk of OSA than those without the risk of OSA. Other

studies have evaluated the association of OSA with emotional stress. Das Santos et al have

reported a higher stress level in moderate and severe apnea patients, which reduced significantly

after receiving medical or surgical treatment.[6] Frequent arousals during the night in OSA alter the

activity of hypothalamic-pituitary-adrenal axis. Repeated and reversible airway obstruction lead to

intermittent hypoxia, sleep fragmentation and decreased sleep duration, causing cortisol and

catecholamine release. [24–27] Cortisol release is associated with higher levels of stress. [28] This

may be contributing to acute stress development in patients of OSA.

Second, we assessed if risk for OSA predicts the quality of sleep. As expected, we found higher

odds of poor-quality sleep in those at high risk of OSA as compared to those at low risk of OSA.

Poor quality of sleep is one the major symptoms of OSA and causes excessive daytime

sleepiness and fatigue. [1] Quality of sleep is closely related to number and extent of arousals.

Frequent arousals are a hallmark of OSA. [1,9]

Third, we assessed if sleep quality can predict acute stress. Using ordinal logistic regression, we

found that poor sleep quality leads to 3.5 times higher chance of having moderate/ severe acute

stress. Some other studies have found that increased perceived stress and sleep quality are

associated with each other. [29,30] Doane and Thurston found that lower sleep efficiency predicted

higher next day stress in adolescents. [31] A few studies have examined the directionality of the

relation between stress and sleep and found it to be bidirectional. [27,31] Sleep quality affects the

quality of replenishment of cognitive processes and energy level. [32] Experimentally produced

sleep restriction or sleep loss produces performance decline the next day. [32,33] Our study is

unable to comment on causality or directionality of effect due to a cross-sectional design. The

importance of our study lies in relating the sleep quality in everyday settings of the participants to

acute stress.

Mediation analysis proves our hypothesis that poor sleep quality mediates the relationship

between OSA and acute emotional stress. Repeated inspiratory airway occlusion in OSA leads to

multiple stressors. The stressors may be frequent episodes of hypoxemia of varying duration and

severity, disrupted sleep due to frequent arousals, reduced total sleep time, and strenuous

respiratory efforts resulting in the generation of severe negative intrathoracic pressures. [35] The

neuro-hormonal, vascular and endocrine responses to OSA include excessive sympathetic

activation which may carry over to the daytime awake state. [36] The powerful sympathetic

excitatory effects may cause increased glucocorticoids and catecholamines and translate into

acute stress. [37]

One of the limitations of our study is the inability to establish the diagnosis of OSA with

polysomnography. Further studies using polysomnography for diagnosis of OSA with objective

measures of stress such as cortisol levels are required to confirm the findings. Secondly, the

duration and onset of acute stress and OSA cannot be determined due to cross sectional design,

and hence the temporal association of acute stress with OSA cannot be established.

Based on the findings of our study, we recommend screening for OSA in young adults presenting

with acute stress, particularly in overweight males. Further studies using polysomnography for

confirmatory diagnosis of OSA and a longitudinal design are recommended to establish causality.

Conclusion

197 Increased risk to obstructive sleep apnea among young adults is associated with acute stress,

198 and the relationship is mediated by sleep quality.

Acknowledgments

200 We would like to acknowledge the Indian Council of Medical Research. This study was conducted

201 under the Short-Term Studentship Award (STS-2019) conferred by the Indian Council of Medical

202 Research.

Disclosure

204 The authors report no conflicts of interest in this work.

205 The abstract of this paper was presented at the 15th World Sleep Congress 2019, September 21-

206 25, at Vancouver, Canada as a poster presentation with interim findings. The poster's abstract

207 was published in "Poster Abstracts" in the journal 'Sleep medicine' and is available at

208 https://www.sciencedirect.com/journal/sleep-medicine/vol/64/suppl/S1.

References

1. Sleep-related breathing disorders in adults: recommendations for syndrome definition and

measurement techniques in clinical research. The Report of an American Academy of

Sleep Medicine Task Force. *Sleep*. 1999;22(5):667-689.

2. Sharma SK, Kumpawat S, Banga A, Goel A. Prevalence and risk factors of obstructive

sleep apnea syndrome in a population of Delhi, India. *Chest*. 2006;130(1):149-156.

3. Al-Barrak M, Shepertycky MR, Kryger MH. Morbidity and mortality in obstructive sleep

apnea syndrome 2: Effect of treatment on neuropsychiatric morbidity and quality of life.

Sleep Biol Rhythms. 2003;1(2):65-74.

4. Millman RP, Fogel BS, McNamara ME, Carlisle CC. Depression as a manifestation of

obstructive sleep apnea: reversal with nasal continuous positive airway pressure. *J Clin*

Psychiatry. 1989;50(9):348-351.

5. Sforza E, Roche F. Sleep Apnea Syndrome and Cognition. *Front Neurol*. 2012;3:87.

6. Santos MAG dos, Nakano T de C, Mendes FA, Duarte BB, Marone SAM. Emotional Stress

Evaluation of Patients with Moderate and Severe Sleep Apnea Syndrome. *Int Arch*

Otorhinolaryngol. 2017;21(1):28-32.

7. Baldwin CM, Griffith KA, Nieto FJ, O'Connor GT, Walsleben JA, Redline S. The association

of sleep-disordered breathing and sleep symptoms with quality of life in the Sleep Heart

Health Study. *Sleep*. 2001;24(1):96-105.

8. Reynolds CF 3rd, Kupfer DJ, McEachran AB, Taska LS, Sewitch DE, Coble PA.

Depressive psychopathology in male sleep apneics. J Clin Psychiatry. 1984 Jul;45(7):287-

90.

9. Macey PM, Woo MA, Kumar R, Cross RL, Harper RM. Relationship between Obstructive

Sleep Apnea Severity and Sleep, Depression and Anxiety Symptoms in Newly-Diagnosed

Patients. *PLoS ONE.* 2010;5(4).

10. Chiang JJ, Bower JE, Irwin MR, Taylor SE, Fuligni AJ. Adiposity moderates links from early

adversity and depressive symptoms to inflammatory reactivity to acute stress during late

adolescence. *Brain Behav Immun.* 2017;66:146-155.

11. Aggarwal K, Akhtar N, Mallick H. Social jet lag, screen time, physical activity and sleep

quality in Indian young adults. *Sleep Med.* 2019;64:S3-S4.

12. Chung F, Subramanyam R, Liao P, Sasaki E, Shapiro C, Sun Y. High STOP-Bang score

indicates a high probability of obstructive sleep apnoea. *BJA Br J Anaesth.*

2012;108(5):768-775.

13. Chiu H-Y, Chen P-Y, Chuang L-P, et al. Diagnostic accuracy of the Berlin questionnaire,

STOP-BANG, STOP, and Epworth sleepiness scale in detecting obstructive sleep apnea:

A bivariate meta-analysis. *Sleep Med Rev.* 2017;36:57-70.

14. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality

Index: a new instrument for psychiatric practice and research. *Psychiatry Res.*

1989;28(2):193-213.

15. Kilpatrick, D. G., Resnick, H. S., & Friedman, M. J. (2013a). Severity of Acute Stress

Symptoms—Adult (National Stressful Events Survey Acute Stress Disorder Short Scale

[NSESSS]) [Measurement instrument].

16. Baron RM, Kenny DA. The moderator–mediator variable distinction in social psychological

research: Conceptual, strategic, and statistical considerations. J Pers Soc Psychol.

1986;51(6):1173-1182.

17. Preacher KJ and Leonardelli GJ. "Calculation for the Sobel Test: An Interactive Calculation

Tool for Mediation Tests [Online software]," 2010. Available at:

http://quantpsy.org/sobel/sobel.htm. Accessed on October 12, 2020.

18. Preacher KJ and Hayes AF. SPSS and SAS Procedures for Estimating Indirect Effects in

Simple Mediation Models. Behaviour Research Methods, Instruments, & Computers. 2004

(36):717-731.

19. Gangwisch JE, Malaspina D, Posner K, et al. Insomnia and Sleep Duration as Mediators of

the Relationship Between Depression and Hypertension Incidence. *Am J Hypertens.*

2010;23(1):62-69.

20. Wing RR, Tate D, Espeland M, et al. Weight gain prevention in young adults: design of the

study of novel approaches to weight gain prevention (SNAP) randomized controlled trial.

BMC Public Health. 2013;13(1):300.

21. Jayawardena R, Ranasinghe P, Wijayabandara M, Hills AP, Misra A. Nutrition Transition

and Obesity Among Teenagers and Young Adults in South Asia. *Curr Diabetes Rev.*

2017;13(5):444-451.

22. Nagappa M, Liao P, Wong J, et al. Validation of the STOP-Bang Questionnaire as a

Screening Tool for Obstructive Sleep Apnea among Different Populations: A Systematic

Review and Meta-Analysis. PloS One. 2015;10(12):e0143697.

23. Aggarwal K, Akhtar N, Mallick HN. High risk of sleep apnoea is associated with acute

stress in young adults. Sleep Med. 2019;64:S7.

24. Miller JN, Berger AM. Screening and assessment for obstructive sleep apnea in primary

care. *Sleep Med Rev.* 2016;29:41-51.

25. Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences.

Lancet Lond Engl. 2009;373(9657):82-93.

26. Young T, Finn L, Peppard PE, et al. Sleep Disordered Breathing and Mortality: Eighteen-

Year Follow-up of the Wisconsin Sleep Cohort. *Sleep.* 2008;31(8):1071-1078.

27. Buckley TM, Schatzberg AF. On the interactions of the hypothalamic-pituitary-adrenal

(HPA) axis and sleep: normal HPA axis activity and circadian rhythm, exemplary sleep

disorders. *J Clin Endocrinol Metab.* 2005;90(5):3106-3114.

28. McEwen BS. Central effects of stress hormones in health and disease: understanding the

protective and damaging effects of stress and stress mediators. *Eur J Pharmacol.*

2008;583(2-3):174-185.

29. Doolin J, Vilches JE, Cooper C, Gipson C, Sorensen W. Perceived stress and worldview

influence sleep quality in Bolivian and United States university students. *Sleep Health.*

2018;4(6):565-571.

30. Roskoden FC, Krüger J, Vogt LJ, et al. Physical Activity, Energy Expenditure, Nutritional

Habits, Quality of Sleep and Stress Levels in Shift-Working Health Care Personnel. *PloS*

One. 2017;12(1):e0169983.

31. Doane LD, Thurston EC. Associations among sleep, daily experiences, and loneliness in

adolescence: Evidence of moderating and bidirectional pathways. *J Adolesc*.

2014;37(2):145-154.

32. Yap Y, Slavish DC, Taylor DJ, Bei B, Wiley JF. Bi-directional relations between stress and

self-reported and actigraphy-assessed sleep: a daily intensive longitudinal study. *Sleep*.

2020;43(3).

33. Pejovic S, Basta M, Vgontzas AN, et al. Effects of recovery sleep after one work week of

mild sleep restriction on interleukin-6 and cortisol secretion and daytime sleepiness and

performance. *Am J Physiol - Endocrinol Metab*. 2013;305(7):E890-E896.

34. Vgontzas AN, Tsigos C, Bixler EO, Stratakis CA, Zachman K, Kales A, Vela-Bueno A,

Chrousos GP. Chronic insomnia and activity of the stress system: a preliminary study. J

Psychosom Res. 1998 Jul;45(1):21-31.

35. Lopez-Jimenez F, Somers VK. Stress measures linking sleep apnea, hypertension and

diabetes--AHI vs arousals vs hypoxemia. *Sleep*. 2006;29(6):743-744.

36. Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in

obstructive sleep apnea. *J Clin Invest*. 1995;96(4):1897-1904.

37. Elenkov IJ, Chrousos GP. Stress hormones, proinflammatory and antiinflammatory

cytokines, and autoimmunity. *Ann N Y Acad Sci*. 2002;966:290-303.

Table 1 Basic descriptives of the study participants

S.No	Characteristic	Total (n=493)	Male (n=256)	Female (n=237)
1	Age (years) Mean ± SD	20.3±1.53	20.04±1.46	20.59±1.57
2	BMI (Kg/m²) Mean ± SD	22.7±3.77	22.97±3.49	20.44±3.91
3 Acute stress				
	None (0)	69 (14%)	43 (8.7%)	26 (5.3%)
	Mild (1)	248(50.3%)	134(27.2%)	114(23.1%)
	Moderate(2)	109(22.1%)	49 (9.9%)	60(12.2%)
	Severe (3)	51(10.3%)	25(5.1%)	26(5.3%)
	Extreme (4)	16 (3.2%)	5 (1.0%)	11(2.2%)
5 Risk of Obstructive Sleep Apnea				
	Low risk	420 (85.2%)	192 (75%)	228 (96.2%)
	Intermediate + High risk	73 (14.8%)	64 (25%)	9 (3.7%)
6 Sleep quality (PSQI)				
	Good (Score 1-5)	315 (63.9%)	173 (35.1%)	142 (28.8%)
	Poor (Score 5- 21)	178 (36.1%)	83 (16.8%)	95 (19.3%)

314 **Notes:**

315 **Abbreviations:** SD, Standard Deviation, PSQI, Pittsburg Sleep Quality Index

317 Table 2: Correlation between acute stress, Sleep Quality and Risk of OSA

	Acute stress	Sleep quality	Risk of OSA
Acute stress	1	.328**	.160**
PSQI		1	.150**
SB			1

Abbreviations: OSA, Obstructive Sleep Apnea

Figure 1 Mediation model

Figure 2 Mediation model with effect size.

Legend: a, Regression coefficient of pathway 'a'; b, Regression coefficient of pathway 'b'; c, Regression
coefficient of pathway 'c'; S_a, Standard error of the coefficient 'a', S_b, Standard error of the coefficient 'b', S_c,
Standard error of the coefficient 'c'
328