**Full Title**: Ancestrally and Temporally Diverse Analysis of Penetrance of Clinical

Variants in 72,434 Individuals

**Running Title**: Penetrance of Clinical Variants

**Authors**: Iain S. Forrest[1,2,3,4], Kumardeep Chaudhary[1,3,4,5], Ha My T. Vy[1,4], Shantanu

Bafna[1,4], Daniel M. Jordan[1,4], Ghislain Rocheleau[1,4], Ruth J.F. Loos[1,4,6], Judy H.

Cho[1,3,4,7], Ron Do[1,3,4].

**Author Affiliations**:

1.  The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine

at Mount Sinai, New York, NY, USA

2.  Medical Scientist Training Program, Icahn School of Medicine at Mount Sinai, New

York, NY, USA

3.  The Bio*Me* Phenomics Center, Icahn School of Medicine at Mount Sinai, New York,

NY, USA

4.  Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount

Sinai, New York, NY, USA

5.  The Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of

Medicine at Mount Sinai, New York, NY, USA

6.  The Mindich Child Health and Development Institute, Icahn School of Medicine at

Mount Sinai, New York, NY, USA

7.  Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY,

USA

**Corresponding Author**:

Ron Do, PhD

Annenberg Building, Floor 18 Room 80A

1468 Madison Ave

New York, NY-10029

Phone Number: 212-241-6206 | Fax Number: 212-849-2643

ron.do@mssm.edu

**Word Count**: 3,840

**No. of Tables and Figures**: 6

**No. of Supplementary Figures**: 5

**No. of Supplementary Tables**: 10

**ABSTRACT**

A major goal of genomic medicine is to quantify the disease risk of genetic variants.
Here, we report the penetrance of 37,772 clinically relevant variants (including those
reported in ClinVar[1] and of loss-of-function consequence) for 197 diseases in an
analysis of exome sequence data for 72,434 individuals over five ancestries and six
decades of ages from two large-scale population-based biobanks (Bio*Me* Biobank and
UK Biobank). With a high-quality set of 5,359 clinically impactful variants, we evaluate
disease prevalence in carriers and non-carriers to interrogate major determinants and
implications of penetrance. First, we associate biomarker levels with penetrance of
variants in known disease-predisposition genes and illustrate their clear biological link to
disease. We then systematically uncover large numbers of ClinVar pathogenic variants
that confer low risk of disease, even among those reviewed by experts, while
delineating stark differences in variant penetrance by molecular consequence.
Furthermore, we ascertain numerous variants present in non-European ancestries and
reveal how increasing carrier age modifies penetrance estimates. Lastly, we examine
substantial heterogeneity of penetrance among variants in known disease-
predisposition genes for conditions such as familial hypercholesterolemia and breast
cancer. These data indicate that existing categorical systems for variant classification
do not adequately capture disease risk and warrant consideration of a more quantitative
system based on population-based penetrance to evaluate clinical impact.

The advent of high-throughput sequencing has led to an exponential increase in

genetic tests for health-related purposes[2,3]. The American College of Medical Genetics

& Genomics (ACMG) recommends clinical action if a pathogenic genetic variant is

found in one of 59 genes (hereafter ACMG59)[4]. This "genetics-first" approach is feasible

if variant pathogenicity is ascertained with high confidence and veracity. A large

database of human variation and phenotypes, ClinVar[1], classifies variant pathogenicity

in a scheme that informs clinical interpretation of genetic test results (e.g., "pathogenic",

"uncertain significance", "benign")[5]. However, most ClinVar variants are of uncertain

clinical significance, and misclassification of 12% of pathogenic and 90% of conflicting

variants has inflated their pathogenicity[6]. Variants implicated in diseases from breast

cancer to cardiomyopathy have been scrutinized for overestimated claims of

pathogenicity[7] and many variants classified as pathogenic were recently downgraded to

lower ClinVar classes[8]. Any unreliability of pathogenicity, already a categorical rather

than quantitative metric of disease risk, further diminishes its clinical value. There is

therefore a critical need to provide accurate information for a variant's effect on disease

to guide patient care.

Penetrance, the probability of a disease phenotype given a particular genotype[9],

is unknown for the vast majority of variants in ClinVar and genetic databases. Well-

known exceptions include ~60% of carriers of certain *BRCA1* and *BRCA2* variants

develop breast cancer by 70 years of age[10] and 73% of individuals heterozygous for

*LDLR* variants present with hypercholesterolemia[11]. Findings of highly penetrant

variants immediately guide treatment, such as statins, and screening, such as

mammograms[12,13]. Penetrance thus gives meaningful and actionable information for

variants by quantifying a carrier's disease risk.

Until recently, penetrance has been primarily derived from family-based or

clinical cohort studies[14–17]. These typically focus on a small number of genes and

inherently maximize penetrance estimates by recruiting patients with disease or family

history of disease, and are therefore susceptible to ascertainment bias[18]. Small sample

size and genetic or environmental modifiers further limit their ability to reliably appraise

penetrance[19]. In contrast, the recent introduction of the UK Biobank (UKB) and other

population-based biobanks has created a trove of genetic and phenotypic data for large

numbers of unrelated individuals[20,21]. Exome sequences coupled to electronic health

records (EHRs) provide an unprecedented opportunity for using a population-based

method to measure penetrance on a large scale with less ascertainment bias than

traditional studies[22]. In addition, multi-ethnic biobanks such as Mount Sinai's Bio*Me*

Biobank (Bio*Me*)[23] enable the analysis of penetrance across diverse ancestries. Recent

studies have begun to use a population-based approach to explore penetrance of genes

in breast cancer[24] and familial hypercholesterolemia (FaH)[25], yet the population-based

penetrance of most variants, genes, and diseases remains uncharacterized.

Here, we perform a comprehensive analysis of variant penetrance observed in

two large-scale EHR-linked population-based biobanks (Bio*Me* and UKB). We

measured the penetrance of 37,772 variants, including a high-quality set of 5,359

clinically impactful variants, for 197 diseases of dominant inheritance using exome

sequences from 72,434 individuals. Penetrance was analyzed in all individuals and

separately in five ancestries across six decades of ages. We investigated determinants

of penetrance, including ClinVar pathogenicity, variant molecular consequence, and

carrier ancestry and age. The clinical manifestation of variants was scrutinized by

examining metabolite levels, biological measurements, and physician notes for carriers,

and clinically significant variants and genes were highlighted. This highly scalable,

population-based approach of generating penetrance estimates for thousands of

clinically relevant variants expands our knowledge of the genetic influences on disease

and has the potential to advance genomic medicine.

**RESULTS**

Analysis of penetrance of clinical variants

A graphical abstract is provided (**Supplementary Figure 1**). Whole-exome

sequencing, quality control, and filtering were performed on 80,773 samples from two

population-based EHR-linked biobanks (**Methods**) to generate a final dataset with

72,434 individuals (**Table 1**). Demographics, self-reported ancestry, biological

measurements, and International Classification of Disease 10 (ICD-10) diagnosis codes

were available for individuals in the final dataset. We defined case-control status for

over 400 non-recessive diseases in ClinVar sourced from the Systematized

Nomenclature of Medicine Clinical Terms (SNOMED CT)[26] using ICD-10 codes[27], of

which 197 had at least one case in the final dataset (**Supplementary Table 1**). We

analyzed 37,772 variants that were either reported in ClinVar for at least one of the

aforementioned diseases or were of predicted loss-of-function (LoF) consequence

(**Supplementary Table 2**). A stringent set of 5,359 clinically impactful (impactful)

variants—defined by pathogenic/likely pathogenic classification in ClinVar or LoF

annotation (splice acceptor/donor, stop gained/lost, frameshift, start lost) with Variant

Effect Predictor[28] in a gene mediating disease via LoF mechanism, and in a gene with

non-recessive inheritance—was used for downstream analyses unless otherwise stated

(**Supplementary Figure 2**). We identified carriers with at least one allele (hereafter

referred to as carriers) for each variant and computed the proportion of carriers affected

with disease to determine penetrance. Variant penetrance was estimated in all carriers

and separately in carriers of five diverse ancestries and six age ranges. The proportion

of non-carriers affected with disease was also quantified for each variant to ascertain

disease prevalence among the population in the absence of a scorable variant.

<u>Validation of phenotyping and penetrance approach</u>

We performed a series of analyses to validate our population-based approach of

phenotyping and computing penetrance. First, we selected nine diseases from a variety

of systems—age-related macular degeneration (AMD), arrhythmogenic right ventricular

cardiomyopathy (ARVC), FaH, familial breast cancer (FBC), type 2 diabetes (T2D),

etc.—and identified cases using both ICD-10 diagnosis codes and previously published

clinical algorithms[29–37]. We then evaluated and compared the penetrance of 208 ClinVar

pathogenic variants corresponding to the diseases in Bio*Me* using both phenotype

approaches (**Supplementary Table 3**). Similar penetrance values were observed for all

diseases. Three diseases (Brugada syndrome, hepatocellular carcinoma, and

pulmonary arterial hypertension) had equivalent mean variant penetrance with both

approaches; mean variant penetrance of the other six diseases varied by 8% or less. A

tabulated list of these results is provided (**Supplementary Table 4**).

Second, we further assessed the accuracy of our phenotyping approach by

manually reviewing physician notes in the problem list (PL) for cases and controls of six

representative diseases: Alzheimer's disease (AD), AMD, FaH, FBC, stroke, and T2D.

We randomly sampled 50 ICD-10-based cases and controls for each disease and

examined their PL for symptoms or findings indicative of a diagnosis while blinded to

ICD-10 disease status (**Supplementary Table 5**). We observed high concordance in

case-control classification between ICD-10- and PL-based phenotyping, ranging from

100% (for AD) to 93% (AMD, FaH). Case concordance ranged from 88% (AMD, FaH)–

100% (AD, FBC, T2D), while control concordance ranged from 94% (T2D)–100% (AD,

stroke).

<u>Biomarker levels linked to variant penetrance</u>

We investigated metabolite and clinical measurements of carriers of impactful

variants with different penetrance in genes with clear biological links to three diseases:

FaH[12], maturity-onset diabetes of the young (MODY)[38], and obesity[39,40] (**Figure 1**).

Carriers of penetrant variants for FaH in *LDLR* had greater low-density lipoprotein

cholesterol (LDL-C) and total cholesterol levels than carriers of incompletely and

nonpenetrant variants (**Figure 1a**). Using linear regression, we observed robust

association between variant penetrance and LDL-C (effect size [β]=0.53 mg/dL per 1%

increase in penetrance, standard error [SE]=0.12; *P*=0.004) and total cholesterol levels

(β=0.98 mg/dL, SE=0.18; *P*=0.002) adjusting for clinical covariates and statin use.

Analogously, carriers of penetrant variants for MODY in hepatocyte nuclear factor-1α

(*HNF1A*) had elevated glucose and hemoglobin A1c (HbA1c) levels compared to

carriers of incompletely penetrant variants (**Figure 1b**). Variant penetrance was

significantly associated with glucose (β=0.76 mg/dL, SE=0.19; *P*=0.008) and HbA1c

levels (β=0.083% [2.4 mg/dL], SE=0.020 [0.57 mg/dL]; *P*=0.01) adjusted for clinical

covariates and diabetes medication use. Lastly, carriers of penetrant variants for obesity

in uncoupling protein 3 (*UCP3*) had higher body mass index (BMI) than carriers of

incompletely and nonpenetrant variants (**Figure 1c**). Variant penetrance was strongly

associated with BMI (β=0.24 kg/m$^2$, SE=0.044; *P*=6x10$^{-4}$) and weight (β=1.3 lbs [0.59

kg], SE=0.26 [0.12 kg]; $P$=8x10$^{-4}$) adjusting for clinical covariates. Together, these

analyses using published phenotype algorithms, physician notes, and biological

measurements support the validity of our population-based penetrance method and

illustrate the clinical presentation of impactful variants.

<u>Distribution of observed variant penetrance and disease risk</u>

We examined the distribution of disease risk conferred by 5,359 impactful

variants and 157 corresponding diseases. Risk difference (RD) is the difference

between the prevalence of disease in carriers and non-carriers, and represents the

excess risk of disease attributed to the variant of interest. A summary of mean variant

RD for all 157 diseases is provided (**Supplementary Table 6**). We observed 565 (11%)

variants with RD exceeding 0.05 for 55 (35%) diseases (**Figure 2**). In contrast, we

detected a very large number of weakly penetrant and nonpenetrant variants that confer

little to no disease risk (4,794 [89%] variants with RD≤0.05), which can be attributed in

part to ClinVar pathogenic variants from traditional studies that maximize penetrance

estimates and have biased ascertainment[18]. Recent studies have demonstrated

overestimation of the disease rates of these variants[6–8], indicating a need for

population-based approaches to more unbiasedly gauge penetrance.

Next, we investigated the prevalence of disease in carriers (i.e., penetrance) and

non-carriers of impactful variants according to ClinVar pathogenicity (pathogenic,

uncertain, conflicting, benign) (**Figure 3a**), ClinVar review status (expert reviewed,

multiple submitters, single submitter, no assertion criteria) (**Figure 3b**), and molecular

consequence (**Figure 3c**). We hypothesized that variants reported as pathogenic in

ClinVar, reviewed by experts, or with LoF consequence have greater *a priori* evidence

of disease risk and are expected to have higher penetrance than variants reported as

benign, without criteria assertion, or of synonymous consequence. In agreement with

these expectations, we observed that mean variant penetrance was highest among

pathogenic (6.9% vs. 0.86% next highest mean for uncertain class; $P=5\times10^{-270}$), expert

reviewed (18% vs. 6.4% next highest mean for multiple submitters; $P=9\times10^{-17}$), and

frameshift variants (10% vs. 4.1% for missense; $P=2\times10^{-13}$). The mean penetrance of

ClinVar variants with conflicting (0.75%), uncertain (0.86%), and benign clinical

significance (0.085%) was similar, warranting caution when interpreting variants that do

not fully meet the ACMG's strict pathogenic criteria[41]. We also note that the mean

penetrance of ClinVar pathogenic variants was only 6.9% and only 18% for those with

the highest review status (expert reviewed), indicating that these broad classifications

do not adequately capture all variants with large effect on disease as measured by

penetrance.

We conducted several sensitivity analyses to account for differing sample sizes

of penetrance estimates. Penetrance distributions according to ClinVar pathogenicity,

review status, and molecular consequence remained similar when stratified by

increasing thresholds of sample sizes (**Supplementary Figure 3**). Singletons (very rare

variants appearing only once) comprised a large portion of the impactful variants

(3,507/5,359 [65%]) and statistically are completely penetrant or nonpenetrant. To test

the validity of singleton penetrance, we evaluated the proportion of penetrant singletons

by ClinVar pathogenicity, review status, and molecular consequence (**Supplementary**

**Figure 4**). If singleton penetrance were accurate, the proportion of penetrant singletons

among pathogenic, expert reviewed, or LoF singletons would be expected to exceed

that of benign, non-expert reviewed, or non-LoF singletons, respectively. We observed

that the proportion of penetrant singletons was in fact greatest among pathogenic (13%

vs. 2.6% in next highest class of conflicting; $P$=2x10$^{-19}$), expert reviewed (41% vs. 12%

in next highest group of multiple submitters; $P$=4x10$^{-11}$), and frameshift singletons (16%

vs. 7.8% for missense; $P$=1x10$^{-3}$). We therefore retained penetrance estimates of

smaller sample sizes in our analyses.

<u>Ancestral and temporal dimensionality of penetrance data</u>

Most penetrance studies to date have focused on one ancestry, typically

European[42,43], with few notable exceptions[11,23,44]. Yet it is well-known that allele

frequency (AF) and disease risk of variants can vary substantially in different

populations[45,46]. Thus, we computed penetrance among multiple self-reported

ancestries, including European, African, Hispanic, and Asian ancestries. To

demonstrate this diversity, we identified numerous ancestry-specific impactful variants.

Out of 2,209 impactful variants in Bio*Me*, we observed 75 (3.4%), 43 (1.9%), 17

(0.77%), and 71 (3.2%) variants exclusively present in African, Hispanic, Asian, and

European ancestry, respectively. Out of 3,408 impactful variants in UKB, we observed 8

(0.23%), 11 (0.32%), and 235 (6.9%) variants specific to African, Asian, and European

ancestry, respectively. Furthermore, we identified ancestry-specific variants that were

highly penetrant, such as an Asian ancestry-specific pathogenic frameshift variant in

*HBB* (NC_000011.10:c.5226994_5226995insC) associated with a greatly increased risk

of thalassemia (RD=0.99; $P$=9x10$^{-6}$) and a European ancestry-specific pathogenic

frameshift variant in *PALB2* (NC_000016.9:c.23647358_23647359del) associated with

a significantly elevated risk of breast cancer (RD=0.92; *P*=0.007).

We also delineated variant penetrance based on different age thresholds of

carriers ranging from at least 20 years (≥20) to at least 70 years (≥70). Age of disease

onset is pertinent for estimating penetrance: congenital or early onset diseases will have

manifested in older carriers of penetrant variants, whereas later onset diseases may not

have presented yet in younger carriers. To address temporality, we characterized the

observed change in variant penetrance with increasing carrier age for 157 diseases

corresponding to the impactful variants, stratified by age of disease onset (**Figure 4**;

**Supplementary Figure 5**). We first annotated and grouped the diseases by age of

onset: Earlier (congenital, childhood, or adolescent), Later (adulthood), or Any

(**Supplementary Table 7**). For each disease group, we then calculated the change in

variant penetrance (ΔPenetrance) between the lowest carrier age threshold and

increasing age thresholds (e.g., variant penetrance for ≥20 versus ≥30, ≥20 versus ≥40,

etc.). As expected, the mean ΔPenetrance increased in the Later disease group with

higher carrier age thresholds in Bio*Me* (**Figure 4a**). The largest mean ΔPenetrance was

+1.8% in the Later disease group when comparing ≥20 and ≥70, while mean

ΔPenetrance remained ~0% in the Earlier and Any disease groups. In UKB, no

significant differences in mean ΔPenetrance were observed in the Later disease group

(**Figure 4b**) likely due to its older age range of carriers (40-69 years) whereby Later

diseases would have manifested. Thus, penetrance estimates of diseases with early or

any age of onset remained stable over time, while penetrance estimates of diseases

with later age of onset predictably increased with higher carrier ages.

<u>Clinical utility of variant penetrance data</u>

A genotype-first approach in genomic medicine must identify penetrant variation in clinically significant genes to inform screening, management, and treatment. The ACMG recommends reporting secondary findings of pathogenic variants in the ACMG59, but acknowledges that insufficient data on penetrance requires ongoing study and revision[49]. *BRCA1* and *BRCA2* are both in the ACMG59, and a recent study estimated the population-based penetrance of pathogenic variants in these genes for FBC using data from the UKB[50]. However, penetrance was estimated at the gene level, and non-ClinVar variants were omitted as were variants in other genes that have been shown to confer significant risk of breast cancer, such as *PALB2*, *CHEK2*, *ATM*, *PTEN*, and others[24]. Thus, we investigated the penetrance of impactful variants in 10 known or suspected breast cancer-predisposition genes (*BRCA1*, *BRCA2*, *PALB2*, *CHEK2*, *ATM*, *PTEN*, *CDH1*, *BARD1*, *BRIP1*, and *RAD51D*) to illustrate the clinical value of a population-based approach of determining penetrance at the variant level (**Figure 5a**). The highest disease risk on average was conferred by impactful variants in *BRCA1* (mean variant penetrance=38%, mean RD=0.32; P=$2\times10^{-6}$), *BRCA2* (mean variant penetrance=38%, mean RD=0.32; *P*=$1\times10^{-10}$), and *PALB2* (mean variant penetrance=26%, mean RD=0.20; *P*=0.009).

These data clarify the disease risk of previously reported variants in ClinVar and novel variants not yet reported in ClinVar or the literature. We highlighted examples of variants that are significantly associated with risk of seven diseases, including FBC, to demonstrate this dual utility (**Supplementary Table 8**). Many impactful variants in

*BRCA2* were strongly associated with FBC, such as a known pathogenic frameshift

variant (NC_000013.11:c.32340301del; penetrance=47%, RD=0.39; *P*=9x10$^{-6}$) and a

previously unreported frameshift variant (NC_000013.11:c.32340630_32340631del;

penetrance=100%, RD=0.96; *P*=4x10$^{-13}$). Evidence of high penetrance strengthens a

candidate variant's case for clinical significance, while assertions of pathogenicity are

refined by quantitative disease risk estimates.

Notably, we observed substantial heterogeneity in the penetrance of variants

even within the same disease-predisposition gene. Impactful variants in *LDLR*, for

instance, exhibited a wide range of penetrance for FaH (**Figure 5b**). Similarly, there was

a large distribution of penetrance associated with impactful variants in *BRCA1* (variant

penetrance standard deviation [SD]=45%, RD SD=0.44), *BRCA2* (variant penetrance

SD=46%, RD SD=0.46), and *PALB2* (variant penetrance SD=42%, RD SD=0.42) for

FBC (**Figure 6a**). Instead of coarsely categorizing variants as simply pathogenic versus

non-pathogenic and collapsing all variants in a gene, these data reveal granularity and

nuance in the penetrance of individual variants.

**DISCUSSION**

A major goal of genomic medicine is to tailor clinical care of patients to their unique genetic composition, especially penetrant variants. Scalability and accuracy are crucial for determining penetrance and employing its information in a clinical setting. Conventional penetrance estimates from family-based or clinical cohort studies typically focus on one variant or gene at a time[14–17] (limited scalability), and have small sample size, ascertainment bias, inconsistent carrier screening, and genetic/environmental confounders[18,19] (limited accuracy). A proliferation of population biobanks, such as UKB and Bio*Me*, has made available thousands of genetic and phenotypic records[20,21,23]. This raises the possibility of using a population-based method to measure penetrance whereby large numbers of unrelated carriers are assessed[22]. Here, we performed a comprehensive assessment of variant penetrance using 72,434 exomes from two large-scale population-based biobanks, with the dataset freely accessible and provided in **Supplementary Tables 9** and **10**.

Reliable penetrance estimates depend on reliable phenotyping of carriers. We performed robust validation analyses to support our phenotyping and penetrance strategy. First, we computed penetrance for nine diseases using both diagnosis codes and previously published clinical algorithms, finding high concordance in all diseases tested. Second, we manually curated physician notes for six diseases to verify ICD-10-based diagnoses. Third, we extracted laboratory and clinical test results (e.g., levels of lipids, glucose, HbA1c, and BMI) for carriers and observed that penetrance was strongly associated with relevant quantitative traits, adjusted for medications and clinical covariates. Hence, by validating our phenotyping and penetrance method, and

evaluating penetrance for thousands of variants in a high-throughput manner, we

ensure both accuracy and scalability. As penetrance studies grow, similar validation

analyses should be implemented to certify fidelity of penetrance estimates.

While several efforts have begun to probe the upward bias of penetrance

estimates in traditional studies, there has not yet been a systematic investigation of the

pervasiveness of overestimated penetrance. In the present study, we examined the

distribution of disease risk associated with a high-quality set of 5,359 impactful variants

and 157 diseases. Persistently lower disease risk was observed for many diseases, with

a few exceptions of highly penetrant variants such as in *BRCA1* and *BRCA2*. This can

be interpreted in light of a few important considerations: 1) small sample size and

stochasticity for rarer variants may contribute to variable penetrance estimates;

however, a number of analyses supported the accuracy of our penetrance

measurements with respect to sample size, including for singletons (**Supplementary**

**Figure 3**, **Supplementary Figure 4**); 2) ascertainment of cases in conventional studies

has been shown to inflate the disease risk of variants[18,19] and recent population-based

studies with lower penetrance estimates for FaH and developmental disorders bolster

this explanation[11,22]; 3) reported pathogenicity does not equate with penetrance, as we

found by interrogating whether existing qualitative classification systems in ClinVar

capture the quantitative disease risk of variants. ClinVar pathogenic variants were more

penetrant than variants of other ClinVar classes yet still weakly penetrant overall, in line

with previous studies[6–8] and consistent with ClinVar's definition of pathogenic that

includes "low penetrance" variants (https://www.ncbi.nlm.nih.gov/clinvar/docs/clinsig/).

We then examined a major driver of variation in penetrance among ClinVar pathogenic

variants: when stratified by ClinVar review status (evidence for clinical significance of a

variant), pathogenic variants that were not expert reviewed had lower penetrance.

In light of these findings, it is incumbent to discuss whether variants reported as

pathogenic but empirically shown to have low penetrance should be classified

differently, or whether categorical systems of disease risk (pathogenic versus non-

pathogenic) should be complemented with a quantitative system based on penetrance[7].

Recent commentary by Khera and Hegele proposed a new paradigm for classifying FaH

based on two parameters: carrier status for a pathogenic *LDLR* variant and severity of

hypercholesterolemia[51]. This advanced schema for disease classification, and others

like it, would greatly benefit from knowledge of penetrance for pathogenic *LDLR* variants

to better stratify disease risk and personalize medical care.

We also emphasize the importance of including diverse ancestries and age

ranges when characterizing variant penetrance. Populations differ by AF and disease

factors[45,46], yet most genetic studies have focused on Europeans[52,53]. Here, we capture

ancestry-specific penetrance in detail, identifying over a hundred variants in non-

European ancestries. While past studies have reported age-dependent penetrance in

age-related diseases such as FBC[17,54], amyotrophic lateral sclerosis[55,56], and obesity[57],

these typically estimate gene-based penetrance whereby all variation in a gene is

aggregated. In contrast, we evaluate age-dependent penetrance at the variant level. As

expected, early onset diseases showed stable penetrance over time while later onset

diseases had increasing penetrance with older carriers. For example, the observed

penetrance of an expert reviewed ClinVar pathogenic frameshift variant in *BRCA2*

(NC_000013.11:c.32340301del) increased from 38% in European ancestry carriers ≥40

years of age to 56% in those ≥60 years of age.

There were several study limitations. First, ICD-10 diagnosis codes from the EHR

were used to define case-control status[26,27] (**Supplementary Table 1**). While commonly

used in EHR-linked biobank studies[61,62], there may be some misclassification[63–66].

Though several validation analyses reinforced the soundness of our phenotyping, these

were completed for a subset of the diseases. Second, we cannot exclude the possibility

of potential bias in our datasets. Bio*Me* is predominantly composed of individuals

recruited from the Mount Sinai Health System and may have a higher burden of

diseases and therefore penetrance estimates. In contrast, the preponderance of healthy

volunteers in UKB may lead to conservative estimates of penetrance. Third, though our

sample size exceeded 72,000 exomes and enabled us to ascertain rare variants, many

penetrance estimates for rare variants (e.g., singletons) are based on low numbers of

carriers and may produce variable estimates. We included raw counts of carriers and

non-carriers for each variant in the dataset so that estimates may be interpreted

accordingly. Fourth, we mapped variants to diseases based on disease genes reported

in ClinVar from pathogenic variant submissions. Inaccuracies in ClinVar submissions,

and therefore our mapping from variant to disease, are possible, though we manually

checked mappings extensively against the literature for accuracy.

In conclusion, we present a large-scale, systematic investigation of variant

penetrance that leveraged thousands of exomes linked to the EHR. While we made

available the penetrance measurements of all 37,772 variants for full transparency, we

utilized a stringent set of 5,359 impactful variants that were ClinVar pathogenic or LoF in

a non-recessive gene for many analyses. We demarcated differences in disease risk

among variants of distinct ClinVar pathogenicity classes, review status, and molecular

consequences. We also accessed a rich resource of clinical phenotypes to thoroughly

explore penetrance: detailed physician notes, medications, biological measurements,

and laboratory results. Critically, we investigated multiple dimensions of penetrance

data spanning five ancestries and six decades of age thresholds. This study provides a

blueprint for future studies to efficiently and accurately determine variant penetrance,

the results of which will greatly improve our understanding of the genetic underpinnings

of human disease.

**METHODS**

Study populations and sample filtering

A flowchart of the study design is provided (**Supplementary Figure 1**). We evaluated penetrance from individuals in two large-scale electronic health record (EHR)-linked population-based biobanks: The BioMe Biobank (BioMe) and UK Biobank (UKB). The study protocols were approved by the Institutional Review Board (IRB) of the Icahn School of Medicine at Mount Sinai. Use of data from UKB was completed and approved using the UK Biobank Resource under Application Number 16218. Informed consent was obtained for all study participants in both BioMe and UKB through the approved IRB protocols. An overview of the demographics and clinical traits for both study populations is provided (**Table 1**). BioMe is an EHR-linked biobank for ~50,000 patients of African, Hispanic, European, and Other (Asian, Native American, and miscellaneous) self-reported ancestry who are recruited from the Mount Sinai Health System in Manhattan, NYC from 2007 onwards. All BioMe participants consented to providing biological and DNA samples linked to de-identified EHRs. A subset of the individuals (n=31,250) was exome sequenced before undergoing quality control. A total of 229 samples with discordance between genetic sex and sex listed in the manifest, low coverage, contamination, low call rate, or duplications were excluded, leaving 30,813 samples. In addition, samples lacking complete demographic data (n=345), younger than 20 years of age (n=610), or without International Classification of Diseases-Clinical Modification 10 (ICD-10) diagnosis data (n=819) were removed to generate the final study set of 29,039 samples.

The UKB is a population-based longitudinal cohort of ~500,000 individuals chiefly

of British self-reported ancestry between 40-69 years of age who were enrolled at

various sites across the United Kingdom between 2006-2010[21,67]. All individuals

consented to providing medical history, demographic data, and DNA samples. A subset

of 49,960 individuals had their exome sequenced and passed standard quality control

(QC) filters, described extensively elsewhere[68]. We further excluded samples that

lacked complete demographic information (n=2), or did not have ICD-10 diagnosis

codes available (n=6,563), leaving a final study set of 43,395 samples for analysis.

<u>Whole-exome sequencing and quality control</u>

In Bio*Me*, variant call files (VCFs) produced by Illumina v4 HiSeq 2500 contained

9,202,884 variants that were called in the samples. Goldilocks Filter (GF) was

implemented on the VCFs. For single nucleotide polymorphisms (SNPs), cells with

depth-normalized quality scores <3 or depth of coverage <7 were set to missing. For

insertions and deletions (indels), cells with depth-normalized quality scores <5 or depth

of coverage <10 were set to missing. Variant sites were then filtered, whereby sites of

heterozygous variation failed the Allele Balance (AB) cutoff and were removed. SNP

sites required ≥1 sample to carry an alternate AB ≥15% and indel sites required ≥1

sample to carry an alternate AB ≥20%. Together, these site filters removed 441,406

sites, leaving 8,761,478 variants after GF. Next, sites with missing genotypes for >2% of

individuals in the dataset (267,955 sites) were removed. AB was calculated for biallelic

SNPs and 320,877 sites with AB <0.3 or >0.8 were removed, leaving 8,172,646 sites.

Lastly, the dataset was filtered to regions within the target regions of the exome capture

platform (4,256,827 sites) and separated into 2 file sets for biallelic and multiallelic sites

(3,948,623 and 308,204, respectively) due to differences in QC procedures.

In UKB, exome data from the first tranche of exome sequence data generated

with the Functional Equivalence pipeline[68] was used. Sequence data and QC for UKB

are described elsewhere[21,67].

<u>Variant curation and annotation</u>

Using exome data from the 72,434 study samples, 37,772 variants associated

with ClinVar[1] diseases were ascertained using PLINK version 2.0[69]. To enrich for

penetrant variants, we selected a strict subset of 5,359 clinically impactful (impactful)

variants for most downstream analyses. These were defined by curating variant

summary information in ClinVar VCF files released in December 2019, functional

annotations from Variant Effect Predictor (VEP)[28] version 99.2, and genic mode of

inheritance from Online Mendelian Inheritance in Man (OMIM)[70]. An overview of variant

selection is provided in **Supplementary Figure 2**. First, we included variants of

pathogenic and/or likely pathogenic classification in ClinVar, and previously unreported

variants with a damaging molecular consequence (splice acceptor/donor, stop

gained/lost, frameshift, or start lost; collectively defined as LoF) annotated by VEP. LoF

variants in a gene were mapped to disease based on prior pathogenic variant

submissions in ClinVar linking genes to diseases (e.g., *BRCA1* LoF variants were

mapped to breast cancer based on prior pathogenic variant submissions in ClinVar

linking *BRCA1* to breast cancer). This filtered out benign variants or variants with

uncertain or conflicting clinical significance in ClinVar and variants with synonymous

molecular consequence. As missense variants have varying and uncertain degrees of

pathogenicity, non-ClinVar missense variants were also excluded. We also noted the

review status for each ClinVar variant, which summarizes the level of evidence

supporting a variant's claim of clinical significance ranging from the lowest level of no

assertion criteria (review status=0), to single submitter or multiple submitters with

conflicting interpretation (review status=1), to multiple submitters with no conflict of

interpretation (review status=2), to reviewed by an expert panel such as ClinGen[71]

(review status=3), to the highest level of practice clinical guidelines (review status=4).

Second, we excluded variants in genes with exclusively recessive mode of inheritance

reported in OMIM. The gene for each variant was retrieved from NCBI reference

sequences (RefSeq[72]) and corroborated with genomic coordinates in OMIM. The mode

of inheritance was mined for all phenotypes reported for each gene in OMIM and genes

were then summarized as Dominant (only dominantly inherited phenotypes), Both (both

dominantly and recessively inherited phenotypes), or Recessive (only recessively

inherited phenotypes).

Phenotyping of carriers

In both UKB and Bio*Me*, we obtained case status using ICD-10 diagnosis codes.

ICD-10 codes are commonly used in genetic population studies to define cases of a

disease[61,62] and they map directly to ClinVar diseases in the Systematized

Nomenclature of Medicine Clinical Terms (SNOMED CT)[27]. All of the 29,039 Bio*Me*

samples and 43,395 UKB samples in the final dataset had ICD-10 diagnosis codes

available. Cases were identified by the presence of a corresponding ICD-10 code while

controls were identified by the absence of all corresponding ICD-10 codes. We thereby

defined case-control status for over 400 SNOMED CT diseases of non-recessive

inheritance in ClinVar (disease inheritance was retrieved from NCBI's MedGen

database at https://www.ncbi.nlm.nih.gov/medgen/); 197 of the diseases were present

with at least one case in the final dataset. A complete list of the cases, controls, and

ICD-10 diagnosis codes for each disease is provided in **Supplementary Table 1**.

Validation of the ICD-10-based phenotyping method was performed in a set of

three analyses in Bio*Me*. First, penetrance estimates using the ICD-10 phenotypes were

compared against results using clinical algorithms from the literature for nine

diseases[29–37]: age-related macular degeneration (AMD), arrhythmogenic right

ventricular cardiomyopathy (ARVC), Brugada syndrome (BrS), familial

hypercholesterolemia (FaH), familial breast cancer (FBC), hepatocellular carcinoma

(HCC), idiopathic pulmonary arterial hypertension (IPAH), prostate cancer (PCa), and

type 2 diabetes (T2D) (**Supplementary Table 3**, **Supplementary Table 4**). Second, we

verified ICD-10 phenotypes by manually reviewing physician notes in the problems list

(PL) for six diseases: Alzheimer's disease (AD), AMD, FaH, FBC, ischemic stroke, and

T2D (**Supplementary Table 5**). For each disease, we extracted physician notes for a

random sample of 50 ICD-10-based cases and 50 ICD-10-based controls. The

presence of symptoms or findings indicative of a diagnosis corroborated ICD-10-based

cases, while the absence of symptoms or findings corroborated ICD-10-based controls.

Third and lastly, we examined the association of biomarker levels with penetrance

estimates for variants in genes with clear biological roles in three diseases: *LDLR*[12] in

FaH (measured by LDL-C and total cholesterol), *HNF1A*[38] in MODY (measured by

glucose and hemoglobin A1c), and *UCP3*[39,40] in obesity (measured by BMI and weight).

We extracted median biological measurements for carriers and medication usage

relevant to FaH (statins) and T2D (insulin, insulin analogs, pramlintide, glucagon-like

peptide 1 agonists, metformin, sulfonylurea, dipeptidyl peptidase 4 inhibitors, glitazone,

sodium-glucose transport protein 2 inhibitors, or alpha-glucosidase inhibitors) to control

for the effect of medications on biological measurements in association analyses.

<u>Statistical analysis</u>

All statistical tests and plots were made using R statistical software version

3.5.3[75]. Differences in categorical variables were assessed with a Fischer's exact test,

while differences in continuous variables were tested with a two-sided t-test.

Significance level was set at $P<0.05$ for comparisons between two groups. Risk

difference (RD) between the prevalence of disease in carriers and non-carriers was

computed, and significance was evaluated with Fisher's exact test. A strict Bonferroni

correction was applied to $P$-values in analyses involving multiple comparisons, including

comparisons of mean penetrance for five ClinVar pathogenicity classes, mean

penetrance for four ClinVar review status levels, and mean penetrance for eight

molecular consequences. Phenotype validation using PL physician notes was

performed by computing the ratio of PL-based cases to ICD-10-based cases and PL-

based controls to ICD-10-based controls to determine case and control concordance,

respectively. Association analyses of variant penetrance with biomarker levels were

performed using a multivariable linear regression, adjusted for clinical covariates of age,

sex, BMI (except in the analysis of obesity), 10 genetic principal components (PCs), and

medication usage where relevant.

<u>Data availability</u>

The UKB data may be browsed at <u>http://biobank.ndph.ox.ac.uk/showcase/</u> and

access to data can be requested at <u>https://www.ukbiobank.ac.uk/register-apply/</u>. More

information about Bio*Me* can be found at

<u>https://icahn.mssm.edu/research/ipm/programs/biome-biobank/researcher-faqs</u>. The

complete penetrance dataset used for all analyses is provided (**Supplementary Tables**

**9** and **10**) with no restrictions on the data released.

**References**

1.  Landrum, M. J. *et al.* ClinVar: public archive of relationships among sequence
variation and human phenotype. doi:10.1093/nar/gkt1113.
2.  Shendure, J., Findlay, G. M. & Snyder, M. W. Genomic Medicine–Progress,
Pitfalls, and Promise. *Cell* vol. 177 45–57 (2019).
3.  Yang, Y. *et al.* Clinical Whole-Exome Sequencing for the Diagnosis of Mendelian
Disorders. *N. Engl. J. Med.* **369**, 1502–1511 (2013).
4.  Watson, M. S. ACMG policy statement: Updated recommendations regarding
analysis and reporting of secondary findings in clinical genome-scale sequencing.
*Genetics in Medicine* vol. 17 68–69 (2015).
5.  Landrum, M. J. & Kattman, B. L. ClinVar at five years: Delivering on the promise.
*Hum. Mutat.* **39**, 1623–1630 (2018).
6.  Shah, N. *et al.* Identification of Misclassified ClinVar Variants via Disease
Population Prevalence. *Am. J. Hum. Genet.* **102**, 609–619 (2018).
7.  Manrai, A. K., Ioannidis, J. P. A. & Kohane, I. S. Clinical genomics: From
pathogenicity claims to quantitative risk estimates. *JAMA - Journal of the
American Medical Association* vol. 315 1233–1234 (2016).
8.  Xiang, J. *et al.* Reinterpretation of common pathogenic variants in ClinVar
revealed a high proportion of downgrades. *Sci. Rep.* **10**, 1–5 (2020).
9.  Griffiths, A. J., Miller, J. H., Suzuki, D. T., Lewontin, R. C. & Gelbart, W. M. *An
Introduction to Genetic Analysis*. (W. H. Freeman, 2000).
10. Mavaddat, N. *et al.* Cancer Risks for BRCA1 and BRCA2 Mutation Carriers:
Results From Prospective Analysis of EMBRACE. **105**, 812–822 (2013).
11. Khera, A. V. *et al.* Diagnostic Yield and Clinical Utility of Sequencing Familial
Hypercholesterolemia Genes in Patients With Severe Hypercholesterolemia. *J.
Am. Coll. Cardiol.* **67**, 2578–2589 (2016).
12. Goldberg, A. C. *et al.* Familial Hypercholesterolemia: Screening, diagnosis and
management of pediatric and adult patients. *J. Clin. Lipidol.* **5**, S1–S8 (2011).
13. Saslow, D. *et al.* American Cancer Society Guidelines for Breast Screening with
MRI as an Adjunct to Mammography. *CA. Cancer J. Clin.* **57**, 75–89 (2007).
14. Hobbs, H. H. *et al.* Evidence for a dominant gene that suppresses
hypercholesterolemia in a family with defective low density lipoprotein receptors.
*J. Clin. Invest.* **84**, 656–664 (1989).
15. Wohlgemuth, M. *et al.* A family-based study into penetrance in
facioscapulohumeral muscular dystrophy type 1. *Neurology* **91**, e444–e454
(2018).
16. Gong, G., Hannon, N. & Whittemore, A. S. Estimating gene penetrance from
family data. *Genet. Epidemiol.* **34**, 373–381 (2010).
17. Kuchenbaecker, K. B. *et al.* Risks of breast, ovarian, and contralateral breast
cancer for BRCA1 and BRCA2 mutation carriers. *JAMA - J. Am. Med. Assoc.* **317**,
2402–2416 (2017).
18. Park, S. *et al.* Adjusting heterogeneous ascertainment bias for genetic association
analysis with extended families. *BMC Med. Genet.* **16**, 62 (2015).
19. Cooper, D. N., Krawczak, M., Polychronakos, C., Tyler-Smith, C. & Kehrer-
Sawatzki, H. Where genotype is not predictive of phenotype: Towards an

understanding of the molecular basis of reduced penetrance in human inherited
disease. *Hum. Genet.* **132**, 1077–1130 (2013).

20. De Souza, Y. G. & Greenspan, J. S. Biobanking past, present and future:
Responsibilities and benefits. *AIDS* vol. 27 303–312 (2013).

21. Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the
Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLOS*
*Med.* **12**, e1001779 (2015).

22. Wright, C. F. *et al.* Assessing the Pathogenicity, Penetrance, and Expressivity of
Putative Disease-Causing Variants in a Population Setting. *Am. J. Hum. Genet.*
**104**, 275–286 (2019).

23. Abul-Husn, N. S. *et al.* Exome sequencing reveals a high prevalence of BRCA1
and BRCA2 founder variants in a diverse population-based biobank. *Genome*
*Med.* **12**, 2 (2019).

24. Hu, C. *et al.* A Population-Based Study of Genes Previously Implicated in Breast
Cancer. *N. Engl. J. Med.* **384**, 440–451 (2021).

25. Fahed, A. C. *et al.* Polygenic background modifies penetrance of monogenic
variants for tier 1 genomic conditions. *Nat. Commun.* **11**, 1–9 (2020).

26. SNOMED CT. https://www.nlm.nih.gov/healthit/snomedct/index.html.

27. SNOMED CT to ICD-10-CM Map. *National Library of Medicine*
https://www.nlm.nih.gov/research/umls/mapping_projects/snomedct_to_icd10cm.
html.

28. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122
(2016).

29. Ning Shang, George Hripcsak, Chunhua Weng, Wendy K. Chung, K. C. Breast
Cancer | PheKB. *PheKB* https://phekb.org/phenotype/breast-cancer (2018).

30. Safarova, M. S., Liu, H. & Kullo, I. J. Rapid identification of familial
hypercholesterolemia from electronic health records: The SEARCH study. *J. Clin.*
*Lipidol.* **10**, 1230–1239 (2016).

31. Papani, R. *et al.* Validation of claims-based algorithms for pulmonary arterial
hypertension: *https://doi.org/10.1177/2045894018759246* (2018)
doi:10.1177/2045894018759246.

32. Ye, J. Z., Delmar, M., Lundby, A. & Olesen, M. S. Reevaluation of genetic variants
previously associated with arrhythmogenic right ventricular cardiomyopathy
integrating population-based cohorts and proteomics data. *Clin. Genet.* **96**, 506–
514 (2019).

33. Goldberg, D. S., Lewis, J. D., Halpern, S. D., Weiner, M. G. & Re, V. Lo.
Validation of a coding algorithm to identify patients with hepatocellular carcinoma
in an administrative database. *Pharmacoepidemiol. Drug Saf.* **22**, 103–107
(2013).

34. Wolf, A. M. D. *et al.* American Cancer Society Guideline for the Early Detection of
Prostate Cancer: Update 2010. *CA. Cancer J. Clin.* **60**, 70–98 (2010).

35. Ritchie, M. D. *et al.* Genome- and phenome-wide analyses of cardiac conduction
identifies markers of arrhythmia risk. *Circulation* **127**, 1377–1385 (2013).

36. Simonett, J. M. *et al.* A validated phenotyping algorithm for genetic association
studies in age-related macular degeneration. *Sci. Rep.* **5**, 1–8 (2015).

37. Kho, A. N. *et al.* Use of diverse electronic medical record systems to identify

genetic risk for type 2 diabetes within a genome-wide association study. *J. Am.*
*Med. Informatics Assoc.* **19**, 212–218 (2012).

38.    Yamagata, K. *et al.* Mutations in the hepatocyte nuclear factor-1α gene in
maturity-onset diabetes of the young (MODY3). *Nature* **384**, 455–458 (1996).

39.    Vidal-Puig, A., Solanes, G., Grujic, D., Flier, J. S. & Lowell, B. B. UCP3: An
uncoupling protein homologue expressed preferentially and abundantly in skeletal
muscle and brown adipose tissue. *Biochem. Biophys. Res. Commun.* **235**, 79–82
(1997).

40.    Musa, C. V. *et al.* Four novel UCP3 gene variants associated with childhood
obesity: Effect on fatty acid oxidation and on prevention of triglyceride storage. *Int.*
*J. Obes.* **36**, 207–217 (2012).

41.    Richards, S. *et al.* Standards and guidelines for the interpretation of sequence
variants: A joint consensus recommendation of the American College of Medical
Genetics and Genomics and the Association for Molecular Pathology. *Genet.*
*Med.* **17**, 405–424 (2015).

42.    Nugent, A. *et al.* Reporting of race in genome and exome sequencing studies of
cancer: a scoping review of the literature. *Genet. Med.* **21**, 2676–2680 (2019).

43.    Fouchier, S. W., Defesche, J. C., Umans-Eckenhausen, M. A. & Kastelein, J. J.
The molecular basis of familial hypercholesterolemia in the Netherlands. *Hum.*
*Genet.* **109**, 602–615 (2001).

44.    Rader, D. J. *et al.* Association of the V122I Hereditary Transthyretin Amyloidosis
Genetic Variant with Heart Failure among Individuals of African or Hispanic/Latino
Ancestry. *JAMA - J. Am. Med. Assoc.* **322**, 2191–2202 (2019).

45.    Cohen, J. *et al.* Low LDL cholesterol in individuals of African descent resulting
from frequent nonsense mutations in PCSK9. *Nat. Genet.* **37**, 161–165 (2005).

46.    Manrai, A. K. *et al.* Genetic Misdiagnoses and the Potential for Health Disparities.
*N. Engl. J. Med.* **375**, 655–665 (2016).

47.    Bonnycastle, L. L. *et al.* Autosomal dominant diabetes arising from a wolfram
syndrome 1 mutation. *Diabetes* **62**, 3943–3950 (2013).

48.    Yang, Y. & Chan, L. Monogenic diabetes: What it teaches us on the common
forms of type 1 and type 2 diabetes. *Endocr. Rev.* **37**, 190–222 (2016).

49.    Green, R. C. *et al.* ACMG recommendations for reporting of incidental findings in
clinical exome and genome sequencing. *Genet. Med.* **15**, 565–574 (2013).

50.    Van Hout, C. V. *et al.* Exome sequencing and characterization of 49,960
individuals in the UK Biobank. *Nature* **586**, 749–756 (2020).

51.    Khera, A. V. & Hegele, R. A. What Is Familial Hypercholesterolemia, and Why
Does It Matter? *Circulation* **141**, 1760–1763 (2020).

52.    Sirugo, G., Williams, S. M. & Tishkoff, S. A. The Missing Diversity in Human
Genetic Studies. *Cell* **177**, 26–31 (2019).

53.    Genetics for all. *Nat. Genet.* **51**, 579 (2019).

54.    Antoniou, A. *et al.* Average risks of breast and ovarian cancer associated with
BRCA1 or BRCA2 mutations detected in case series unselected for family history:
A combined analysis of 22 studies. *Am. J. Hum. Genet.* **72**, 1117–1130 (2003).

55.    De Belleroche, J., Orrell, R. & King, A. Familial amyotrophic lateral
sclerosis/motor neurone disease (FALS): A review of current developments. *J.*
*Med. Genet.* **32**, 841–847 (1995).

56. Murphy, N. A. *et al.* Age-related penetrance of the C9orf72 repeat expansion. *Sci.*
*Rep.* **7**, (2017).
57. Stutzmann, F. *et al.* Prevalence of melanocortin-4 receptor deficiency in
europeans and their age-dependent penetrance in multigenerational pedigrees.
*Diabetes* **57**, 2511–2518 (2008).
58. Kobayashi, Y. *et al.* Pathogenic variant burden in the ExAC database: An
empirical approach to evaluating population data for clinical variant interpretation.
*Genome Med.* **9**, 13 (2017).
59. Claussnitzer, M. *et al.* A brief history of human disease genetics. *Nature* **577**,
179–189 (2020).
60. Bomba, L., Walter, K. & Soranzo, N. The impact of rare and low-frequency
genetic variants in common disease. *Genome Biol.* **18**, 1–17 (2017).
61. Xue, A. *et al.* Genome-wide association analyses identify 143 risk variants and
putative regulatory mechanisms for type 2 diabetes. *Nat. Commun.* **9**, 1–14
(2018).
62. Choquet, H. *et al.* A multiethnic genome-wide association study of primary open-
angle glaucoma identifies novel risk loci. *Nat. Commun.* **9**, 1–14 (2018).
63. Bradley, W. G. Inaccuracy of the ICD-9-CM in identifying the diagnosis of
ischemic cerebrovascular disease. *Neurology* **51**, (1998).
64. Öhman, L., Johansson, M., Jansson, J. H., Lind, M. & Johansson, L. Positive
predictive value and misclassification of diagnosis of pulmonary embolism and
deep vein thrombosis in swedish patient registries. *Clin. Epidemiol.* **10**, 1215–
1221 (2018).
65. Doktorchik, C. *et al.* Validation of a case definition for depression in administrative
data against primary chart data as a reference standard 11 Medical and Health
Sciences 1117 Public Health and Health Services. *BMC Psychiatry* **19**, 1–8
(2019).
66. Cozzolino, F. *et al.* A diagnostic accuracy study validating cardiovascular ICD-9-
CM codes in healthcare administrative databases. The Umbria Data-Value
Project. *PLoS One* **14**, e0218919 (2019).
67. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic
data. *Nature* **562**, 203–209 (2018).
68. Hout, C. V. Van *et al.* Whole exome sequencing and characterization of coding
variation in 49,960 individuals in the UK Biobank. *bioRxiv* 572347 (2019)
doi:10.1101/572347.
69. Chang, C. C. *et al.* Second-generation PLINK: Rising to the challenge of larger
and richer datasets. *Gigascience* **4**, 7 (2015).
70. Amberger, J. S., Bocchini, C. A., Schiettecatte, F., Scott, A. F. & Hamosh, A.
OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of
human genes and genetic disorders. *Nucleic Acids Res.* **43**, D789–D798 (2014).
71. Rehm, H. L. *et al.* ClinGen — The Clinical Genome Resource. *N. Engl. J. Med.*
**372**, 2235–2242 (2015).
72. Pruitt, K. D., Tatusova, T. & Maglott, D. R. NCBI reference sequences (RefSeq):
A curated non-redundant sequence database of genomes, transcripts and
proteins. *Nucleic Acids Res.* **35**, D61 (2007).
73. Ferrer, J. *et al.* Mitochondrial glycerol-3-phosphate dehydrogenase: Cloning of an

alternatively spliced human islet-cell cDNA, tissue distribution, physical mapping, and identification of a polymorphic genetic marker. *Diabetes* **45**, 262–266 (1996).

74. Meyre, D. *et al.* Variants of ENPP1 are associated with childhood and adult obesity and increase the risk of glucose intolerance and type 2 diabetes. *Nat. Genet.* **37**, 863–867 (2005).

75. R Core Team. *R: a language and environment for statistical computing.* (R Foundation for Statistical Computing, 2018).

**Author Contributions**

IF and RD designed the study. IF, RL, JC, and RD obtained the data. IF, KC, HV, SB, DJ, and GR analyzed and/or interpreted the data. IF and RD drafted and revised the manuscript. All authors read and provided critical comments on the manuscript.

**Acknowledgements**

The Bio*Me* healthcare delivery cohort at Mount Sinai was founded and maintained with a generous gift from the Andrea and Charles Bronfman Philanthropies. We thank the individuals who were involved in the quality control and/or file handling for the exome sequencing and genome-wide genotyping data, including Aayushee Jain, Lisheng Zhou, Michael Preuss, Quingbin Song, Stephane Wenric, and Steve Ellis. We also thank the thesis advisory committee of IF, including Bruce D. Gelb, Sander Houten, Paz Polak, and Stuart Scott, for their critical feedback and expertise. Research reported in this paper was supported by the Office of Research Infrastructure of the National Institutes of Health under award numbers S10OD018522 and S10OD026880. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

**Funding**

IF is supported by T32GM007280 the Medical Scientist Training Program Training Grant from the National Institute of General Medical Sciences of the National Institutes of Health. RD is supported by R35GM124836 from the National Institute of General Medical Sciences of the National Institutes of Health, and R01HL139865 from the

846    **Competing interests**

847    RD received grants from AstraZeneca, grants and nonfinancial support from Goldfinch

848    Bio, is a scientific co-founder, consultant and equity holder for Pensieve Health, and is a

849    consultant for Variant Bio.

 **FIGURES**

866 **Figure 1**. Biomarker levels in carriers of clinically impactful variants with varying

867 penetrance for three diseases in the Bio*Me* Biobank.

868
869
Metabolite and clinical measurements for carriers of clinically impactful variants with

varying penetrance for three diseases in the Bio*Me* Biobank. Clinically impactful

variants are reported as pathogenic/likely pathogenic in ClinVar or are loss-of-function

in a gene that mediates disease via loss-of-function mechanism, and in a gene with

non-recessive inheritance. Mean of measurements is compared between penetrance

strata with two-tailed t-test: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. **a**, Variants in low-

density lipoprotein receptor (*LDLR*) are stratified by penetrance for familial

hypercholesterolemia (Low [0%], Medium [greater than 0 and less than 100%], and

High [100%]) and shown as box plots with measurements of low-density lipoprotein

cholesterol (LDL) and total cholesterol (cholesterol) levels in carriers. **b**, Variants in

hepatocyte nuclear factor 1-α (*HNF1A*) are stratified by penetrance for maturity-onset

diabetes of the young (Low [0%], Medium [greater than 0 and less than or equal to

50%], and High [greater than 0 and less than 100%]) and depicted as box plots with

levels of glucose and hemoglobin A1c (HbA1c) in carriers. No impactful variants in

*HNF1A* were of Low penetrance. **c**, Variants in uncoupling protein 3 (*UCP3*) are

stratified by penetrance for obesity (Low [0%], Medium [greater than 0 and less than

100%], and High [100%]) and displayed as box plots with body mass index (BMI) and

weight of carriers.

894 **Figure 2**. Distribution of observed disease risk for 5,359 clinically impactful variants in 72,434 individuals.

895
896

Distribution of observed disease risk for 5,359 clinically impactful variants in 72,434 individuals. Risk difference is the

difference in disease prevalence between carriers and non-carriers of a variant of interest. Clinically impactful variants are

reported as pathogenic/likely pathogenic in ClinVar or are loss-of-function in a gene that mediates disease via loss-of-

function mechanism, and in a gene with non-recessive inheritance. Diseases are sorted by descending mean variant risk

difference (diamonds), with all variant risk difference estimates plotted (points) along with the standard deviation (error

bars) per disease. A complete tabulated list of results is provided in **Supplementary Table 6**.

915    **Figure 3**. Penetrance of 34,301 variants stratified by ClinVar pathogenicity, ClinVar review status, and molecular

916    consequence.

917
918

Penetrance of 34,301 variants in non-recessive genes stratified by ClinVar pathogenicity, ClinVar review status, and

molecular consequence. Penetrance distributions are shown as violin plots in red, purple, and green color on a base-10

logarithmic scale with the mean penetrance overlaid as points, alongside disease prevalence in non-carriers shown as

violin plots in grey color with the mean disease prevalence superimposed as points. Pathogenic/likely pathogenic variants

are grouped as pathogenic and benign/likely benign variants are grouped as benign. *, two-tailed t-test *P*<0.05; **,

*P*<0.01; ***, *P*<0.001; ****, *P*<0.0001. **a**, Penetrance is stratified by classification of variant pathogenicity in ClinVar.

Pathogenic variants on average have the highest penetrance (6.9% vs. 0.86% next highest mean penetrance in uncertain

class; $P$=5x10$^{-270}$). Only pathogenic variants confer a significantly increased risk of disease on average to carriers

compared to baseline disease risk in non-carriers (risk difference [RD]=0.046; $P$=1x10$^{-22}$). Other pathogenicity, variants

with a ClinVar pathogenicity classification other than pathogenic, benign, conflicting, or uncertain. **b**, Penetrance is

stratified by variant review status in ClinVar as reviewed by experts (review status=3), multiple submitters (review

status=2), single submitter (review status=1), or no assertion criteria (review status=0). Variants reviewed by experts

(review status=3) have the highest penetrance on average (18% vs. 6.3% next highest mean penetrance in review

status=2 group; $P$=9x10$^{-17}$). Pathogenic variants of review status=3, =2, and =1 confer a significantly increased risk of

disease on average, with RD=0.15 ($P$=1x10$^{-13}$), RD=0.042 ($P$=3x10$^{-10}$), and RD=0.020 ($P$=4x10$^{-3}$), respectively. **c**,

Penetrance is stratified by molecular consequence annotated with Variant Effect Predictor (VEP). Frameshift variants on

average have the highest penetrance (10% vs. 4.1% for missense; $P$=2x10$^{-13}$). There was a significantly elevated risk of

disease on average with frameshift (RD=0.064; $P$=3x10$^{-19}$), splice acceptor (RD=0.020; $P$=5x10$^{-3}$), splice donor

(RD=0.011; $P$=4x10$^{-2}$), stop gained (RD=0.018; $P$=3x10$^{-7}$), and missense variants (RD=0.022; $P$=4x10$^{-4}$).

**Figure 4**. Association between age of disease onset and age-dependent change in

penetrance for 157 diseases.

940
941

Association between age of disease onset and age-dependent change in penetrance for

157 diseases. Diseases correspond to the 5,359 clinically impactful variants and are

grouped according to their age of onset: Earlier, Later, or Any. Change in penetrance is

displayed as a violin plot for each age of onset group when comparing two carrier age

thresholds with the mean change in penetrance superimposed as a point. ΔPenetrance

(%), change in variant penetrance represented as a percent (+ values indicate greater

penetrance estimate with the older age threshold and - values indicate greater

penetrance estimate with the younger age threshold); age comparison, two carrier age

thresholds for which penetrance is compared (e.g., 20-70 compares penetrance with

carriers ≥20 years old and penetrance with carriers ≥70 years old); onset, disease

groups according to age of onset; *, two-tailed t-test *P*<0.05; **, *P*<0.01; ***, *P*<0.001;

****, *P*<0.0001. A complete plot including diseases with ΔPenetrance >5% or

ΔPenetrance <-5% is provided (**Supplementary Figure 5**). **a**, The Bio*Me* Biobank has

carriers of age 20-90 years old and ΔPenetrance is assessed over five carrier age

thresholds: Later diseases have greater ΔPenetrance on average than Earlier diseases

for 20-40 (*P*=0.006), 20-50 (*P*=0.002), 20-60 (*P*=0.02), and 20-70 (*P*=0.02) age

comparisons. **b**, UK Biobank has carriers of age 40-69 years old and ΔPenetrance is

evaluated over two carrier age thresholds: there is no difference in ΔPenetrance

between Later and Earlier diseases for either 40-50 (*P*=0.2) or 40-60 (*P*=0.2) age

comparisons.

976  **Figure 5**. Disease risk associated with clinically impactful variants in disease-

977  predisposition genes for breast cancer and hypercholesterolemia.

978
979

Disease risk associated with clinically impactful variants in known breast cancer- and

hypercholesterolemia-predisposition genes. Clinically impactful (impactful) variants are

reported as pathogenic/likely pathogenic in ClinVar or are loss-of-function in a gene that

mediates disease via loss-of-function mechanism, and in a gene with non-recessive

inheritance. **a**, Penetrance of impactful variants in 10 known or suspected breast

cancer-predisposition genes is displayed as pink violin plots with the mean penetrance

overlaid as a point, alongside disease prevalence in non-carriers shown as grey violin

plots with the mean prevalence superimposed as a point. Violin plots are sorted by

genes with descending mean penetrance, with the highest mean variant penetrance of

38% in *BRCA1* (n=48 variants, mean risk difference [RD]=0.32; $P$=2x10$^{-6}$) and 38% in

*BRCA2* (n=92 variants, mean RD=0.32; $P$=1x10$^{-10}$). *, two-tailed t-test $P$<0.05; **,

$P$<0.01; ***, $P$<0.001; ****, $P$<0.0001. **b**, Heterogeneity in the penetrance of 59 impactful

992 low-density lipoprotein receptor (*LDLR*) variants is shown. The number of variants is

993 plotted as a histogram across four quartiles of penetrance. [0, 25), penetrance less than

994 25%; [25, 50), penetrance greater than or equal to 25% and less than 50%; [50, 75),

995 penetrance greater than or equal to 50% and less than 75%; [75, 100], penetrance

996 greater than or equal to 75%.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

 **TABLES**

1016

1017 **Table 1.** Overview of baseline demographic and clinical traits for 72,434 individuals

1018 from two population-based biobanks.

| Trait | BioMe (n=29,039) | UKB (n=43,395) |
|---|---|---|
| Male, n (%) | 11,684 (40%) | 19,330 (45%) |
| Age, mean (SD) | 59 (16) | 57 (8) |
| European ancestry, n (%) | 9,376 (32%) | 40,447 (93%) |
| African ancestry, n (%) | 7,190 (25%) | 916 (2%) |
| Hispanic ancestry, n (%) | 8,528 (29%) | -- |
| Asian ancestry, n (%) | 1,349 (5%) | 1,088 (3%) |
| Mixed ancestry, n (%) | 2,028 (7%) | 344 (0.8%) |
| BMI in kg/m2, mean (SD) | 28 (7) | 28 (5) |
| SBP in mmHg, mean (SD) | 127 (20) | 140 (19) |
| DBP in mmHg, mean (SD) | 73 (12) | 82 (11) |
| LDL-C in mg/dL, mean (SD) | 98 (36) | 63 (16) |
| Total cholesterol in mg/dL, mean (SD) | 177 (45) | 102 (21) |
| HbA1c in %, mean (SD) | 6 (2) | 6 (3) |
| Glucose in mg/dL, mean (SD) | 106 (47) | 93 (20) |
| HTN, n (%) | 12,962 (45%) | 11,094 (26%) |
| T2D, n (%) | 7,044 (24%) | 2,820 (6%) |
| FBC, n (%) | 1,488 (5%) | 1,581 (4%) |
| AMD, n (%) | 1,041 (4%) | 352 (0.8%) |

1019

Overview of baseline demographic and clinical traits for 72,434 individuals from two

population-based biobanks: BioMe Biobank (BioMe) and UK Biobank (UKB). n, number;

SD, standard deviation; SBP, systolic blood pressure; DBP, diastolic blood pressure;

LDL-C, low-density lipoprotein cholesterol; HbA1c, hemoglobin A1c; HTN, hypertension;

T2D, type 2 diabetes; FBC, familial breast cancer; AMD, age-related macular

degeneration; --, no Hispanic ancestry individuals are in UKB.