Epicardial adipose tissue thickness is associated with increased severity and mortality related to SARS-CoV-2 infection

Roopa Mehta*[1,2], Omar Yaxmehen Bello-Chavolla*[3], Leonardo Mancillas-Adame[4], Marcela Rodriguez-Flores[5], Natalia Ramírez Pedraza[6], Bethsabel Rodríguez Encinas[6] , Carolina Isabel Pérez Carrión[2], María Isabel Jasso Ávila[2], Jorge Carlos Valladares García[2] , Pablo Esteban Vanegas-Cedillo[2], Diana Hernández Juárez[2] , Arsenio Vargas-Vázquez[1,7], Neftali Eduardo Antonio-Villa[1,7], Monica Chapa-Ibarguengoitia[4], Paloma Almeda-Valdés[1,2], Daniel Elias-Lopez[1,2], Arturo Galindo-Fraga[8], Alfonso Gulias-Herrero[9], Alfredo Ponce de Leon[8], José Sifuentes-Osornio[8,9] Carlos A. Aguilar-Salinas[1,10,11]

*Both authors contributed equally

[1]Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán (INCMNSZ). [2]Department of Endocrinology and Metabolism, INCMNSZ, [3]Research Division, Instituto Nacional de Geriatría, [4]Facultad de Medicina, Universidad Autonoma de Nuevo Leon, [5]Department of Obesity, INCMNSZ, [6]Department of Radiology, INCMNSZ, [7]MD/PhD (PECEM) program, Faculty of Medicine, National Autonomous University of Mexico, [8]Department of Infectology, INCMNSZ, [9]Internal Medicine Division, [10]Instituto Tecnologico y de Estudios Superiores de Monterrey Tec Salud, [11]Division of Nutrition, INCMNSZ,

Short title: Epicardial fat thickness and COVID-19 mortality

Corresponding authors

Roopa Mehta, FRCP. Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán. E-mail: roopamehta@yahoo.com

Carlos Aguilar-Salinas, MD. Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán. E-mail: caguilarsalinas@yahoo.com

CONFLICT OF INTERESTS: Nothing to disclose.

FUNDING: This research received no specific funding

Word count: 2859 words; 4 figures, 4 tables, 35 references

ABSTRACT (250 WORDS)

BACKGROUND: Obesity, in particular, visceral adipose tissue has been linked to adverse COVID-19 outcomes. The amount of epicardial adipose tissue (EAT) may have relevant implications given its proximity to the heart and lungs. Here, we explored the role of EAT in increasing the risk for COVID-19 adverse outcomes.

METHODS: We included 748 patients with COVID-19 attending a reference centre in Mexico City. EAT thickness, sub-thoracic and pericardial fat were measured using thoracic CT scans. We explored the association of each thoracic adipose tissue compartment with critical COVID-19 and mortality according to the presence or absence of obesity. Mediation analyses evaluated the role of EAT in facilitating the effect of age, body mass index and cardiac troponin levels with adverse COVID-19 outcomes.

RESULTS: EAT thickness was associated with increased risk of COVID-19 mortality (HR 1.18, 95%CI 1.01-1.39) independent of age, gender, comorbid conditions and BMI. Increased EAT was associated with lower SpO2 and PaFi index and higher levels of cardiac troponins, D-dimer, C-reactive protein, and 4C severity score, independent of obesity. EAT mediated 13.1% (95%CI 3.67-28.0%) and 5.1% (95%CI 0.19-14.0%) of the effect of age and 19.4% (95%CI 4.67-63.0%) and 12.8% (95%CI 0.03-46.0%) of the effect of BMI on requirement for intubation and mortality, respectively. EAT also mediated the effect of increased cardiac troponins on myocardial infarction during COVID-19.

CONCLUSION: EAT is an independent risk factor for severe COVID-19 and mortality independent of obesity. EAT partly mediates the effect of age and BMI and increased cardiac troponins on adverse COVID-19 outcomes.

Keywords: COVID-19, Obesity, Mortality, Epicardial fat, Visceral fat

INTRODUCTION

Obesity is a major risk factor for disease severity related to severe acute coronavirus 2 (SARS-CoV-2) infection[1]. Patients with obesity have a higher risk for, hospitalization, intensive care unit (ICU) admission, invasive mechanical ventilation (IMV) and death [1,2]. Obesity is associated with a state of low-grade inflammation, which in the setting of coronavirus disease (COVID-19) can result in further inflammation, endothelial dysfunction and subsequent cytokine storm, increasing the risk for severe disease [2–4]. In Mexico, obesity is highly prevalent; in particular, the accumulation of abdominal visceral adipose tissue (VAT) has been shown to influence COVID-19 outcomes, independent of additional comorbid conditions and risk factors.[5-8]

Visceral obesity is associated with increased cardiometabolic burden, including risk of arterial hypertension, type 2 diabetes mellitus and cardiovascular disease[5]. Accumulation of VAT results in adipose tissue dysfunction, morphological changes in adipocytes, fibrosis, altered secretion of adipokines and increased inflammation[6]. VAT has proven to be related to increased risk for severe COVID-19 and need for intensive care.[11, 12] The engagement of SARS-CoV-2 with angiotensin converting enzyme 2 (ACE 2) in visceral fat is thought to impair the function of this enzyme, resulting in increased production of angiotensin II, thus enhancing the production of inflammatory cytokines.[11] Levels of ACE 2 mRNA in visceral fat correlate with body mass index; this is not the case for subcutaneous fat. Epicardial adipose tissue (EAT) is an ectopic fat deposit, located between the heart and the pericardium and is thought to contribute to adverse cardiovascular outcomes[7]. EAT may play a significant role in mediating the effects of obesity on COVID-19 outcomes due to its thoracic localization and may influence the cardiovascular complications of COVID-19[8–10]. In this study, we examined the association of EAT with adverse COVID-19 outcomes and explored potential pathways which may link the accumulation of EAT was mediators of well-established risk factors related to severe COVID-19.

METHODS

Study population

This study included consecutive patients evaluated at the Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán (INCMNSZ), a COVID-19 reference centre in Mexico City between 17th March and 31st May 2020. Subjects were initially assessed at triage and required either ambulatory or in-hospital care for COVID-19, confirmed with computerized tomography (CT) and/or via RT-qPCR test from nasopharyngeal swabs. All patients had moderate to severe disease as defined by National Institute of Health criteria. *Moderate* illness was defined as evidence of lower respiratory disease during clinical assessment or imaging and who have saturation of oxygen (SpO$_2$) ≥94% on air, and s*evere illness as* SpO$_2$ <94% on air, a ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO$_2$/FiO$_2$) <300 mm Hg, respiratory frequency >30 breaths/min, or lung infiltrates >50%). All patients underwent a chest CT, and a radiologist determined the degree of pulmonary parenchymal disease and assessed epicardial fat thickness. In addition, a medical history, anthropometric measurements and laboratory tests were obtained. The electronic files of each patient were retrospectively reviewed to document the outcomes during hospitalization. All proceedings were approved by the research and ethics committee of the INCMNZ (Ref. 3383) and informed consent was waived due to the nature of the study.

Laboratory and clinical measurements

Clinical variables and laboratory measures were obtained at the time of initial evaluation. Physical examination included: weight, height, body mass index (BMI, calculated as weight in kilograms divided by squared height in meters), pulse oximeter saturation (SpO2), respiratory rate (RR), temperature and arterial blood pressure (BP). Laboratory measurements included: full blood count, chemistry panel including liver function tests, C-reactive protein (CRP), fibrinogen, D-dimer, ferritin, troponin I (TPNI), erythrocyte

sedimentation rate (ESR) and procalcitonin levels. The blood samples were processed at the central laboratory of the INCMNSZ.

Epicardial adipose tissue measurements

All patients underwent unenhanced CT scans including low-dose CT and two ultra-low-dose CT protocols and commonly reported imaging features of COVID-19 pneumonia were captured. Thorax CT was performed using a 64-slice scanner (GE MEDICAL SYSTEMS Revolution EVO). Epicardial adipose tissue (EAT), the adipose tissue located between the visceral layer of the pericardium and the surface of the heart was measured. EAT thickness was measured at 3 points (right atrioventricular fossa, left atrioventricular fossa, and anterior interventricular fossa) in the reformatted 4-chamber view using the multiplanar reconstruction (MPR) tool on the workstation (19, 20, 21). The maximum thickness of the EAT was recorded, from the surface of the myocardium to the pericardium (measured perpendicular to the pericardium). The measurements were made on 2 different occasions, obtaining a total of 6 measurements; the average of them was used in the statistical analysis.

Pericardial adipose tissue (PAT) was quantified with the volume measurement tool with the Carestream system of the workstation. The thickness of the thoracic subcutaneous adipose tissue (TscAT) was measured from the anterior border of the sternum to the skin, at the level of the mitral valve in the axial plane of the tomography. The 80[th] gender-specific percentile of EAT thickness was obtained and used as the threshold to define increased EAT thickness.

In addition, chest CT findings were recorded and used to evaluate severity of COVID infection.

COVID-19 outcomes

Outcomes included mortality and requirement for mechanical intubation. Furthermore, a composite of death and requirement for invasive mechanical ventilation was used as a

definition for critical COVID-19. For time-to-event analyses, time from self-reported symptom onset prior to evaluation until last follow-up (censoring) or death, whichever occurred first was estimated. The COVID-19 specific 4C mortality score was estimated; this reflects increased risk of mortality in this population[11].

Statistical analyses

The clinical characteristics of subjects with visceral obesity (EAT > 80th percentile) were compared according to presence or absence of obesity, as assessed by BMI, using Student's t-test or Mann-Whitney U according to variable distribution for continuous variables and using chi-squared tests for categorical variables. To ensure variable symmetry prior to modeling, the *bestNormalize* R package was used and identified the better transformation as out-of-sample normalization via 10-fold cross-validation with 5 repeats for all CT-derived measurements. All statistical analyses were conducted using R 4.0.2.

Assessment of risk based on thoracic fat distribution

The CT-derived adipose tissue measures and body-mass index (BMI) were compared between cases who eventually required mechanical ventilation or died during hospitalization and those who did not, using Student's t-test. Linear regression analysis was used to assess which clinical features were independently associated with EAT thickness in COVID-19 patients. To assess the predictive ability of each thoracic adipose tissue segment, Cox proportional risk regression models were generated including an unadjusted model (Model 1) and models adjusted for age, gender and number of comorbid conditions (Model 2) and further adjustments by BMI to assess independent risk after adjustment (Model 3). In addition, the construct of increased EAT using the definition of >80th gender-specific percentile of EAT thickness was assessed using the same models. To assess the impact of thoracic adipose tissue compartments on the risk of mechanical ventilation, similar models were adjusted using fixed effects logistic regression models. To

assess the independent effect of BMI, Models 1 and 2 were fitted for normalized BMI values for both outcomes.

Post-estimation simulation of mortality risk

The 1-unit incremental risk associated with normalized EAT measures was evaluated using Cox regression models adjusted for age, gender, number of comorbid conditions and BMI- This was further stratified by BMI-determined obesity status to evaluate the independent effect of EAT thickness in predicting mortality risk related to COVID-19 using the *simPH* and *survival* R packages.

Mediation analyses

The first hypothesis was that EAT was partially mediating the mortality risk related to age, given the association of increases in visceral and epicardial adipose tissue with aging, and BMI[12]. The secondary hypothesis was that EAT may induce cardiac damage and that this could increase COVID-19 severity and   mortality risk[8]. To test these hypotheses, casual mediation analyses were conducted using the *mediation* R package, adjusting logistic regression models by age, gender, number of comorbid conditions or BMI as required. Confidence intervals were estimated using White's heteroskedasticity-consistent estimator for the covariance matrix, derived from quasi-Bayesian Monte Carlo simulation based on normal approximation.

RESULTS

<u>Study population</u>

We included 748 patients with confirmed COVID-19, with a mean age of 51.22±13.62 years, male predominance (n=470, 63.1%), and a mean BMI of 30.25±5.71. Median follow-up was 14.0 days (IQR 10.0-20.0) and 629 patients required hospitalization (85.5%). Overall, 138 patients received invasive mechanical ventilation (18.5%) and 164 in-hospital deaths (21.9%) were recorded. Type 2 diabetes (T2D) was present in 191 patients (26.3%), 300 patients had obesity (44.4%) and 282 were overweight (41.8%).

Median EAT thickness was 9.67mm (IQR 7.67-12.0), median sub thoracic fat thickness was 16mm (IQR 11.0-23.0) and pericardial fat was present in 451 patients (60.5%).

Thoracic fat distribution and COVID-19 severity

Cases that required mechanical ventilation had greater EAT thickness, pericardial fat volume and BMI with no significant difference in subthoracic adipose tissue thickness. In fatal cases, EAT thickness and pericardial fat volume were significantly higher; BMI was greater but failed to reach statistical significance and sub thoracic fat was no different between fatal and non-fatal cases (**Figure 1**). When stratifying cases according to BMI-defined obesity, those who had increased EAT were older, had worsened respiratory parameters, altered coagulation and overt inflammation, and had higher rates of severe outcomes and intubation as well as higher mortality scores independent of BMI status (**Table 1**). EAT thickness was independently associated with male gender (β=-0.236, 95%CI -0.382,-0.090), oxygen saturation (β=-0.014, 95%CI -0.019,-0.008) and higher 4C mortality scores (β=0.063, 95%CI 0.039, 0.087, adjusted R^2=0.103). In a second model, EAT thickness was associated with increased levels of ultrasensitive cardiac troponins (β=0.0002, 95%CI 0.00005, 0.0004), BMI (β=0.034, 95%CI 0.021, 0.048) and age (β=0.017, 95%CI 0.011, 0.023, adjusted R^2=0.081), even after adjustments for C-reactive protein levels.

EAT thickness is an independent predictor of COVID-19 mortality

Using Cox regression analysis, EAT thickness was a predictor of COVID-19 mortality, even after adjustment for age, gender, number of comorbid conditions and BMI (**Table 2**). Using post-estimation simulation, we observed monotonic increases in mortality risk with increasing normalized EAT thickness both in obese and non-obese subjects (**Figure 2**). Subjects who had increased EAT had higher risk of both mortality and requirement for invasive mechanical ventilation. Similarly, after assessing the independent effect of other adipose tissue segments, pericardial fat volume was associated with increased mortality in

univariate analyses but failed to retain significance after adjustments for age, gender, number of comorbid conditions and BMI. Similarly, sub thoracic fat thickness was associated with increased risk of mortality and mechanical ventilation after adjustment for age, gender, and number of comorbid conditions but after adjustment for BMI this association failed to maintain statistical significance. In contrast, BMI was associated with both mortality and invasive mechanical ventilation even after adjustment for covariates (**Table 2**)

EAT partially mediates the effect of age and BMI on COVID-19 mortality

Given the association of BMI, increased cardiac troponins and chronological age on COVID-19 mortality levels and EAT thickness, two causal mediation models were explored. We explored EAT thickness as a mediator of both age and BMI and in increasing risk of COVID-19 mortality and critical COVID-19 (**Figure 2**). EAT thickness mediates 13.1% (95%CI 3.67-28.0%) of the effect of age and 19.4% (95%CI 4.67-63.0%) of the effect of BMI on the risk of critical COVID-19. Similarly, EAT thickness mediates 5.91% (95%CI 0.19-14.0%) of the effect of age and 12.8% (95%CI 0.03-46.0%) of the effect of BMI on risk of mortality, after adjustment of age, gender and number of comorbid conditions (**Table 4**).

EAT-induced heart damage mediates COVID-19 adverse outcomes

The second causal mediation model explored whether increased risk for COVID-19 outcomes was partially mediated by EAT-induced heart damage (**Figure 3**). Cardiac troponins mediate 24.2% (95%CI 4.8-62.0%) of the effect of EAT on the risk of critical COVID-19 and 29.6% (95%CI 3.9-100.0%) of the effect of EAT on the risk of COVID-19 mortality adjusted for age and gender. Regarding cardiovascular events, there were 19 registered myocardial infarctions, amongst whom 13 cases died. EAT thickness is a significant predictor of myocardial infarction in COVID-19 (OR 1.78, 95%CI 1.09-2.97);

increased EAT thickness increased risk for myocardial infarction in COVID-19 (OR 2.83, 95%CI 1.07-7.31), after adjustment for age, gender and comorbid conditions.

DISCUSSION

Here, we showed that EAT thickness is a significant predictor of COVID-19 severity and mortality independent of BMI, age, and comorbid conditions. Similarly, subjects with increased EAT had higher levels of COVID-19 severity markers independent of BMI. Overall, EAT thickness was associated with decreased oxygen saturation, male gender, and increased COVID-19 severity; similarly, EAT thickness was associated with increased ultrasensitive cardiac troponin levels. Sub thoracic and pericardial adipose tissue did not show an increased risk for adverse outcomes or mortality independent of BMI. Notably, EAT thickness partly mediates the effect of age and BMI related risk on COVID-19 adverse outcomes. These findings confirm that adipose tissue distribution, in particular the presence of visceral adiposity more so than total adipose tissue, may play a significant role in modulating the adaptive responses to SARS-CoV-2 infection, and thus increase disease severity in COVID-19[13].

The metabolic and inflammatory role of VAT and its role in cardiovascular and metabolic health is well documented[14]. Beyond the risk associated with whole-body adipose tissue as assessed by BMI, VAT depots significantly contribute to the increased cardiovascular risk profile and low-grade pro-inflammatory state observed in patients with obesity[15]. Increased accumulation of VAT has been associated with increased deposition of ectopic fat, particularly in the liver and in the epicardium as EAT[16]. Notably, EAT shares microcirculation with the myocardium, favoring the rapid exchange of metabolic and inflammatory products[17]; the implication of this anatomical relationship suggests that, under acute or chronic inflammatory conditions, these interactions might prove deleterious to cardiovascular function. Increased epicardial adipogenesis induces the secretion of pro-inflammatory adipokines which contribute to atrial and ventricular damage[7,12,17]. In the

setting of COVID-19, EAT is postulated to play a role in generating the imbalance between the production of anti and pro-inflammatory adipokines, thus contributing to the enhanced cytokine storm and multisystem involvement documented in severe COVID-19[8]. In this study, EAT thickness was a significant predictor of adverse biochemical profiles, critical COVID-19 and increased mortality independent of BMI.

Using mediation analyses, EAT thickness was shown to play a significant role in mediating the risk of critical COVID-19 and mortality attributable to increasing age and BMI. Other authors have shown that increased VAT deposition has a greater predictive value for adverse COVID-19 outcomes compared to BMI. This depot-specific adipose tissue dysfunction may be particularly relevant in younger cases[18–21]. These results are consistent with previous findings in this population and further contribute to the characterization of the mechanisms underlying the risk attributable to aging and obesity in severe COVID-19 [25–28].

Finally, increased EAT accumulation was shown to be associated with increased cardiac damage and this may contribute to COVID-19 severity and mortality. Increased expression of ACE II in whole-body and visceral adipose tissue in persons with obesity has been proposed as a potential mechanism for increasing the risk of COVID-19 by promoting viral spread and shedding and by amplifying the inflammatory responses [22–24]. Increased risk of cardiovascular complications in COVID-19 has been associated with the pro-thrombotic and inflammatory adaptive responses to SARS-CoV-2 infection as well as direct damage to the myocardium[9,25,26]. Increased EAT thickness was associated with increased risk of in-hospital myocardial infarction and was associated with increased cardiac troponin levels, possibly indicating direct cardiovascular injury. Since EAT is only a measure of ectopic fat deposition around the myocardium, its measurement can be considered complementary to BMI. Hence, it may be most useful to assess risk for potential cardiac involvement in COVID-19.

This work has certain strengths and limitations. The study analyses a large number of patients in whom EAT thickness was measured at admission and who were followed throughout hospitalization to record COVID-19 relevant outcomes. Furthermore, this work is novel as it assesses the different types of intrathoracic adipose tissue, including EAT, subthoracic and pericardialm in addition to whole-body adipose tissue measures such as BMI. Using mediation analyses, the degree of contribution of EAT in risk of critical disease and mortality was also quantified. Limitations include the use of EAT thickness instead of volume; this was because the CT scans on admission were simple scans in which volume could not be assessed. There were only a small number of recorded cardiovascular events, this means the full impact of EAT on COVID-19 cardiovascular complications could not be evaluated. The results only apply to moderate or severe infection as mild cases were not included dut to the nature of case recruitment. The study was carried out at a single COVID-19 centre, and the findings may not be representative at a population level. Finally, since abdominal CT scans were not available for most patients, the contribution of thoracic vs. abdominal fat could not be assessed, an area which should be evaluated in future studies.

In conclusion, EAT thickness was associated with increased risk for critical COVID-19 and mortality compared to other types of intrathoracic adipose tissue. EAT was also associated with increased cardiac troponin levels, suggesting it may play a role in the risk for cardiovascular complications in COVID-19. Furthermore, EAT appears to be a mediator of the effect of age and BMI on adverse disease outcomes. Hence, this work adds to the body of evidence suggesting that EAT may be an important protagonist in adverse metabolic and inflammatory responses to SARS-CoV-2.

CONFLICT OF INTEREST/FINANCIAL DISCLOSURE: Nothing to disclose.

FUNDING: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

ACKNOWLEDGMENTS

All authors approved the submitted version. AVV, and NEAV are enrolled in the PECEM program at the Faculty of Medicine of UNAM; AVV and NEAV are supported by CONACyT.

CONTRIBUTIONS: research idea and study design: RM, LMA, MRF; data acquisition: RM, NRM, BRE, IJA, JCVG, CIPC, PEVC, DHJ, CAAS; data analysis/interpretation: OYBC, RM, CAAS, NEAV, AVV; statistical analysis: OYBC; manuscript drafting: RM, OYBC, NEAV, AVV, PAV, DEL, CAAS; supervision or mentorship: RM, AGF, AGH, APLG, JSO, CAAS. Each author contributed important intellectual content during manuscript drafting or revision and accepts accountability for the overall work by ensuring that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

DATA AVAILABILITY: Data is available from the corresponding author upon reasonable request. Code for reproducibility of results available at:

https://github.com/oyaxbell/covid_metabolism

REFERENCES

1.   Huang Y, Lu Y, Huang Y-M, et al. Obesity in patients with COVID-19: a systematic review and meta-analysis. *Metabolism*. 2020;113:154378. doi:10.1016/j.metabol.2020.154378
2.   Vepa A, Bae JP, Ahmed F, Pareek M, Khunti K. COVID-19 and ethnicity: A novel pathophysiological role for inflammation. *Diabetes Metab Syndr*. 2020;14(5):1043-1051. doi:10.1016/j.dsx.2020.06.056
3.   Chiappetta S, Sharma AM, Bottino V, Stier C. COVID-19 and the role of chronic inflammation in patients with obesity. *Int J Obes 2005*. 2020;44(8):1790-1792. doi:10.1038/s41366-020-0597-4
4.   Korakas E, Ikonomidis I, Kousathana F, et al. Obesity and COVID-19: immune and metabolic derangement as a possible link to adverse clinical outcomes. *Am J Physiol Endocrinol Metab*. 2020;319(1):E105-E109. doi:10.1152/ajpendo.00198.2020
5.   Bello-Chavolla OY, Antonio-Villa NE, Vargas-Vázquez A, et al. Metabolic Score for

Visceral Fat (METS-VF), a novel estimator of intra-abdominal fat content and cardio-metabolic health. *Clin Nutr Edinb Scotl*. 2020;39(5):1613-1621. doi:10.1016/j.clnu.2019.07.012

6.   Unamuno X, Gómez-Ambrosi J, Rodríguez A, Becerril S, Frühbeck G, Catalán V. Adipokine dysregulation and adipose tissue inflammation in human obesity. *Eur J Clin Invest*. 2018;48(9):e12997. doi:10.1111/eci.12997

7.   Sato F, Maeda N, Yamada T, et al. Association of Epicardial, Visceral, and Subcutaneous Fat With Cardiometabolic Diseases. *Circ J Off J Jpn Circ Soc*. 2018;82(2):502-508. doi:10.1253/circj.CJ-17-0820

8.   Malavazos AE, Goldberger JJ, Iacobellis G. Does epicardial fat contribute to COVID-19 myocardial inflammation? *Eur Heart J*. 2020;41(24):2333. doi:10.1093/eurheartj/ehaa471

9.   Dhakal BP, Sweitzer NK, Indik JH, Acharya D, William P. SARS-CoV-2 Infection and Cardiovascular Disease: COVID-19 Heart. *Heart Lung Circ*. 2020;29(7):973-987. doi:10.1016/j.hlc.2020.05.101

10.   Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the COVID-19 Pandemic. *J Am Coll Cardiol*. 2020;75(18):2352-2371. doi:10.1016/j.jacc.2020.03.031

11.   Knight SR, Ho A, Pius R, et al. Risk stratification of patients admitted to hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ*. 2020;370:m3339. doi:10.1136/bmj.m3339

12.   Silaghi A, Piercecchi-Marti M-D, Grino M, et al. Epicardial adipose tissue extent: relationship with age, body fat distribution, and coronaropathy. *Obes Silver Spring Md*. 2008;16(11):2424-2430. doi:10.1038/oby.2008.379

13.   Márquez-Salinas A, Fermín-Martínez CA, Antonio-Villa NE, et al. Adaptive metabolic and inflammatory responses identified using accelerated aging metrics are linked to adverse outcomes in severe SARS-CoV-2 infection. *medRxiv*. Published online November 24, 2020:2020.11.03.20225375. doi:10.1101/2020.11.03.20225375

14.   Neeland IJ, Ross R, Després J-P, et al. Visceral and ectopic fat, atherosclerosis, and cardiometabolic disease: a position statement. *Lancet Diabetes Endocrinol*. 2019;7(9):715-725. doi:10.1016/S2213-8587(19)30084-1

15.   Stefan N. Causes, consequences, and treatment of metabolically unhealthy fat distribution. *Lancet Diabetes Endocrinol*. 2020;8(7):616-627. doi:10.1016/S2213-8587(20)30110-8

16.   Smith U. Abdominal obesity: a marker of ectopic fat accumulation. *J Clin Invest*. 2015;125(5):1790-1792. doi:10.1172/JCI81507

17.   Packer M. Epicardial Adipose Tissue May Mediate Deleterious Effects of Obesity and Inflammation on the Myocardium. *J Am Coll Cardiol*. 2018;71(20):2360-2372. doi:10.1016/j.jacc.2018.03.509

18.   Favre G, Legueult K, Pradier C, et al. Visceral fat is associated to the severity of COVID-19. *Metabolism*. 2020;115:154440. doi:10.1016/j.metabol.2020.154440

19.   Watanabe M, Caruso D, Tuccinardi D, et al. Visceral fat shows the strongest association with the need of intensive care in patients with COVID-19. *Metabolism*. 2020;111:154319. doi:10.1016/j.metabol.2020.154319

20.   Deng M, Qi Y, Deng L, et al. Obesity as a Potential Predictor of Disease Severity in Young COVID-19 Patients: A Retrospective Study. *Obes Silver Spring Md*. 2020;28(10):1815-1825. doi:10.1002/oby.22943

21.   Iacobellis G, Malavazos AE, Ferreira T. COVID-19 Rise in Younger Adults with Obesity: Visceral Adiposity Can Predict the Risk. *Obes Silver Spring Md*. 2020;28(10):1795. doi:10.1002/oby.22951

22.   Iacobellis G, Secchi F, Capitanio G, et al. Epicardial Fat Inflammation in Severe

COVID-19. *Obesity*. 2020;28(12):2260-2262. doi:https://doi.org/10.1002/oby.23019

23.   Malavazos AE, Corsi Romanelli MM, Bandera F, Iacobellis G. Targeting the Adipose Tissue in COVID-19. *Obes Silver Spring Md*. 2020;28(7):1178-1179. doi:10.1002/oby.22844

24.   Ryan PM, Caplice NM. Is Adipose Tissue a Reservoir for Viral Spread, Immune Activation, and Cytokine Amplification in Coronavirus Disease 2019? *Obes Silver Spring Md*. 2020;28(7):1191-1194. doi:10.1002/oby.22843

25.   Long B, Brady WJ, Koyfman A, Gottlieb M. Cardiovascular complications in COVID-19. *Am J Emerg Med*. 2020;38(7):1504-1507. doi:10.1016/j.ajem.2020.04.048

26.   Levi M, Thachil J, Iba T, Levy JH. Coagulation abnormalities and thrombosis in patients with COVID-19. *Lancet Haematol*. 2020;7(6):e438-e440. doi:10.1016/S2352-3026(20)30145-9

TABLES

Table 1. Comparison of demographic and clinical markers of COVID-19 patients comparing epicardial adipose tissue thickness (EAT) (increased >80th sex-specific percentile or not) stratified by obesity status as defined by body-mass index (BMI).

Parameter	Overall	Non-Obese			Obese		
		EAT not-increased (n=368)	EAT increased (n=74)	P-value	EAT not-increased (n=230)	EAT increased (n=76)	P-value
Age (years)	51 (41-61)	51.5 (13.7)	56.9 (15.2)	0.006	48.4 (12.9)	52.5 (13.8)	0.002
Male Sex (%)	470 (63.1)	227 (62.2)	49 (66.2)	0.602	149 (64.7)	45 (59.2)	0.944
Low-Socioeconomic Status (%)	530 (71.2)	262 (71.9)	53 (71.6)	0.682	161 (71.8)	54 (71.1)	0.682
Critical outcome (%)	236 (31.5)	91 (24.7)	34 (45.9)	0.003	78 (33.9)	33 (43.5)	0.025
Intubation (%)	138 (18.4)	46 (12.5)	19 (12.5)	0.006	54 (23.4)	19 (25)	0.002
Mortality (%)	164 (21.9)	67 (18.2)	22 (29.8)	0.036	48 (20.8)	27 (35.5)	0.182

Arterial Hypertension (%)	212 (29.2)	88 (24.9)	23 (31.9)	0.277	73 (32.2)	28 (37.3)	0.039
Type 2 diabetes (%)	191 (26.3)	87 (24.6)	23 (32.3)	0.226	55 (24.2)	26 (34.7)	0.857
Time since DM2 diagnosis (years)	7 (3-14.8)	8 (5-14.5)	6 (2-12.5)	0.370	6 (1-15)	6 (5-10)	0.773
Smoking status (%)	47 (7.7)	21 (7.26)	6 (7.25)	0.792	16 (8.2)	4 (6.2)	0.826
CKD (%)	26 (3.6)	14 (3.2)	4 (5.5)	0.780	3 (1.3)	5 (6.7)	0.345
CVD (%)	19 (2.6)	9 (2.6)	1 (1.4)	0.868	6 (2.6)	3 (4.0)	0.774
Cirrhosis (%)	6 (0.82)	2 (0.05)	1 (1.4)	0.999	2 (0.8)	1 (1.3)	0.998
BMI (kg/m2)	30.4 (11.2)	27 (25.1-28.7)	27.5 (25-29)	0.061	33.2 (31-37)	34.1 (31-36)	0.013
Respiratory Rate (rpm)	28 (11.2)	25 (20-30)	30 (26-36)	<0.001	26 (22-33)	28 (24-39)	<0.001
Heart Rate (bpm)	101 (18.6)	102 (90-113)	103 (90-115)	0.529	102 (90-115)	107 (90-115)	0.385

Systolic Arterial Pressure (mmHg)	121 (110-132)	120 (110-130)	120 (110-130)	0.587	127 (113-135)	128 (111-134)	0.771
Diastolic Arterial Pressure (mmHg)	75 (69-80)	73 (69-80)	70 (66-80)	0.397	78 (70-82)	76 (68-80)	0.343
Oxygen saturation (%)	86 (77-89)	88 (80-90)	79 (63-87)	<0.001	85 (79-89)	82 (60-87)	<0.001
C-reactive protein	14.3 (7.1-22.4)	14.93 (9.02-23.20)	19.3 (10.8-28.5)	<0.001	14.4 (7.9-21.2)	19 (9-28.6)	<0.0001

TABLE 2. Cox proportional risk regression models to predict mortality related to COVID-19 using CT-scan derived fat measures and BMI transformed using repeated out-of sample 10-fold cross-validation and the definition of visceral obesity as epicardial fat thickness values >80th sex-adjusted percentiles for the population. Model 1: Univariate, Model 2: Ajusted for age, gender and comorbid conditions, Model 3: Adjusted for age, gender, comorbid conditions and BMI

Model	Model 1		Model 2		Model 3	
	HR (95%CI)	P-value	HR (95%CI)	P-value	HR (95%CI)	P-value
Epicardial fat thickness	1.290 (1.108-1.501)	0.001	1.212 (1.033-1.423)	0.019	1.180 (1.003-1.389)	0.047
Extrapericardial fat volume	1.223 (1.036-1.445)	0.018	1.108 (0.924-1.328)	0.269	1.049 (0.870-1.265)	0.616
Subthoracic fat volume	1.067 (0.907-1.256)	0.434	1.230 (1.019-1.483)	0.031	1.117 (0.889-1.404)	0.343
Body-mass index	1.091 (0.931-1.280)	0.283	1.262 (1.042-1.529)	0.018	-	-
Increased EAT (>p80, sex-based)	1.570 (1.123-2.196)	0.008	1.433 (1.023-2.009)	0.037	1.409 (1.006-1.975)	0.046

TABLE 3. Logistic regression models to predict requirement for invasive ventilation in COVID-19 cases, using CT-scan derived fat measures and BMI transformed using repeated out-of sample 10-fold cross-validation and the definition of increased epicardial fat thickness (EAT) values >80th sex-adjusted percentiles for the population. Model 1: Univariate, Model 2: Ajusted for age, gender and comorbid conditions, Model 3: Adjusted for age, gender, comorbid conditions and BMI.

Model	Model 1		Model 2		Model 3	
	OR (95%CI)	P-value	OR (95%CI)	P-value	OR (95%CI)	P-value
Epicardial fat thickness	1.205 (1.001-1.454)	0.049	1.272 (1.048-1.548)	0.015	1.210 (0.993-1.478)	0.060
Extrapericardial fat volume	1.054 (0.855-1.294)	0.617	0.989 (0.791-1.234)	0.926	0.874 (0.691-1.102)	0.258
Subthoracic fat volume	1.093 (0.908-1.316)	0.346	1.242 (1.007-1.535)	0.044	0.999 (0.763-1.306	0.995
Body-mass index	1.408 (1.167-1.705)	<0.001	1.418 (1.149-1.766)	<0.001	-	-
Increased EAT	1.690 (1.094-2.572)	0.016	1.791 (1.148-	0.009	1.689 (1.078-	0.020

| obesity (>p80, sex-based) | 2.759) | 2.614) |

TABLE 4. Causal mediation models. Firstly, evaluating the effect of age and BMI, on COVID-19 outcomes via epicardial fat thickness. Secondly, the role of epicardial adipose tissue thickness on COVID-19 outcomes via cardiac troponins. Abbreviation: ACME: average causal mediation effects; ADE: average direct effects

Causal Mediation Model	Outcome (Y)	Effector (E)	Mediator (M)	ACME (95%CI)	ADE (95%CI)	Total Effect (95%CI)	Proportion Mediated (95%CI)
1	Critical COVID-19	Age (years)	Epicardial fat thickness	0.000336 (0.000077-0.001)	0.00207 (0.00160-0.001)	0.00240 (0.00188-0.001)	13.1% (3.67-28.0%)
		BMI (kg/m^2)		0.0109 (0.0018-0.020)	0.0457 (0.0080-0.090)	0.0566 (0.0186-0.090)	19.4% (4.67-63.0%)
		Epicardial fat thickness	Cardiac troponins	0.0148 (0.0030-0.030)	0.0465 (0.010-0.800)	0.0613 (0.0232-0.100)	24.2% (4.8-62.0%)

2	COVID-19 mortality	Age (years)	Epicardial fat thickness	0.0000494 (0.0000013-0.001)	0.000731 (0.000377-0.001)	0.000780 (0.000394-0.001)	5.91% (0.19-14.0%)
		BMI (kg/m^2)		0.00710 (0.00014-0.020)	0.048 (0.010-0.090)	0.0554 (0.0173-0.100)	12.8% (0.03-46.0%)
		Epicardial fat thickness	Cardiac troponins	0.0115 (0.0022-0.0200)	0.0274 (-0.0033, 0.060)	0.0389 (0.0061-0.0700)	29.6% (3.9-100.0%)

Figure 1. Boxplots comparing transformed levels of epicardial fat thickness, pericardial fat, subthoracic fat and body-mass index

(BMI), according to the need for invasive mechanical ventilation (IVM, A-D) or COVID-19 mortality(E-H).

Figure 2. Post-estimation simulation of the risk of epicardial fat thickness transformed by repeated out-of sample 10-fold cross-validation adjusted for age, gender, comorbid conditions and BMI (A) stratified in non-obese (B) and obese (C) cases with confirmed COVID-19.

Figure 3. Diagrams of causal mediation models evaluating the role of epicardial adipose tissue (EAT) thickness as a mediating factor of the impact of age and body-mass index (BMI) on severe COVID-19 and mortality (COVID-19 outcome). A third causal mediation model was developed to investigate the association of ultrasensitive cardiac troponins as mediators of the relationship between EAT and adverse COVID-19 outcomes via direct heart damage