1

COVID-19 and Cognitive Impairment: Severity, Evolution, and Functional Impact during Inpatient Rehabilitation

Ruchi Patel MA OTR/L[1], Irene Savrides MA OTR/L[1], Christine Cahalan MS OTR/L[1], Gargi

Doulatani MA[2], Michael W. O'Dell MD[1,2], Joan Toglia PhD OTR/L[1,2,4], & Abhishek Jaywant,

PhD[1,2,3]

[1]Department of Rehabilitation Medicine, NewYork-Presbyterian Hospital/Weill Cornell Medical

Center

[2]Department of Rehabilitation Medicine, Weill Cornell Medicine

[3]Department of Psychiatry, Weill Cornell Medicine

[4]School of Health and Natural Sciences, Mercy College

Manuscript Word Count: 1196 words

Corresponding Author:

Abhishek Jaywant, PhD

525 E 68th St, F-1232, New York NY 10065

Email: abj2006@med.cornell.edu

Tel: 857-225-5783

Abstract

Objective: To determine the frequency, magnitude, and change in cognitive impairment in patients with COVID-19 undergoing acute inpatient rehabilitation. We secondarily evaluated correlates of cognitive impairment and the relationship between cognition and functional gain.

Design: Cross-sectional observational study with assessments at admission and discharge

Setting: Acute inpatient rehabilitation unit within a large, urban academic medical center

Participants: 77 patients hospitalized for COVID-19 and subsequently admitted to an inpatient rehabilitation unit between March-August 2020, 45 of whom were re-assessed at discharge.

Interventions: N/A

Main Outcome Measures: Montreal Cognitive Assessment (MoCA) scores on admission and discharge (when available) and Quality Indicator for Self-Care (QI-SC) scores on admission and discharge.

Results: 62/77 (80.5%) of patients demonstrated cognitive deficits on the MoCA at admission: 39/77 (50.6%) were mildly impaired, 20/77 (26%) moderately impaired, and 3/77 (3.9%) severely impaired. Cognitive impairment was associated with a prior history of delirium, but not age or length of acute care hospitalization. 32/45 (71.1%) patients with discharge scores improved and met the MoCA minimally clinically important difference (MCID); however, 35/45 (77.8%) continued to score in the impaired range. Patients who met the MoCA MCID demonstrated significantly greater QI-SC score gains than those that did not meet the MCID (p=.02).

Conclusion: Cognitive impairment is common among hospitalized COVID-19 patients requiring acute inpatient rehabilitation. Cognitive impairment improves over the course of inpatient

42   rehabilitation, and is associated with functional gain. Nonetheless, cognitive deficits frequently

43   remain present at discharge, indicating the need for systematic assessment and follow-up,

44   especially given the association with functional outcome.

45   **Key Words:** Coronavirus, SARS-CoV-2, Cognition, Activities of Daily Living, Rehabilitation

1.Introduction

Acute and critical illness frequently result in cognitive impairment, affecting function and quality of life[1]. COVID-19 can result in respiratory distress, hypoxia, ICU admission, intubation/ventilation, delirium, and vascular complications, all associated with cognitive impairment[2]. Cognitive deficits lead to dependence in activities of daily living[3] (ADLs), prompting the need for assessing and treating cognitive dysfunction to optimize function.

Studies on COVID-19 show a high frequency of cognitive impairment in mild to severe disease expressions. Among community-dwelling patients with mild cases of COVID-19, 21% of patients experienced a decline in cognitive functioning[4]. 81% of hospitalized but post-acute COVID-19 patients score abnormally on neuropsychological measures of attention, processing speed, and executive function[5]. There is still limited knowledge on how cognitive deficits evolve and impact functional outcome. One study found 80% of COVID-19 patients on a rehabilitation unit had cognitive deficits, with continued impairment in over half of the patients at 1 month follow-up[6]; however, the relationship to functional gain was not evaluated. This study aims to determine the magnitude and change in cognitive impairment, correlates of cognitive impairment, and the relation between cognition and function, in a relatively large sample of COVID-19 patients admitted to inpatient rehabilitation.

2.Methods

94 patients hospitalized for COVID-19 were admitted to an inpatient rehabilitation unit (IRU) from March-August 2020. Seventeen patients were not administered our cognitive measure as they were either deemed as not exhibiting cognitive impairment during instrumental ADLs or

underwent other cognitive testing, resulting in a final sample of 77. Patients were medically

stable, able to tolerate 3 hours/day of therapy, and had potential to gain function. Infection

control guidelines were followed, allowing for admission of positive COVID-19 patients and in-

person administration of our cognitive measure. All data were extracted from retrospective chart

reviews under approval of the Weill Cornell Medicine Institutional Review Board. 19/77 patients

underwent a separate neuropsychological evaluation during their admission, the results of which

have been published[5]; the results described here are novel.

Demographic/clinical data extracted included age, gender, prior level of function, intubation

history and length, delirium history, and IRU and acute care lengths of stay (LOS) (Table 1).

The Montreal Cognitive Assessment (MoCA) was administered by occupational therapists (OT)

72 hours before/after admission to the IRU. The minimal clinical important difference (MCID)

of 3 points was used[7]. Scores range from 0-30 and are stratified as follows: $\geq$26=normal, 18-

25=mild impairment, 11-17=moderate impairment, and $\leq$10=severe impairment[8]. Admission

MoCA assessments were administered on all 77 subjects and discharge MoCA assessments on

45/77. Three patients received the Blind MoCA, which was converted to 30-point scale per

established guidelines[8]. 32 patients did not receive discharge assessment as they either scored

normal on the initial assessment, were unavailable, and/or declined.

The Quality Indicator for Self-Care (QI-SC) is a standard of care functional assessment for

inpatient rehabilitation facilities. Items scored on a scale of 1 to 6 include eating, grooming,

toileting, bathing, upper body dressing, lower body dressing, and donning/doffing footwear.

Total score ranges from 7, indicating total assistance, to 42, indicating independence[9]. The QI-

SC is reliable and discriminates between the quality of care among rehabilitation facilities[9]. OTs

assigned admission and discharge QI-SC scores to all participants.

Statistical analyses used non-parametric tests to evaluate group differences (Mann-Whitney U),

within-group change from admission to discharge (Wilcoxon Signed-Ranks Test), association

between continuous variables (Spearman rank-order correlations), and chi-square tests for

categorical variables. Effect sizes are reported as Cohen's d.

3.Results

80.5% of patients demonstrated cognitive deficits on the MoCA at admission, of whom 15/77

were in the normal range, 39/77 had mild deficits, 20/77 had moderate deficits, and 3/77 had

severe deficits (Figure 1). Admission MoCA scores were higher for patients who did not

previously have delirium (*M*=22.03, *SD*=4.64) than patients who previously had delirium

(*M*=18.95, *SD*=5.8), *U*=466.5, *p*=.02, Cohen's d=.58. Admission MoCA did not differ based on

whether a patient was intubated (*U*=454.5, *p*=.82), age (r_s=-.15, *p*=.19) or acute hospitalization

length of stay (r_s=-.08 *p*=.50).

Patients with discharge MoCA scores (n=45) did not differ from those who were not

administered the MoCA at discharge (n=32) in age, gender, prior neurologic history, prior

delirium, prior intubation, or prior hypoxia (all p's>.05). These 45 patients with admission and

discharge MoCA scores improved on the MoCA (*M*=4.02, *SD*=3.67, Z=5.08, p<.001, Cohen's

d=1.10) and QI-SC (*M*=12.2, *SD*=6.18, Z=5.47, p<.001, Cohen's d=1.97). 32/45 patients

improved and met the MoCA MCID at discharge. This group of patients had a greater increase in

QI-SC scores (*M*=13.88, *SD*=5.01) versus patients who did not meet the MCID or declined

(*M*=9.46, *SD*=7.79), Mann-Whitney *U*=113.0, *p*=.02, Cohen's d=0.75, a relationship that

persisted in a linear regression model accounting for age and delirium. Despite overall

improvement, 28/45 patients still had mild deficits on the MoCA at discharge, 7 had moderate

deficits, none had severe deficits, and 10 scored in the normal range.

4.Discussion

Among 77 COVID-19 patients undergoing acute inpatient rehabilitation, 80.5% of patients

demonstrated cognitive impairment, the majority of whom had mild deficits. This number is

consistent with studies using neuropsychological assessments to evaluate cognition in

hospitalized COVID-19 patients[5-6] and expands on these findings with a larger sample than

previous studies. Consistent with the literature, delirium during acute hospitalization is

associated with greater cognitive impairment at admission[10], emphasizing the importance of

early detection and management of delirium in COVID-19. The lack of association between

admission cognitive impairment and intubation is also consistent with prior work.[5]

COVID-19 patients exhibited significant improvement in cognition with a large effect size.

32/45 (71.1%) percent exhibited clinically meaningful improvement, possibly reflecting natural

recovery, improvement in residual acute mental status changes, and/or rehabilitation intervention

effects. Mean QI-SC scores for this subgroup improved with a large effect size, signifying that

patients required less assistance for basic ADLs. Notably, significantly greater functional

improvement was seen in patients who exhibited a clinically meaningful improvement in

cognition than those that did not. Cognitive ability is associated with short- and long-term

independence with self-care tasks in older adults[3] and our results reinforce the importance of

targeting cognition to improve functional gains in COVID-19. COVID-19 patients warrant

further follow-up as a majority continued to demonstrate mild deficits at discharge.

4.1.Study Limitations

Discharge cognitive data was not available for 32/77 patients, limiting the ability to determine

the true prevalence of cognitive deficits in our sample, although there were no

demographic/cliincal differences between the 45 with and 32 without discharge cognitive data.

Further research is necessary regarding the underlying etiology of the cognitive deficits as

imaging results and the effects of psychiatric illness[5] were not addressed in this sample. Finally,

this sample included a relatively sick group of patients from the first wave of COVID-19;

findings may not apply to milder disease presentations or from subsequent waves of illness.

5.Conclusion

This study confirms the presence of cognitive impairment in COVID-19 patients with prolonged

hospitalization requiring inpatient rehabilitation and provides new insights into the evolution of

cognitive status. As most patients demonstrated mild/moderate cognitive deficits on admission,

routine cognitive screening should be conducted for COVID-19 patients. Despite improvement

during inpatient rehabilitation, a majority demonstrated mild cognitive deficits at discharge,

indicating ongoing cognitive needs. As cognitive improvement is associated with functional

gain, interventions targeting cognitive dysfunction in the acute inpatient rehabilitation setting

may be important in optimizing independence.

Acknowledgements: The authors received no funding to complete this study. All authors have

no conflicts or competing interests to declare. AJ is funded through a Career Development

Award from the National Center of Medical Rehabilitation Research/ Eunice Kennedy Shriver

National Institute of Child Health and Human Development (1K12HD093427-04), via a

subaward from Georgetown University. We thank the staff of the IRU for assisting in cognitive

and functional assessments, and for caring for, our COVID-19 patients.

References

1. Ehlenbach WJ, Hough CL, Crane PK, et al. Association between acute care and critical illness

hospitalization and cognitive function in older adults. JAMA. 2010;303(8):763–770.

doi:10.1001/jama.2010.167

2. Wilcox ME, Brummel NE, Archer K, Ely EW, Jackson JC, Hopkins RO. Cognitive

dysfunction in ICU patients: risk factors, predictors, and rehabilitation interventions. Crit Care

Med. 2013 Sep;41(9 Suppl 1): S81-98. doi: 10.1097/CCM.0b013e3182a16946. PMID:

23989098.

3. Millán-Calenti JC, Tubío, J., Pita-Fernández, S., Rochette, S., Lorenzo, T., Maseda, A.

Cognitive impairment as predictor of functional dependence in an elderly sample. Archives of

Gerontology and Geriatrics. 2012;54(1):197-201. https://doi.org/10.1016/j.archger.2011.02.010.

4. Del Brutto, O.H., Wu, S., Mera, R.M., Costa, A.F., Recalde, B.Y. and Issa, N.P. Cognitive

decline among individuals with history of mild symptomatic SARS-CoV-2 infection: A

longitudinal prospective study nested to a population cohort. Eur J Neurol. 2021

https://doi.org/10.1111/ene.14775 Online ahead of print.

5. Jaywant A, Vanderlind WM, Alexopoulos GS, Fridman CB, Perlis RH, Gunning FM.

Frequency and profile of objective cognitive deficits in hospitalized patients recovering from

COVID-19. Neuropsychopharmacology. 2021 Feb 15:1–6. doi: 10.1038/s41386-021-00978-8.

Epub ahead of print. PMID: 33589778; PMCID: PMC7884062.

6. Alemanno F, Houdayer E, Parma A, Spina A, Del Forno A, Scatolini A, Angelone S,

Brugliera L, Tettamanti A, Beretta L, Iannaccone S. COVID-19 cognitive deficits after

respiratory assistance in the subacute phase: A COVID-rehabilitation unit experience. PLoS One.

2021 Feb 8;16(2):e0246590. doi: 10.1371/journal.pone.0246590. PMID: 33556127; PMCID:

PMC7870071. eCollection 2021.

7. Wu CY, Hung SJ, Lin KC, Chen KH, Chen P, Tsay PK. Responsiveness, minimal clinically

important difference, and validity of the MoCA in stroke rehabilitation. Occup Ther Int. 2019

Apr 14:2517658. doi: 10.1155/2019/2517658. PMID: 31097928; PMCID: PMC6487084.

8. www.mocatest.org Accessed March 1st, 2021.

9. Pardasaney PK, Deutsch A, Iriondo-Perez J, Ingber MJ, McMullen T. Measuring inpatient

rehabilitation facility quality of care: Discharge self-care functional status quality measure. Arch

Phys Med Rehabil. 2018 Jun;99(6):1035-1041. doi: 10.1016/j.apmr.2017.02.023. Epub 2017

Mar 28. PMID: 28363701.

10. Hopkins RO, Jackson JC. Short- and long-term cognitive outcomes in intensive care unit

survivors. Clin Chest Med. 2009 Mar;30(1):143-53, ix. doi: 10.1016/j.ccm.2008.11.001. PMID:

19186286.

	N=77	N= 45	N=32
Table 1. Demographics and clinical characteristics of main sample and subsample with discharge MoCA. Patients without discharge MoCA are included for comparison.			
	(Full Sample)	**(Patients with discharge MoCA)**	**(Patients without a discharge MoCA)**
Gender			
Male	49 (63.6%)	30 (66.7%)	19 (59.4%)
Female	28 (36.4%)	15 (33.3%)	13 (40.6%)
Age	61.03 (15.67)	61.98 (14.11)	59.69 (17.77)
Acute hospitalization length of stay	37.03 (31.8)	39.4 (34.11)	33.69 (28.4)
Inpatient rehabilitation unit length of stay	13.43 (6.47)	14.44 (6.06)	12 (6.84)
Documented delirium	44 (57.1%)	28 (62.2%)	16 (50%)
Intubation	59 (76.6%)	35 (77.8%)	24 (75%)
Intubation length	13.31 (10.53)	13.75 (10.03)	12.67 (11.37)
Documented hypoxia	69 (89.6%)	39 (86.7%)	30 (93.8%)
History of neurological diagnosis	2 (2.6%)	1 (2.2%)	1 (3.1%)
Initial MoCA scores	20.29 (5.47)	17.96 (4.71)	23.47 (5.06)
Discharge MoCA scores		22.02 (4.23)	
Initial QI-SC score	25.75 (6.21)	24.96 (4.94)	26.88 (7.59)
Discharge QI-SC score	36.57 (8.37)	37.11 (6.97)	35.81 (10.09)

199   Note. MoCA = Montreal Cognitive Assessment; QI-SC = Quality Indicator for Self-Care; All

200   values are N (%) or Mean (SD).

202   **Figure 1.** Classification of cognitive impairment by score on the Montreal Cognitive Assessment

203   (MoCA) at admission in the full sample, admission in the subsample with discharge scores, and

204   discharge in the latter sample.