Title: Factors preventing SARS-CoV-2 transmission during unintentional exposure in a GP

practice, Germany, 2020

Authors: T. Sonia Boender[1,2#], Jennifer K. Bender[3,4#*], Angelika Krüger[5], Kai Michaelis[6], and

Udo Buchholz[7]

[#]These authors contributed equally to this article

[1]Postgraduate Training for Applied Epidemiology (PAE), Department of Infectious Disease

Epidemiology, Robert Koch Institute, Germany, BoenderS@rki.de

[2]European Programme for Intervention Epidemiology Training (EPIET), European Centre for

Disease Prevention and Control, (ECDC), Sweden

[3]Department of Infectious Diseases, Nosocomial Pathogens and Antibiotic Resistances Unit,

Robert Koch Institute, Germany, BenderJ@rki.de

[4]European Programme for Public Health Microbiology Training (EUPHEM), European Centre

for Disease Prevention and Control, (ECDC), Sweden

[5]Public Health Office Nuremberg, Germany

[6]Department of Infectious Disease Epidemiology, Respiratory Infections Unit, Robert Koch

Institute, Germany

[7]Department of Infectious Disease Epidemiology, Respiratory Infections Unit, Robert Koch

Institute, Germany

Word count: 1471

Summary

Two general practitioners (GPs) with SARS-CoV-2 infection provided in-person patient care to

patients of their joint medical practice before and after symptom onset, up until SARS-CoV-2

laboratory confirmation. Through active contact tracing, the local public health authorities

recruited the cohort of patients that had contact with either GP in their putative infectious period.

In this cohort of patient contacts, we assess the frequency and determinants of SARS-CoV-2-

transmission from GPs to patients. We calculated incidence rate ratios (IRR) to explore the type

of contact as explanatory variable for COVID-19 cases. Among the cohort of 83 patient contacts,

we identified 22 (27%) COVID-19 cases; 17 (21%) possible, 3 (4%) probable, and 2 (2%)

confirmed cases. All 22 cases had contact with a GP when they did not wear a mask, and/or

when contact was ≥10 minutes. Importantly, patients who had contact <10 minutes with a GP

wearing a facemask were at reduced risk (IRR 0.21; 95%CI 0.01-0.99) of COVID-19. This

outbreak investigation adds to the body of evidence in supporting current guidelines on measures

at preventing transmission of SARS-CoV-2 in an outpatient setting.

Abstract

Two general practitioners (GPs) with a SARS-CoV-2 infection provided in-person patient

care to patients of their joint medical practice before and after symptom onset, up until SARS-

CoV-2 laboratory confirmation. In a retrospective cohort study of patient contacts, we assessed

the risk (frequency and determinants) of SARS-CoV-2 transmission from the GPs to their

patients. Our findings support the use of facemasks for GPs, and short consultation time, to

minimize the risk of transmission.

Text

Background

In Germany, the first case of COVID-19 was confirmed on 27 January 2020 (1). This was followed by several introductions and clusters of cases of COVID-19. On 27 February, Germany reported a total of 26 cases (2). By 3 March 2020, 196 cases were reported, of which the majority (n=101) belonged to a large cluster in North-Rhine Westphalia or were linked to importation from high risk areas (n=35, from Italy, China, Iran) (3). Three days later, on 6 March 2020, a total of 639 confirmed COVID-19 cases were reported in Germany (4). Extensive contact tracing by the responsible health authorities was ongoing.

In this field report, we report about one of the first of ten cases of COVID-19 that was notified to a local public health authority in Germany, early March 2020, in the context of a GP practice.

Epidemiology of the incident

Two GPs, working in their joint medical practice, had fallen ill with COVID-19 end of February / early March 2020. To ensure confidentiality, we report an anonymised timeline, counting days 1 to 22, starting two days before the first day of symptom onset of one of the GPs. Both GPs had been attending to patients shortly before and after onset of respiratory symptoms, until laboratory confirmation of SARS-CoV-2 infection on day nine (**Figure 1**). Of note, because of the limited number of cases of COVID-19 in the area at that time, exposure to SARS-CoV-2 outside of the GP practice was deemed unlikely.

Response & outbreak investigation

Through active contact tracing, the local public health authorities recruited the cohort of

patients that had contact with either GP in their putative infectious period (starting two days

before and ending 10 days after the date of symptom onset). The cohort of patients that was

exposed to SARS-CoV-2 during their GP visit provided the unique opportunity to investigate the

likelihood of transmission in a GP setting. Therefore, and in accordance with national contact

tracing guidelines (5), we included patients in our cohort who were exposed to either GP two

days before their respective date of symptom onset until laboratory confirmation of SARS-CoV-

2 infection of the GP. Thus, the study period included all valid patient contacts from day one to

day nine of the anonymised timeline. The GPs were self-isolating from day nine onwards. Both

GPs made a full recovery. Close contacts of either GP were ordered a 14-day home quarantine.

Quarantine included active daily monitoring of COVID-19 symptoms and testing for SARS-

CoV-2 in the event of symptom onset and/or at the end of the quarantine period. The full referent

period can therefore be defined as the day one (two days before the symptom onset of the first

GP) up to and including day 22 (last day of quarantine). Work-related contacts included six

practice assistants; all reported distant contact, tested negative for SARS-CoV-2 and were not

considered part of the cohort.

In addition to routine contact tracing phone calls, we interviewed all patient contacts by

phone at the end of their putative incubation period (14 days after exposure) and again four

weeks later, using a standardised questionnaire. For each GP visit, the (medical) purpose,

duration and type of contact with the GP was extracted, including which GP was seen, the

distance to the GP, if the patient was physically examined, and whether or not the GP wore a

medical facemask. At the time, wearing a facemask by GPs was not yet recommended as a basic

infection prevention and control (IPC) measure for all situations of direct contact to all patients

(regardless of infection status) in Germany.

We collected information on potential symptom onset in the 14 days after risk contact

with the GP, including the type and timing of general and COVID-19-associated symptoms (6).

New onset was defined as a symptom that started the day after the risk contact up until 14 days

after the GP visit. COVID-19-compatible symptoms were separated into respiratory and non-

respiratory symptoms. Respiratory symptoms were defined as cough, sore throat, common-cold

(runny nose), shortness of breath and/or pneumoniae. Non-respiratory symptoms were defined as

fever, anosmia (loss of the sense of smell), ageusia (loss of the sense of taste), dysgeusia

(distortion of the sense of taste), headache, back-pain, muscle ache, joint-pain, loss of appetite

and/or weight, nausea, vomiting, diarrhoea, conjunctivitis, skin rash, swollen lymph nodes and/or

apathy.

Case definitions were based on self-reported symptoms and laboratory criteria. Possible

cases reported onset of at least two COVID-19-compatible symptoms. Probable cases met the

criteria for a possible case, but also had pneumonia, anosmia, ageusia or dysgeusia. Confirmed

cases tested positive for SARS-COV-2, regardless of symptoms (i.e. national COVID-19 case

definition).

We used negative binomial regression to calculate incidence rate ratios (IRR) to explore

the risk (frequency and determinants) of SARS-CoV-2 transmission from GPs to patients using

the MASS and fmsb packages in R Version 3.6.3.

We conducted this outbreak investigation as part of the official tasks of the local public

health authorities of the respective district supported by the RKI upon official request in

accordance to §4 of the German Protection against Infection Act. Therefore, this investigation

was exempt from additional institutional review.

<u>Results & discussion</u>

We reached 91 (70%) of 131 patient contacts as provided by the GPs. Of these, four

contacts did not meet our inclusion criteria --they were not exposed to or were exposed outside

the infectious period of either GP--and four did not consent. Thus, 83 (91%) of 91 were included

in the cohort study.

The median age was 45 [IQR 7-87] years, 56 (68%) were female. There were 25 (30%)

current smokers (tobacco) and 54 (65%) reported underlying conditions. Some patients visited

their GPs more than once during the infectious period of the GPs, we therefore recorded 89

contact-events in our study (GP_X n=43; GP_Y n=46). Patients spent a median of 10 minutes (IQR

10-20) with their GP. The GPs wore a medical facemask during 31 (35%) of 89 contact-events,

more often during physical examination (25/56; 45%) compared to no physical examination

(6/32; 19%; Fisher's exact P=.046).

In total, 28 participants (33.7%) reported onset of COVID-19 associated symptoms after

exposure to either GP, 22 (26.5%) reported onset of respiratory symptoms including fever, 21

(25.3%) reported onset respiratory symptoms, irrespective of fever. Onset of sore throat (n=17;

61%), cough (n=10; 36%), common cold (n=10; 36%), headache (n=9; 32%) and apathy (n=7;

25%) were the most frequently reported symptoms (**Figure 1**). We identified 22 (27%) COVID-

19 cases, including 17 (21%) possible, 3 (4%) probable, and 2 (2%) confirmed cases (**Figure 2**

& Table 1).

No single type of contact was significantly associated with the risk of getting COVID-19

(**Table 2**); however, the trends indicated that contact with GP X, contact in the pre-symptomatic

phase (data included for GP X only, as GP Y has not seen any patient in their pre-symptomatic

phase), and contact ≥10 minutes increased the risk, while contact with a GP wearing a facemask

decreased the risk (**Table 2**). All 22 identified COVID-19 cases had contact with a GP when they

did not wear a mask, and/or when contact was ≥10 minutes. Importantly, contact <10 minutes

with a GP wearing a facemask was significantly associated with a reduced risk (IRR 0.21;

95%CI 0.01-0.99) of COVID-19. When assessing the risk using a cumulative score (1 point for

each: duration ≥10 minutes; distance <1 meter and/or physical examination; contact without a

facemask or contact ≥10 minutes with facemask), trends indicated that increased exposure

increased risk of COVID-19 (IRR per point 1.23; 95%CI 0.76-2.15).

Our study has a few limitations. First, and due to the observational nature of our study,

case definitions rely on self-reported symptoms and the availability of PCR test results (64/83

were tested), and are therefore an estimation of the true number of cases. Also, not all potentially

confounding factors could be considered in the statistical analysis, because of the limited sample

size. Second, we cannot exclude that the patients were exposed to another source for SARS-

CoV-2 infection, even though community transmission in Germany at the time was limited (3,

4). Last, case definition for possible cases included symptoms that could also have been caused

by others such as seasonal influenza which was still circulating by the time the patients were

exposed to the infectious GPs.

    Conclusion

In conclusion, the use of facemasks for GPs and short consultation times helped limiting the

spread of SARS-CoV-2 from GPs to their patients. Should a GP or other healthcare workers

outside of a hospital setting with COVID-19 be shedding SARS-CoV-2 at the peak of

infectiousness (i.e. shortly before or after symptom onset) then these types of IPC measures may

prevent transmission to susceptible patients. Our investigation lends support to current guidelines

from WHO and the ECDC, both recommending all healthcare workers to continuously wear a

medical facemask at work especially in areas of community transmission (7-10).

Acknowledgments

We thank the Robert Koch Institute's team supporting the interviews: Katharina Alpers,

Nicu Anghel, Mona Askar, Eva Feuerhahn, Christina Frank, Jane Hecht, Alexandra Holzer, Anja

Klingeberg, Teresa Nygren, Andreas Reich, Viktoria Schönfeld, Doreen Staat, Jan Walter, and

especially Felix Moek for the organization. We further thank Merle Böhmer from the federal

state public health authority and the local public health authorities, especially Clara Vicén

Renner for carrying out several interviews. Last but not least, we very much appreciate and thank

all participants of the study.

Disclaimers

No conflict of interest reported. TSB and JB are fellows of the ECDC Fellowship

Programme, supported financially by the European Centre for Disease Prevention and Control

(ECDC). The views and opinions expressed herein do not state or reflect those of ECDC. ECDC

is not responsible for the data and information collation and analysis and cannot be held liable

for conclusions or opinions drawn.

Data availability statement

Detailed data are confidential and protected by German law. Anonymized data are

available from the corresponding author upon reasonable request.

References

1.   Bohmer MM, Buchholz U, Corman VM, Hoch M, Katz K, Marosevic DV, et al.

Investigation of a COVID-19 outbreak in Germany resulting from a single travel-associated

primary case: a case series. Lancet Infect Dis. 2020.

2.   Coronavirus Disease 2019 (COVID-19) Daily Situation Report of the Robert Koch

Institute 27/02/2020. Robert Koch Institute 27 February 2020.

3.   Coronavirus Disease 2019 (COVID-19) Daily Situation Report of the Robert Koch

Institute 03/03/2020. Robert Koch Institute 3 March 2020.

4.   Coronavirus Disease 2019 (COVID-19) Daily Situation Report of the Robert Koch

Institute 06/03/2020. Robert Koch Institute 6 March 2020.

5.   Kontaktpersonennachverfolgung bei respiratorischen Erkrankungen durch das

Coronavirus SARS-CoV-2: Robert Koch Institute; 2020 [2 July 2020]. Available from:

https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Kontaktperson/Management.h

tml.

6.   Tong JY, Wong A, Zhu D, Fastenberg JH, Tham T. The Prevalence of Olfactory and

Gustatory Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis.

Otolaryngol Head Neck Surg. 2020;163(1):3-11.

7.   Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schunemann HJ, et al. Physical

distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-

CoV-2 and COVID-19: a systematic review and meta-analysis. Lancet. 2020;395(10242):1973-

87.

8.   Howard J, Huang A, Li Z, Tufekci Z, Zdimal V, van der Westhuizen H, et al. Face Masks

Against COVID-19: An Evidence Review. Preprints. 2020(2020040203).

9.   Mitze T, Kosfeld R, Rode J, Wälde K. Face Masks Considerably Reduce COVID-19

Cases in Germany:A Synthetic Control Method Approach. Bonn, Germany: IZA Institute of

Labor Economics June 2020. Contract No.: IZA DP No. 13319.

10.   ECDC. COVID-19 infection prevention and control measures for primary care, including

general practitioner practices, dental clinics and pharmacy settings: first update. European Centre

for Disease Prevention and Control 19 October 2020.

Figure 1. Percentage of self-reported onset of COVID-19-associated symptoms after contact with

either GP, presented as a percentage of the full cohort, as recalled by the 28/83 study

participants.

Figure 2. Date of symptom (following the anonymized timeline) onset of possible, probable and

confirmed COVID-19 cases within the cohort of patient contacts of two SARS-CoV-2-positive

GPs in a medical practice in Germany, February-March 2020. Symptom onset of either GP, the

days they attended to patients, and closure of the GP medical practice are marked by blue (GP

X), pink (GP Y) and black arrows, respectively.

217   Table 1. Summary of characteristics of the five confirmed and probable COVID-19 cases within the cohort of patient contacts of two

218   SARS-CoV-2-positive GPs in a GP practice in Germany, 2020.

	Case 1	Case 2	Case 3	Case 4	Case 5
Case classification	Confirmed	Confirmed	Probable	Probable	Probable
Age group (years)	Between 20-25	Between 40-45	Between 20-25	Between 45-50	Between 80-85
Sex	Female	Female	Female	Female	Female
Incubation period (days)	3	14	3	1	5
Underlying conditions	No	No	No	Yes	Yes
GP visited	X	X	Y	Y	Y
Time with GP (minutes)	10	10	13	12	5
Distance to GP (meter)	<1	<1	1-2	<1	<1
Close contact[a]	Close	Close	Close	Close	Close
Physical examination by GP	Yes	No	Yes	Yes	Yes
GP wore a face mask	No	No	Yes	Yes	No
Risk contact score[b]	2	2	2	2	2

219   [a]Combination of distance <1m and/or physical examination/blood drawn/injection; [b]Risk contact score (scale 0-3), one point for each:

220   duration (1 point if ≥10 minutes), distance (1 point for <1 meter and/or physical examination), GP with facemask (1 point for contact

221   without a facemask, or contact ≥10 minutes with a facemask)

222 Table 2. Incidence and incidence rate ratios (IRR) of COVID-19 cases (possible, probable and confirmed), by risk contact-event with

223 SARS-CoV-2-positive general practitioners (GP).

Type of risk contact	Exposure	Total (n=89)	%	No case (n=67)	%	Case (n=22)	%	Incidence rate ratios	95%CI lower	95%CI upper
GP	GP Y	46	51.7	36	53.7	10	45.5	1 (ref)	(ref)	(ref)
	GP X	43	48.3	31	46.3	12	54.5	1.28	0.55	3.04
Phase of infection of GPs	Symptomatic	20	22.5	16	23.9	4	18.2	1 (ref)	(ref)	(ref)
	Pre-symptomatic	69	77.5	51	76.1	18	81.8	1.30	0.49	4.51
Duration	≥10 minutes	70	78.7	52	77.6	18	81.8	1 (ref)	(ref)	(ref)
	<10 minutes	18	20.2	15	22.4	3	13.6	0.65	0.15	1.91
Facemask worn by GP	No facemask	53	59.6	39	58.2	14	63.6	1 (ref)	(ref)	(ref)
	Facemask	31	34.8	24	35.8	7	31.8	0.85	0.32	2.05
	Unknown	5	5.6	4	6.0	1	4.5	-	-	-
Duration and facemask worn by GP combined[a]	Exposed (contact ≥10 minutes and/or GP not wearing facemask)	75	84.3	53	79.1	22	100	1 (ref)	(ref)	(ref)
	Unexposed (GP wore facemask and <10 minutes contact)	14	15.7	14	20.9	0	0.0	0.21	0.01	0.99
Risk contact score[b]	Per point (*continuous*)	NA	NA	NA	NA	NA	NA	1.23	0.76	2.15

224 [a] IRR calculated using Haldane-Anscombe correction; [b] Risk contact score (scale 0-3), one point for each: duration (1 point if ≥10

225 minutes), distance (1 point for <1 meter and/or physical examination), GP with facemask (1 point for contact without a facemask, or

226 contact ≥10 minutes with a facemask).

14