1   **Title**: Background rates of all-cause mortality, hospitalizations, and emergency department
2   visits among nursing home residents in Ontario, Canada to inform COVID-19 vaccine safety
3   assessments
4

5   **Authors**: Maria Sundaram[1,2], Sharifa Nasreen[2], Andrew Calzavara[1], Siyi He[1], Hannah Chung[1],
6   Susan E. Bronskill[1,2,3], Sarah A. Buchan[1,2,4], Mina Tadrous[1,3], Peter Tanuseputro[1,5], Kumanan
7   Wilson[5], Sarah Wilson[1,2,4], Jeffrey C. Kwong[1,2,4,6,7], on behalf of the Canadian Immunization
8   Research Network (CIRN) investigators
9
Affiliations:
[1] ICES, Toronto, ON
[2] Dalla Lana School of Public Health, University of Toronto, Toronto, ON
[3] Women's College Research Institute, Toronto, ON
[4] Public Health Ontario, ON
[5] Department of Medicine, University of Ottawa and Bruyere and Ottawa Hospital Research
Institutes, Ottawa, ON
[6] Department of Family and Community Medicine, University of Toronto, Toronto, ON
[7] University Health Network, Toronto, ON
19
Corresponding author:
Jeff Kwong
Senior Scientist, ICES
G1 06, 2075 Bayview Avenue
Toronto, Ontario, Canada, M4N 3M5
jeff.kwong@utoronto.ca
Phone: (416) 480-4055 x1-7665
Fax: (416) 480-6048

28
29

Abstract

Background. Nursing home (NH) residents are prioritized for COVID-19 vaccination. We report

monthly mortality, hospitalizations, and emergency department (ED) visit incidence rates (IRs)

during 2010-2020 to provide context for COVID-19 vaccine safety assessments.

Methods. We observed outcomes among NH residents using administrative databases. IRs

were calculated by month, sex, and age group. Comparisons between months were assessed

using one-sample t-tests; comparisons by age and sex were assessed using chi-squared tests.

Results. From 2010-2019, there were 83,453 (SD: 652.4) NH residents per month, with an

average of 2.3 (SD: 0.28) deaths, 3.1 (SD: 0.16) hospitalizations, and 3.6 (SD: 0.17) ED visits

per 100 residents per month. From March to December 2020, mortality IRs were increased, but

hospitalization and ED visit IRs were reduced (p<0.05).

Conclusion. We identified consistent monthly mortality, hospitalization, and ED visit IRs during

2010-2019. Marked differences in these rates were observed during 2020, coinciding with the

COVID-19 pandemic.

Keywords: long-term care, mortality, hospitalization, baseline rates, COVID-19 vaccine safety
49

Introduction[1]

Nursing home (NH) residents have been disproportionately affected by the coronavirus disease

2019 (COVID-19) pandemic, comprising a substantial proportion of total COVID-19 deaths in

many countries, including over 80% of COVID-19 deaths in Canada during the first epidemic

wave [1, 2]. The vaccine rollout has prioritized, among others, NH residents because of their

high risk of poor outcomes from COVID-19 [3].

However, NH residents are generally frail and are at comparatively higher risk of poor

outcomes, even outside the context of a global pandemic [4, 5]. As a result, deaths and

healthcare use within the expected range of event numbers for NH residents could create an

erroneous impression that vaccine rollout is associated with increased risk of such events [6].

There is a lack of data on poor outcome rates among NH residents in Canada to facilitate

comparisons of observed versus expected outcome rates during vaccination rollout. Therefore,

we sought to quantify background rates of all-cause mortality, hospitalizations, and emergency

department (ED) visits among all NH residents in Ontario, the most populous province in

Canada. Since COVID-19 and related restrictions may also impact these outcomes, we report

these rates separately during 2020.

Methods

We conducted an observational analysis using population-level health administrative databases

encompassing all residents of Ontario, Canada (2020 population: approximately 14.7 million, of

which 10% live in rural areas; median age: 40.4 years) [7, 8]. Ontario has a single-payer health

system that provides universal access to physician and hospital services for all Ontario

residents, including NH residents, who receive personal and nursing care and subsidized

---

[1] **Abbreviations**: NH, nursing home. COVID-19, coronavirus disease 2019. ED, emergency
department.

housing [9]. In Ontario, NH may be not-for-profit (managed either by public, municipal, or private

companies), or for-profit institutions managed by private companies [9].

These datasets were linked using unique encoded identifiers and analyzed at ICES (formerly

called the Institute for Clinical Evaluative Sciences). ICES is a not-for-profit research institute

that is a prescribed entity under Ontario's Personal Health Information Protection Act (PHIPA).

Section 45 of PHIPA authorizes ICES to collect personal health information, without consent, for

the purpose of analysis or compiling statistical information with respect to the management of,

evaluation, or monitoring of, the allocation of resources to or planning for all or part of the health

system.

Cohort definitions

We identified all Ontarians aged 18 years or older living in NHs using the physician billing

database Ontario Health Insurance Plan (encompassing most outpatient interactions with the

healthcare system), Ontario Drug Benefit (encompassing prescriptions), and the Continuing

Care Reporting System (encompassing information about individuals in NH) databases.

Information on age, sex and death were collected from the Registered Persons Database. We

identified individual cohorts beginning at the start of each month from January 1, 2010 to

December 31, 2019 inclusive, and from January 1, 2020 to December 31, 2020. Individuals

were included in these cohorts if they were alive at the beginning of a month and recorded as

living in a NH at least one day in that month or in the previous 90 days. We defined NH

residents in this way because administrative datasets in Ontario may have incomplete capture

of individual entry in, and exit from, NH care; but interactions with the healthcare system that

would identify the individual as a NH resident are expected to happen at least once every 90

days. As a result, ICES uses a multi-component definition of NH resident that incorporates

information from healthcare system interactions in a 90-day period that can identify individuals

as NH residents.

Statistical analysis

We calculated monthly rates per 100 NH residents overall, by sex, and by age group (18-69

years, 70-79 years, 80-89 years, and ≥90 years). We also conducted analyses comparing adults

aged ≥80 years with adults younger than 80 years. Estimates from 2010-2019 were combined

over years by month, and by sex and age group separately. Direct comparisons were conducted

either with one-sample t-tests, in the case of comparisons by month from 2010-2019 vs. 2020,

or with chi-squared tests, in the case of comparisons by sex or age. We also calculated the

relative change in mortality in 2020, compared with mortality in 2010-2019, using a simple ratio

(mortality in 2020 divided by average mortality in 2010-2019). As the first presumed SARS-CoV-

2 positive case was identified in Canada on January 25, 2020, and Health Canada did not report

that the primary source of infections was "local" transmission until March 24, 2020 [10], we

defined the pandemic period in 2020 as beginning approximately on March 1, 2020.

Results

From January 1, 2010 through December 31, 2019, there were an average of 83,453 (SD:

652.4) NH residents in Ontario in any given month (**Figure 1A**). Of these, an average of 57,104

(SD: 291.6; ~68% of the total in any given month) were women, and an average of 58,733 (SD:

550.6; ~70% of the total in any given month) were aged ≥80 years.

During this period, approximately 2.3 (SD: 0.28) deaths per 100 NH residents occurred per

month (**Figure 1B; Table 1**). These rates were higher for men (incidence rate [IR]: 2.6; SD:

0.31) than women (IR: 2.2; SD: 0.27) (p<0.001; **Figure 1E**), and for residents aged ≥80 years

(IR: 5.6; SD: 0.68) than residents aged <80 years (IR: 2.6; SD: 0.33) (p<0.001; **Figure 1H**). An

average of 0.4 (SD: 0.04) more monthly deaths per 100 residents occurred during winter

months (December, January, February, and March) than during other months during 2010-2019

(p<0.001; **Table 1**).

We observed monthly averages of 3.1 (SD: 0.16) hospitalizations per 100 residents and 3.6

(SD: 0.17) ED visits per 100 residents during 2010-2019 (**Figures 1C and 1D; Table 1**).

Hospitalization rates were higher for men (IR: 3.8; SD: 0.20) than women (IR: 2.7; SD: 0.15)

(p<0.001) (**Figure 1F; Table 1**) and were lower for older residents (IR: 5.8; SD: 0.34) than

younger residents (IR: 6.8; SD: 0.35) (p<0.001; **Figure 1I**). Similarly, ED visit rates were higher

for men (IR: 4.3; SD: 0.21) than women (IR: 3.3; SD: 0.18) (p<0.001; **Figure 1G**) and lower for

older residents (IR: 6.7; SD: 0.33) than younger residents (IR: 8.6; SD: 0.45) (p<0.001; **Figure**

1J).

We observed a steady decline in the total number of NH residents during 2020 (non-parametric

test for trend p<0.01; **Figure 1A**). No statistically significant differences were observed in

mortality rates in January or February of 2020 compared to those months in previous years.

Mortality rates rose during March 2020, and were higher in all months of the pandemic period

(March – December 2020) compared to corresponding months in 2010-2019 (p<0.05 for each

month) (**Figure 1B, Table 2**); these differences were most pronounced in April and May 2020.

Mortality was higher for men (**Figure 1E**) and older residents (**Figure 1H**). The ratio of excess

deaths in 2020 to average mortality incidence in 2010-2019 was highest in April and May 2020,

across sexes and age groups (**Table 3**).

There was a substantial reduction in hospitalizations during 2020 compared to previous years

(p<0.001 for each month during the pandemic period; **Figure 1C** and **Table 2**). Hospitalization

rates were higher for men than women (p<0.001; **Figure 1F**) and lower with increasing age

throughout the pandemic period (**Figure 1I**). Similar to hospitalizations, there was a substantial

reduction in ED visits during 2020 compared to previous years (p<0.001 for each month during

2020; **Figure 1D** and **Table 2**). Patterns by sex and age were similar as for hospitalizations

(**Figures 1G** and **1J**).

Discussion

This report presents rates of all-cause mortality, hospitalizations, and ED visits in NH residents

in Ontario, Canada during a 10-year pre-pandemic period (2010-2019) and during 2020.

Mortality rates in 2010-2019 are similar to rates reported by others in Canada [4], with rates

being higher in men compared to women, and higher with increasing age. As expected,

mortality rates spiked during the early phase of the COVID-19 pandemic, with high mortality

rates among all age groups and sexes in April and May of 2020, similar to overall estimates for

the province of Ontario [11]. Although mortality rates returned to values closer to pre-pandemic

rates in August 2020, they rose again in November and December 2020, in keeping with the

timing of Ontario's second wave [12].

The decrease in mortality rates during summer 2020 is likely multifactorial. First, changes in

provincial-level testing guidelines may have enabled earlier identification of COVID-19 cases.

On April 8, 2020, Ontario officially expanded its testing guidelines beyond symptomatic NH

residents to all residents, including all newly admitted individuals and asymptomatic residents

who had contact with a confirmed-positive case [13]. Earlier identification of COVID-19 cases

may have contributed to a subsequent reduction in mortality by preventing potential outbreaks

and by providing needed care to residents earlier in the course of their illness. Second,

enhanced infection prevention measures were implemented in NHs province-wide.[14] Third,

reduced mortality rates in the summer months likely reflect a reduction in community-level

transmission of SARS-CoV-2 experienced throughout Ontario [11].

Our analysis additionally identified a decline in hospitalizations and ED visits during 2020,

compared to 2010-2019. This finding may initially seem counterintuitive but, taken in context

with simultaneous increases in NH setting mortality and decreases in the total number of NH

residents, these reductions likely indicate a reduction in hospitalized deaths, which has been

reported previously [15]. There may have also been an increased unwillingness to seek

hospital-based care for non-COVID-19-related issues during the beginning of the first pandemic

wave; future analyses should also consider the reasons noted for ED visits and hospitalizations

to identify COVID-19-related vs. non-COVID-19-related incidents. Our analysis identified that

older adults (≥90 years) had the lowest hospitalization and ED visit rates both in pre-pandemic

and pandemic periods; and that younger adults (18-64 years) had the highest hospitalization

and ED visits in both periods. We believe these patterns reflect different goals of care for

different age groups and populations.

This analysis uses large-scale health administrative datasets, which are not yet currently

updated to the end of December 2020 with vital statistics. As a result, we are only able to report

data up to the end of 2020, an important limitation of this work. If feasible, we will update this

report as more data become available.

We intend for these findings to be used to contextualize potential increases in adverse events of

special interest (AESI) [16] during the COVID-19 vaccine rollout in NH residents. Whereas

mortality, hospitalization, and ED visit rates were highly stable over 10 pre-pandemic years, the

marked changes in rates observed during 2020 indicate that caution is needed in determining

the appropriate baseline to use when interpreting rates of AESIs following the COVID-19

vaccine rollout. Existing investigations on mortality in elderly and frail individuals who received

COVID-19 vaccines has identified that mortality rates were consistent with high background

rates of death in this population [17]; our data will help contextualize Canada- and Ontario-

specific rates in a similar way.

Conclusions

This report identified relatively consistent monthly mortality, hospitalization, and ED visit rates

for Ontario NT residents over a ten-year pre-pandemic period. During 2020, we identified a

substantial increase in mortality rates during April and May, followed by a regression during

summer 2020 and subsequent increase in November and December 2020. In contrast,

hospitalization and ED visit rates declined markedly in 2020 and (as of the latest data available)

have not yet returned to baseline levels. These results provide important context for future

analyses to identify potential AESI in NH residents.

Funding

This work was supported by the Canadian Immunization Research Network (CIRN) through a

grant from the Public Health Agency of Canada and the Canadian Institutes of Health Research

(CNF 151944). This study was also supported by ICES, which is funded by an annual grant

from the Ontario Ministry of Health (MOH). No endorsement by ICES or MOH is intended or

should be inferred.

Conflict of interest

The authors declare no conflicts of interest.

Authors' contributions

JCK conceived of the study design and oversaw the study. AC obtained the data and calculated

incidence rates. MS conducted additional statistical analysis and drafted the manuscript. PT,

SEB, KW, and MT provided methodological input. All authors interpreted the results, critically

revised the manuscript, and have approved the final version for publication.

References

[1] Pandemic experience in the long-term care sector: how does Canada compare with other

countries? Ottawa, ON: Canadian Institute for Health Information; 2020.

[2] Sepulveda ER, Stall NM, Sinha SK. A comparison of COVID-19 mortality rates among long-

term care residents in 12 OECD countries. JAMDA. 2020;21:1572-4 e1.

[3] WHO SAGE roadmap for prioritizing uses of COVID-19 vaccines in the context of limited

supply. Geneva, Switzerland: World Health Organization; 2020.

[4] Tanuseputro P, Chalifoux M, Bennett C, Gruneir A, Bronskill SE, Walker P, et al.

Hospitalization and Mortality Rates in Long-Term Care Facilities: Does For-Profit Status Matter?

J Am Med Dir Assoc. 2015;16:874-83.

[5] Gruneir A, Bell CM, Bronskill SE, Schull M, Anderson GM, Rochon PA. Frequency and

pattern of emergency department visits by long-term care residents--a population-based study.

J Am Geriatr Soc. 2010;58:510-7.

[6] Branswell H. CDC advisory panel's lone dissenter on why long-term care residents shouldn't

receive COVID-19 vaccine first. STATNews2020.

[7] Table 17-10-0005-01: Population estimates on July 1st, by age and sex. Canada: Statistics

Canada; 2020.

[8] Sundaram ME, Calzavara A, Mishra S, Kustra R, Chan AK, Hamilton MA, et al. Individual

and social determinants of SARS-CoV-2 testing and positivity in Ontario, Canada: a population-

wide study. CMAJ. 2021.

[9] Stall NM, Jones A, Brown KA, Rochon PA, Costa AP. For-profit long-term care homes and

the risk of COVID-19 outbreaks and resident deaths. CMAJ. 2020;192:E946-E55.

[10] COVID-19: A Canadian timeline. Canada: HealthcareNetwork; 2020.

[11] All Ontario: Case numbers and spread. Ontario: Ontario Ministry of Health; 2021.

[12] COVID-19: Modelling Update. COVID-19 Fall Preparedness. Ontario: Government of

Ontario; 2020.

[13] COVID-19 Provincial Testing Guidance Update: April 8, 2020. Ontario: Ministry of Health;

2020.

[14] COVID-19 Long-Term Care Resources. Ontario: Public Health Ontario; 2021.

[15] Brown KA, Daneman N, Buchan SA, Chan AK, Stall NM. Temporal variations in the

intensity of care provided to community and nursing home residents who died of COVID-19 in

Ontario, Canada. medRxiv. 2020.

[16] Law B, Sturkenboom M. D2.3: Priority List of Adverse Events of Special Interest: COVID-

19. SPEAC: Safety Platform for Emergency vACcines CEPI; 2020.

[17] GACVS COVID-19 Vaccine Safety subcommittee meeting to review reports of deaths of

very frail elderly individuals vaccinated with Pfizer BioNTech COVID-19 vaccine, BNT162b2.

Geneva, Switzerland: World Health Organization; 2021.

 Table 1. Mean incidence rates per 100 nursing home residents, monthly from 2010-2019.

	Overall: IR (SD)	Women: IR (SD)	Men: IR (SD)	18-69: IR (SD)	70-79: IR (SD)	80-89: IR (SD)	90+: IR (SD)
Mortality rates							
January	2.8 (0.28)	2.7 (0.29)	3.2 (0.29)	1.1 (0.18)	2.2 (0.19)	2.8 (0.26)	4.0 (0.47)
February	2.4 (0.19)	2.3 (0.20)	2.7 (0.20)	0.9 (0.09)	1.8 (0.21)	2.5 (0.19)	3.4 (0.25)
March	2.5 (0.13)	2.4 (0.16)	2.9 (0.11)	1.0 (0.10)	1.8 (0.14)	2.6 (0.12)	3.6 (0.25)
April	2.3 (0.11)	2.2 (0.10)	2.6 (0.14)	0.9 (0.13)	1.8 (0.14)	2.4 (0.11)	3.2 (0.18)
May	2.2 (0.07)	2.1 (0.07)	2.5 (0.09)	0.9 (0.08)	1.7 (0.12)	2.3 (0.12)	3.1 (0.13)
June	1.9 (0.10)	1.8 (0.10)	2.20 (0.12)	0.8 (0.08)	1.4 (0.13)	2.0 (0.07)	2.7 (0.19)
July	2.0 (0.07)	1.9 (0.07)	2.3 (0.10)	0.8 (0.09)	1.6 (0.09)	2.1 (0.06)	2.8 (0.18)
August	2.1 (0.11)	1.9 (0.08)	2.4 (0.18)	0.8 (0.09)	1.5 (0.08)	2.1 (0.15)	2.9 (0.16)
September	2.1 (0.11)	1.9 (0.11)	2.4 (0.14)	0.8 (0.05)	1.6 (0.11)	2.1 (0.16)	2.9 (0.15)
October	2.3 (0.16)	2.2 (0.13)	2.7 (0.22)	0.9 (0.16)	1.8 (0.16)	2.4 (0.16)	3.3 (0.21)
November	2.3 (0.14)	2.2 (0.14)	2.7 (0.18)	0.9 (0.11)	1.8 (0.13)	2.3 (0.14)	3.3 (0.22)
December	2.5 (0.14)	2.4 (0.13)	2.9 (0.18)	1.0 (0.11)	1.9 (0.12)	2.6 (0.13)	3.5 (0.25)
Hospitalization rates							
January	3.3 (0.16)	3.0 (0.17)	4.1 (0.17)	3.4 (0.40)	3.8 (0.19)	3.5 (0.23)	2.9 (0.20)
February	3.0 (0.07)	2.6 (0.06)	3.7 (0.13)	3.1 (0.16)	3.4 (0.21)	3.0 (0.10)	2.5 (0.13)
March	3.3 (0.10)	2.9 (0.10)	4.1 (0.13)	3.4 (0.24)	3.8 (0.24)	3.4 (0.15)	2.8 (0.14)
April	3.1 (0.14)	2.7 (0.13)	3.8 (0.21)	3.3 (0.31)	3.6 (0.22)	3.2 (0.17)	2.6 (0.15)
May	3.1 (0.10)	2.8 (0.09)	3.8 (0.15)	3.4 (0.22)	3.5 (0.14)	3.2 (0.13)	2.6 (0.10)
June	2.9 (0.08)	2.6 (0.08)	3.6 (0.13)	3.2 (0.21)	3.3 (0.17)	3.0 (0.16)	2.5 (0.08)
July	3.1 (0.10)	2.8 (0.10)	3.7 (0.15)	3.4 (0.13)	3.5 (0.17)	3.1 (0.13)	2.6 (0.12)
August	3.0 (0.08)	2.7 (0.08)	3.7 (0.14)	3.3 (0.22)	3.5 (0.24)	3.1 (0.15)	2.6 (0.10)

September	3.0 (0.06)	2.7 (0.07)	3.7 (0.09)	3.2 (0.19)	3.4 (0.14)	3.1 (0.10)	2.5 (0.13)
October	3.1 (0.07)	2.8 (0.09)	3.9 (0.09)	3.4 (0.16)	3.6 (0.19)	3.2 (0.09)	2.6 (0.08)
November	3.0 (0.07)	2.6 (0.07)	3.7 (0.12)	3.2 (0.21)	3.4 (0.23)	3.0 (0.09)	2.5 (0.14)
December	3.1 (0.12)	2.7 (0.13)	3.9 (0.13)	3.4 (0.18)	3.6 (0.10)	3.2 (0.15)	2.6 (0.18)

ED visit rates

January	3.7 (0.10)	3.4 (0.08)	4.3 (0.21)	4.7 (0.20)	3.9 (0.21)	3.6 (0.14)	3.2 (0.13)
February	3.3 (0.09)	3.0 (0.08)	3.9 (0.22)	4.2 (0.23)	3.6 (0.26)	3.2 (0.07)	2.8 (0.16)
March	3.7 (0.09)	3.4 (0.12)	4.3 (0.12)	4.7 (0.22)	3.9 (0.13)	3.6 (0.11)	3.2 (0.18)
April	3.6 (0.08)	3.3 (0.12)	4.2 (0.09)	4.7 (0.19)	3.9 (0.19)	3.5 (0.12)	3.1 (0.13)
May	3.7 (0.08)	3.4 (0.08)	4.4 (0.15)	4.8 (0.17)	4.0 (0.14)	3.6 (0.11)	3.3 (0.15)
June	3.7 (0.10)	3.4 (0.09)	4.3 (0.18)	4.8 (0.21)	3.9 (0.25)	3.6 (0.11)	3.2 (0.12)
July	3.8 (0.10)	3.5 (0.14)	4.4 (0.11)	5.0 (0.20)	4.1 (0.10)	3.8 (0.13)	3.2 (0.18)
August	3.8 (0.09)	3.5 (0.09)	4.4 (0.19)	5.0 (0.26)	4.1 (0.15)	3.7 (0.10)	3.2 (0.14)
September	3.6 (0.05)	3.3 (0.09)	4.2 (0.15)	4.6 (0.14)	3.8 (0.19)	3.5 (0.09)	3.2 (0.10)
October	3.7 (0.10)	3.4 (0.13)	4.3 (0.16)	4.9 (0.19)	3.9 (0.20)	3.6 (0.15)	3.1 (0.12)
November	3.4 (0.09)	3.1 (0.11)	4.0 (0.15)	4.4 (0.16)	3.7 (0.19)	3.4 (0.11)	2.9 (0.10)
December	3.6 (0.11)	3.3 (0.15)	4.2 (0.15)	4.6 (0.19)	3.9 (0.15)	3.6 (0.14)	3.1 (0.18)

IR=incidence rate; SD=standard deviation

Table 2. Incidence rates per 100 nursing home residents during 2020.

	Overall	Women	Men	18-69	70-79	80-89	90+
Mortality rates							
January	2.8	2.8	2.9	1.1	2.3	2.8	4.0
February	2.4	2.2	2.8	0.9	1.7	2.4	3.4
March	2.6	2.4	3.2	1.1	1.8	2.8	3.6
April	4.0	3.7	4.7	1.7	3.0	4.1	5.5
May	3.3	3.0	3.9	1.8	2.7	3.4	4.2
June	2.2	2.0	2.6	1.1	1.7	2.2	2.8
July	2.1	2.0	2.4	0.9	1.5	2.1	3.0
August	2.2	2.0	2.6	0.9	1.6	2.1	3.1
September	2.2	2.1	2.5	0.9	1.7	2.3	3.0
October	2.6	2.5	2.9	1.0	1.7	2.7	3.7
November	2.9	2.6	3.5	0.8	2.1	3.0	4.1
December	3.3	3.1	3.8	1.3	2.5	3.4	4.6
Hospitalization rates							
January	3.2	2.8	3.8	3.6	3.5	3.4	2.5
February	2.8	2.5	3.5	3.3	3.4	2.9	2.3
March	2.2	1.9	2.8	2.9	2.5	2.2	1.7
April	1.7	1.5	2.1	2.6	2.1	1.7	1.1
May	1.9	1.7	2.3	2.7	2.5	1.8	1.3
June	1.9	1.6	2.4	2.4	2.2	1.8	1.5
July	2.0	1.7	2.5	2.5	2.2	2.0	1.6
August	2.2	1.9	2.8	2.6	2.5	2.2	1.9
September	2.2	1.9	2.9	2.6	2.8	2.3	1.7

October	2.3	2.0	3.0	2.7	2.6	2.6	1.8
November	2.4	2.0	3.2	3.0	2.9	2.4	1.8
December	2.4	2.0	3.2	3.0	3.1	2.2	1.9

ED visit rates

January	3.5	3.2	4.1	4.8	3.8	3.2	3.0
February	3.1	2.9	3.5	4.1	3.3	3.0	2.7
March	2.3	2.1	2.6	3.3	2.5	2.2	1.8
April	1.2	1.0	1.5	2.1	1.3	1.2	0.8
May	1.6	1.4	2.0	2.3	1.6	1.6	1.4
June	1.9	1.8	2.3	2.9	2.0	1.8	1.6
July	2.2	1.9	2.6	3.2	2.3	2.1	1.8
August	2.3	2.1	2.9	3.3	2.4	2.3	2.0
September	2.4	2.1	2.9	3.0	2.6	2.3	2.0
October	2.3	2.0	3.0	3.2	2.5	2.4	1.8
November	2.3	2.0	2.8	3.2	2.5	2.4	1.7
December	2.4	2.1	3.0	3.7	2.7	2.2	1.9

276
277

278 **Table 3. Ratio of excess deaths in 2020 to average mortality incidence in previous years**
279 **for COVID-19.**

	Overall	Women	Men	18-69	70-79	80-89	90
Mortality rates							
January	1.01	1.05	0.93	1.04	1.05	0.99	0.99
February	0.99	0.96	1.05	0.98	0.95	0.99	1.00
March	1.05	1.01	1.10	1.14	0.99	1.07	1.02
April	1.74	1.71	1.78	1.99	1.67	1.73	1.71
May	1.48	1.44	1.55	2.02	1.61	1.49	1.35
June	1.11	1.08	1.16	1.36	1.20	1.11	1.05
July	1.04	1.05	1.01	1.09	0.98	1.01	1.06
August	1.05	1.03	1.08	1.05	1.03	1.01	1.06
September	1.07	1.09	1.04	1.11	1.05	1.09	1.05
October	1.11	1.12	1.09	1.09	0.95	1.13	1.13
November	1.24	1.20	1.30	0.85	1.20	1.30	1.23
December	1.31	1.30	1.34	1.29	1.31	1.31	1.31

280 Estimates are calculated with the following formula: (incidence rate in 2020)/(average incidence
281 rate in 2010-2019).
282
283
284

Figure 1. Nursing home residents (A) and monthly incidence rates of mortality, hospitalization, and emergency department visits per 100 nursing home residents (panels B-D), by sex (panels E-G) and by age (panels H-J).