Title: Background rates of all-cause mortality, hospitalizations, and emergency department visits among nursing home residents in Ontario, Canada to inform COVID-19 vaccine safety assessments

Authors: Maria Sundaram[1,2], Sharifa Nasreen[2], Andrew Calzavara[1], Siyi He[1], Hannah Chung[1], Susan E. Bronskill[1,2,3], Sarah A. Buchan[1,2,4], Mina Tadrous[1,4], Peter Tanuseputro[1, 5], Kumanan Wilson[5], Sarah Wilson[1,2,4], Jeffrey C. Kwong[1,2,4,6,7], on behalf of the Canadian Immunization Research Network (CIRN) investigators

Affiliations:
[1] ICES, Toronto, ON
[2] Dalla Lana School of Public Health, University of Toronto, Toronto, ON
[3] Women's College Research Institute, Toronto, ON
[4] Public Health Ontario, ON
[5] Department of Medicine, University of Ottawa and Bruyere and Ottawa Hospital Research Institutes, Ottawa, ON
[6] Department of Family and Community Medicine, University of Toronto, Toronto, ON
[7] University Health Network, Toronto, ON

Corresponding author:
Jeff Kwong
Senior Scientist, ICES
G1 06, 2075 Bayview Avenue
Toronto, Ontario, Canada, M4N 3M5
jeff.kwong@utoronto.ca
Phone: (416) 480-4055 x1-7665
Fax: (416) 480-6048

Abstract word count: 284

Manuscript word count: 1,533

Abstract

Background. Nursing home (NH) residents have been disproportionately affected by the coronavirus disease 2019 (COVID-19) pandemic and are prioritized for vaccination. NH residents are generally at increased risk of poor outcomes due to advanced age, frailty, and complex health conditions. We report monthly incidence rates of deaths, hospitalizations, and emergency department (ED) visits during ten pre-pandemic years (2010-2019) and 2020 to provide context for assessments of COVID-19 vaccine safety in NH residents.

Methods. We observed deaths, hospitalizations, and ED visits among all Ontarians living in NHs using health administrative databases. Monthly incidence rates were calculated by month, by sex, and by age group. Direct comparisons between months were assessed using one-sample t-tests; direct comparisons by age and sex were assessed using chi-squared tests.

Results. From January 1, 2010 through December 31, 2019, there were, on average, 83,453 (SD: 652.4) Ontarians living in NHs in any given month, with an average of 2.3 (SD: 0.28) deaths, 3.1 (SD: 0.16) hospitalizations, and 3.6 (SD: 0.17) ED visits per 100 residents per month. Mortality rates were higher for men ($p<0.001$) and residents aged ≥80 years ($p<0.001$). Hospitalization and ED visit rates were higher for men but were lower for residents aged ≥80 years. From January to October 2020, the number of NH residents declined markedly. Mortality rates were increased in 2020 compared to 2010-2019, but hospitalization and ED visit rates were reduced in 2020 compared to 2010-2019 ($p<0.001$).

Conclusion. We identified relatively consistent monthly mortality, hospitalization, and ED visit rates during ten pre-pandemic years. Marked differences in these rates were observed during 2020, coinciding with heightened COID-19 infection rates and restrictions. These results provide context to the assessment of COVID-19 vaccine safety outcomes in this high-risk population.

Introduction

Nursing home (NH) residents have been disproportionately affected by the coronavirus disease 2019 (COVID-19) pandemic, comprising a substantial proportion of total COVID-19 deaths in many countries, including over 80% of COVID-19 deaths in Canada during the first epidemic wave.[1,2] The vaccine rollout has prioritized, among others, NH residents because of their high risk of poor outcomes from COVID-19.[3]

However, NH residents are generally frail and are at comparatively higher risk of poor outcomes, even outside the context of a global pandemic.[4,5] As a result, deaths and healthcare use within the expected range of event numbers for NH residents could create an erroneous impression that vaccine rollout is associated with increased risk of such events.[6] There is a lack of data on poor outcome rates among NH residents in Canada to facilitate comparisons of observed versus expected outcome rates during vaccination rollout. Therefore, we sought to quantify background rates of all-cause mortality, hospitalizations, and emergency department (ED) visits among NH residents in Ontario, the most populous province in Canada. Since COVID-19 and related restrictions may also impact these outcomes, we also report these rates during 2020.

Methods

We conducted an observational analysis using population-level health administrative databases from Ontario, Canada. These datasets were linked using unique encoded identifiers and analyzed at ICES. Ontario has a single-payer health system that provides universal access to physician and hospital services for all Ontario residents.

Cohort definitions

We identified Ontarians aged 18 years or older living in NHs using the physician billing database Ontario Health Insurance Plan, Ontario Drug Benefit, and the Continuing Care Reporting System databases. Information on age, sex and death were collected from the Registered Persons Database. We identified individual cohorts beginning at the start of each month from January 1, 2010 to December 31, 2019 inclusive, and from January 1 2020 to the most recent available data: October 31, 2020 for mortality or September 30, 2020 for hospitalizations and ED visits. Individuals were included in these cohorts if they were alive at the beginning of a month and recorded as living in a NH at least one day in that month or in the three previous months.

Statistical analysis

We calculated monthly rates per 100 NH residents overall, by sex, and by age group (18-69 years, 70-79 years, 80-89 years, and ≥90 years). We also conducted analyses comparing adults aged ≥80 years with adults younger than 80 years. Estimates from 2010-2019 were combined over years by month, and by sex and age group separately. Direct comparisons were conducted either with one-sample t-tests, in the case of comparisons by month from 2010-2019 vs. 2020, or with chi-squared tests, in the case of comparisons by sex or age. We also calculated the relative increase in mortality in 2020 likely due to COVID-19, compared with mortality in 2010-2019, using the following formula:

$$\frac{monthly\ mortality\ incidence\ in\ 2020 - average\ monthly\ mortality\ incidence\ in\ 2010 - 2019}{average\ monthly\ mortality\ incidence\ in\ 2010 - 2019}$$

Results

From January 1, 2010 through December 31, 2019, there were an average of 83,453 (SD: 652.4) NH residents in Ontario in any given month (**Figure 1A**). Of these, an average of 57,104 (SD: 291.6; ~68% of the total in any given month) were women, and an average of 58,733 (SD: 550.6; ~70% of the total in any given month) were aged ≥80 years.

During this period, approximately 2.3 (SD: 0.28) deaths per 100 NH residents occurred per month (**Figure 1B; Supplemental Table 1**). These rates were higher for men (incidence rate [IR]: 2.6; SD: 0.31) than women (IR: 2.2; SD: 0.27) (p<0.001; **Figure 2A**), and for residents aged ≥80 years (IR: 5.6; SD: 0.68) than residents aged <80 years (IR: 2.6; SD: 0.33) (p<0.001; **Figure 2D**). An average of 0.4 (SD: 0.04) more monthly deaths per 100 residents occurred during winter months (December, January, February, and March) than during other months during 2010-2019 (p<0.001; **Supplemental Table 1**).

We observed monthly averages of 3.1 (SD: 0.16) hospitalizations per 100 residents and 3.6 (SD: 0.17) ED visits per 100 residents during 2010-2019 (**Figures 1C and 1D; Supplemental Table 1**). Hospitalization rates were higher for men (IR: 3.8; SD: 0.20) than women (IR: 2.7; SD: 0.15) (p<0.001) (**Figure 2B; Supplemental Table 1**) and were lower for older residents (IR: 5.8; SD: 0.34) than younger residents (IR: 6.8; SD: 0.35) (p<0.001; **Figure 2E**). Similarly, ED visit rates were higher for men (IR: 4.3; SD: 0.21) than women (IR: 3.3; SD: 0.18) (p<0.001; **Figure 2C**) and lower for older residents (IR: 6.7; SD: 0.33) than younger residents (IR: 8.6; SD: 0.45) (p<0.001; **Figure 2F**).

We observed a steady decline in the total number of NH residents during 2020 (non-parametric test for trend p<0.01; **Figure 1A**). Mortality rates rose during the early pandemic period, and were higher in March, April, May, June, July, and September of the pandemic period compared to corresponding months in 2010-2019 (p<0.05 for each month) (**Figure 1B, Supplemental Table 2**); these differences were most pronounced in April and May 2020. Mortality was higher for men (**Figure 2A**) and older residents (**Figure 2D**). The ratio of excess deaths in 2020 to average mortality incidence in 2010-2019 was highest in April and May 2020, across sexes and age groups (**Supplemental Table 3**).

There was a substantial reduction in hospitalizations during 2020 compared to previous years (p<0.001 for each month during the pandemic period; **Figure 1C** and **Supplemental Table 2**). Hospitalization rates were higher for men than women (p<0.001; **Figure 2B**) and lower with increasing age throughout the pandemic period (**Figure 2E**). Similar to hospitalizations, there was a substantial reduction in ED visits during 2020 compared to previous years (p<0.001 for each month during 2020; **Figure 1D** and **Supplemental Table 2**). Patterns by sex and age were similar as for hospitalizations (**Figures 2C** and **2F**).

Discussion

This report presents rates of all-cause mortality, hospitalizations, and ED visits in NH residents in Ontario, Canada during a 10-year pre-pandemic period (2010-2019) and during January-October 2020. Mortality rates in 2010-2019 are similar to rates reported by others in Canada,[4] with rates being higher in men compared to women, and higher with increasing age. As expected, mortality rates spiked during the early phase of the COVID-19 pandemic, with high mortality rates among all age groups and sexes in April and May of 2020, similar to overall estimates for the province of Ontario.[7] However, mortality rates returned to values closer to pre-pandemic rates from August to October 2020.

The decrease in mortality rates is likely multifactorial. First, changes in provincial-level testing guidelines may have enabled earlier identification of COVID-19 cases. On April 8, 2020, Ontario officially expanded its testing guidelines beyond symptomatic NH residents to all residents, including all newly admitted individuals and asymptomatic residents who had contact with a confirmed-positive case.[8] Earlier identification of COVID-19 cases may have contributed to a subsequent reduction in mortality by preventing potential outbreaks and by providing needed care to residents earlier in the course of their illness. Second, enhanced infection prevention measures were implemented in NHs province-wide.[9] Third, reduced mortality rates in the summer months likely reflect a reduction in community-level transmission of SARS-CoV-2 experienced throughout Ontario.[7]

Our analysis additionally identified a decline in hospitalizations and ED visits during 2020, compared to 2010-2019. This finding may initially seem counterintuitive but, taken in context with simultaneous increases in NH setting mortality and decreases in the total number of NH residents, these reductions likely indicate a reduction in hospitalized deaths, which has been reported previously.[10] There may have also been an increased unwillingness to seek hospital-based care for non-COVID-19-related issues during the beginning of the first pandemic wave; future analyses should also consider the reasons noted for ED visits and hospitalizations to identify COVID-19-related vs. non-COVID-19-related incidents.

This analysis uses large-scale health administrative datasets, which are not yet currently updated to the end of December 2020 with vital statistics. As a result, we are not yet able to report more recent data. This is an important limitation of this work. If feasible, we will update this report as more data become available.

We intend for these findings to be used to contextualize potential increases in adverse events of special interest (AESI)[11] during the COVID-19 vaccine rollout in NH residents. Whereas mortality, hospitalization, and ED visit rates were highly stable over 10 pre-pandemic years, the marked changes in rates observed during 2020 indicate that caution is needed in determining the appropriate baseline to use when interpreting rates of AESIs following the COVID-19 vaccine rollout. Existing investigations on mortality in elderly and frail individuals who received COVID-19 vaccines has identified that mortality rates were consistent with high background rates of death in this population;[12] our data will help contextualize Canada- and Ontario-specific rates in a similar way.

Conclusions

This report identified relatively consistent monthly mortality, hospitalization, and ED visit rates for Ontario NT residents over a ten-year pre-pandemic period. During 2020, we identified a substantial increase in mortality rates during April and May, followed by a return to baseline rates for the remainder of 2020. In contrast, hospitalization and ED visit rates declined markedly in 2020 and (as of the latest data available) have not yet returned to baseline levels. These results provide important context for future analyses to identify potential AESI in NH residents.

Acknowledgments

This work was supported by the Canadian Immunization Research Network (CIRN) through a grant from the Public Health Agency of Canada and the Canadian Institutes of Health Research (CNF 151944). This study was also supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health (MOH). No endorsement by ICES or MOH is intended or should be inferred.

Conflict of interest

The authors declare no conflicts of interest.

Authors' contributions

JCK conceived of the study design and oversaw the study. AC obtained the data and calculated incidence rates. MS conducted additional statistical analysis and drafted the manuscript. PT, SEB, KW, and MT provided methodological input. All authors interpreted the results, critically revised the manuscript, and have approved the final version for publication.

References

1. *Pandemic experience in the long-term care sector: how does Canada compare with other countries?* Ottawa, ON: Canadian Institute for Health Information;2020.
2. Sepulveda ER, Stall NM, Sinha SK. A comparison of COVID-19 mortality rates among long-term care residents in 12 OECD countries. *JAMDA.* 2020;21(11):1572-1574 e1571. PMC7522741.
3. *WHO SAGE roadmap for prioritizing uses of COVID-19 vaccines in the context of limited supply.* Geneva, Switzerland: World Health Organization;2020.
4. Tanuseputro P, Chalifoux M, Bennett C, Gruneir A, Bronskill SE, Walker P, Manuel D. Hospitalization and Mortality Rates in Long-Term Care Facilities: Does For-Profit Status Matter? *J Am Med Dir Assoc.* 2015;16(10):874-883.
5. Gruneir A, Bell CM, Bronskill SE, Schull M, Anderson GM, Rochon PA. Frequency and pattern of emergency department visits by long-term care residents--a population-based study. *J Am Geriatr Soc.* 2010;58(3):510-517.
6. Branswell H. CDC advisory panel's lone dissenter on why long-term care residents shouldn't receive COVID-19 vaccine first. *STATNews*2020.
7. All Ontario: Case numbers and spread. 2021; https://covid-19.ontario.ca/data. Accessed February 5, 2021.
8. *COVID-19 Provincial Testing Guidance Update: April 8, 2020.* Ontario: Ministry of Health;2020.
9. COVID-19 Long-Term Care Resources. 2021; https://www.publichealthontario.ca/en/diseases-and-conditions/infectious-diseases/respiratory-diseases/novel-coronavirus/long-term-care-resources. Accessed March 10, 2021.
10. Brown KA, Daneman N, Buchan SA, Chan AK, Stall NM. Temporal variations in the intensity of care provided to community and nursing home residents who died of COVID-19 in Ontario, Canada. *medRxiv.* 2020.
11. Law B, Sturkenboom M. *D2.3: Priority List of Adverse Events of Special Interest: COVID-19.* CEPI; May 25, 2020 2020.
12. GACVS COVID-19 Vaccine Safety subcommittee meeting to review reports of deaths of very frail elderly individuals vaccinated with Pfizer BioNTech COVID-19 vaccine, BNT162b2 [press release]. Geneva, Switzerland: World Health Organization, 22 January 2021.

Figure 1. Total nursing home residents and monthly incidence rates of mortality, hospitalization, and emergency department visits per 100 nursing home residents.

Figure 2. Monthly incidence rates of mortality, hospitalization, and emergency department visits per 100 nursing home residents, by sex (panels A-C) and by age (panels D-F).