Estimating the increased transmissibility of the B.1.1.7 strain over previously circulating strains in England using fractions of GISAID sequences and the distribution of serial intervals

Chayada Piantham[1], Kimihito Ito[2*],

[1]*Graduate School of Infectious Diseases, Hokkaido University; chayada@czc.hokudai.ac.jp*
[2]*Research Center for Zoonosis Control, Hokkaido University; itok@czc.hokudai.ac.jp*

Abstract

The B.1.1.7 strain, a variant strain of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is thought to have higher transmissibility than previously circulating strains in England. The fraction of the B.1.1.7 strain among SARS-CoV-2 viruses in England have grown rapidly. In this paper, we propose a method to estimate the selective advantage of a mutant strain over previously circulating strains using the time course of the fraction of B.1.1.7 strains. Our approach is based on the Maynard Smith's model of allele frequencies in adaptive evolution, which assumes that the selective advantage of a mutant strain over previously circulating strains is constant over time. Applying this method to the sequence data in England using serial intervals of COVID-19, we found that the transmissibility of the B.1.1.7 strain is 40% (with a 95% confidence interval (CI) from 40% to 41%) higher than previously circulating strains in England. The date of the emergence of B.1.1.7 strains in England was estimated to be September 20, 2020 with its 95% CI from September 11 to September 20, 2020. The result indicated that the control measure against the B.1.1.7 strain needs to be strengthened by 40% from that against previously circulating strains. To get the same control effect, contact rates between individuals need to be restricted to 0.71 of the contact rates that have been achieved form the control measure taken for previously circulating strains.

Keywords: COVID-19, B.1.1.7, selective advantage, adaptive evolution, serial interval, GISAID, England, SARS-CoV-2

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of COVID-19, has been rapidly evolving since its emergence in 2019. In December 2020, the Public Health England detected a new cluster of SARS-CoV-2 viruses phylogenetically distinct from the other strains circulating in the United Kingdom (Chand et al., 2020). These viruses belong to the lineage B.1.1.7 according to the PANGO nomenclature (Rambaut, Loman, et al., 2020) and the World Health Organization (WHO) designated them as Variant of Concern, year 2020, month 12, variant 01 (VOC 202012/01) (World Health Organization, 2020).

The B.1.1.7 strain was first detected in England in September 2020, and the numbers of infections with this strain increased in October and November in 2020 (Chand et al., 2020). By February 2021, B.1.1.7 strain occupied 95% of the SARS-CoV-2 circulating in England (Davies et al., 2021).

Several studies have compared the transmissibility of the B.1.1.7 strain to that of previously circulating strains. Davies et al. estimated that the reproduction number of the B.1.1.7 strain is 43–90% (with a 95% credible interval of 38–130%) higher than preexisting strains using data from England (Davies et al., 2021). However, different models resulted in different ranges of estimates in their multiplicative increase in reproduction number (R). Grabowski et al. estimated a 83–118% increase with a confidence interval of 71–140% compared to previously circulating strains in England (Grabowski et al., 2021). Volz et al. estimated 40–75% increase in R using data from England (Volz et al., 2021). Their methods use linear regression of log odds ratio between B.1.1.7 strain and previously circulating strains and estimate increase of transmissibility under assumption that R is constant over time during the target period of analysis. Washington et al. estimated a 35–45% increase using data from the United States of America using Volz's method (Washington et al., 2021). Chen et al. estimated 49–65% increase in reproduction number using data from Switzerland also under the constant R assumption (Chen et al., 2021). Due to the high transmissibility the B.1.1.7 strain, strong control measure such as lockdown was taken when the strain was introduced (Davies et al., 2021). Thus, the constant R assumption is questionable when analyzing the increase in the reproduction number of B.1.1.7 compared to that of previously circulating strains.

In this paper, we propose a method to estimate the selective advantage of a mutant strain over previously circulating strains. Our approach is based on the Maynard Smith's model of allele frequencies in adaptive evolution, which assumes that the selective advantage of a mutant strain over previously circulating strains is constant over time (Maynard Smith & Haigh, 1974). Appling the developed method to the sequence data in England using the serial interval distribution of COVID-19 estimated by Nishiura et al. (Nishiura et al., 2020), we estimate the increase in the instantaneous reproduction number of B.1.1.7 strains compare to that of previously circulating strains. Based on the estimate, we discuss its implication to control measures for COVID-19.

Materials and Methods

Sequence data

Nucleotide sequences of SARS-CoV-2 viruses was downloaded from GISAID EpiCoV database (Shu & McCauley, 2017) on March 1, 2021. Nucleotide sequences that determined from viruses detected in England were selected and aligned to the reference amino acid sequence of S protein of SARS-CoV-2 virus (YP_009724390) using DIAMOND (Buchfink et al., 2015). The aligned nucleotide sequences were translated into amino acid sequences, then were aligned with the reference amino acid sequence of S protein using MAFFT (Katoh et al., 2002). Amino acid sequences having either an ambiguous amino acid or more than ten gaps were excluded from the rest of analyses. Table 1 shows amino acids on S protein of characterizing B.1.1.7 strain, retrieved from the PANGO database (Rambaut, Loman, et al., 2020).

Table 1. Amino acids on S protein which are used to define B.1.1.7 strains

Position on S protein	Amino acid
69	Deletion
70	Deletion
144	Deletion
501	Tyrosine (Y)
570	Aspartic acid (D)
681	Histidine (H)
716	Isoleucine (I)
982	Alanine (A)
1118	Histidine (H)

We divided amino acid sequences into three groups based on amino acids shown in Table 1. The first group is sequences having all of B.1.1.7-defining amino acid substitutions in Table 1. We call a virus in this group a "B.1.1.7 strain". The second group are sequences which have none of the B.1.1.7-defining substitutions. We call a virus in the this group a "non-B.1.1.7 strain". The third group are sequences which have at least one but incomplete set of the B.1.1.7-defining amino acid substitutions. We called a strain in the third group a "B.1.1.7-like strain". Table 2 shows the number of sequences categorized into each group. We used the number of B.1.1.7 strains and non-B.1.1.7 strains for the rest of the analyses. B.1.1.7-like strains were excluded from the analyses, since we do not know whether they had the same transmissibility as B.1.1.7 strains or not. Figure 1 shows the daily numbers of GISAID sequences of B.1.1.7 strains, non-B.1.1.7 strains and B.1.1.7-like strains detected in England from September 1, 2020 to February 19, 2021. These numbers are provided in Supplementary Table 1.

Table 2. Number of GISAID sequences in England from September 1, 2020 to February 19, 2021

B.1.1.7 strains	Non-B.1.1.7	B.1.1.7-like	Total
71692	65840	2227	139759

Figure 1. Number of sequences in England from September 1, 2020 to February 19, 2021. The nucleotide sequences were retrieved from GISAID on March 1, 2020.

Serial interval distribution

The method we propose in this paper uses discrete distributions of serial intervals. Function $g(i)$ gives the probability that the onset day of a secondary infection is at i days after the onset day of its primary case. We obtained the values of $g(i)$ by discretizing the lognormal distribution of serial intervals of COVID-19 estimated by Nishiura et al. (Nishiura et al., 2020). Thus, the probability mass function of serial intervals is given by

$$g(i) = \int_i^{i+1} f(t)dt \,,$$

where $f(t)$ is the probability density function of a lognormal distribution with a mean of 4.7 days and a standard deviation of 2.9 days.

Model of Advantageous Selection

Consider we have a large population of viruses consisting of strains of two genotypes A and a, of which fraction in the viral population at a calendar date t are $q_A(t)$ and $q_a(t)$, respectively. Suppose also that genotype A is mutant of a that emerges at time t_0.

111 We assume that a virus of genotype A generates $1 + s$ times as many secondary transmissions as those of
112 genotype a. Then, s can be considered as the coefficient of selective advantage in adaptive evolution. As
113 described by Maynard Smith and Haigh (1974), the fraction of viruses of allele A after the n
114 transmissions, q_n, satisfies

$$q_{n+1} = \frac{(1 + s)q_n}{(1 + s)q_n + (1 - q_n)} = \frac{1 + s}{1 + sq_n} q_n. \tag{1}$$

116 Let $g(i)$ be a discrete distribution of serial intervals defined in the previous subsection. Since the
117 expected fraction of allele A in the population at calendar time t can be represented as s and $q_A(t - i)$ for
118 $0 \le i \le t$ with a probability of $g(i)$, the value of $q_A(t)$ can be represented as follows.

$$q_A(t) = \sum_{i=0}^{t} g(i) \frac{1 + s}{1 + sq_A(t - i)} q_A(t - i). \tag{2}$$

121 Assuming $g(0) = 0$, we can approximate the formula.

$$q_A(t) \approx \sum_{i=1}^{t} g(i) \frac{1 + s}{1 + sq_A(t - i)} q_A(t - i). \tag{3}$$

Likelihood Function

124 Let $n(t)$ be the number of sequences of either genotype A or a observed at calendar date t. Let $d_1, \dots, d_k$
125 be calendar dates such that $n(d_i) > 0$ for $1 \le j \le k$. Suppose that we have $n_A(d_j)$ samples of genotype A
126 at calendar date d_j. Since genotype A emerged at time t_0, $q_A(d_j) = 0$ and $q_a(d_j) = 1$ for $d_j < t_0$. Let
127 q_0 be initial frequency of genotype A, i.e., $q_0 = q_A(t_0)$. Then the following equation gives the likelihood
128 function of s, t_0, and q_0 for observing $n_A(d_j)$ samples of viruses of genotype A at calendar date d_j.

$$L\left(s, t_0, q_0 ; n_A(d_j)\right) = \binom{n(d_j)}{n_A(d_j)} q_A(d_j)^{n_A(d_j)} \left(1 - q_A(d_j)\right)^{n(d_j) - n_A(d_j)}, \tag{4}$$

131 for $1 \le j \le k$. The likelihood function of s, t_0, and q_0 for observing $n_A(d_1), \dots, n_A(d_k)$ sequences of
132 genotype A at calendar dates $d_1, \dots, d_k$ is given by the following formula.

$$L\left(s, t_0, q_A(t_0) ; n_A(d_1), \dots, n_A(d_k)\right) = \prod_{j=1}^{k} L\left(s, t_0, q_A(t_0) ; n_A(d_j)\right) \tag{5}$$

Parameter estimation from sequence data

The B.1.1.7 strain was first detected in England on September 20, 2020. We assume that t_0 is this day or
someday before this day. Parameters s, t_0, and q_0 were estimated by maximizing likelihood of

observations on September 1, 2020 and later on. B.1.1.7 strains, viruses having complete subset of
B.1.1.7-difining substitutions on its S protein were considered as genotype A. The non-B.1.1.7 strains,
viruses having none of B.1.1.7-defining substitutions were considered to be genotype a. The B.1.1.7-liike
strains, viruses having an incomplete set of B.1.1.7 substitutions on the S protein, were excluded from the
analysis. Parameters of s, t_0, and q_0 were estimated by maximizing log likelihood defined in Equation (5).
The 95% confidence intervals of parameters were estimated by profile likelihood (Pawitan, 2013). The
optimization of likelihood function was done by the nloptr package in R (Johnson; Rowan, 1990).

Results

The selective advantage of B.1.1.7 strains over non-B.1.1.7 strains, s, was estimated to be 0.40 with its
95% confidence intervals from 0.40 to 0.41 (Table 2). The date of emergence of B.1.1.7 strains in
England, t_0, was estimated to be September 20, 2020 with its 95% confidence interval from September
11, 2020 to September 20, 2020. The initial fraction of B.1.1.7 among non-B.1.1.7 and B.1.1.7 strains at
the emergence in England, q_0, was estimated to be 0.0030 with its 95% confidence intervals from 0.0014
to 0.0031.
151
152   **Table 2. Maximum likelihood estimations of parameters**

Parameter	Estimate	95% CI
s	0.40	[0.40, 0.41]
t_0	September 20, 2020	[September 11, 2020, September 20, 2020]
q_0	0.0030	[0.0014, 0.0031]

153
Figure 2 shows the time course of the fraction of B.1.1.7 strains among all strains except B.1.1.7-like
strains detected in the England from September 1, 2020 to February 19, 2021. White circles indicate daily
fractions of B.1.1.7 strains among all strains except B.1.1.7-like strains. Solid line indicates the time
course of fraction of B.1.1.7 strains calculated using parameters estimated from the data. Dashed lines
indicate its lower and upper bound of its 95% CI.

Figure 2. Time course of the fraction of B.1.1.7 strains among all strains except B.1.1.7-like strains

detected in England from September 1, 2020 to February 19, 2021. White circles indicate the

fraction of B.1.1.7 strains among B.1.1.7 and non-B.1.1.7 strains. The nucleotide sequences were

retrieved from GISAID on March 1, 2021. Solid line indicates the time course of fraction of B.1.1.7

strains calculated using parameters estimated from the data. Dashed lines indicate its lower and

upper bound of its 95% CI.

Discussion

In this paper, the selective advantage of the B.1.1.7 strain in England over non-B.1.1.7 strains was

estimated to be 0.40 with a 95% CI from 0.40 to 0.41. The date of emergence of B.1.1.7 strains in

England was estimated to be September 20, 2020 with its 95% confidence interval from September 11,

2020 to September 20, 2020. The initial fraction of B.1.1.7 among all sequences except B.1.1.7-like

strains at the time of emergence in England was estimated to be 0.0030 with its 95% confidence intervals

from 0.0014 to 0.0031.

The estimated selective advantage of 0.40 indicates that the B.1.1.7 strain is 40% more transmissible than

that of previously circulating strains in England. This means that the control measures for the B.1.1.7

strain needs to be strengthened by 40% compared to that for the previously circulating strains. To get the

same control effect, contact rates between individuals needs to be restricted below $1/1.40 = 0.71$ of the

contact rates to achieve the same control measure for the previously circulating strains.

Our estimation method is based on the principle that the expected fraction of a mutant strain among all strains can be determined from those in the previous generation using the serial interval distribution of infections. The method assumes that the selective advantage of a mutant strain over previously circulating strains is constant over time, which is based on Maynard Smith's model of allele frequencies in adaptive evolution. The method is related to Fraser's method for estimating the instantaneous reproduction number (Fraser, 2007). However, it is our future work to mathematically clarify the relationship between the selective advantage estimated using our method and the instantaneous reproduction numbers of two strains.

The selective advantage estimated in this study relies on the serial interval distribution, and thus the results may change depending on serial interval distribution used in the analysis. In this paper, we used the serial interval distribution estimated by Nishiura et al. (Nishiura et al., 2020). The distribution is a lognormal distribution with a mean serial interval of 4.7 days with a standard deviation of 2.9 days. Several groups have estimated the serial intervals of SARS-CoV-2 using different datasets. Some variations are observed among these estimated values (Rai et al., 2021). Volz et al. (Volz et al., 2021) assumes a fixed serial interval of 6.5 days based on results by Bi et al. (Bi et al., 2020). However, Ali et al. have reported that the serial interval estimated using data from China during before January 22, 2020 was longer than estimates after January 22, 2020 (Ali et al., 2020). The serial interval estimated by Bi et al. contains data before January 22, 2020 and there might be some possibility that the estimated serial interval does not reflect the current situation. This is the reason why we did not use serial interval estimated by Bi et al.

As of March 17, 2021, the B.1.1.7 strain has now been observed in 93 countries (Rambaut, Holmes, et al., 2020). The selective advantage of the B.1.1.7 strains over previously circulating strains in other countries remains to be our future work. Variant strains originated in Brazil and South Africa also show higher transmissibility than previously circulating strains (World Health Organization, 2021). There is an urgent need to estimate the selective advantage of these strains over previously circulating strains. We hope that the developed methodology provides important information for countries in the world to establish control measures against variants strains such as the B.1.1.7 strain.

Acknowledgement

We gratefully acknowledge the laboratories responsible for obtaining the specimens and the laboratories where genetic sequence data were generated and shared via the GISAID Initiative, on which this research is based. This work was supported by CREST (grant number JPMJCR1413) from Japan Science and Technology Agency (http://www.jst.go.jp/), and the World-leading Innovative and Smart Education (WISE) Program (1801) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan (http://www.mext.go.jp/). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of Interest

We declare that there is no conflict of interest.

References

1. Ali, S. T., Wang, L., Lau, E. H. Y., Xu, X. K., Du, Z., Wu, Y., Leung, G. M., & Cowling, B. J. (2020). Serial interval of SARS-CoV-2 was shortened over time by nonpharmaceutical interventions. *Science, 369*(6507), 1106-1109. doi:10.1126/science.abc9004

2. Bi, Q., Wu, Y., Mei, S., Ye, C., Zou, X., Zhang, Z., Liu, X., Wei, L., Truelove, S. A., Zhang, T., Gao, W., Cheng, C., Tang, X., Wu, X., Wu, Y., Sun, B., Huang, S., Sun, Y., Zhang, J., Ma, T., Lessler, J., & Feng, T. (2020). Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis, 20*(8), 911-919. doi:10.1016/S1473-3099(20)30287-5

3. Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nat Methods, 12*(1), 59-60. doi:10.1038/nmeth.3176

4. Chand, M., Hopkins, S., Dabrera, G., Achison, C., Barclay, W., Ferguson, N., Volz, E., Loman, N., Rambaut, A., & Barrett, J. (2020). *Investigation of novel SARS-COV-2 variant. Variant of Concern 202012/01* (PHE gateway number: GW-1824). Retrieved from https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/959438/Technical_Briefing_VOC_SH_NJL2_SH2.pdf

5. Chen, C., Nadeau, S., Topolsky, I., Manceau, M., Huisman, J. S., Jablonski, K. P., Fuhrmann, L., Dreifuss, D., Jahn, K., Beckmann, C., Redondo, M., Kobel, O., Noppen, C., Risch, L., Risch, M., Wohlwend, N., Kas, S., Bodmer, T., Roloff, T., Stange, M., Egli, A., Eckerle, I., Denes, R., Feldkamp, M., Nissen, I., Santacroce, N., Burcklen, E., Aquino, C., de Gouvea, A. C., Moccia, M. D., Grüter, S., Sykes, T., Opitz, L., White, G., Neff, L., Popovic, D., Patrignani, A., Tracy, J., Schlapbach, R., Dermitzakis, E. T., Harshman, K., Xenarios, I., Pegeot, H., Cerutti, L., Penet, D., Blin, A., Elies, M., Althaus, C., Beisel, C., Beerenwinkel, N., Ackermann, M., & Stadler, T. (2021). Quantification of the spread of SARS-CoV-2 variant B.1.1.7 in Switzerland. *medRxiv*, 2021.2003.2005.21252520. doi:10.1101/2021.03.05.21252520

6. Davies, N. G., Abbott, S., Barnard, R. C., Jarvis, C. I., Kucharski, A. J., Munday, J. D., Pearson, C. A. B., Russell, T. W., Tully, D. C., Washburne, A. D., Wenseleers, T., Gimma, A., Waites, W., Wong, K. L. M., van Zandvoort, K., Silverman, J. D., Group, C. C.-W., Consortium, C.-G. U., Diaz-Ordaz, K., Keogh, R., Eggo, R. M., Funk, S., Jit, M., Atkins, K. E., & Edmunds, W. J. (2021). Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*. doi:10.1126/science.abg3055

7. Fraser, C. (2007). Estimating individual and household reproduction numbers in an emerging epidemic. *PLoS One, 2*(8), e758. doi:10.1371/journal.pone.0000758

8. Grabowski, F., Preibisch, G., Giziński, S., Kochańczyk, M., & Lipniacki, T. (2021). SARS-CoV-2 Variant of Concern 202012/01 Has about Twofold Replicative Advantage and Acquires Concerning Mutations. *Viruses, 13*(3), 392. Retrieved from https://www.mdpi.com/1999-4915/13/3/392

9. Johnson, S. G. The NLopt nonlinear-optimization package. Retrieved from http://ab-
initio.mit.edu/nlopt
10. Katoh, K., Misawa, K., Kuma, K., & Miyata, T. (2002). MAFFT: a novel method for
rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res,*
30(14), 3059-3066. doi:10.1093/nar/gkf436
11. Maynard Smith, J., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene.
Genet Res, 23(1), 23-35. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/4407212
12. Nishiura, H., Linton, N. M., & Akhmetzhanov, A. R. (2020). Serial interval of novel
coronavirus (COVID-19) infections. *Int J Infect Dis, 93*, 284-286.
doi:10.1016/j.ijid.2020.02.060
13. Pawitan, Y. (2013). *In All Likelihood: Statistical Modelling and Inference Using*
Likelihood. Croydon: Oxford University Press.
14. Rai, B., Shukla, A., & Dwivedi, L. K. (2021). Estimates of serial interval for COVID-19:
A systematic review and meta-analysis. *Clin Epidemiol Glob Health, 9*, 157-161.
doi:10.1016/j.cegh.2020.08.007
15. Rambaut, A., Holmes, E. C., O'Toole, A., Hill, V., McCrone, J. T., Ruis, C., du Plessis,
L., & Pybus, O. G. (2020). A dynamic nomenclature proposal for SARS-CoV-2 lineages
to assist genomic epidemiology. *Nat Microbiol, 5*(11), 1403-1407. doi:10.1038/s41564-
020-0770-5
16. Rambaut, A., Loman, N., Pybus, O., Barclay, W., Barrett, J., Carabelli, A., Connor, T.,
Peacock, T., Robertson, D. L., & Volz, E. (2020, 5 February 2021). Preliminary genomic
characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of
spike mutations. Retrieved from https://virological.org/t/preliminary-genomic-
characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-
spike-mutations/563
17. Rowan, T. (1990). *Functional Stability Analysis of Numerical Algorithms.* (Ph.D. Thesis).
University of Texas, Austin.
18. Shu, Y., & McCauley, J. (2017). GISAID: Global initiative on sharing all influenza data -
from vision to reality. *Euro Surveill, 22*(13). doi:10.2807/1560-
7917.ES.2017.22.13.30494
19. Volz, E., Mishra, S., Chand, M., Barrett, J. C., Johnson, R., Geidelberg, L., Hinsley, W.
R., Laydon, D. J., Dabrera, G., O'Toole, Á., Amato, R., Ragonnet-Cronin, M., Harrison,
I., Jackson, B., Ariani, C. V., Boyd, O., Loman, N. J., McCrone, J. T., Gonçalves, S.,
Jorgensen, D., Myers, R., Hill, V., Jackson, D. K., Gaythorpe, K., Groves, N., Sillitoe, J.,
Kwiatkowski, D. P., Flaxman, S., Ratmann, O., Bhatt, S., Hopkins, S., Gandy, A.,
Rambaut, A., & Ferguson, N. M. (2021). Transmission of SARS-CoV-2 Lineage B.1.1.7
in England: Insights from linking epidemiological and genetic data. *medRxiv*,
2020.2012.2030.20249034. doi:10.1101/2020.12.30.20249034
20. Washington, N. L., Gangavarapu, K., Zeller, M., Bolze, A., Cirulli, E. T., Schiabor
Barrett, K. M., Larsen, B. B., Anderson, C., White, S., Cassens, T., Jacobs, S., Levan, G.,
Nguyen, J., Ramirez, J. M., Rivera-Garcia, C., Sandoval, E., Wang, X., Wong, D.,
Spencer, E., Robles-Sikisaka, R., Kurzban, E., Hughes, L. D., Deng, X., Wang, C.,
Servellita, V., Valentine, H., De Hoff, P., Seaver, P., Sathe, S., Gietzen, K., Sickler, B.,

Antico, J., Hoon, K., Liu, J., Harding, A., Bakhtar, O., Basler, T., Austin, B., Isaksson,
M., Febbo, P., Becker, D., Laurent, M., McDonald, E., Yeo, G. W., Knight, R., Laurent,
L. C., de Feo, E., Worobey, M., Chiu, C., Suchard, M. A., Lu, J. T., Lee, W., &
Andersen, K. G. (2021). Genomic epidemiology identifies emergence and rapid
transmission of SARS-CoV-2 B.1.1.7 in the United States. *medRxiv.*
doi:10.1101/2021.02.06.21251159

21. World Health Organization. (2020). SARS-CoV-2 Variants. *Disease Outbreak News.*
    Retrieved from https://www.who.int/csr/don/31-december-2020-sars-cov2-variants/en/

22. World Health Organization. (2021). *COVID-19 Weekly Epidemiological Update.*
    Retrieved from https://www.who.int/publications/m/item/weekly-epidemiological-
    update---9-february-2021