Modelling the impact of rapid tests, tracing and distancing in lower-income countries suggest that optimal policies vary with rural-urban settings

Xilin Jiang [1,2,3], Wenfeng Gong [4 *], Zlatina Dobreva[5], Ya Gao [6], Matthew Quaife[5], Christophe Fraser [1], Chris Holmes[1,2,7 *]

[1] Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford
[2] Department of Statistics, University of Oxford
[3] Wellcome Centre for Human Genetics, University of Oxford
[4] Bill & Melinda Gates Foundation, Seattle WA, USA
[5] Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, UK
[6] Department of International Health, Johns Hopkins University
[7] The Alan Turing Institute, London, UK

*Corresponding authors:

Wenfeng.Gong@gatesfoundation.org

Address: Unit 1901, Tower B, Ping An International Financial Center, No.1-3 Xinyuan South Road, Chaoyang District , Beijing 100027, PRC

cholmes@stats.ox.ac.uk
Address: Department of Statistics, University of Oxford, 24–29 St Giles', Oxford OX1 3LB, UK

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Bill and Melinda Gates Foundation (BMGF).

Financial & competing interests disclosure:
Wenfeng Gong is an employee at the Bill and Melinda Gates Foundation. Xilin Jiang and Ya Gao served as consultants to the BMGF. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

Abstract

Low- and middle-income countries (LMICs) remain of high potential for hotspots for COVID-19 deaths and emerging variants given the inequality of vaccine distribution and their vulnerable healthcare systems. We aim to evaluate containment strategies that are sustainable and effective for LMICs. We constructed synthetic populations with varying contact and household structures to capture LMIC demographic characteristics that vary across communities. Using an agent-based model, we explored the optimal containment strategies for rural and urban communities by designing and simulating setting-specific strategies that deploy rapid diagnostic tests, symptom screening, contact tracing and physical distancing. In low-density rural communities, we found implementing either high quality (sensitivity > 50%) antigen rapid diagnostic tests or moderate physical distancing could contain the transmission. In urban communities, we demonstrated that both physical distancing and case finding are essential for containing COVID-19 (average infection rate < 10%). In high density communities that resemble slums and squatter settlements, physical distancing is less effective compared to rural and urban communities. Lastly, we demonstrated contact tracing is essential for effective containment. Our findings suggested that rapid diagnostic tests could be prioritised for control and monitor COVID-19 transmission and highlighted that contact survey data could guide strategy design to save resources for LMICs. An accompanying open source R package is available for simulating COVID-19 transmission based on contact network models.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes the COVID-19, has rapidly spread worldwide since the end of 2019. [1] Inadequate vaccine distribution and insufficient healthcare resources have contributed to many low- and middle- income countries (LMICs) suffering more from the pandemic than affluent countries. [2] Many LMICs rely on testing, tracing and physical distancing to control and monitor COVID-19, [3,4] such as test, travel restriction, and testing and tracing programs, given the inequitable distribution of vaccines and medication. While contributing to COVID-19 control and surveillance, these non-pharmaceutical interventions (NPIs) may cause considerable damage to the economies as people put their life and work on hold. [5] Therefore, identifying effective NPIs for LMICs could prepare the government for new variants and peaks while easing the burden on the economy and society. [4]

Emerging immunity-escaping variants from LMICs alert us to the importance of consistent surveillance and containment of COVID-19. [6,7] Affordable and effective testing methods are crucial to reduce the economic burden of LMICs. Many COVID-19 containment policies rely on the real-time reverse transcription-polymerase chain reaction (RT-PCR), which can be challenging to implement in LMIC because of the limited clinical resources [8,9] and laboratory capacities. [10,11] The financial and lab personnel cost of maintaining large scale PCR testing would put a heavy burden on the vulnerable healthcare systems of LMICs. Therefore, it is essential to leverage alternative testing methods, such as rapid diagnostic tests (RDT) based on antigen detection, [12–14] which is cheaper and less resource-demanding. Additionally, community healthcare workers could perform contact tracing and quarantine to further mitigate the SARS-CoV-2 transmission, which has been implemented successfully during HIV and Tuberculosis outbreaks. [15] Combining RDT, contact tracing and quarantine, we can increase the testing capacity, ensure early and accurate detection of infected cases, contain disease transmission, and reduce the speed of viral mutation accumulation.

In this study, we evaluated the dynamics of COVID-19 transmission in younger populations that are typical of LMICs and the effectiveness of testing, tracing and distancing. We constructed synthetic contact networks using contact numbers and demographic structures in lower-income

communities. To capture the different infection rates across geographical locations, [16,17], we

simulated SARS-CoV-2 transmission in various contact structures and captured the uncertainty

of the epidemic sizes using bootstrap samples of contact networks. We evaluated the realistic

combined strategy involving test and tracing using PCR and RDT, advising patients to self

isolate, and physical distancing. Our simulation results suggest the level of containment required

to contain COVID-19 depends on contact frequency in the communities. We also showed that

antigen RDT and symptom-based diagnosis could be deployed in several settings for better

containment outcomes. To help researchers replicate results with demographics that are not

included in our analysis and to update the results using properties of new variants, we released an

open-source R package for contact network based transmission simulation which accommodates

customised parameter settings.

Material and methods

Processing demographic data

We cited the age distribution and household sizes from the United nation. [18,19] (Supplementary

Table 1) We considered three age groups (0-14, 15-25 and 25+ years old) for simplicity and to

capture the susceptibility of the younger population and contact structure differences between

younger and older groups. We cited the contact survey data from a study in Uganda to compute

age-specific contact matrices and contact distribution within households.[20] To compare

geographies that have different contact numbers, we simulated contact networks that correspond

to LMICs using PERC survey data from Africa CDC. [21] When using the contact matrices in

simulation, we scaled the contact matrix to match the average daily contact number under each

setting. (Table 1)

Agent-based modelling

We used an agent-based model and configured the communities based on the age distributions

and household sizes cited from a Uganda study. [20] We simulated populations using contact

numbers that span a range of African countries. Additionally, we abstracted three representative

community settings that have distinct demographics: (1) a rural community which represent

geographies with low contact rates and medium household sizes; (2) an urban community which

represent geographies with high contact rates and medium household sizes; (3) a high density

community which represent high contact number and large household sizes, which aim to

capture communities such as slums and squatter settlements. (Table 1) For each setting, we

simulated 20 synthetic populations, where each household was composed of at least one adult

above age 25, and each household was assigned to a location on a two-dimensional plane. The

contact networks for the synthetic population were configured for the corresponding household

structure, age-specific contact matrices, and geographical clustering, using an Exponential

Random Graph Model (ERGM). [22] See Supplemental Methods for details on the ERGM.

After constructing the synthetic populations with contact and household structure, we randomly

choose one of the 20 synthetic populations and simulated COVID-19 transmission initiated by

importing 2 cases. At each time step of a simulated outbreak, one realised contact network of

ERGM was sampled to represent the daily dynamic of contacts. (Figure 1) Infection events were

sampled among the contacts proportional to the transmissibility multiplied by generation time,

which is a Weibull distribution that has a mode on the day of onset. [23] The day of onset for each

infected individual was sampled from the cited incubation time distribution [24]. The susceptibility

of the younger group (0-14 years old) was set to 50% of that of the older group (15+ years old).

[25,26] The death rate is set as 1% for the older group (25+ years old) and 0.2% for the younger

group (0-25 years old). [27] Using the age-dependent susceptibility and basic reproduction number,

we computed the next generation matrix and the transmissibility. (Supplementary Methods) At

each infection event, we sampled whether an infected individual is asymptomatic using a

Bernoulli distribution of probability equal to 30%. (Table 1) For detailed parameterization of

transmission dynamics, see Supplementary Methods.

We tested if our model reflects the cited contact data and transmission dynamics by comparing

the simulated contact distribution and generation time with the empirical data. (Figure 2) The age

mixing contact matrix generated from ERGM is also compared to the survey data.

(Supplementary Figure 1) We performed simulations that covered the range of contact numbers

collected from Africa CDC. (Figure 3) We cautioned these contact numbers are sampled from a

small proportion of the populations and should not be considered as representing the contact

number for each entire nation.

NPI simulation

We evaluated the impact of testing alone and in combination with other NPIs, which include

contact tracing, quarantine, and physical distancing, for LMICs. (Figure 1B) The testing methods

considered were PCR and antigen RDT. We simulated that only individuals with symptoms and

agree to be tested will receive testing. In our simulation, an infected individual with symptoms

will be discovered at a rate which is the product of four parameters: healthcare seeking rate, test

consent rate, sampling success rate and lab sensitivity of the test. Values and sources of testing

parameters were summarised in Table 2.

Our simulation assumes a test is always performed with close contact tracing and quarantine,

unless specified otherwise. We assumed only close contacts of confirmed cases who consent to

be isolated will be traced. Therefore, the overall quarantine rate is determined by the product of

three parameters: isolation consent rate of the infector, tracing success rate (we assumed a

tracing success rate equal to 100% for members within the same household of the infector), and

quarantine consent rate of the contacts. Physical distancing was modelled by reducing the

number of contacts outside the households. We simulated the effect of physical distancing by

reducing the non-household contact number by 20% to 80% with 20% increments. We

approximate the effective reproduction number Re under physical distancing by scaling Re

proportionally to the average daily contact numbers.

In addition to individual containment measures, we also evaluated combined strategies based on

current practice of COVID-19 testing. The combined strategies evaluated included 1) symptom-

based isolation + PCR testing, 2) symptom-based isolation + antigen RDT, 3) and symptom-

based isolation + PCR testing + antigen RDT. (Supplementary Table 3) In practice, it does not

introduce extra cost to suggest individuals who show up at the test centre with symptoms to self

isolate. Therefore we assumed all three containment strategies are accompanied by symptom

based isolation, which has a low compliance rate (20%). All three strategies were replicated with

and without physical distancing for 200 outbreak trajectories to estimate the average daily

incidence number. We simulated each trajectory for 100 days to guarantee that the transmission

extincts within the synthetic population.

Sensitivity analysis

We performed sensitivity analyses for a list of parameters that could vary (Supplementary Table

4) and used regression to evaluate the impact of changing these parameters on the epidemic

sizes. For the baseline conditions, we simulated outbreaks over grids of values for transmission

and demographic variables that are assumed or could not be determined from literature. We

computed R^2 using the total infected number as the dependent variable to capture the proportion

of variance explained by each variable. (Table 1). For the intervening conditions, we performed

the same simulations over grids of values for testing and NPI variables. We computed the R^2

using infected numbers under three different intervention strategies as the dependent variables

and testing/NPI variables in Table 2 as independent variables. In total we have 2,028 simulation

settings (3 community settings, 4 containing strategies and 169 values for the variables) which

are listed in Supplementary table 5. For each simulation setting, we simulated 200 simulated

trajectories of 100 days to obtain the bootstrap confidence interval (BCI) of the estimated

infection numbers.

The code generated during this study is bundled up, with an R-package available at:

https://github.com/Xilin-Jiang/NetworkCOVID19. Please refer to the GitHub page for installing

the package and setting parameters for replicating our results or performing other contact

network based analysis.

Role of the funding source

Results

Effectiveness of testing and physical distancing

To investigate the effectiveness of testing and physical distancing under different contact

numbers based on survey data from African CDC. [21] We found moderate physical distancing

(permitting 60% of non-household contacts) could contain the outbreaks for geographies when

the contact number is below 11, with infection rate equal to 15.4% [95% BCI 13.5% - 17.2%].

(Figure 3B) Notably, for majority of LMIC countries considered, which have more than 20 close

contacts per day, even strong physical distancing alone (permitting only 40% of non-household

contacts) could not protect more than half of the population (infection rate = 53.5% [95% BCI

50.9% - 56.0%]). Only strict lockdown that reduces non-household contact number to 20% of

that of normal could contain the outbreaks. (contact number = 20: infection rate = 5.1% [95%

BCI 4.4% - 5.9%])

RDT or PCR alone could only mitigate the transmission in communities with less than 20

contacts, while combined with physical distancing, the mitigation effect increases, with 34.8%

[95% BCI 33.3% - 36.3%] infection rate under moderate physical distancing. (contact number =

20, non-household contact rate= 60%) The mitigating effect increases as physical distancing gets

stronger, with RDT slightly outperforming PCR tests. When simulating with a contact rate of 13,

we found that using PCR or antigen RDT alone will isolate or quarantine more than 25% of the

population and reduce the proportion of population infected as the isolation compliance rate

increases. (Supplementary Figure 2A) When isolation compliance rates are at optimistic levels

(90%), antigen RDT testing outperformed PCR and reduced the infection rate to 47.4% [95%

BCI 45.8% - 49.1%].

Combined strategies have varying performance in rural and urban communities

Motivated by the improved containment effect of combining tests with social distancing, we

further evaluated combined strategies based on practice. Firstly, we assume patients who show

up at the test centre with symptoms will be advised to isolate and told their contact to isolate

(symptome-based isolation), which is modelled at a low compliance rate (Table 2); Secondly, we

assumed a patient could provide sample for both a RDT and a PCR test. To contrast the contact

structure between populations, we focused on three representative communities abstracted from

contact data. We found that the symptom+PCR has approximately similar performance as

symptom+RDT, while symptom+PCR+RDT has better performance in all three community

settings. (Figure 4A, Supplementary Figure 3) In rural settings, either symptom+PCR or

symptom+RDT could contain the outbreaks (symptom+PCR: 4.3% infected [95% BCI: 3.6% -

4.9%], Supplementary Table 3). In urban settings, symptom+PCR+RDT could suppress infection

rate to 32.6% [95% BCI: 31.4% - 33.7%], while with physical distancing reduces non-household

contact number by 40%, either symptom+PCR or RDT could contain the outbreaks. Notably,

physical distancing could effectively flatten the curve in urban communities, (Supplementary

Figure 3D) while the effect is less prominent in high density communities. Combined, these

results show that affordable options such as RDT could be sufficient for low-density geographies

similar to the rural settings, whereas testing and tracing programmes need to be combined with

physical distancing to achieve containment in more densely populated areas similar to the urban

settings.

The distributions of infection rates show substantial uncertainty for rural communities and for

urban communities when physical distancing is in place. (Figure 4A, Supplementary Table 3)

These uncertainties suggest the outbreaks might have substantial variation in the epidemic sizes

even for communities with similar demographics. To quantify the probability that imported cases

do not start an outbreak, we defined the stochastic extinction events as the trajectories that

infected less than 5% of the population. Under the three settings considered, the probability of

stochastic extinction increases when testing, isolation and quarantine are implemented. In rural

settings, symptom+RDT would increase the probability of stochastic extinction from 29% to

70%. In urban settings, the stochastic extinction events happened in 62% of the trajectories

when physical distancing and symptom+RDT+PCR are implemented, which is a much larger

proportion compared to 0.5% when no NPI is implemented. In the high density settings, the

stochastic extinction probability remains low even when physical distancing and

symptom+RDT+PCR are all implemented (13.5%). We verified that the probability of stochastic

extinction does not depend on simulated population size. (Supplementary figure 4) The

possibility of achieving high stochastic extinction rates in rural and urban settings suggest

consistent testing, tracing, and physical distancing could reduce the probability of full scale

outbreaks resulting from imported cases.

The impact of RDT sensitivity, tracing and assumed parameters

Sampling success rate, antigen loading, and kit technology could all impact on the sensitivity of antigen RDT. [28,29] When simulating combined strategies with varying antigen RDT sensitivity, our simulations show that the containment effect increases as the RDT sensitivity increases. (Figure 4B) To reach a similar containment effect as in Figure 4A, which assumes an antigen RDT sensitivity of 56% (70% sampling success rate and 80% kit sensitivity), the kits deployed should have a sensitivity above 40%.

For the role of contact tracing, we found when the quarantine compliance rate increased from 30% to 100% under the symptom+RDT+PCR strategy, the average infection rate reduced from 15.7% to 9.5% in rural settings, from 53.2% to 36.5% in urban settings, and from 66.5% to 49.5% in high density settings. (Figure 4C) Note only contacts that are of isolated infectors that are successfully traced are subject to the quarantine. Without close contact quarantine measures, the combination strategies show no significant effect on reducing infection numbers (Supplementary Figure 5). Moreover, we found that the number of infected individuals discovered by testing is less than those discovered by contact tracing when the contact quarantine compliance rate is above 25%. (Supplementary Figure 6A)

The impact of other testing parameters on the containment outcomes are summarised in Supplementary table 3 using R^2 and P-values (Supplementary Methods Section 4). Among these parameters, strong correlation between isolation consent rates and containing effectiveness are observed for all strategies and settings. Days from symptom onset to isolation, quarantine consent rate of contacts and tracing success rate are correlated with containing effectiveness in urban and high density settings, while less impactful in the rural settings. Additionally, we found that starting testing and quarantine after 10% of the population are infected could protect 31% of the healthy population under urban settings and 21% of the healthy population under the high density settings (Supplementary Figure 4B).

Discussion

After two years since the beginning of the COVID-19 pandemic, special attention needs to be

paid to LMICs as they suffer the most from global health inequality. Low vaccination rates and

vulnerable health care systems in these countries could lead to overwhelming outbreaks and

harbour new variants. [6,7,30] Our primary goal is to prioritise tools for LMICs to protect themselves

against COVID-19, where we highlighted four interconnected elements: community knowledge,

RDT, physical distancing, and contact tracing. These elements could be utilised to address two

pressing challenges in LMICs. Firstly, how to design a cost-effective strategy for LMICs to

reduce the financial burden of controlling the transmission of COVID-19? Secondly, how to

design setting-specific surveillance or containment programmes that account for geographical

variability in LMICs [17,31,32]? We argue that community knowledge about demographics and

contact structure is a valuable resource that could facilitate government decision making. For

example, if a community is known to be densely populated and have crowded households, our

results suggest that physical distancing is expected to be less effective there. As another example,

if survey data suggested some communities have less indoor contacts than average, we would

expect a light containment strategy such as mild physical distancing alone could work

effectively, thus saving resources from an over-stringent strategy. In practice, policy makers

could perform contact surveys in representative communities and use them to construct a

classification system. Our analyses present the trends of transmission dynamics and containment

effectiveness across different contact numbers and demographics. When used in combination

with community knowledge, our results could guide policy design that achieves the best

outcomes while saving resources.

Antigen RDT has several desirable properties for application in LMICs. It is cheap, easy to

distribute, has quick turnarounds, and doesn't require lab facilities. LMICs require consistent

containment to protect their health systems from being overwhelmed, which makes RDT an ideal

choice to avoid huge financial costs. Specifically, RDT might be prioritised in resource-limited

regions where contact number is low and performing PCR tests is prohibitive. Though containing

power subjects to the test sensitivity and community scenario, our results suggested high quality

kits that have a sensitivity above 50% will provide similar containment power shown in the

simulation. However, the detrimental effect of false negative and false positive results might

negate the containment effects. [33] We suggest real world RDT application should be

accompanied by careful sensitivity and false positivity evaluation.

We also want to emphasise that policy design should serve its goals. If aiming at a zero-COVID

policy, such as those deployed by China (as of March 5th, 2022), our results show that contact

tracing is an essential element to implement, which relies on abundant personnel and quarantine

capacities. However, if the goal is to flatten the curve to avoid overwhelmed health systems,

mild-physical distancing is an effective measure which saves resources and might serve the

young population of LMICs better. If the goal is to monitor the transmission for early alarm of

new variants and prepare for outbreaks, mass RDT tests could detect abnormal outbreaks that

link to new variants and auxiliary PCR tests could monitor prevalence of each variant.

Successful containment strategy should apply community knowledge to serve the goal, whether

it is suppressing, delaying, or monitoring the transmission of COVID-19.

There are some limitations in our study. Firstly, our simulated population is an approximation to

the communities in LMICs and is limited by the available empirical evidence from LMICs.

Therefore, to apply our results to specific geography requires knowledge of the target population,

including demographic and contact information. Secondly, like many simulation studies, we had

to choose our parameters from empirical studies that are not consistent with each other. [34] This

difficulty is most pronounced when we are setting the parameters for age-dependent

susceptibility and asymptomatic rate of the infected. Lastly, Compared to simulations on larger

networks that involve more sophisticated configuration such as schools and shopping malls, [35]

our strategy might not capture superspreading events that were reported. Future development of

approximation methods might provide comparable accuracy while scaling to large populations.

Contributors

356 All authors contributed to study design. XJ designed the model, XJ and WG implemented the

357 model, and XJ, WG, CF and CH analysed and verified data. All authors interpreted results, XJ,

358 CF and CH drafted the manuscript, XJ made the figures, and all authors critically revised and

359 approved the manuscript.

Code availability

The code generated during this study is available at:

https://github.com/Xilin-Jiang/NetworkCOVID19

Acknowledgement

Computation used the Oxford Biomedical Research Computing (BMRC) facility, a joint development between the Wellcome Centre for Human Genetics and the Big Data Institute supported by Health Data Research UK and the NIHR Oxford Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. We thank Microsoft China Co. Ltd. for computation power during the exploration stage of the study. We thank Yang Liu, Jiayao Lei and Mark Jit for discussion and comments on the manuscript.

373 **Tables and Figures**

Parameter	Sensitivity analysis range			Coefficient of determination			References
	Rural	Urban	High density	Rural	Urban	High density	
Population size	1000			0.005 (P = 0.222)	0.001 (P = 0.604)	0.009 (P = 0.109)	[36]
Number of initial infected patients	2			0.24 (P = 0)	0.064 (P = 0)	0.022 (P = 0)	Assumed
Daily close contact number	7	13	14	0.879 (P = 0)	0.806 (P = 0)	0.855 (P = 0)	[20,37]
Average household size and distribution	5 (Uganda)	5 (Uganda)	15 (India)	NA			[18,38]
Percentage of contact between members of the same household	50%	23%	50%	0.001 (P = 0.381)	0.173 (P = 0)	0.199 (P = 0)	[20,37]
Asymptomatic proportion	30% (10%-90%)			0.001 (P = 0.22)	0 (P = 0.634)	0 (P = 0.887)	[39,40]
Asymptomatic infection rate	20% (10%-90%)			0 (P = 0.4)	0.001 (P = 0.119)	0.002 (P = 0.066)	Assumed
Healthcare seeking rate	70%		60%	0.001 (P = 0.403)	0 (P = 0.649)	0.002 (P = 0.176)	Assumed
Basic reproduction number (R0)	1.45	2.7	2.9	NA			[41]
Susceptibility for younger population (< 15 years old)	50%			NA			[26]

Death rate for each age group	0.2% (0-14 years old); 0.2% (15-24 years old); 1% (25+ years old);	NA	[27]

Table 1: Summary of population characteristics and transmission parameters and their impact on transmission. For parameters where the source is not available for at least one of the three community settings, we simulated models with different choices of values and used regression to evaluate their impact on model outputs. (Supplementary Table 4, Supplementary Methods Section 4) The coefficient of determination (R^2) shows the proportion of variance in the infected number that is captured by the parameter; The P-values in the parentheses show the rate of type one error of rejecting the null hypothesis that the parameter is not correlated with infection number. Last column shows the sources for cited parameters.

Parameter	Symptom	PCR	Antigen RDT	Data source
Test consent rate	100%	80%		Assumption
Isolation consent rate	20%	90%		Assumption
Days from symptom onset to isolation (days)	1	5	1	Assumption
Duration that close contacts are traced (days)	3	5	3	Assumption
Tracing success rate (non-household contacts)	85%	80%	85%	Assumption
Quarantine consent rate of contacts	50%	70%		Assumption
Sampling success rate	NA	70%		[29,42,43]
Lab sensitivity	NA	100%	80% (10%-90%)	[28,29,44]
Transmission rate after quarantine	Rural: 10%; Urban: 10%; High density: 20%			Assumption

Table 2: Summary of testing and NPI parameters. Last column shows the sources for cited parameters; values of assumed parameters are used in the containment simulations in Figure 3 and Figure 4, while separate sensitivity analyses are performed for each assumed parameter (Supplementary Table 3 and Supplementary Table 4). For rationales behind the chosen value for assumed parameters, see Supplementary Methods.

Figure 1: Schematic representation of methodology. A) For each parameter setting we simulated 20 synthetic populations and inferred the contact network with ERGM, using age- and household-structured contact data. At each simulation step, a contact network is sampled from the ERGM, where we simulate the transmission of SARS-CoV-2. B) We evaluated the effect of physical distancing by reducing the sampling probability of non-household contacts. Testing, isolation, contact tracing, and quarantine are also simulated using the contact networks.

Figure 2: Properties of simulated outbreaks from the agent-based model. A) Example outbreak in a network composed of 1000 individuals, simulated under the urban setting without NPI. The left, middle and right panel show the distributions of infected individuals (red dots) on day 20, day 30 and day 40 from the day that initial cases are imported. For visualisation purposes we fix the network structure across days here, while in our analysis a different contact network is sampled each day. B) Comparison of the simulated contact number distribution with those computed from the cited Uganda survey data.[24] The simulated contact number distribution is computed by sampling 100 daily contact realisations from each of the 20 synthetic populations. C) Comparison of generation time from simulation (red histogram) and that from cited Uganda survey data [24] (blue dotted curve).

Figure 3: Evaluation of containing strategies. A) Example of case finding and quarantine in one simulated outbreak trajectory with NPIs, where a moderate physical distancing reduces 40% of the contact outside the household and symptom+PCR testing is performed. The network structure is fixed for visualisation, while in simulation a contact network is different each day. B) Social distancing and tests have different containing effects over different geographies. Synthetic populations were constructed using a grid of contact numbers that covers several reference LMICs. Left panel shows simulation with different levels of physical distance, indicated by the percentage of non-household contacts that are still permitted (no physical distancing: 100%; strongest physical distancing: 20%). The middle and right panel shows physical distancing combined with PCR or RDT tests respectively. The size and colour of the circle shows the mean

Figure 4: The impacts of testing strategies, sensitivity, and compliance rate on epidemic size

vary with community settings. A) Infection rates under different combined strategies and in three

abstracted community settings. Each dot represents one 100-day simulated outbreak and each

box presents 200 simulated outbreaks, throughout all panels. Green boxes represent simulation

with physical distancing (60% non-household contact permitted) and orange boxes show those

without physical distancing. The quintiles of infection rate and mortality rate are summarised in

Supplementary Table 3. B) Infection rates for each community setting when applying antigen

RDT of various sensitivity. The blue boxes show the infection rate when applying the most

effective testing strategy identified in the upper panel, with different antigen RDT sensitivity. C)

Infection rates for each community setting when different quarantine compliances rates of traced

contacts are simulated. The blue boxes show the infection rates when simulating containment

using the most effective testing strategy identified in the first panel. In all three panels, one dot

represents the epidemic size of one bootstrap trajectory. The red dashed line shows an infection

rate at 5%, dots below which are defined as stochastic extinction events.

Supplementary Tables and Figures

Age group (in years)	Uganda	
0-14	48.10%	
15-24	20.30%	
25+	31.60%	
Household Size (members)	Uganda	High density community
1	11.00%	0.00%
2-3	22.00%	6.00%
4-5	27.00%	17.00%
6+	40.00%	77.00%

Supplementary Table 1: Summary of the demographic information used in the study. We used the age and household size distributions from Uganda, except for the household size distribution in high density communities, where we used data from Afghanistan, which has the largest average household size recorded by the United Nations. [18,19]

Contact number	Physical distancing					Physical distancing + PCR					Physical distancing + RDT				
	100%	80%	60%	40%	20%	100%	80%	60%	40%	20%	100%	80%	60%	40%	20%
8	31.8% (0.5% - 51.1%)	9.0% (0.3% - 25.9%)	2.0% (0.3% - 5.1%)	0.8% (0.2% - 1.6%)	0.4% (0.2% - 0.8%)	8.1% (0.5% - 17.4%)	2.9% (0.3% - 7.6%)	1.4% (0.3% - 3.0%)	0.6% (0.2% - 1.3%)	0.5% (0.2% - 0.9%)	5.9% (0.4% - 15.2%)	2.8% (0.4% - 7.3%)	1.2% (0.3% - 2.6%)	0.7% (0.2% - 1.4%)	0.4% (0.2% - 0.7%)
11	66.7% (66.6% - 75.0%)	48.1% (1.3% - 62.2%)	15.4% (0.5% - 34.2%)	2.5% (0.3% - 7.2%)	0.7% (0.2% - 1.4%)	29.9% (13.3% - 39.0%)	14.0% (1.0% - 26.6%)	4.4% (0.6% - 10.2%)	1.4% (0.3% - 3.1%)	0.6% (0.2% - 1.1%)	26.2% (7.7% - 37.7%)	10.7% (0.6% - 23.1%)	3.5% (0.4% - 8.0%)	1.1% (0.3% - 2.6%)	0.6% (0.2% - 1.1%)
14	82.5% (80.2% - 85.7%)	72.4% (70.1% - 78.3%)	45.5% (0.5% - 62.2%)	9.1% (0.4% - 21.4%)	1.2% (0.3% - 2.8%)	46.5% (41.7% - 51.6%)	32.5% (20.1% - 43.4%)	12.0% (0.8% - 25.2%)	2.7% (0.3% - 5.9%)	0.9% (0.2% - 2.0%)	43.4% (37.9% - 50.4%)	29.4% (5.1% - 41.3%)	10.4% (1.0% - 24.0%)	2.7% (0.4% - 6.1%)	0.8% (0.2% - 1.6%)
17	88.5% (87.3% - 90.9%)	83.5% (81.1% - 85.8%)	69.4% (66.4% - 75.8%)	26.5% (1.0% - 47.9%)	2.3% (0.3% - 5.8%)	52.2% (47.3% - 56.9%)	44.6% (39.0% - 51.0%)	25.7% (6.1% - 38.5%)	6.1% (0.7% - 14.0%)	1.3% (0.3% - 2.8%)	51.2% (46.0% - 55.7%)	40.6% (34.1% - 47.7%)	19.4% (1.9% - 34.1%)	3.9% (0.4% - 8.4%)	1.1% (0.3% - 2.3%)
20	92.8% (91.3% - 94.2%)	88.8% (86.8% - 90.6%)	79.6% (78.0% - 83.8%)	53.5% (27.4% - 67.4%)	5.1% (0.3% - 13.2%)	58.7% (54.3% - 63.2%)	52.2% (47.6% - 56.6%)	39.3% (29.5% - 47.1%)	13.0% (1.5% - 25.7%)	1.9% (0.3% - 4.5%)	57.0% (52.0% - 61.8%)	49.9% (45.5% - 55.0%)	34.8% (22.0% - 44.7%)	10.1% (1.0% - 20.5%)	1.9% (0.4% - 4.2%)
23	95.1% (94.1% - 96.2%)	92.1% (90.7% - 93.5%)	86.2% (83.8% - 88.5%)	69.4% (65.3% - 76.8%)	10.5% (0.4% - 25.8%)	63.5% (60.1% - 67.3%)	57.8% (53.1% - 61.6%)	48.8% (43.6% - 53.4%)	24.1% (15.8% - 38.6%)	3.4% (0.5% - 7.9%)	61.6% (56.7% - 65.7%)	55.1% (50.8% - 60.0%)	44.6% (37.1% - 51.7%)	17.7% (1.3% - 31.0%)	2.7% (0.5% - 6.8%)
26	96.6% (95.6% - 97.5%)	94.2% (93.1% - 95.3%)	89.0% (88.0% - 91.7%)	78.1% (76.1% - 82.7%)	22.7% (0.7% - 44.3%)	68.1% (64.8% - 71.7%)	62.3% (58.9% - 65.9%)	54.3% (50.4% - 59.1%)	34.5% (15.8% - 46.2%)	5.0% (0.5% - 11.3%)	65.2% (61.1% - 68.8%)	60.0% (56.3% - 63.6%)	51.6% (46.4% - 56.9%)	27.2% (7.5% - 41.2%)	4.3% (0.5% - 9.9%)
29	97.6% (96.9% - 98.3%)	95.8% (94.9% - 96.9%)	92.3% (91.0% - 93.8%)	85.0% (82.6% - 87.1%)	33.1% (1.1% - 57.9%)	72.0% (69.0% - 74.9%)	66.2% (62.3% - 69.8%)	59.5% (55.4% - 63.5%)	44.4% (36.3% - 52.4%)	8.0% (1.0% - 17.3%)	68.6% (64.8% - 72.2%)	63.5% (59.3% - 67.9%)	56.8% (51.5% - 61.2%)	38.9% (22.1% - 48.8%)	6.8% (0.8% - 13.7%)

Supplementary Table 2: Summary of the distributions of infection rates for that of Figure 3B. The mean of infection rates is computed from 200 simulated trajectories of 100 days for each setting. The range in the bracket shows the 10% and 90% quantiles of the infection rates from the 200 bootstrap sample trajectories.

Strategies	Infection rate			Mortality rate		
	Rural	Urban	High density	Rural	Urban	High density
No intervention	18.0% (0.5% - 38.2%)	79.5% (77.3% - 82.4%)	87.0% (85.2% - 89.4%)	0.1% (0.0% - 0.2%)	0.4% (0.2% - 0.6%)	0.4% (0.1% - 0.7%)
Symptom + PCR	4.3% (0.4% - 11.5%)	41.0% (36.0% - 47.5%)	51.8% (47.1% - 56.9%)	0.0% (0.0% - 0.1%)	0.2% (0.0% - 0.4%)	0.3% (0.1% - 0.5%)
Symptom + RDT	4.0% (0.4% - 10.3%)	39.7% (34.1% - 46.9%)	51.6% (47.5% - 56.8%)	0.0% (0.0% - 0.1%)	0.2% (0.0% - 0.4%)	0.3% (0.1% - 0.5%)
Symptom + PCR + RDT	3.0% (0.4% - 7.6%)	32.6% (25.4% - 39.8%)	45.3% (40.9% - 50.3%)	0.0% (0.0% - 0.1%)	0.2% (0.0% - 0.4%)	0.2% (0.0% - 0.4%)
Reducing 40% of non-household contact	2.5% (0.3% - 6.9%)	33.7% (0.8% - 55.5%)	74.1% (71.6% - 78.9%)	0.0% (0.0% - 0.1%)	0.2% (0.0% - 0.4%)	0.4% (0.2% - 0.6%)
Symptom + PCR + reducing 40% of non-household contact	1.3% (0.3% - 3.2%)	8.4% (0.7% - 19.7%)	31.1% (7.0% - 43.0%)	0.0% (0.0% - 0.0%)	0.0% (0.0% - 0.1%)	0.1% (0.0% - 0.3%)
Symptom + RDT + reducing 40% of non-household contact	1.2% (0.3% - 2.7%)	6.5% (0.6% - 14.4%)	29.8% (6.1% - 42.4%)	0.0% (0.0% - 0.0%)	0.0% (0.0% - 0.1%)	0.1% (0.0% - 0.3%)
Symptom + PCR + RDT + reducing 40% of non-household contact	1.1% (0.3% - 2.2%)	5.0% (0.6% - 11.8%)	21.6% (3.2% - 34.4%)	0.0% (0.0% - 0.0%)	0.0% (0.0% - 0.1%)	0.1% (0.0% - 0.2%)

Supplementary Table 3: Summary of the distributions of infection and mortality rate for three community settings. The mean of infection rate and mortality rates is computed from 200 simulated trajectories of 100 days for each strategy. The range in the bracket shows the 10% and 90% quantiles of the infection rates from 200 bootstrap samples.

A)

Parameter	Symptom	PCR	Antigen RDT	Antibody RDT	Coefficient of determination		
					Strategy 1	Strategy 2	Strategy 3
Test consent rate	100%	80%		70%	0.001 (P = 0.289)	0.007 (P = 0.022)	0.065 (P = 0)
Isolation consent rate	20%	90%			0.218 (P = 0); symptom	0.245 (P = 0); symptom	0.107 (P = 0); antigen
Antibody testing day (days from onset)	NA			8	0 (P = 0.63)	0.005 (P = 0.012)	0 (P = 0.97)
Days from symptom onset to isolation (days)	1	5	1	8	0.031 (P = 0)	0.036 (P = 0)	0.079 (P = 0)
Duration that close contacts are traced (days)	3	5	3	8	0.001 (P = 0.361)	0.149 (P = 0)	0.004 (P = 0.065)
Tracing success rate (non-household contacts)	85%	80%	85%	75%	0.021 (P = 0)	0.04 (P = 0)	0.078 (P = 0)
Quarantine consent rate of contacts	50%	70%			0.05 (P = 0)	0.072 (P = 0)	0.093 (P = 0)
Sampling success rate	NA	70%		100%	NA		
Lab sensitivity	NA	100%	80% (10%-90%)	90%	NA		
Half of patient develop detectable serological response (Days from infection)	NA			12	NA		
Transmission rate after quarantine	Rural: 10%; Non-slum urban: 10%; Slum: 20%				0.004 (P = 0.042)	0.004 (P = 0.04)	0.002 (P = 0.115)
Test kit availability	NA	Rural: none Non-slum urban: unlimited Slum: 2 per week per 1000 people	Unlimited		NA		

Setting: Rural   Strategy 1: Symptom;   Strategy 2: Symptom + Antibody RDT;   Strategy 3: Antigen RDT ;

B)

Parameter	Symptom	PCR	Antigen RDT	Antibody RDT	Coefficient of determination		
					Strategy 1	Strategy 2	Strategy 3
Test consent rate	100%	80%		70%	0.028 (P = 0)	0.056 (P = 0)	0.187 (P = 0)
Isolation consent rate	20%	90%			0.438 (P = 0); symptom	0.388 (P = 0); symptom	0.19 (P = 0); antigen
Antibody testing day (days from onset)	NA			8	0 (P = 0.868)	0 (P = 0.946)	0 (P = 0.839)
Days from symptom onset to isolation (days)	1	5	1	8	0.16 (P = 0)	0.173 (P = 0)	0.287 (P = 0)
Duration that close contacts are traced (days)	3	5	3	8	0.002 (P = 0.166)	0.317 (P = 0)	0 (P = 0.787)
Tracing success rate (non-household contacts)	85%	80%	85%	75%	0.181 (P = 0)	0.235 (P = 0)	0.246 (P = 0)
Quarantine consent rate of contacts	50%	70%			0.214 (P = 0)	0.248 (P = 0)	0.221 (P = 0)
Sampling success rate	NA	30%		0	NA		
Lab sensitivity	NA	100%	80%	90%	NA		
Half of patient develop detectable serological response (Days from infection)	NA			7	NA		
Transmission rate after quarantine	Rural: 10%; Non-slum urban: 10%; Slum: 20%				0.033 (P = 0)	0.03 (P = 0)	0.096 (P = 0)
Test kit availability	NA	Rural: none Non-slum urban: unlimited Slum: 2 per week per 1000 people	Unlimited		NA		

Setting: Non-slum urban   Strategy 1: Symptom + PCR;   Strategy 2: Symptom + PCR + Antibody RDT;   Strategy 3: Antigen RDT + PCR;

C)

Parameter	Symptom	PCR	Antigen RDT	Antibody RDT	Coefficient of determination		
					Strategy 1	Strategy 2	Strategy 3
Test consent rate	100%	80%		70%	0.037 (P = 0)	0.055 (P = 0)	0.341 (P = 0)
Isolation consent rate	20%	90%			0.448 (P = 0); symptom	0.445 (P = 0); symptom	0.32 (P = 0); antigen
Antibody testing day (days from onset)	NA			8	0 (P = 0.886)	0.016 (P = 0)	0 (P = 0.847)
Days from symptom onset to isolation (days)	1	5	1	8	0.226 (P = 0)	0.282 (P = 0)	0.257 (P = 0)
Duration that close contacts are traced (days)	3	5	3	8	0.064 (P = 0)	0.408 (P = 0)	0.05 (P = 0)
Tracing success rate (non-household contacts)	85%	80%	85%	75%	0.289 (P = 0)	0.365 (P = 0)	0.367 (P = 0)
Quarantine consent rate of contacts	50%	70%			0.354 (P = 0)	0.471 (P = 0)	0.267 (P = 0)
Sampling success rate	NA	70%		100%	NA		
Lab sensitivity	NA	100%	80% (10%-90%)	90%	NA		
Half of patient develop detectable serological response (Days from infection)	NA			12	NA		
Transmission rate after quarantine	Rural: 10%; Non-slum urban: 10%; Slum: 20%				0.101 (P = 0)	0.095 (P = 0)	0.074 (P = 0)
Test kit availability	NA	Rural: none Non-slum urban: unlimited Slum: 2 per week per 1000 people	Unlimited		NA		

Setting: Slum Strategy 1: Symptom + limited PCR; Strategy 2: Symptom + limited PCR + Antibody RDT; Strategy 3: Antigen RDT + limited PCR;

Supplementary Table 4: The impact of NPI parameter setting on the effectiveness of containment. Panel A-C shows sensitivity analysis under rural, urban and high density settings. The coefficient of determination (R^2) shows the proportion of variance in the final infected number (under each containing strategy) that is captured by the testing parameters; The P-values in the bracket show the rate of type one error of rejecting the null hypothesis that the parameter is not correlated with outcome.

ID	Description	Values								
1	Basic reproduction number	1.5	2	2.5	3	3.5				
2	Number of daily contact	2	4	7	10	14				
3	Number of intial inporting cases	1	2	5	8	15				
4	Healthcare seeking rate	30%	50%	70%	90%	100%				
5	Asymptomatic rate	10%	20%	30%	40%	50%	60%	70%	80%	90%
6	Transmission ratio for isolated patients	5%	10%	15%	20%	25%				
7	Daily PCR tests per 1000 population for slum	Not available	1	2	10	20	50	1000		
8	Community size	100	1000	2000						
9	Infection number in 1000 to trigger containment	No delay	5	20	50	100	250	500		
10	Percentage of contacts that are with a member of household, fixing the number of total contacts	20%	40%	60% (did not converge for slum settings)	80% (did not converge for three settings)					
11	Social distancing, which is computed as the proportion of non-household contact that is stopped.	0%	20%	40%	60% (did not converged for rural setting)	80% (did not converged for rural and slum settings)				
12	Non-household contact trace successful rate	0%	15%	35%	55%	75%	85%	95%		
13	Delay after tested postive to isolation (days)	0	1	3	5					
14	Duration of contact tracing, counted from the day of isolation (days)	Symptom screening/Antigen RDT = 1; Antibody RDT = 6; PCR=3	Symptom screening/Antigen RDT = 3; Antibody RDT = 8; PCR = 5	Symptom screening/Antigen RDT = 5; Antibody RDT = 10; PCR = 7	Symptom screening/Antigen RDT = 7; Antibody RDT = 12; PCR = 9					
15	Delay after onset for PCR testing results (days); contact tracing duration is set to be same as the delay, as we assume healthcare workers will start contact tracing once they identify an individual with onset.	0	1	3	5	7	10	14		
16	Delay after onset for antibody sampling (days); contact tracing duration is set to be same as the delay, as we assume healthcare workers will start contact tracing once they identify an individual with onset.	5	6	7	8	9	10	11		
17	Seroconversion rate	opimistic: half of individual has detectable antibody 7 days after onset	opimistic: half of individual has detectable antibody 12 days after infection	conservative: half of individual has detectable antibody 9 days after onset						
18	Test acceptance rate (conditional on an individual seek healthcare)	Antibody (blood sample) = 30%; PCR/Antigen (swab sample) = 40%	Antibody (blood sample) = 50%; PCR/Antigen (swab sample) = 60%	Antibody (blood sample) = 70%; PCR/Antigen (swab sample) = 80%	90%					
19	Sympotom-based isolation consent rate, simulated with other testing measures	10%	20%	30%	40%	50%	60%	70%	80%	90%
20	Sympotom-based isolation consent rate (simulation with only symptom detection, no other detecting measures)	10%	20%	30%	40%	50%	60%	70%	80%	90%
21	PCR isolation consent rate (simulation with only PCR testing, no other detecting measure)	10%	20%	30%	40%	50%	60%	70%	80%	90%
22	Antibody RDT isolation consent rate (simulation with only antibody RDT, no other detecting measure)	10%	20%	30%	40%	50%	60%	70%	80%	90%
23	Antigen RDT isolation consent rate (simulation with only antigen RDT, no other detecting measure)	10%	20%	30%	40%	50%	60%	70%	80%	90%
24	close contact quarantine compliance rate (conditional on the infector consent to be isolated)	Symptom screen = 10%; Testing = 30%	Symptom screen = 30%; Testing = 50%	Symptom screen = 50%; Testing = 70%	Symptom screen = 70%; Testing = 90%	Symptom screen = 90%; Testing = 100%				

Supplementary Table 5: Parameter settings for sensitivity analysis. The Description explains what variables are changed and rows contain the values of each variable used for simulation. For each value we performed the simulation for baseline and three types of containing strategies under the rural, urban and high density settings.

ID	Description	Values				
2	**Number of daily contact**	**2**	**4**	**7**	**10**	**14**
	Rural	0.99	1.00	1.00	1.00	0.14
	Non-slum urban	1.00	1.00	0.99	1.00	1.00
	Slum	1.00	0.94	0.99	1.00	1.00
9	**Community size**	**100 (did not converge for rural setting)**	**1000**	**2000**		
	Rural	0.000002 *	1.00	1.00		
	Non-slum urban	1.00	1.00	1.00		
	Slum	1.00	1.00	1.00		
11	**Percentage of contacts that are with a member of household, fixing the number of total contacts**	**20%**	**40%**	**60% (did not converge for rural setting)**	**80% (did not converge for three settings)**	
	Rural	1.00	1.00	1.00	0 *	
	Non-slum urban	1.00	1.00	0.006*	0 *	
	Slum	0.99	1.00	1.00	0 *	
12	**Social distancing, which is computed as the proportion of non-household contact that is stopped.**	**0%**	**20%**	**40%**	**60%**	**80%**
	Rural	1.00	1.00	0.99	1.00	1.00
	Non-slum urban	1.00	1.00	1.00	1.00	1.00
	Slum	1.00	1.00	1.00	1.00	1.00

Supplementary Table 6: Lack-of-fit test for the ERGM inference. We performed lack-of-fit tests of the target statistics for the four variables that alter the ERGMs. IDs correspond to Supplementary Table 3. For each value (bold), we compute the P-value by comparing the 20 fitted networks with the input to the ERGMs. * shows the cases when the networks are rejected as good fits.

Supplementary Figure 1: Comparison of fitted age-mixing contact matrix with those cited from survey data. The color scale shows the total number of contacts for each mixing within the populations.

500 **Supplementary Figure 2** A) Proportion of populations that are infected (at the end of the

501 trajectories) and under quarantine (max value throughout the trajectories) using PCR and antigen

502 RDT as detection methods. The dots with bars show the mean value with 95% confidence

503 interval from 200 simulated trajectories for each isolation compliance rate. B) Proportion of the

population infected when physical distancing blocks different proportions of non-household

contacts. The box shows the corresponding effective reproductive number for the level of

physical distancing. The combined effect of physical distancing with each testing method shown

in B is plotted with different colours.

Supplementary Figure 3: Summary of averaged outbreak trajectories in rural, urban and high

density settings. A-B) Averaged trajectories of outbreaks under rural setting and those with three

NPI containing strategies. Left panel shows the simulation without physical distancing and the

right panel shows simulation with a physical distancing that blocks 40% of non-household

contacts. C-D) Same averaged trajectories under urban setting; Left panel shows simulation

without physical distancing and right panel shows simulation with a physical distancing that

blocks 40% of non-household contacts. E-F) Same averaged trajectories under high density

community setting; Left panel shows simulation without physical distancing and right panel

shows simulation with a physical distancing that blocks 40% of non-household contacts. The

curves are the mean value of daily infected numbers for a synthetic population of 1000; The

shaded area indicated the 95% confidence interval of the estimation; Both mean and confidence

intervals are computed from 200 simulated outbreaks.

Supplementary Figure 4: Probability of stochastic extinction against simulated population size under the rural setting. 200 bootstrap outbreaks were simulated for each population size, with stochastic extinction probability computed as the proportion of outbreaks that have less than 50 cases at the end of the 100th day.

Supplementary Figure 5: Averaged trajectories when no close contact tracing is performed under rural, urban and high density settings. The curves are the mean value of daily infected numbers for a synthetic population of 1000; The shaded area indicated the 95% confidence

 interval of the estimation; Both mean and confidence interval are computed from 200 simulated

outbreaks.

Supplementary Figure 6: Evaluation of close contact tracing and delayed responses. A)

Proportion of total infected individuals who are either isolated by testing (red dots) or

quarantined by tracing (green dots) varies as the compliance rate of quarantine a close contact of

a confirmed case changes. B) Implementation of containing measures after a certain proportion

of the population was infected could still protect a proportion of the population, compared to the

circumstances with no intervention. All simulations are performed for rural, urban and high

desnity settings.

References

1.   Zhu, N. *et al*. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med*.

382, 727–733 (2020).

2.   Tatar, M., Shoorekchali, J. M., Faraji, M. R. & Wilson, F. A. International COVID-19 vaccine

inequality amid the pandemic: Perpetuating a global crisis? *J. Glob. Health* **11**, 03086 (2021).

3.   Nkengasong, J. N., Ndembi, N., Tshangela, A. & Raji, T. COVID-19 vaccines: how to ensure Africa

has access. (2020).

4.   Brauner, J. M. *et al.* Inferring the effectiveness of government interventions against COVID-19.

Science **371**, (2021).

5.   Hodgins, S. & Saad, A. Will the Higher-Income Country Blueprint for COVID-19 Work in Low-

and Lower Middle-Income Countries? *Glob Health Sci Pract* **8**, 136–143 (2020).

6.   Collie, S., Champion, J., Moultrie, H., Bekker, L.-G. & Gray, G. Effectiveness of BNT162b2

Vaccine against Omicron Variant in South Africa. *N. Engl. J. Med.* **386**, 494–496 (2022).

7.   Pulliam, J. R. C. *et al.* Increased risk of SARS-CoV-2 reinfection associated with emergence of the

Omicron variant in South Africa. *MedRxiv* (2021).

8.   Bong, C.-L. *et al.* The COVID-19 Pandemic: Effects on Low- and Middle-Income Countries.

Anesth. Analg. **131**, 86–92 (2020).

9.   Murthy, S., Leligdowicz, A. & Adhikari, N. K. J. Intensive care unit capacity in low-income

countries: a systematic review. *PLoS One* **10**, e0116949 (2015).

10. Hasell, J. *et al.* A cross-country database of COVID-19 testing. *Sci Data* **7**, 345 (2020).

11. Nkengasong, J. Let Africa into the market for COVID-19 diagnostics. *Nature* **580**, 565 (2020).

12. Porte, L., Legarraga, P., Vollrath, V. & Aguilera, X. Evaluation of novel antigen-based rapid

detection test for the diagnosis of SARS-CoV-2 in respiratory samples. *International Journal of*

(2020).

13. Krueger, L. J., Gaeddert, M., Koeppel, L. & Bruemmer, L. Evaluation of the accuracy, ease of use

and limit of detection of novel, rapid, antigen-detecting point-of-care diagnostics for SARS-CoV-2.

medRxiv (2020).

14. Jacobs, J. *et al*. Implementing COVID-19 (SARS-CoV-2) Rapid Diagnostic Tests in Sub-Saharan

Africa: A Review. *Front*. *Med*. **7**, 557797 (2020).

15. Shapiro, A. E. *et al*. Community-based targeted case finding for tuberculosis and HIV in household

contacts of patients with tuberculosis in South Africa. *Am*. *J*. *Respir*. *Crit*. *Care Med*. **185**, 1110–

1116 (2012).

16. Baqui, P., Bica, I., Marra, V. & Ercole, A. Ethnic and regional variations in hospital mortality from

COVID-19 in Brazil: a cross-sectional observational study. *The Lancet Global* (2020).

17. Mohanan, M., Malani, A., Krishnan, K. & Acharya, A. Prevalence of SARS-CoV-2 in Karnataka,

India. *JAMA* **325**, 1001–1003 (2021).

18. United Nation. Household Size and Composition Around the World 2017. (2017).

19. Department of Economic and Social Affairs. *World Population Prospects 2019 - Volume II:*

Demographic Profiles. (United Nations, 2020).

20. le Polain de Waroux, O. *et al*. Characteristics of human encounters and social mixing patterns

relevant to infectious diseases spread by close contact: a survey in Southwest Uganda. *BMC Infect*.

Dis. **18**, 172 (2018).

21. Dobreva, Z, Gimma, A, Rohan, H, Djoudalbaye, B, Tshangela, A, Quaife, M. Characterising social

contacts under COVID-19 control measures in 18 African countries. in *1st International Conference*

on Public Health in Africa (CPHIA2021).

22. Robins, G., Pattison, P., Kalish, Y. & Lusher, D. An introduction to exponential random graph (p*)

models for social networks. *Soc*. *Networks* **29**, 173–191 (2007).

23. Sun, K. *et al*. Transmission heterogeneities, kinetics, and controllability of SARS-CoV-2. *Science*

(2020) doi:10.1126/science.abe2424.

24. Ferretti, L. *et al*. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital

contact tracing. *Science* **368**, (2020).

25. Zhang, J. *et al*. Age profile of susceptibility, mixing, and social distancing shape the dynamics of the

novel coronavirus disease 2019 outbreak in China. *medRxiv* (2020)

doi:10.1101/2020.03.19.20039107.

26. Davies, N. G. *et al*. Age-dependent effects in the transmission and control of COVID-19 epidemics.

MedRxiv (2020).

27. Verity, R. *et al*. Estimates of the severity of coronavirus disease 2019: a model-based analysis.

Lancet Infect. Dis. **20**, 669–677 (2020).

28. Scohy, A. *et al*. Low performance of rapid antigen detection test as frontline testing for COVID-19

diagnosis. *J. Clin. Virol.* **129**, 104455 (2020).

29. Pilarowski, G. *et al*. Performance characteristics of a rapid SARS-CoV-2 antigen detection assay at a

public plaza testing site in San Francisco. *medRxiv* (2020) doi:10.1101/2020.11.02.20223891.

30. Daria, S., Asaduzzaman, M., Shahriar, M. & Islam, M. R. The massive attack of COVID-19 in India

is a big concern for Bangladesh: The key focus should be given on the interconnection between the

countries. *Int. J. Health Plann. Manage.* **36**, 1947–1949 (2021).

31. Nordling, L. The pandemic appears to have spared Africa so far. Scientists are struggling to explain

why. *Science* (2020) doi:10.1126/science.abe2825.

32. Makoni, M. COVID-19 in Africa: half a year later. *Lancet Infect. Dis.* **20**, 1127 (2020).

33. Bock, P. *et al*. Understanding low sensitivity of community-based HIV rapid testing: experiences

from the HPTN 071 (PopART) trial in Zambia and South Africa. *J. Int. AIDS Soc.* **20**, 21780 (2017).

34. Omori, R., Matsuyama, R. & Nakata, Y. The age distribution of mortality from novel coronavirus

disease (COVID-19) suggests no large difference of susceptibility by age. *Sci. Rep.* **10**, 16642

(2020).

35. Hinch, R., Probert, W. J. M., Nurtay, A., Kendall, M. & Wymatt, C. OpenABM-Covid19-an agent-

based model for non-pharmaceutical interventions against COVID-19 including contact tracing.

medRxiv (2020).

36. Food and Agriculture Organization. *Guidelines on defining rural areas and compiling indicators for*

development policy. (2018).

37.   Mossong, J., Hens, N., Jit, M., Beutels, P. & Auranen, K. POLYMOD social contact data. (2017).

38.   Kumar, S. *et al*. Who interacts with whom? Social mixing insights from a rural population in India.

628   *PLoS One* **13**, e0209039 (2018).

629   39.   Nishiura, H. *et al*. Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-

630   19). *Int. J. Infect. Dis.* **94**, 154–155 (2020).

631   40.   Nogrady, B. What the data say about asymptomatic COVID infections. *Nature* **587**, 534–535 (2020).

632   41.   Anderson, R. M., Heesterbeek, H., Klinkenberg, D. & Hollingsworth, T. D. How will country-based

633   mitigation measures influence the course of the COVID-19 epidemic? *Lancet* **395**, 931–934 (2020).

634   42.   Wölfel, R. *et al*. Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**,

635   465–469 (2020).

636   43.   Wu, F. *et al*. Neutralizing Antibody Responses to SARS-CoV-2 in a COVID-19 Recovered Patient

637   Cohort and Their Implications. (2020) doi:10.2139/ssrn.3566211.

638   44.   Li, Z. *et al*. Development and clinical application of a rapid IgM-IgG combined antibody test for

639   SARS-CoV-2 infection diagnosis. *J. Med. Virol.* **92**, 1518–1524 (2020).