SARS-CoV-2 spike protein gene variants with N501T and G142D mutation dominated

infections in minks in the US

Hugh Y. Cai[1], Allison Cai[2]

Animal Health Laboratory, University of Guelph, Guelph, Ontario, Canada (H. Cai)

Faculty of Science, University of British Colombia, BC, Canada (A. Cai)

*Corresponding author: Hugh Cai, Animal Health Laboratory, University of Guelph, 419

Gordon Street, Bldg 89, Guelph, Ontario, Canadda. N1G 2W1. hcai@uoguelph.ca

Running title: SARS-CoV-2 N501T-G142D variants dominated infections in minks in the US

Abstract

Large number of minks were infected with SARS-CoV-2 virus containing the spike protein

Y453F mutation in Europe, causing zoonosis concerns. To evaluate the genetic characteristics of

the US and Canadian mink-derived SARS-CoV-2 sequences, we analyzed all animal-derived

(977), all Canadian (19,529) and US (173,277) SARS-CoV-2 sequences deposited in GISAID

from December 2019 to March 12, 2021, and identified 2 dominant novel variants, the N501T-

G142D variant and N501T-G142D-F486L variant, in the US mink-derived SARS-CoV-2

sequences. These variants were not found in minks from Canada or other countries. The Y453F

mutation was not identified in the mink-derive sequences in the US and Canada. The N501T

mutation occurred two months earlier in the human than in the minks in the US, and the novel

N501T-G142D variant and N501T-G142D-F486L variant were found in human prior to minks.

The result of this study indicates that the novel variants may have evolved during human

infection and then transmitted to mink populations in the US.

Keywords

SARS-CoV-2, COVID-19, spike protein, S gene, mutation, variant, mink

Genomic surveillance of circulating SARS-CoV-2 variants is critical for epidemiologic

tracking and infection control. Rapid global spread of a variant harbouring a mutation located on

the SARS-CoV-2 spike protein (S protein) has been reported.[6,7] In the GISAID database of

SARS-CoV-2 genomes as of March 17, 2021, 95.7% (758,684 of 792,388) of the SARS-CoV-2

genome has the D614G mutation in the spike protein gene; and 22% (178,186 of 792.388) had

N501Y mutation which occurred in the more virulent UK variant (lineage B.1.1.7) with

increased transmissibility by 50% and is potentially more virulent;[1] a similar phenomenon was

observed with the South African variant lineage B.1.351[8] and the Brazil variant lineage

B.1.525,[4,9] both have the N501Y mutation. Large numbers of mutants with Y453F mutations

were found in infected mink in Denmark, which might have been spread from human to animal

and back to human.[5] We analyzed the sequences of the SARS-CoV-2 S protein gene collected

from mink in the USA and Canada, and identified 2 dominant novel variants with 2 and 3

mutations each.

From the GISAID (Global Initiative on Sharing All Influenza Data, https://www.gisaid.org),
we downloaded the following SARS-CoV-2 sequences collected from December 2019 to the
date of writing (March 12, 2021): all animal-derived sequences (977) including those from *Canis
lupus familiaris, Chlorocebus sabaeus, Felis catus, Manis javanica, Manis pentadactyla, Mus
musculus, Mustela lutreola, Neovison vison, Panthera leo, Panthera tigris jacksoni, Rhinolophus
affinis, Rhinolophus malayanus and Rhinolophus shameli*; all sequences collected in Canada
(19,529) and the US (173,277). The SARS-CoV-2 isolate Wuhan-Hu-1 collected in December
19, 2019 (GenBank accession NC045512) was used as a reference for mutation analysis. All
nucleotide position labeling in our study was based on alignment with this sequence. SARS-
CoV-2 full genome sequences were multiple aligned with the bioinformatic software, Geneious
v.11 (Auckland, New Zealand) using "Geneious Multiple Alignment" and "Map to a Reference
Assembly" function. The aligned sequences were visually examined to confirm that they were
aligned properly. The variants/SNP were identified by the software automatically and verified by
visual confirmation. Short fragments (30 nt) containing the novel mutations identified in our
study were used as queries to blast search the local databases consisting of the downloaded
sequences to determine the incidence of the novel mutations using the "BLAST" function of the
Geneious software. The global incidence of "N501Y" was determined using the "search"
function with substitution "N501Y" in GISAID. The global incidence of "N501T" was not
determined since the query of   substitution "N501T" was not available in GISAID.

The N501Y mutation that was common to UK, South African, and Brazil variants, and the
Y453F mutation that widely prevailed in COVID-19 positive mink in Denmark and the
Netherlands were absent in all the US and Canadian mink SARS-CoV-2 sequences examined.
Interestingly, an N501T mutation was found in almost all US mink SARS-CoV-2 sequences (100
of 101; 99%); furthermore, all N501T mutant sequences, except 2 of poor quality, had the
G142D mutation (Supplementary Table 1). Based on GISAID records, all US mink SARS-CoV-
2 sequences were collected from Michigan and Wisconsin in October, 2020. We found that,
except 1 sequences with poor quality, all (54) Michigan mink sequences had the F486L
mutation, and all (55) Michigan mink sequences had the Y144 deletion, in addition to the N501T
and G142D mutations (Supplementary Table 1). Hereafter, we designated the sequences with
N501T and G142D, but without F486L, mutations as the N501T-G142D variant (WI mink
variant); and the ones with all 3 novel mutations as the N501T-G142D-F486L variant (MI mink

variant). In addition to the above 3 mutations and 1 deletion in Spike protein, G614D, the

mutation that dominated the SARS-CoV-2 sequences, existed in all US and Canadian mink-

derived sequences (Supplementary Table 1, Supplementary Table 2). No N501T-G142D nor

N501T-G142D-F486L variants were found in mink from Canada or other countries.

Outside the US, among 335 Netherlands mink-derived SARS-CoV-2 S protein gene

sequences, 5 collected from April to June 2020 had the N501T mutation without G142D or

F486L mutations (Supplementary Table 3), which is consistent with a previous report on N501T

on Netherlands mink-derived sequences.[2] We also found F486L mutation in 60% (204 of 335)

Netherlands mink-derived SARS-CoV-2 S protein gene sequences, none of which with N501T-

G142D variants. The F486L mutation was unique to Netherlands mink-derived sequences

causing by nucleotide mutation T23018C, different from the F486L mutation found in the US

mink-derived sequences due to nucleotide T23020G mutation. Among 454 Denmark mink

SARS-CoV-2 sequences, 3 had N501T mutations as described earlier, none had G142D or

F486L mutations (Supplementary Table 3).

In order to determine the origin of the 2 novel variants, we analyzed all human-derived

SARS-CoV-2 sequences collected from the US and Canada. Three WI N501T-G142D variant

and 2 MI N501T-G142D-F486L variant sequences were identified from the US human-derived

sequences collected from Wisconsin on October 3, 2020 and Michigan on October 6, 2020, and

they had all the mutations as those described above for the mink sequences except lacking of the

Y144 deletion in S gene. Furthermore, we identified N501T mutants without G142D and F486L

in US human-derived sequences collected as early as August 2020, which was earlier than the

mink-derived sequences collected in October, 2020 (Supplementary Table 4, Fig. 1). In Canada,

the N501T mutation only existed in human-derived sequences (Supplementary Table 2). The

N501T mutation was described in 2 Italian human-derived sequences collected in August 2020.[3]

We therefore speculate that the N501T mutation may have evolved during human infection and

was then transmitted to mink populations in the US, and it is less likely that the mutations

occurred in mink population after infected with human strains of the virus. The fact that the two

novel variants were found in almost all infected minks in the US in short period of times within

days also support this speculation.

Fig. 1. Emerging timeline of human-derived (above the timeline) and mink-derived (below the timeline) SARS-CoV-2 spike protein N501T-G142D and N501T-G142D-F486L variants in the US. Timeline scale is not in proportion.

To evaluate the transmissibility of N501T mutants, the incidence of N501T and N501Y mutants was enumerated from US and Canadian GISAID SARS-CoV-2 sequences collected up to March 12, 2021: 2.4% (4,097 of 173,277) and 0.77% (1,339 of 173,277) of US sequences had the N501Y and N501T maturation, respectively; and 1.6% (315/19,529) and 0.06% (12/19,529) of Canadian sequences had the N501Y and the N501T mutation, respectively (Supplementary Table 5). Comparatively, the incidence of N501Y is much lower in the US and Canada than globally. Our analysis results indicate that the N501T mutants had weaker transmissibility compared to the N501Y mutants that include UK B.1.17, South Africa B.1.351, and Brazil P.1. However, it cannot be ruled out that the different time of introduction of N501Y and N501T (April 2020 vs August 2020) may have affected the occurrences of the 2 mutants (Fig. 1).

It has been described that the mutations at S protein N501 can significantly increase the

transmissibility, and likely virulence of SARS-CoV-2; the UK variant N501Y is an example. It

was predicted that a single N501T mutation might significantly enhance the binding affinity

between 2019-nCoV RBD and human ACE2.[10] Our analysis revealed that the N501T and

G142DL mutations occurred in 99% of mink-derived sequences collected in the US; all, except

one with poor quality, mink sequences collected from Michigan also contained the F486L

mutation. The large number of new variants occurring in the mink population warrants further

study on how these changes affected their interaction with the ACE2 receptor, and thereby the

transmissibility, virulence, and immunogenicity in humans and mink. It is important to monitor

the emerging new variants and determine their impact on human and animal health.

Acknowledgement

We gratefully acknowledge the authors, originating and submitting laboratories of the sequences

from GISAID's hCov-19 Database on which this research is based. See supplementary

acknowledgement for details.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship,

and/or publication of this article.

Funding Support:

The authors received no financial support for the research, authorship, and/or publication of this

article.

Supplementary material

Supplementary material for this article is available online.

References

1. ECDC (21 January 2021). "Risk related to the spread of new SARS-CoV-2 variants of concern in the EU/EEA - first update". European Centre for Disease Prevention and Control. Retrieved 2 February 2021.

2. Elaswad A, Fawzy M, Basiouni S, and ShehataAA. Mutational spectra of SARS-CoV-2 isolated from animals PeerJ. 8: e10609. Published online 2020 Dec 18. doi: 10.7717/peerj.10609

3. Fiorentini S et al. First detection of SARS-CoV-2 spike protein N501 mutation in Italy in August, 2020. Lancet Infect Dis 2021 Published Online, January 12, 2021, https://doi.org/10.1016/S1473-3099(21)00007-4 (Retrieved March 12, 2021)

4. Kupferschmidt K. New mutations raise specter of 'immune escape. Science. 371 (6527): 329–330. doi:10.1126/science.371.6527.329. PMID 33479129. Retrieved 25 January 2021.

5. Munnink BBO, et al. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. Science. 2021; 371:172-177 DOI: 10.1126/science.abe5901

6. Na W, Moon H and Song DA. Comprehensive review of SARS-CoV-2 genetic mutations and lessons from animal coronavirus recombination in one health perspective. J Microbiol. 2021. 59, 332–340 https://doi.org/10.1007/s12275-021-0660-4

7. Phan T. Genetic diversity and evolution of SARS-CoV-2. Infect Genet Evol. 2020; 81:104260

8. Tegally H et al. Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa medRxiv 2020.12.21.20248640; doi: https://doi.org/10.1101/2020.12.21.20248640

9. Voloch, CM et al. Genomic characterization of a novel SARS-CoV-2 lineage from Rio de Janeiro, Brazil . Retrieved 15 January 2021. doi:10.1101/2020.12.23.20248598 – via medRxiv.

10. Wan Y, et al. Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. J Virol 2020; 94:e00127-20. https://doi.org/10.1128/JVI.00127-20.