The Preventive Treatment of Migraine with Low-Dose Naltrexone/Acetaminophen Combination: Findings of a Small, Randomized, Double-Blind, and Placebo-Controlled Clinical Trial with an Open-Label Extension for None-Responders

Annette C. Toledano[1]

[1] Sponsor/investigator. Founder/medical director, Allodynic Therapeutics, LLC, North Miami, Florida.

Corresponding author:

Annette Toledano, M.D.

E-mail: annette.toledano@allodynic.com

Phone: 305 895 6808

ABSTRACT

We tested low-dose naltrexone/acetaminophen combination for episodic migraine prevention. We randomly assigned patients to naltrexone/acetaminophen (n=6) or placebo (n=6) for a 12-week double-blind treatment. Non-responders continued into open-label treatment with naltrexone/acetaminophen (n=5) for additional 12 weeks. Patients were adults who experienced 5 to 17 (average 9.7) migraine days at baseline. The primary endpoint was the mean change in the monthly migraine days during the last 4 weeks of the double-blind treatment period. The key secondary endpoint was the mean change in the monthly migraine days from the 4-week double-blind follow-up (2nd baseline) to the last 4 weeks of the open-label treatment period.

The magnitude of the treatment effect for naltrexone/acetaminophen observed in the double-blind period was 2.2 fewer monthly migraine days than placebo (p=0.43).

Four out of 6 (66.7%) naltrexone/acetaminophen-treated patients experienced 75% reduction in migraine days compared to 1 out of 6 (16.7%) placebo-treated patients (p=0.09).

In the open-label phase, treatment with naltrexone/acetaminophen (n=5) led to 8.2 fewer mean monthly migraine days (from 11.8 to 3.6), representing 69.5% improvement (p=0.03), and 100% of the patients experienced a 50% reduction in monthly migraine days.

Adverse events were mild/moderate and transient, included dry mouth, fatigue, sedation, nausea, and feeling jittery.

We postulate that naltrexone's toll-like receptor (TLR4) antagonism properties prevent pro-inflammatory cytokines' production in the trigeminal ganglion averting "overactive nerves" and migraine. Although this trial used low-dose naltrexone (defined as 1 – 5 mg/day), we postulate mid-dose naltrexone (MDN) (defined as 6 – 10 mg/day) may offer a greater migraine prevention control.

Keywords: Pain; Migraine prevention; Headache; Naltrexone; Analgesia; Acetaminophen; Low-Dose Naltrexone; Mid-Dose Naltrexone; Toll-Like Receptor 4 Antagonist; TLR4 Antagonist; Neuropathic Pain; Cytokine Storm.

INTRODUCTION

The estimated global prevalence of migraine is 14.7% (that's around 1 in 7 people).[1] In the global burden of disease study, updated in 2013, migraine was the sixth highest

cause worldwide of years lost due to disability (YLD).[2] In the United States, approximately 38 million Americans are afflicted by migraine, and available treatments do not adequately meet the needs of many. Hence, there is a need for new treatments for preventing migraine. Oral naltrexone/acetaminophen combination, if proven effective, may provide greater efficacy/tolerability ratio than existing migraine prevention treatments.

Naltrexone,[3] an opioid antagonist approved for addiction treatment, is also an analgesic due to recently-discovered toll-like receptor 4 (TLR4) antagonism properties. Inhibiting the TLR4 with naltrexone in nerve cells of the dorsal root ganglia (DRG) and trigeminal ganglion led to reduction in pro-inflammatory cytokines' production and reversal of neuropathic pain and migraine in animal studies.[4–10] Naltrexone can prevent a "localized cytokine storm" (our term) in nerve cells averting pain. We postulate that naltrexone's toll-like receptor (TLR4) antagonism properties prevent pro-inflammatory cytokines' production in the trigeminal ganglion averting "overactive nerves" and migraine.

The Interagency Pain Research Coordinating Committee (IPRCC) voted a study that used naloxone[5] (a opioid/TLR4 antagonist similar to naltrexone) as one of 2009-2013 pain research advances that represent significant progress in the field. "This research supports TLR4 as a potential therapeutic target for treating chronic pain in patients, and, as the establishment of a completely new class of pain-relieving medication, would be a remarkable advance in pain treatment."[11]

TLR4 is an innate immune system receptor that usually detects invasion of foreign agents such as viruses and initiates a cascade leading to cytokines' production to eliminate them. However, the TLR4 can also be triggered by endogenous damage

molecules originating from injured tissues (e.g., a herniated intervertebral disc) and lead to cytokines' production and pain.[12] The inborn, innate immune system is distinct from the learned, adaptive immune system.

Naltrexone's analgesic properties are due to its inhibition of the production of pro-inflammatory cytokines, e.g., interleukin (IL)-6, tumor necrosis factor (TNF)-α, interferon-β, calcitonin gene-related peptide (CGRP), nitric oxide (NO), and reactive oxygen species (ROS) in nerve cells, [5,8,13,14] preventing a "localized cytokine storm" (our term) and pain generation. Naltrexone exerts its action at the beginning of the cascade leading to the production of many pro-inflammatory cytokines blocking their creation. Drugs that neutralize individual cytokines after their formation include adalimumab (Humira), a TNF-α blocker; anakinra (Kineret), an IL-1 blocker; Tocilizumab (Actemra), an IL-6 blocker; and rimegepant, (Nurtec), a CGRP blocker. Naltrexone can prevent the production of multiple cytokines eliminating the need to neutralize them. But once created, naltrexone, cannot stop these cytokines.

We hypothesize that the pathophysiological event underlying migraine is the excessive production of pro-inflammatory cytokines in the trigeminal ganglion creating a neuro-inflammatory response resulting in a "localized cytokine storm." Similarly, in the dorsal root ganglions, a "localized cytokine storm" results in neuropathic back pain. We postulate that a localized "cytokine storm" is the underlying event leading to "overactive nerves" (layman's term) and nerve pain.

Pro-inflammatory cytokines exaggerate neuronal excitability, contributing to neuropathic pain and migraine. Activation of TLR4 has been implicated in the pathogenesis of

migraine[9,15] and (+)-Naltrexone blocked the development of facial allodynia in modeled migraine in rats.[9]

Dr. Bernard Bihari invented Low-Dose Naltrexone (LDN) (a daily dose of 1 to 5 mg) in the mid-1980s for "normalizing the immune system function."[16,17] However, scientists discovered the innate immune system and TLRs in humans in the 1990s (a Nobel Prize was awarded in 2011).[18] The prevailing theory for LDN's mechanism of action was that it increases endorphin production, systemically upregulating endogenous opioid signaling by a transient opioid-receptor blockade.[19] later research attributed the analgesic properties of naltrexone to TLR4 antagonism leading to pro-inflammatory cytokines' production inhibition. Currently, LDN is widely accepted as an alternative medicine modality and is used by its proponents to treat various medical conditions. It is almost sold as an everyday supplement by certain pharmacies.[19,20]

Grassroots interest in off-label LDN sprang clinical trials for fibromyalgia, multiple sclerosis, Crohn's disease, and complex regional pain syndrome (CRPS).[21–25]

Although LDN is reportedly used as an off-label treatment for various medical conditions, there are no confirmatory studies for these off-label uses. Addiction specialists are the primary prescribers of naltrexone.

Acetaminophen potentially enhances the combination in several ways. Acetaminophen created synergy for analgesia in combination drugs such as Vicodin (hydrocodone/ acetaminophen).[26] Similarly, combining naltrexone with acetaminophen could attain synergy for analgesia. Acetaminophen 1000 mg was already established as an effective acute migraine medication and can therefore enhance the combination's treatment

effect on migraine.[27,28] Acetaminophen has the public's trust as the world's most used drug. Acetaminophen was found to be also an emotional pain reducer. Acetaminophen significantly reduced hurt feelings in human studies.[29,30] Acetaminophen's emotional pain reducing properties could potentially be augmented by naltrexone[31] and enhance the combination's effect on the overall sense of well-being.

Naltrexone's established cytokine production inhibition properties in the trigeminal and dorsal root ganglia averting pain could theoretically reduce pain symptoms associated with COVID-19 and COVID-19 vaccination. Three experimental TLR4 antagonists FP7, Eritoran, and retrocyclin 101, were significantly better than placebo in treating lethal influenza.[32–34] TLR4 signaling is a key disease pathway controlling the severity of acute lung injury.[35] Naltrexone, a readily available TLR4 antagonist, needs to be evaluated for the treatment of pain symptoms in COVID-19.

Interestingly, the innate immune system's pathway that mounts an initial response to viral infections is the same one that leads to neuropathic pain when triggered intrinsically by damage-molecules.

Naltrexone/acetaminophen combines naltrexone's unutilized analgesic properties with the established and well-trusted analgesic properties of acetaminophen.

METHODS

Study Population

Patients were recruited to our single site in Miami, Florida, through billboards and social media advertising. The study was conducted between August 25, 2017, and July 26, 2018. ANODYNE-3, (the study's name) enrolled patients 18 to 75 years of age with a

history of migraine with or without aura for at least one year that was consistent with the diagnosis criteria of the International Classification of Headache Disorders (ICHD)-3rd edition (beta version)[36] and have experienced 5 to 17 migraine days in the 4-week baseline. The study population included 11 patients with episodic migraine and one with chronic migraine. Patients needed to have at least 5 migraine days during the baseline. Patients using opioid medications or who had a history within the previous 3 years of drug abuse were excluded.

Our goal was to enroll 48 patients, but we were unable to meet that goal due to hurricane Irma's (September 2017) impact on our community.

Trial Oversight

The sponsor/investigator – Annette Toledano, M.D., was responsible for all the trial elements, including design, execution, data collection, analysis, and interpretation.

The trial's protocol and informed consent were approved by the Schulman Associates IRB (now Advarra). All patients provided written informed consent before starting the study procedures. The informed consent informed the patients of the conflict of interest of the sponsor/investigator. The study patients were compensated for participation.

This study was conducted under an Investigational New Drug (IND) application with the United States Food and Drug Administration (FDA).

Trial Design

The ANODYNE-3 trial was a phase 2, single-site, randomized, double-blind, and placebo-controlled proof-of-concept study. We evaluated the patients in seven site visits during the double-blind phase and four visits during the open-label phase. The visits

included: baseline (week -4), randomization (week 0), and double-blind treatment

(weeks 1, 4, 8, and 12), follow-up (week 16), open-label treatment (weeks 20, 24, and

28), and open-label follow-up (week 32) (Figure 1).

4-week Baseline	12-week Double-blind (DB) treatment with naltrexone/acetaminophen or placebo	4-week Follow-up	12-week Open-label treatment with naltrexone/acetaminophen for DB none-responders	4-week Open-label Follow-up

Figure 1: Study design flow-chart

We randomly assigned patients in 1:1 ratio to naltrexone 2.25 mg/acetaminophen 325

mg (n=6) or placebo (n=6) BID for a 12-week double-blind treatment period. Non-

responders (defined as ≤50% reduction in monthly migraine days in the last 4 weeks of

treatment) continued into open-label naltrexone/acetaminophen treatment (n=5) for

additional 12 weeks. The open-label phase intended to gain insight into a "real world"

response to the drug.

Patients recorded in a daily paper diary the duration and headache severity. They also

recorded photophobia, phonophobia, nausea, vomiting, and any acute migraine

medication use.

The study pharmacist prepared the study medication from marketed tablets placed in

two single-ingredient capsules backfilled with microcrystalline cellulose. The study

pharmacist devised the randomization schedule using a block size of 12. The study

medication kits were sequentially numbered, and the investigator assigned the kits to

patients consecutively. Patients took the study medication at approximately 8 am and 8

pm every day. At each site visit, the patients received enough medication to cover until

the next visit, plus three extra days in case of a delayed appointment. Patients returned
bottles and extra capsules at each site visit for drug accountability.

Efficacy Assessments and Endpoint

The primary efficacy endpoint was the mean change in monthly migraine days from
baseline to the last 4 weeks of the double-blind treatment period. The key secondary
efficacy endpoint was the mean change in monthly migraine days from the double-blind
follow-up (2nd baseline) to the last 4 weeks of the open-label treatment period.

Additional secondary endpoints were the proportion of patients achieving at least 50%,
75%, and 100% reduction in monthly migraine days, acute migraine medication
treatment days, migraine headache-hours, at least moderate migraines days, and
severe migraine days from baseline to the last 4 weeks of treatment. Other endpoints
were Headache Impact Test (HIT-6), Pittsburgh Insomnia Rating Scale-20 (PIRS-20),
and Patient Global Impression of Change (PGIC). We assessed treatment-emergent
adverse events and serious adverse events. We performed a physical examination,
obtained vital signs (systolic and diastolic blood pressure, pulse, and weight), and
recorded concomitant medications during each site visit. The patients had a 12-lead
electrocardiogram at the screening visit and monthly clinical laboratory tests (serum
chemistry and hematology).

Statistical Analysis

The efficacy and safety analyses included all randomized patients who received at least
one dose of the study medication, recorded a baseline migraine severity rating, and
reported at least one post-dose assessment. All statistical tests were 2-sided hypothesis

tests performed at the 5% significance level. Statistical analyses were conducted using MedCalc statistical software.[37]

RESULTS

Patients

We assessed 94 patients for eligibility; 12 patients met the inclusion criteria and were randomized. Six patients received naltrexone 2.25 mg/acetaminophen 325 mg and 6 received placebo. All the 12 randomized patients completed the double-blind treatment period, and 5 of 6 non-responders completed the open-label treatment period (Figure 2).

The baseline characteristics of the patients were similar in the two groups (Table 1). The mean age was 40.3 years, and the trial population was predominantly female (83.3%) and Caucasian (91.7%). Migraine histories were similar in terms of migraine duration, the number of migraine days per month in the previous 3 months, and the monthly migraine days with acute medication use over the preceding 3 months. Acute migraine medication classes, specifically triptans – were identical in both groups. Migraine characteristics were similar among the treatment groups during the baseline period. The mean number of headache days in the 4-week baseline was 9.7 for both groups and the days of acute migraine medication use were 8.2 and 7.7 for naltrexone/acetaminophen and placebo, respectively. Headache impact test (HIT) and Pittsburgh Insomnia Rating Scale-20 (PIRS-20) scores were comparable at the screening visit.

Figure 2: Patients' distribution

Table 1: Baseline characteristics

	Total (n=12)	Naltrexone 2.25 mg/ acetaminophen 325 mg (n=6)	Placebo (n=6)
Characteristics			
Age - years	40.3±12.0	41.5±14.3	39.0±10.4
Female gender - no. (%)	10 (83.3)	4 (66.7)	6 (100)
Race or ethnic group - no. (%)†			
Caucasian, no. (%)	11 (91.7)	6 (100.0)	5 (83.3)
African American, no. (%)	1 (8.3)	0 (0.0)	1 (16.7)
Hispanic ethnic group, no. (%)	8 (66.7)	4 (66.7)	4 (66.7)
Body mass index‡	26.8±7.0	26.9±6.0	26.7±8.4
Migraine History			
Duration of migraine illness - year	17.0±13.1	20.3±12.4	13.7±13.8
Migraine days/mth in past 3 months – no.	8.9±1.9	8.8±2.3	9.0±1.5
Migraine days/month with acute medication use in past 3 months, no.	8.5±2.2	8.2±2.6	8.8±1.9
Previous use of preventive medications - no. %	3 (25.0)	1 (16.7)	2 (33.3)
Acute migraine medication classes			
Combination analgesics - no. (%)	5 (41.7)	2 (33.3)	3 (50.0)
Non-steroidal anti-inflammatory drugs - no. (%)	2 (16.7)	2 (33.3)	0 (0.0)
Acetaminophen - no. (%)	1 (8.3)	0 (0.0)	1 (16.7)
Triptans - no. (%)	4 (33.3)	2 (33.3)	2 (33.3)
Migraine characteristics during baseline			
Migraine days per month§	9.7±3.3	9.7±4.2	9.7±2.9
Acute migraine medication days/month	8.2±2.7	7.7±2.1	8.7±3.5
Monthly migraine hours, avg./patient	62.6±41.2	70.8±47.6	54.4±36.2
Headache Impact Test (HIT-6) score at screening¶	70.2±5.2	70.0±6.3	70.2±4.9
Pittsburgh Insomnia Rating Scale-20 (PIRS-20) score at screening**	32.0±13.0	35.7±13.9	27.2±13.3

*Plus–minus values are means ±SD
†Race and ethnic group were reported by the patient.
‡Calculated as weight in kilograms divided by height in meters squared.
§ A migraine day was defined as a calendar day in which a patient reported a headache of ≥ 30-minute duration and met criteria for migraine with or without aura.
¶ Headache Impact Test (HIT-6). Scores range from 36 to 78, with a score of over 50 indicating disability. **Pittsburgh Insomnia Rating Scale-20 (PIRS-20). PIRS-20 scores range from 0 to 60, with scores above 20 suggesting insomnia.

Efficacy Results

Comparing the baseline to the last 4-weeks of the double-blind period, treatment with naltrexone/acetaminophen (n=6) led to 5.7 fewer monthly migraine days, whereas treatment with placebo led to 3.5 fewer monthly migraine days, both groups improving from a baseline rate of 9.7 monthly migraine days. The magnitude of the treatment effect observed in the double-blind period was 2.2 fewer monthly migraine days (p=0.43) (Table 2 and Figure 3). In the double-blind period, 66.7% of the naltrexone/acetaminophen patients experienced a 50% reduction in migraine headache days from baseline to the last 4 weeks of the double-blind period compared to 33.3% in placebo patients (p=0.27). 66.7% of naltrexone/acetaminophen patients experienced 75% reduction in migraine days compared to 16.7% of placebo patients, (p=0.09), and 50.0% of the naltrexone/acetaminophen-treated patients experienced 100% reduction in monthly migraine days compared to 16.7% of placebo patients, (p=0.27) (Table 2).

The magnitude of the treatment effect from baseline to the last 4 weeks of the double-blind treatment for Headache Impact Test (HIT) score was -9.8 points (p=0.29), and for Pittsburgh Insomnia Rating Scale-20, it was -11.7 (p=0.27). The magnitude of the treatment effect in monthly migraine hours in the double-blind period was a 28.8 hours reduction (p=0.35).

In the open-label period, treatment with naltrexone/acetaminophen (n=5) led to 8.20 fewer monthly migraine days (from 11.80 to 3.60), amounting to 69.5% improvement, (p=0.03) (Figure 4). Also, 100% of the patients experienced a 50% reduction in the monthly migraine days (Table 3). The treatment effect from the double-blind follow-up to the last 4 weeks of the open-label period for Headache Impact Test and Pittsburgh

Insomnia Rating Scale-20 was -25.8 (p=0.0002) and -17.0 (p=0.09) points, respectively. The treatment effect in monthly migraine hours in the open-label period was -36.2 hours (p=0.09). In the open-label period, there were statistically significant reductions from baseline to the last 4 weeks of treatment in the days of use of any acute headache medications, Headache Impact Test (HIT), at least moderate migraine days, and severe headache days.

The improvements in Headache Impact Test (HIT) and Pittsburgh Insomnia Rating Scale-20 (PIRS-20) indicate enhancement of quality of life measures.

Acetaminophen-overuse head could be expected in patients having more than 15 headaches per month who consume acetaminophen more than 15 days per month for more than 3 months.[36]

Since the study administered acetaminophen 650 mg daily, we were concerned that acetaminophen may induce medication overuse headaches. The study provided data on 11 three-month treatment courses with naltrexone/acetaminophen for nine unique patients and two patients who received two consecutive courses separated by 4 weeks. During each of these courses, migraine days were lower than the baseline (Figure 5 and Figure 6). This study provided preliminary evidence that a daily dose of naltrexone 4.5 mg/acetaminophen 650 mg did not induce medication overuse headache after three months of treatment.

Table 2: Endpoints from baseline to month-3 of the double-blind treatment period*

	Naltrexone 2.25 mg/ acetaminophen 325 mg (n=6)	Placebo (n=6)	Diff. v. placebo	p-value
Migraine days per month§ no.	-5.7±3.2	-3.5±5.6	-2.2	0.43
≥50% reduction in monthly migraine days, no. (%)	4 (66.7)	2 (33.3)	2 (33.3)	0.27
≥75% reduction in monthly migraine days, no. (%)	4 (66.7)	1 (16.7)	3 (50.0)	0.09
100% reduction in monthly migraine days, no. (%)	3 (50.0)	1 (16.7)	2 (33.3)	0.29
Days of any acute headache medication use per month, no.	-5.0±5.4	-3.2± 5.5	-1.8	0.57
≥75% reduction in days of any acute headache medication use, no (%)	4 (66.7)	1 (16.7)	3 (50.0)	0.09
Headache Impact Test (HIT-6) score¶	-20.7±19.1	-10.8±13.0	-9.8	0.30
≥50% reduction in HIT-6, no. (%)	4 (66.7)	1 (16.7)	3 (50.0)	0.09
Pittsburgh Insomnia Rating Scale-20 (PIRS-20)** score	-22.7± 22.6	-11.0±8.9	-11.7	0.27
≥50% reduction in PIRS-20 score, no. (%)	5 (83.4)	2 (33.3)	3 (50.0)	0.27
Patient Global Impression of Change (PGIC)‡ change ≥2 points, (%)	4 (66.7)	2 (33.3)	2 (33.3)	0.27
Patient Global Impression of Change (PGIC)‡ change=3 points, (%)	4 (66.7)	2 (33.3)	2 (33.3)	0.27
Monthly migraine hours, avg./patient	-46.1±59.4	-17.4±41.1	-28.8	0.35
≥70% reduction in monthly migraine hours, no. (%)	4 (66.7)	1 (16.7)	3 (50.0)	0.09
At least moderate headache days, no.	-4.8±4.4	-2.8±5.)	-2.0	0.52
≥75% reduction in at least moderate migraine days, no. (%)	4 (66.7)	1 (16.7)	3 (50.0)	0.09
Severe headache days, no.	-2.2±2.8	-1.8±3.9	-0.3	0.86

*Plus–minus values are means ±SD. The analysis included all the 12 patients who were randomized (all of them completed the study).
§ A migraine day was defined as a calendar day in which a patient reported a headache of ≥ 30-minute duration and met criteria for migraine with or without aura.
¶ Headache Impact Test (HIT-6). Scores range from 36 to 78, with a score of over 50 indicating disability.
** Pittsburgh Insomnia Rating Scale-20 (PIRS-20). Scores range from 0 to 60, with scores above 20 suggesting insomnia.
‡ Patient Global Impression of Change PGIC: 1=Very much worse. 2=Much worse. 3=minimally worse. 4=no change. 5=minimally improved 6=Much improved. 7=very much improved.
p-value - 95% confidence intervals

Table 3: Endpoints from double-blind follow-up (2nd baseline) to month-3 of open-label treatment period*

	Double-blind follow-up (baseline) (n=5)	last 4 weeks of open-label treatment (n=5)	Difference from baseline	p-value
Migraine days per month, § no.	11.8±6.0	3.6±3.7	-8.2±3.9	0.03
≥50% reduction in monthly migraine days, no. (%)		5 (100)	5 (100)	n/a
≥75% reduction in monthly migraine days, no. (%)		3 (60)	3 (60)	n/a
100% reduction in monthly migraine days, no. (%)		3 (60)	3 (60)	n/a
Days of use of any acute headache medications per month, no.	9.8±4.3	1.6± 2.5	-8.2±2.8	0.0063
Headache Impact Test (HIT-6) score¶	68.2±5.7	42.4± 7.1	-25.8±4.4	0.0002
HIT-6 Mean % response (adjusted for score of 36 being 100% improvement)			80% (15%)	0.0002
** Pittsburgh Insomnia Rating Scale-20 (PIRS-20) score	25.0±18.5	8.0±7.0	-17.0±17.2	0.09
Patient Global Impression of Change (PGIC)‡ change ≥2 points, (%)		5, (100)	5, (100)	n/a
Patient Global Impression of Change (PGIC)‡ change=3 points, (%)		4, (80)	4, (80)	n/a
Monthly migraine hours, no.	55.3±34.5	19.2±24.6	-36.2±32.4	0.09
At least moderate migraine days, no.	10.8±5.0	2.6±2.7	-8.1±4.2	0.01
Severe headache days, no.	6.8±4.8	1.0±1.2	-5.8±3.4	0.03

* Plus–minus values are means ±SD. The analysis included 5 of the 6 patients who were non-responders in the double-blind period. Non-responders were defined as ≤50% reduction in monthly migraine days in the last 4 weeks of treatment.
§ A migraine day was defined as a calendar day in which a patient reported a headache of ≥ 30-minute duration and met criteria for migraine with or without aura.
¶ Headache Impact Test (HIT-6). Scores range from 36 to 78, with a score of over 50 indicating disability.
** Pittsburgh Insomnia Rating Scale-20 (PIRS-20). Scores range from 0 to 60, with scores above 20 suggesting insomnia.
‡ Patient Global Impression of Change (PGIC): 1=Very much worse. 2=Much worse. 3=minimally worse. 4=no change. 5=minimally improved 6=Much improved. 7=very much improved.
p-value - 95% confidence intervals

Figure 3: Change in mean monthly migraine days from baseline in the double-blind phase.

Figure 4: Mean monthly migraine days from baseline in the open-label phase.

Figure 5: Monthly migraine days for individual naltrexone/acetaminophen-treated patients in the double-blind phase.

Figure 6: Monthly migraine days in individual naltrexone/acetaminophen-treated patients in the open-label phase. BL=Baseline.

Safety

Treatment-emergent adverse events were reported by 66.7% (4 of 6) of patients in the naltrexone 2.25 mg/acetaminophen 325 mg group, 33.3% (2 of 6) in the placebo group, and 60.0% (3 of 5) in the open-label period. (Table 4).

Dry mouth, fatigue, sedation, nausea, and feeling jittery were the most commonly reported events. All the adverse events were mild/moderate in severity and resolved within 1 to 3 days. No serious adverse events were reported. No deaths or discontinuations due to an adverse event were reported. No changes in monthly liver transaminases or bilirubin were observed.

Table 4: Treatment-emergent adverse events*

	Naltrexone 2.25 mg/ acetaminophen 325 mg (n=6)	Placebo (n=6)	Un-blinded Naltrexone 2.25 mg/ acetaminophen 325 mg (n=5)
Death — no. (%)	0	0	0
Serious adverse event — no. (%)	0	0	0
Discontinuation due to adverse event - no. (%)	0	0	0
Patients ≥1 Adverse event during the trial — no. (%)	4 (66.7)	2 (33.3)	3 (60)
All adverse events*			
Dry Mouth — no. (%)	1 (16.7)		1 (20.0)
Fatigue — no. (%)	1 (16.7)	1 (16.7)	1 (20.0)
Feeling Jittery			1 (20.0)
Gastroesophageal reflux — no. (%)	1 (16.7)		
Heavy breathing — no. (%)	1 (16.7)	1 (16.7)	
Nausea — no. (%)	1 (16.7)		
Palpitations, no. (%)	1 (16.7)		
Sedation, no. (%)	1 (16.7)	1 (16.7)	1 (20.0)
Tinnitus, no. (%)			1 (20.0)
Vomiting, no. (%)	1 (16.7)		
Total adverse events, no.	8	3	5
* All the reported adverse events.			

DISCUSSION

In the preventive treatment of migraine, naltrexone/acetaminophen demonstrated higher response rates than placebo in several key endpoints. The double-blind and open-label phase results together suggest naltrexone/acetaminophen may be useful in the preventive treatment of migraine.

There were fewer migraine days per month in the double-blind phase and higher proportions of patients experienced a 50%, 75%, and 100% reduction in monthly migraine days in the naltrexone/acetaminophen-treated group. Naltrexone/acetaminophen improved the quality of life compared with placebo, as indicated by the Headache Impact Test, Pittsburgh Insomnia Rating Scale-20, and the Patient Global Impression of Change.

The open-label treatment phase, representing "real world" response, demonstrated statistically significant reductions in migraine days per month, days of any acute headache medications use, Headache Impact Test score, at least moderate migraine days, and severe headache days.

The daily dose of naltrexone in this study was approximately 1/10th of the approved addiction treatment dose. The daily acetaminophen dose was 650 mg.

Although this trial used low-dose naltrexone (defined as 1 – 5 mg/day), we postulate mid-dose naltrexone (MDN) (defined as 6 – 10 mg/day) may offer a greater migraine prevention control.

Limitations of this trial included its tiny sample size and treatment periods of just 3 months. The treatment period needed to be longer than 3 months to assess the acetaminophen-overuse headache risk.

Targeting the toll-like receptor-4 represents a novel approach to treating migraine. Adding acetaminophen, the world's most used drug, could enhance the combination's analgesic effect and boost the public's trust conferring an advantage over naltrexone alone. This trial provides preliminary evidence for the potential benefits of naltrexone as a migraine prevention treatment.

This study was the first to assess a naltrexone product for migraine prevention.

Adequately powered clinical trials are needed to confirm these findings. The study's sample size was too small to achieve statistical significance for the primary endpoint.

PATENTS

The naltrexone/acetaminophen combination received two U.S. patents for treating pain and a U.S. patent for treating emotional pain.

Author Contribution: Concept, design, statistical analysis, and data interpretation: Annette Toledano.

Dr. Toledano had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study Pharmacist: Ayman Mohamed PharmD.

Conflict Of Interest Disclosures: Dr. Annette Toledano reports she is the founder/medical director of Allodynic Therapeutics, LLC and the inventor/patent holder

of several naltrexone/acetaminophen patents. Allodynic Therapeutics is a clinical-stage, specialty Biopharmaceutical Company focused on painful conditions with high unmet needs.

Funding/Support: This trial was fully funded by Allodynic Therapeutics, LLC.

Disclaimer: The opinions expressed in this article are those of the author.

Grants: None

Trial Registration: ClinicalTrials.gov Identifier**:** NCT03194555.

Institutinal Review Board (IRB): This trial's protocol and informed consent were approved by the Schulman Associates IRB (now Advarra).

Investigational New Drug Application: This study was conducted under an Investigational New Drug (IND) application with the United States Food and Drug Administration (FDA).

Acknowledgments: We acknowledge the immense contribution of the study patients, the study pharmacist, and the clinical research coordinators, whose hard work made this trial possible.

REFERENCES

1. Steiner TJ, Stovner LJ, Birbeck GL. Migraine: the seventh disabler. *J Headache Pain.* 2013;14(1):1. doi:10.1186/1129-2377-14-1

2. Headache disorders. Accessed December 9, 2020. https://www.who.int/news-room/fact-sheets/detail/headache-disorders

3. FDA. REVIA® (naltrexone hydrochloride tablets USP) 50 mg Opioid Antagonist. FDA. Accessed August 29, 2020. https://www.accessdata.fda.gov/drugsatfda_docs/label/2013/018932s017lbl.pdf

4. Hutchinson MR, Zhang Y, Brown K, et al. Non-stereoselective reversal of neuropathic pain by naloxone and naltrexone. *Eur J Neurosci*. 2008;28(1):20-29. doi:10.1111/j.1460-9568.2008.06321.x

5. Nicotra L, Loram LC, Watkins LR, Hutchinson MR. Toll-like receptors in chronic pain. *Exp Neurol*. 2012;234(2):316-329. doi:10.1016/j.expneurol.2011.09.038

6. Ellis A, Wieseler J, Favret J, et al. Systemic Administration of Propentofylline, Ibudilast, and (+)-Naltrexone Each Reverses Mechanical Allodynia in a Novel Rat Model of Central Neuropathic Pain. *J Pain Off J Am Pain Soc*. 2014;15(4):407-421. doi:10.1016/j.jpain.2013.12.007

7. Lewis SS, Loram LC, Hutchinson MR, et al. (+)-naloxone, an opioid-inactive toll-like receptor 4 signaling inhibitor, reverses multiple models of chronic neuropathic pain in rats. *J Pain Off J Am Pain Soc*. 2012;13(5):498-506. doi:10.1016/j.jpain.2012.02.005

8. Wang X, Zhang Y, Peng Y, et al. Pharmacological characterization of the opioid inactive isomers (+)-naltrexone and (+)-naloxone as antagonists of toll-like receptor 4. *Br J Pharmacol*. 2016;173(5):856-869. doi:10.1111/bph.13394

9. Wieseler J, Ellis A, McFadden A, et al. Supradural inflammatory soup in awake and freely moving rats induces facial allodynia that is blocked by putative immune modulators. *Brain Res*. 2017;1664:87-94. doi:10.1016/j.brainres.2017.03.011

10. Selfridge BR, Wang X, Zhang Y, et al. Structure-Activity Relationships of (+)-Naltrexone-Inspired Toll-Like Receptor 4 (TLR4) Antagonists. *J Med Chem*. 2015;58(12):5038-5052. doi:10.1021/acs.jmedchem.5b00426

11. 2009-2013 Pain Research Advances | Interagency Pain Research Coordinating Committee. Accessed October 28, 2020. https://www.iprcc.nih.gov/Pain-Research/Pain-Research-Advances/2009-2013-Research-Advances

12. Kato J, Svensson CI. Role of extracellular damage-associated molecular pattern molecules (DAMPs) as mediators of persistent pain. *Prog Mol Biol Transl Sci*. 2015;131:251-279. doi:10.1016/bs.pmbts.2014.11.014

13. Parkitny L, Younger J. Reduced Pro-Inflammatory Cytokines after Eight Weeks of Low-Dose Naltrexone for Fibromyalgia. *Biomedicines*. 2017;5(2). doi:10.3390/biomedicines5020016

14. Cant R, Dalgleish AG, Allen RL. Naltrexone Inhibits IL-6 and TNFα Production in Human Immune Cell Subsets following Stimulation with Ligands for Intracellular Toll-Like Receptors. *Front Immunol*. 2017;8. doi:10.3389/fimmu.2017.00809

15. Su M, Ran Y, He Z, et al. Inhibition of toll-like receptor 4 alleviates hyperalgesia induced by acute dural inflammation in experimental migraine. *Mol Pain*. 2018;14:1744806918754612. doi:10.1177/1744806918754612

16. *Low Dose Naltrexone (LDN) Pioneer Dr. Bernard Bihari Talks About His Life.*; 2011. Accessed October 26, 2020. https://www.youtube.com/watch?v=x54Jccr8GT8

17. Bihari B. Bernard Bihari, MD: Low-dose Naltrexone for Normalizing Immune System Function. *Altern Ther Health Med 2013;19:56–65.*

18. The Nobel Prize in Physiology or Medicine 2011. NobelPrize.org. Accessed October 26, 2020. https://www.nobelprize.org/prizes/medicine/2011/popular-information/

19. Toljan K, Vrooman B. Low-Dose Naltrexone (LDN)—Review of Therapeutic Utilization. *Med Sci.* 2018;6(4). doi:10.3390/medsci6040082

20. Brown N, Panksepp J. Low-dose naltrexone for disease prevention and quality of life. *Med Hypotheses.* 2009;72(3):333-337. doi:10.1016/j.mehy.2008.06.048

21. Younger J, Mackey S. Fibromyalgia symptoms are reduced by low-dose naltrexone: a pilot study. *Pain Med Malden Mass.* 2009;10(4):663-672. doi:10.1111/j.1526-4637.2009.00613.x

22. Younger J, Noor N, McCue R, Mackey S. Low-dose naltrexone for the treatment of fibromyalgia: findings of a small, randomized, double-blind, placebo-controlled, counterbalanced, crossover trial assessing daily pain levels. *Arthritis Rheum.* 2013;65(2):529-538. doi:10.1002/art.37734

23. Cree BAC, Kornyeyeva E, Goodin DS. Pilot trial of low-dose naltrexone and quality of life in multiple sclerosis. *Ann Neurol.* 2010;68(2):145-150. doi:10.1002/ana.22006

24. Smith JP, Stock H, Bingaman S, Mauger D, Rogosnitzky M, Zagon IS. Low-dose naltrexone therapy improves active Crohn's disease. *Am J Gastroenterol.* 2007;102(4):820-828. doi:10.1111/j.1572-0241.2007.01045.x

25. Chopra P, Cooper MS. Treatment of Complex Regional Pain Syndrome (CRPS) Using Low Dose Naltrexone (LDN). *J Neuroimmune Pharmacol.* 2013;8(3):470-476. doi:10.1007/s11481-013-9451-y

26. Beaver WT. Combination Analgesics. *Am J Med.* 1984;77(3):38-53. doi:10.1016/S0002-9343(84)80101-1

27. Marmura MJ, Silberstein SD, Schwedt TJ. The acute treatment of migraine in adults: the american headache society evidence assessment of migraine pharmacotherapies. *Headache.* 2015;55(1):3-20. doi:10.1111/head.12499

28. Lipton RB, Baggish JS, Stewart WF, Codispoti JR, Fu M. Efficacy and Safety of Acetaminophen in the Treatment of Migraine: Results of a Randomized, Double-

blind, Placebo-Controlled, Population-Based Study. *Arch Intern Med*. 2000;160(22):3486-3492. doi:10.1001/archinte.160.22.3486

29. Dewall CN, Macdonald G, Webster GD, et al. Acetaminophen reduces social pain: behavioral and neural evidence. *Psychol Sci*. 2010;21(7):931-937. doi:10.1177/0956797610374741

30. Mischkowski D, Crocker J, Way BM. A Social Analgesic? Acetaminophen (Paracetamol) Reduces Positive Empathy. *Front Psychol*. 2019;10. doi:10.3389/fpsyg.2019.00538

31. Wardle MC, Bershad AK, de Wit H. Naltrexone alters the processing of social and emotional stimuli in healthy adults. *Soc Neurosci*. 2016;11(6):579-591. doi:10.1080/17470919.2015.1136355

32. Perrin-Cocon L, Aublin-Gex A, Sestito SE, et al. TLR4 antagonist FP7 inhibits LPS-induced cytokine production and glycolytic reprogramming in dendritic cells, and protects mice from lethal influenza infection. *Sci Rep*. 2017;7:40791. doi:10.1038/srep40791

33. Shirey KA, Lai W, Scott AJ, et al. The TLR4 antagonist Eritoran protects mice from lethal influenza infection. *Nature*. 2013;497(7450):498-502. doi:10.1038/nature12118

34. Prantner D, Shirey KA, Lai W, et al. The θ-defensin retrocyclin 101 inhibits TLR4- and TLR2-dependent signaling and protects mice against influenza infection. *J Leukoc Biol*. 2017;102(4):1103-1113. doi:10.1189/jlb.2A1215-567RR

35. Imai Y, Kuba K, Neely GG, et al. Identification of oxidative stress and Toll-like receptor 4 signaling as a key pathway of acute lung injury. *Cell*. 2008;133(2):235-249. doi:10.1016/j.cell.2008.02.043

36. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. 2013;33(9):629-808. doi:10.1177/0333102413485658

37. Schoonjans F. MedCalc's Comparison of proportions calculator. MedCalc. Accessed August 31, 2020. https://www.medcalc.org/calc/comparison_of_proportions.php