Title: Disparities in SARS-CoV-2 seroprevalence among individuals presenting for care in central North Carolina over a six-month period

Authors: Cesar A. Lopez[1*], Clark H. Cunningham[2*] Sierra Pugh[3], Katerina Brandt[4], Usaphea P. Vanna[1], Matthew J. Delacruz[1], Quique Guerra[1], Samuel Jacob Goldstein[5], Yixuan J. Hou[6], Margaret Gearhart[7], Christine Wiethorn[8], Candace Pope[8], Carolyn Amditis[9], Kathryn Pruitt[10], Cinthia Newberry-Dillon[10], John Schmitz[11], Lakshmanane Premkumar[1], Adaora A. Adimora[6,12], Michael Emch[4], Ross Boyce[12], Allison E. Aiello[6], Bailey K. Fosdick[3], Daniel B. Larremore[13], Aravinda M. de Silva[1], Jonathan J Juliano[6,12], Alena J. Markmann[12#]

*co-first authors
#corresponding author

1. Department of Microbiology and Immunology, University of North Carolina School of Medicine, Chapel Hill NC 27599, USA
2. Department of Genetics, University of North Carolina School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill NC 27599, USA
3. Department of Statistics, Colorado State University, Fort Collins, CO, 80523, USA
4. Department of Geography, University of North Carolina at Chapel Hill, Chapel Hill, NC 27514, USA; Carolina Population Center, Chapel Hill, NC 27516, USA
5. Department of Environmental Sciences and Engineering, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
6. Department of Epidemiology, University of North Carolina at Chapel Hill School of Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
7. McLendon Clinical Laboratories, UNC Healthcare, Chapel Hill, NC 27599, USA
8. Johnston Health Laboratories, Johnston Health, Smithfield, NC 27577
9. Rex Healthcare Laboratory, UNC Healthcare, Chapel Hill, NC 27607, USA
10. Chatham Clinical Laboratory, Chatham Hospital, Siler City, NC 27344, USA
11. Department of Pathology & Laboratory Medicine, University of North Carolina School of Medicine, Chapel Hill NC 27599, USA
12. Department of Medicine, Division of Infectious Diseases, University of North Carolina School of Medicine, Chapel Hill NC 27599, USA
13. Department of Computer Science & BioFrontiers Institute, University of Colorado Boulder, Boulder, CO, 80303, USA

Alena J Markmann, MD, PhD; 160 Dental Circle, Chapel Hill, NC 27514; 585-880-5812

Email: alena.markmann@unchealth.unc.edu

ABSTRACT

Background
Robust community-level SARS-CoV-2 prevalence estimates have been difficult to obtain in the American South and outside of major metropolitan areas. Furthermore, though some previous studies have investigated the association of demographic factors such as race with SARS-CoV-2 exposure risk, fewer have correlated exposure risk to surrogates for socioeconomic status such as health insurance coverage.

Methods
We used a highly specific serological assay utilizing the receptor binding domain of the SARS-CoV-2 spike-protein to identify SARS-CoV-2 antibodies in remnant blood samples collected by the University of North Carolina Health system. We estimated the prevalence of SARS-CoV-2 in this cohort with Bayesian regression, as well as the association of critical demographic factors with higher prevalence odds.

Findings
Between April 21st and October 3rd of 2020, a total of 9,624 unique samples were collected from clinical sites in central NC and we observed a seroprevalence increase from 2·9 (1·7, 4·3) to 9·1 (7·2, 11·1) over the study period. Individuals

57 who identified as Latinx were associated with the highest odds ratio of SARS-CoV-2 exposure at 7·77 overall (5·20,
58 12·10). Increased odds were also observed among Black individuals and individuals without public or private health
59 insurance.
60
61 **Interpretation**
62 Our data suggests that for this care-accessing cohort, SARS-CoV-2 seroprevalence was significantly higher than
63 cumulative total cases reported for the study geographical area six months into the COVID-19 pandemic in North
64 Carolina. The increased odds of seropositivity by ethnoracial grouping as well as health insurance highlights the urgent
65 and ongoing need to address underlying health and social disparities in these populations.
66
67 **RESEARCH IN CONTEXT**
68
Evidence before this study
We searched PubMed for studies published through March 21st, 2021. We used search terms that included "COVID-
19", "SARS-CoV-2", "prevalence" and "seroprevalence". Our search resulted in 399 papers, from which we identified
58 relevant studies describing SARS-CoV-2 seroprevalence at sites around the United States from March 1 to
December 9, 2020, 12 of which utilized remnant clinical samples and three of which overlapped with our study area.
Most notably, one study of 4,422 asymptomatic inpatients and outpatients in central NC from April 28-June 19, 2020
found an estimated seroprevalence of 0·7 - 0·8%, and another study of 177,919 inpatients and outpatients (3,817 from
NC) from July 27-September 24, 2020 found an estimated seroprevalence of 2·5 - 6·8%.
77
Added value of this study
This is the largest SARS-CoV-2 seroprevalence cohort published to date in NC. Importantly, we used a Bayesian
framework to account for uncertainty in antibody assay sensitivity and specificity and investigated seropositivity by
important demographic variables that have not yet been studied in this context in NC. This study corroborates other
reports that specific demographic factors including race, ethnicity and the lack of public or private insurance are
associated with elevated risk of SARS-CoV-2 infection. Furthermore, in a subset of serum samples, we identify other
SARS-CoV-2 antibodies elicited by these individuals, including functionally neutralizing antibodies.
85
Implications of all the available evidence
It is difficult to say the exact seroprevalence in the central North Carolina area, but a greater proportion of the
population accessing healthcare has been infected by SARS-CoV-2 than is reflected by infection cases confirmed by
molecular testing. Furthermore, local governments need to prioritize addressing the many forms of systemic racism
and socioeconomic disadvantage that drive SARS-CoV-2 exposure risk, such as residential and occupational risk, and
an urgent need to provide access to SARS-CoV-2 testing and vaccination to these groups.

INTRODUCTION

In December 2019, a cluster of pneumonia cases in China's Hubei province heralded the beginning of what would become a global pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Despite attempts to contain the virus, SARS-CoV-2 has spread around the world, causing over 100 million infections and over 2 million deaths due to the respiratory disease it causes, COVID-19.[1] Serological testing complements molecular testing for evaluating the spread of SARS-CoV-2 and can be deployed efficiently at the population level.[2] Recently, large prevalence studies around the United States using remnant samples from healthcare settings have reported substantial geographic variation in prevalence by state: around 30% in New York but less than 2% in North Carolina (NC), the focus of the present study.[3,4] Notably, two other studies overlap with the present cohort both temporally and geographically. One study of 4,422 asymptomatic inpatients and outpatients in central NC from April 28-June 19, 2020 found an estimated seroprevalence of 0·7 - 0·8%, and another study of 177,919 remnant clinical laboratory samples from routine screening (3,817 from NC) from July 27-September 24, 2020 found an estimated seroprevalence of 2·5 - 6·8%.[5,6] While overall seroprevalence estimates of a given study depend on sampling method, assay characteristics, geography, and temporal factors, seroprevalence studies can provide information on the spread of COVID-19 that is missed by looking at the number of confirmed acute cases alone.

Seroprevalence studies are also useful for identifying demographic factors such as racial, ethnic and socioeconomic disparities among those exposed to SARS-CoV-2.[4,7,8] The COVID-19 pandemic has been shaped by the deep and historic impacts of structural racism on disease disparities in US society as identified by serologic studies as well as hospitalization and mortality rates.[9,10] For example, COVID-19 case and hospitalization rates among Black, Hispanic and Native American populations in the US, according to the Centers for Disease Control and Prevention, are 2.5-4.5 times higher than those in white populations[11]. Structural and occupational factors previously identified as drivers of race and ethnic disparities in health include unequal labor market opportunities and higher representation in essential work positions that lack job security, access to infection prevention control, benefits, and sick leave.[12–17] Here, we confirm the findings of disparate SARS-CoV-2 exposure among racial and ethnic groups in the US by measuring seroprevalence in a large southern US health-care seeking cohort using remnant blood samples.

The following results are from the first six months (April 21-October 3, 2020) of an ongoing seroprevalence study using convenience remnant samples from clinical laboratories in central NC. The study catchment area covers Wake, Orange, Chatham, Johnston, Durham and Alamance counties and includes the county of the first confirmed case in NC,[18] which occurred on March 3rd, 2020. On October 3rd, 2020, the cumulative total PCR and antigen-confirmed SARS-CoV-2 cases in the study catchment area was 52,722 (2·7% of the population), with 1,266 confirmed deaths.[19,20] We used an in-house enzyme-linked immunoassay (ELISA) against the receptor-binding domain (RBD) of the spike protein of SARS-CoV-2[21] and applied Bayesian inference[22] to estimate seroprevalence and demographic risk factors of SARS-CoV-2 infection in a healthcare-seeking cohort over a six-month period.

METHODS

Sampling Strategy and Data Collection

Remnant plasma and serum samples were collected from four hospital-based clinical laboratories affiliated with the University of North Carolina (UNC) Health system. These laboratories receive and process clinical samples from inpatient units as well as outpatient clinics in NC. Each week, up to 300 remnant samples belonging to individuals 5-99 years of age were arbitrarily selected by the clinical laboratory for testing from each location. Samples were collected between April 21st, 2020 – October 3rd, 2020. Medical record numbers were recorded for each sample and duplicates were discarded. We abstracted the following demographic and clinical data from electronic medical records (EMR, Epic): age, legal sex, ethnicity, race, address including city, state and ZIP code, insurance coverage, insurance type, inpatient or outpatient status, encounter diagnosis (ICD-10 code), inpatient date of discharge, and whether or not COVID-19 testing was performed within a 30-day window prior to study sample collection. Written informed consent was not required due to the use of routinely collected samples. All data for this study were collected under UNC IRB #20-0791, which is conducted under Good Clinical Research Practices (GCP) and compliant with institutional IRB oversight. De-identified samples used for assay validation were collected under UNC IRBs #20-0913 and #08-0895.

Enzyme-Linked Immunosorbent Assays

A total Ig and IgM SARS-CoV-2 RBD ELISA that does not react with common endemic human coronaviruses was
used in this study as previously described.[21] The spike protein N-terminal domain (NTD) antigen (16–305 amino
acids, Accession: P0DTC2.1) was cloned into the pαH mammalian expression vector and purified using nickel-
nitrilotriacetic acid agarose in the same manner. Each measurement was conducted in duplicate and duplicate values
with variance > 25% and/or one value above assay cutoff were repeated. A correlation plot shows using 140 COVID-
19 PCR-confirmed cases between our RBD Ig P/N ratios and the neutralization assay described below (**Figure S1**).
175
176 *Nucleocapsid protein ELISA*
177
178 Detection of IgG antibody to SARS-CoV-2 N antigen was performed with the EUA approved Abbott SARS-CoV-2
179 IgG assay (Abbott Laboratories) on the Abbott Architect i2000SR immunoassay analyzer as previously described.[23]
180
181 *SARS-CoV-2 Neutralization Assays*
182
To further characterize the SARS-CoV-2 antibody responses of this study, viral neutralization assays were obtained
for 110 ELISA-positive samples that were selected randomly using the sample_n() function of the dplyr R package.
Luciferase-expressing, full-length SARS-CoV-2 isolate WA1 strain (GenBank Accession#: MT020880) was
engineered and recovered via reverse genetics and used to titer serially diluted sera on Vero E6 USAMRID cell as
described previously.[24] The sample dilution at which a 50% reduction in RLU was observed relative to that of the
virus control wells was used as the 50% neutralization titer (NT50) for that sample.
189
190 *Statistical Methods and Analyses*
191
192 To account for plate-to-plate variability, we used positive to negative (P/N) ratios defined as the average optical
193 density (OD) of the sample divided by the average OD of the negative control in the respective ELISA plate. Following
194 the CDC recommendation to set specificity to 99·5%, we chose the 0.995 quantile of the P/N ratio for the negative
195 validation samples as the P/N cutoff.[25]
196
We fit two statistical models to estimate seroprevalence. First, we fit a Bayesian autoregressive logistic model to
estimate weekly prevalence across the six-month study period while accounting for uncertainty in the assay specificity
and sensitivity due to finite lab validation samples. Second, we fit a Bayesian logistic regression model to estimate
prevalence and conditional odds ratios by subpopulation with main effects for gender, race/ethnicity, age, in/out-
patient status, and health insurance payor, while again accounting for uncertainty in the assay test characteristics
(**Table S1**). Each group was compared to females, non-Latinx white, ages 5-17, outpatient, and private payor health
insurance status as respective baseline categories. Details are given in Supplementary Methods: Bayesian
seroprevalence models with unknown sensitivity and specificity. These Bayesian hierarchical models (BHM)
simultaneously model study data and validation data to produce prevalence estimates and credible intervals that reflect
both uncertainty due to the finite study sample as well as the uncertainty in the sensitivity and specificity of the ELISA,
with statistical uncertainty represented by 95% credible intervals.
208
209
210 **RESULTS**
211
212 *Cohort Characteristics*
213
From April 21, 2020 – October 3, 2020, after excluding duplicate samples, 9,624 remnant samples were analyzed
from four UNC Health hospitals in central North Carolina. The six counties most heavily sampled were Orange,
Johnson, Chatham, Wake, Durham and Alamance, with 6,946 (72·2%) of individuals residing in these counties
(**Figure 1**). The study consists of 5,417 females (56·3%) and 4,206 males (43·7%) which is similar to the
demographics of this region (**Table 1**). Less than 6% of individuals were in the youngest age group (5-17 years old),
though this age group represents over 18% of the study area's population. Approximately 90% of study individuals
were insured, with 8% falling into the self-pay category. The majority of sampled individuals were seen at UNC
Memorial Hospital, ~3% were acute or trauma cases and ~5% had a visit diagnosis of fever or respiratory symptoms
(**Table S2**). Overall, approximately 1% of patients had an associated COVID-19 visit diagnosis, with a significant
difference between inpatients (2·8%) and outpatients (0·3%) (Chi-squared test; p<0·0001) (**Table S3**).
224

Overall seroprevalence estimates

The six-month period of the study was divided into three, two-month cohorts. The BHM-derived seroprevalence estimates increased from around 3% in April/May to around 9% in August/September **(Table 2)**. Raw seroprevalence estimates also showed a similar increasing trend over the study period, but because they do not take into account assay performance uncertainty, they are slightly higher at ~5% and ~11%. Furthermore, seroprevalence estimates peaked in early August following a hospitalization peak in mid-July **(Figure 2A, 2C)**. Cumulative PCR-positive COVID-19 cases reported by the state for these six counties increased over the study period **(Figure 2B)** with the most rapid accumulation of cases occurring from June to August. Unexpectedly, seroprevalence peaks followed by a slight decline, related to raw seroprevalence estimates at Johnston County hospital which surged from 7·81% in the first two months to 18·00% in the second two months coinciding with a peak in PCR-confirmed cases in the region, followed by a measured decline in raw seroprevalence to 14·80% in the final two-month period **(Table S6)**. This peak and decline was not affected by the removal of cases with ICD-10 visit codes for "COVID-19" or those we identify as "respiratory disease" (data not shown).

Clinical and demographic differences in seroprevalence estimates

Latinx-identifying individuals have higher SARS-CoV-2 seroprevalence at 15-33% compared to non-Latinx individuals which have only 1-11% seroprevalence over the study period **(Table 2)**. Individuals with "Other/Unknown" or "Self-pay" insurance status had a higher estimated seroprevalence (~20-40% or ~1-18%, respectively) than those with private or public health insurance (~3-9%). Approximately 30% of Latinx individuals in this study were either in the other/unknown or self-pay health categories, disproportionately comprising ~27% of these two categories but only accounting for ~8% of our study population **(Table S5)**.

To better compare the relative odds of SARS-CoV-2 seroprevalence for each clinical and/or demographic characteristic, we calculated conditional odds ratios for each variable we collected using the BHM **(Table 3)**. Latinx individuals had the highest odds of SARS-CoV-2 exposure throughout the study period compared to non-Latinx white individuals, OR 7·77 overall (5·20, 12·10), ranging from 14·53 (6·47, 36·72) in the first two months to 4·34 (2·61, 7·41) in the last two months of the study. Individuals with unknown insurance status also had an elevated odds ratio of seropositivity at 3·81 (2·23, 6·54) compared to those with private insurance status. Over the entire period of the study, non-Latinx Black individuals, individuals aged 50-64 years, and inpatients, also had increased odds ratios of approximately two-fold compared to non-Latinx white individuals, individuals aged 0-17, and outpatients, respectively. The overall difference in odds ratios by age appears to be driven primarily by increased odds ratios in the first two months.

SARS-CoV-2 RBD positive subset analysis

To determine the SARS-CoV-2 antibody repertoire in a subset of RBD Ig seropositive individuals, we randomly selected 110 participants and tested their sera for: RBD IgM, NTD IgG, and SARS-CoV-2 neutralizing antibodies. About 75% of individuals were positive for RBD IgM, 60% had NTD IgG antibodies, and about 50% had detectable neutralizing antibodies **(Figure 3A)**. Of the participants with detectable functionally neutralizing antibodies, 23% had a high titer > 1:1280, 47% had a moderate titer of 1:160-1:1279, and 30% had a lower titer of 1:10-1:159. Furthermore, RBD Ig P/N antibody signal correlated more strongly with functionally neutralizing antibody levels **(Figure 3B)**, than NTD IgG signal **(Figure 3C)**. We also found that 36% (29/80) of those in this subset with an ICD-10 code binned as "Other" had detectable neutralizing antibodies, while 83% (25/30) of individuals with an ICD-10 code of "COVID-19" or what we identify as "respiratory disease" had neutralizing antibodies **(Figure 3D)**. There was substantial agreement between the RBD Ig ELISA results reported here and 150 study individuals for which a clinical SARS-CoV-2 nucleocapsid IgG (Abbott assay) was available (Cohen's kappa=0·685) **(Table S4).**

DISCUSSION

Here we describe SARS-CoV-2 seroprevalence in a total of 9,624 unique healthcare-seeking individuals in central North Carolina using clinical remnant samples from four regional hospitals between April and October 2020. Employing a Bayesian framework[22] to capture assay uncertainty in both field and lab validation data, we estimate a significant increase in overall seroprevalence from 2·9% (95% CI 1·7% - 4·3%) at the start of the study period, to 9·1% (95% CI 7·2% - 11·1%) at the end of the study period, approximately six months after the first case in the state.

The end-of-study prevalence identified here is significantly higher than the cumulative number of cases identified by
PCR or antigen testing in the same county region at the same date, though determining the degree to which the
identified cases undercount true infections requires more representative sampling.

A previous study from central North Carolina that overlaps with the first two months of our study period found
seroprevalence in an asymptomatic healthcare-seeking cohort below 1% using the Abbott nucleocapsid IgG assay.[5]
This is much lower than the ~3% seropositive estimate in our cohort over this time period, and may be due to under-
sampling of Latinx individuals in that study and/or preferential sampling of asymptomatic individual. There is also
growing concern about the use and performance of nucleocapsid IgG assays in individuals with asymptomatic or mild
disease.[26] The nationwide CDC study that used remnant clinical samples from inpatients and outpatients found a
seroprevalence of 6·8% in NC in September 2020, which is closer to our estimate of 9·1% during the final two months
of this analysis.

The conditional odds ratios we calculated assume that all other variables are held constant while estimating the effect
of one demographic variable at a time. We found that Latinx individuals had the highest odds of SARS-CoV-2
seropositivity, and that non-Latinx Black individuals also had high odds of SARS-CoV-2 seropositivity, corroborating
previous observations.[4,7,8] The high odds ratios by race and ethnicity decrease over time, consistent with the virus
spreading first among individuals with high exposure risk and later to the rest of the population. Residential
segregation, crowded households, socioeconomic disadvantage, mass incarceration, and inequities in access to
insurance, health care, and access to testing, vaccination, and treatments have all been cited as factors that have
contributed to the large and sustained racial and ethnic disparities in COVID-19 in the US.[13,15,27–29] We also observed
that individuals that fell into the "self-pay" category for their healthcare or otherwise had unknown healthcare status
had higher SARS-CoV-2 seropositivity and odds ratios. The significant overlap in the Latinx population and these
insurance categories is concerning because the high odds ratios and seroprevalence in these categories can lead to
much higher exposure risk among the significant number of underinsured Latinx individuals[30].

Studies of PCR-positive symptomatic COVID-19 cases have reported good neutralizing antibody responses in these
individuals.[31] Thus, it was surprising that we observed 51% of individuals in our RBD-positive subset analysis did not
have detectable neutralizing antibodies. Though we do not know what proportion of individuals in our study had
asymptomatic infections, low neutralizing antibody titers may be explained by short duration of viral replication in
respiratory compartments and low to no viral replication in the serum or blood of those with mild or asymptomatic
disease. Not surprisingly, when we looked at our neutralizing antibody results by ICD-10 code, the majority of all
individuals with a "respiratory disease" or "COVID-19" diagnosis had developed neutralizing antibodies. Reports of
mild disease COVID-19 cohorts support the idea that detectable neutralizing antibody titers are not necessarily
identified after mild COVID-19.[23,32] In this subset analysis we also found that 75% had RBD IgM antibodies,
indicating that their infections likely occurred within the past three months.[23] Furthermore, a majority of individuals
in this subset had detectable NTD IgG antibodies; the NTD has recently been found to be an important target for the
B.1.1.7, B.1.351, and B.1.1.28.1 SARS-CoV-2 variants.[33]

The primary limitation of this study is that the study population, composed of individuals accessing care at UNC area
hospitals and clinics may differ from the overall population in central North Carolina in ways that are not captured in
demographic data (e.g., overall health status). Accordingly, we have chosen to not weight our dataset to county
demographics and therefore do not provide overall estimates of seroprevalence in the six-county area as that would
require more representative sampling methodology.[34] Furthermore, many clinics and hospital elective procedures were
closed or only seeing patients virtually during the first few months of the study period.

The unexpected seroprevalence peak observed at the Johnston County hospital suggests that the population accessing
care at these clinical sites did not have consistent exposure risk over time. As expected, seroprevalence estimates in
this cohort track closely with COVID-19 hospitalizations in the four hospitals in this study with a two-week lag which
could be due to time to seroconvert. Declining antibody over this time period to undetectable levels is unlikely, as the
length of the study is shorter than it takes for significant antibody decline to undetectable levels, although little is
known about antibody levels over time in the asymptomatic population.[31]

Other limitations of the study include that we could not break down odds ratios by all races and/or by race and ethnicity
at the same time, or by multiracial categories because the number of individuals became too small to allow broad
interpretation. Finally, though the "self-pay" insurance category includes the uninsured, we cannot confidently state

that everyone in this category was uninsured because lack of insurance is not a specific category that is captured in
the EMR. Although SARS-CoV-2 seroprevalence of healthcare-seeking individuals is an imperfect comparison to the
general population, we maintain that it is a useful sentinel population to understand overall trends, especially when
attempting to surveil rural populations residing in areas without strong public health systems and spread over a large
geographic area.
342
Based on our estimates of seroprevalence in the population accessing healthcare, cumulative case numbers confirmed
by molecular diagnostics are likely under-representing the true number of cases. Public health distancing measures,
mask wearing, and vaccination should continue to be prioritized in order to lower the transmission of SARS-CoV-2
and subsequent loss of lives. Our findings of a significantly higher odds of SARS CoV-2 seropositivity among Latinx
and non-Latinx Black populations corroborate numerous studies describing large racial and ethnic disparities in
SARS-CoV-2 infection, morbidity and mortality in the US.[4,7,8] Vaccination programs should address structural and
occupational factors that drive race and ethnic disparities in health outcomes in the US to ensure that individuals at
particularly high exposure risk of SARS-CoV-2 have timely access to SARS-CoV-2 vaccination.
351

352   **Acknowledgements**
353   We would like to thank Jennifer Dan, Alessandro Sette and Shane Crotty's Laboratories at La Jolla Institute of
354   Immunology for supplying some of the serum controls in our validation cohort.
355
356   **Data Sharing**
357   Deidentified individual data will be shared beginning 9 to 36 months following publication provided the investigator
358   who proposes to use the data has approval from an Institutional Review Board (IRB), Independent Ethics Committee
359   (IEC), or Research Ethics Board (REB), as applicable, and executes a data use/sharing agreement with UNC.
360
361   **Funding Sources**
362   This study was supported by funds from the NC Department of Health and Human Services, Division of Public Health
363   (Contract 00041877) and the SeroNet program of the National Cancer Institute (1U01CA261277-01). Funding sources
364   played no role in study design, in the collection, analysis, and interpretation of data, in the writing of the report, or in
365   the decision to submit the paper for publication.
366

367   **Declaration of interests**
368   The authors declare no conflicts of interest.
369

Author contributions
CL, CHC: Conceptualization, Data curation, Investigation, Project Administration, Writing – original draft. SP, KB:
Formal analysis, Software, Visualization. MD, QG, UPV: Data curation, Investigation, Project Administration. SJG:
Formal analysis, Software, Visualization. YJH: Investigation. PL: Methodology, Resources. RR, MG, CW, KP, CA:
Resources. JS: Conceptualization, Resources. ME: Conceptualization, Formal analysis, Resources, Supervision. RB,
AA: Conceptualization, Funding acquisition. BKF: Conceptualization, Formal analysis, Resources, Software,
Supervision. DBL: Conceptualization, Formal analysis, Software. ADS: Conceptualization, Funding acquisition,
Methodology, Project Administration, Resources, Supervision. JJJ: Conceptualization, Funding acquisition, Project
Administration, Supervision. AJM: Conceptualization, Funding acquisition, Investigation, Project Administration,
Supervision, Writing – original draft. CL, CHC, SP, KB, MD, QG, UPV, SJG, YJH, PL, JS, ME, RB, AA, BKF, DBL,
ADS, JJJ, AJM: Writing- reviewing & editing.

381
382

383   **REFERENCES**
384
1   Home - Johns Hopkins Coronavirus Resource Center. 2020. https://coronavirus.jhu.edu/ (accessed Feb 26,
2021).
2   Population-based age-stratified seroepidemiological investigation protocol for coronavirus 2019 (COVID-
19) infection. https://www.who.int/emergencies/diseases/novel- (accessed June 21, 2020).
3   Havers FP, Reed C, Lim T, *et al.* Seroprevalence of Antibodies to SARS-CoV-2 in 10 Sites in the United
States, March 23-May 12, 2020. *JAMA Intern Med* 2020. DOI:10.1001/jamainternmed.2020.4130.
4   Anand S, Montez-Rath M, Han J, *et al.* Prevalence of SARS-CoV-2 antibodies in a large nationwide sample
of patients on dialysis in the USA: a cross-sectional study. *Lancet* 2020; **0**. DOI:10.1016/S0140-

6736(20)32009-2.

5   Barzin A, Schmitz JL, Rosin S, *et al.* SARS-CoV-2 seroprevalence among a southern U.S. population
indicates limited asymptomatic spread under physical distancing measures. *MBio* 2020; **11**: 1–8.

6   Bajema KL, Wiegand RE, Cuffe K, *et al.* Estimated SARS-CoV-2 Seroprevalence in the US as of
September 2020. *JAMA Intern Med* 2020. DOI:10.1001/jamainternmed.2020.7976.

7   Rogawski McQuade ET, Guertin KA, Becker L, *et al.* Assessment of Seroprevalence of SARS-CoV-2 and
Risk Factors Associated With COVID-19 Infection Among Outpatients in Virginia. *JAMA Netw open* 2021;
4: e2035234.

8   Feehan AK, Velasco C, Fort D, *et al.* Racial and workplace disparities in seroprevalence of SARS-CoV-2,
Baton Rouge, Louisiana, USA. *Emerg Infect Dis* 2021; **27**: 314–7.

9   Bailey ZD, Feldman JM, Bassett MT. How Structural Racism Works — Racist Policies as a Root Cause of
U.S. Racial Health Inequities. *N Engl J Med* 2020; **384**: 768–73.

10   Khazanchi R, Evans CT, Marcelin JR. Racism, Not Race, Drives Inequity Across the COVID-19
Continuum. JAMA Netw. open. 2020; **3**: e2019933.

11   Centers for Disease Control. Risk for COVID-19 Infection, Hospitalization, and Death By Race/Ethnicity.
2021; published online Feb 18. https://www.cdc.gov/coronavirus/2019-ncov/covid-data/investigations-
discovery/hospitalization-death-by-race-ethnicity.html (accessed Feb 23, 2021).

12   Kerwin D, Nicholson M, Alulema D, Warren R. US Foreign-Born Essential Workers by Status and State,
and the Global Pandemic. 2020 DOI:10.14240/cmsrpt0520.

13   Bailey ZD, Robin Moon J. Racism and the political economy of COVID-19: Will we continue to resurrect
the past? *J Health Polit Policy Law* 2020; **45**: 937–50.

14   Alberti PM, Lantz PM, Wilkins CH. Equitable Pandemic Preparedness and Rapid Response: Lessons from
COVID-19 for Pandemic Health Equity. *J Health Polit Policy Law* 2020; **45**: 921–35.

15   Laster Pirtle WN. Racial Capitalism: A Fundamental Cause of Novel Coronavirus (COVID-19) Pandemic
Inequities in the United States. *Heal Educ Behav* 2020; **47**: 504–8.

16   McClure ES, Vasudevan P, Bailey Z, Patel S, Robinson WR. Racial Capitalism Within Public Health—How
Occupational Settings Drive COVID-19 Disparities. *Am J Epidemiol* 2020; **189**: 1244–53.

17   Dyal JW, Grant MP, Broadwater K, *et al.* COVID-19 Among Workers in Meat and Poultry Processing
Facilities ― 19 States, April 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**.
DOI:10.15585/mmwr.mm6918e3.

18   NCDHHS: North Carolina Identifies First Case of COVID-19.
https://web.archive.org/web/20200407230739/https://www.ncdhhs.gov/news/press-releases/north-carolina-
identifies-first-case-covid-19 (accessed May 22, 2020).

19   US Census Bureau. American Community Survey 5-Year Data (2014-2018). 2019
https://www.census.gov/data/developers/data-sets/acs-5year.html (accessed Dec 3, 2020).

20   Data Behind the Dashboards: NC DHHS COVID-19. https://covid19.ncdhhs.gov/dashboard/data-behind-
dashboards (accessed Feb 13, 2021).

21   Premkumar L, Segovia-Chumbez B, Jadi R, *et al.* The receptor binding domain of the viral spike protein is
an immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. *Sci Immunol* 2020; **5**:
eabc8413.

22   Larremore DB, Fosdick BK, Bubar KM, *et al.* Estimating SARS-CoV-2 seroprevalence and epidemiological
parameters with uncertainty from serological surveys. *Elife* 2021; **10**. DOI:10.7554/eLife.64206.

23   Markmann AJ, Giallourou N, Ryan Bhowmik D, *et al.* Sex disparities and neutralizing antibody durability to
SARS-CoV-2 infection in convalescent individuals. *medRxiv* 2021; : 2021.02.01.21250493.

24   Hou YJ, Okuda K, Edwards CE, *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection
Gradient in the Respiratory Tract. *Cell* 2020; published online May 27. DOI:10.1016/j.cell.2020.05.042.

25   Centers for Disease Control and Prevention. Interim Guidelines for COVID-19 Antibody Testing | CDC.
2020; published online Aug 1. https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antibody-tests-
guidelines.html (accessed Feb 26, 2021).

26   Ripperger TJ, Uhrlaub JL, Watanabe M, *et al.* Orthogonal SARS-CoV-2 Serological Assays Enable
Surveillance of Low-Prevalence Communities and Reveal Durable Humoral Immunity. *Immunity* 2020; **53**:
925-933.e4.

27   Hagan LM, Williams SP, Spaulding AC, *et al.* Mass Testing for SARS-CoV-2 in 16 Prisons and Jails — Six
Jurisdictions, United States, April–May 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 1139–43.

28   Webb Hooper M, Nápoles AM, Pérez-Stable EJ. COVID-19 and Racial/Ethnic Disparities. JAMA - J. Am.
Med. Assoc. 2020; **323**: 2466–7.

29   Figueroa JF, Wadhera RK, Lee D, Yeh RW, Sommers BD. Community-Level Factors Associated With
Racial And Ethnic Disparities In COVID-19 Rates In Massachusetts. *Health Aff* 2020; **39**: 1984–92.
30   Berchick ER, Barnett JC, Upton RD. Health Insurance Coverage in the United States: 2018 Current
Population Reports. 2019.
31   Long QX, Tang XJ, Shi QL, *et al.* Clinical and immunological assessment of asymptomatic SARS-CoV-2
infections. *Nat Med* 2020; **26**: 1200–4.
32   Dan JM, Mateus J, Kato Y, *et al.* Immunological memory to SARS-CoV-2 assessed for up to 8 months after
infection. *Science (80-)* 2021; **371**. DOI:10.1126/science.abf4063.
33   Diamond M, Chen R, Xie X, *et al.* SARS-CoV-2 variants show resistance to neutralization by many
monoclonal and serum-derived polyclonal antibodies. *Res Sq* 2021; published online Feb 10.
DOI:10.21203/rs.3.rs-228079/v1.
34   Shook-Sa BE, Boyce RM, Aiello AE. Estimation Without Representation: Early Severe Acute Respiratory
Syndrome Coronavirus 2 Seroprevalence Studies and the Path Forward. *J Infect Dis* 2020; **222**: 1086–9.

462
463
464   **FIGURE LEGENDS**
465
466   **Figure 1. Catchment area for hospital remnant sample collection for UNC Health hospitals.** Remnant samples
467   were collected from hospital clinical laboratories from each of the four sites indicated by the red dots. (A) Number of
468   samples collected by count as well as (B) the rate of sampling.[19]
469
470
471   **Figure 2. Trends in seroprevalence estimates.** (A) Weekly posterior mean seroprevalence estimates and 95%
472   credible intervals for the study period of 4/21-10/3 of the hospital samples by ELISA plotted over time over the course
473   of the study period. (B) Cumulative daily COVID-19 PCR+ cases from the six-county area 4/19-10/3, and (C) weekly
474   COVID-19 hospitalizations in the six-county area 4/19-10/3 from NC Department of Health and Human Services.
475
476
Figure 3. Antibody repertoires in an RBD Ig positive subset. 110 RBD Ig positive samples were chosen at random
to undergo SARS-2 antibody repertoire analysis. (A) Percent of individuals with RBD IgM, NTD IgG and functionally
neutralizing antibodies (NT50). (B) Correlation plot of NT50 and RBD Ig. (C) Correlation plot of NTD IgG and RBD
Ig, r_s = Spearman correlation coefficient displayed in the top left of panels (B) and (C). (D) NT50 values for each
diagnosis binning category based on ICD-10 codes. Medians shown in blue. Two-tailed Mann-Whitney,
****p<0.0001, **p=0.0078.

505　**TABLES**
506

Table 1. Study participants by demographic factors of interest.							
	4/19-6/13		6/14-8/08		8/09-10/03		6-county Demographics (%)
	N	(%)	N	(%)	N	(%)	
Sex							
Female	1947	56·2	2020	57·3	1450	55·1	51·8
Male	1515	43·7	1508	42·7	1183	44·9	48·2
Unreported	1	0·0	0	0·0	0	0·0	—
Age							
5-17	259	7·5	163	4·6	150	5·7	18·4
18-49	1311	37·9	1052	29·8	830	31·5	48·7
50-64	926	26·7	1030	29·2	725	27·5	19·7
65-99	967	27·9	1283	36·4	928	35·2	13·1
Race/Ethnicity							
NL White	2113	61·0	2267	64·3	1628	61·8	59·7
NL Black	845	24·4	803	22·8	603	22·9	21·0
NL Other	210	6·1	195	5·5	194	7·4	8·2
Latinx	295	8·5	263	7·5	208	7·9	11·1
In/Out patient							
Inpatient	1057	30·5	961	27·2	839	31·9	—
Outpatient	2394	69·1	2562	72·6	1792	68·1	—
Unknown	12	0·3	5	0·1	2	0·1	—
Payor							
Public	1825	52·7	2050	58·1	1509	57·3	—
Private	1249	36·1	1172	33·2	920	34·9	—
Self-Pay	326	9·4	254	7·2	181	6·9	—
Other/Unknown	63	1·8	52	1·4	23	0·8	—

507
508
509
510　**Table 1**. **Study participants by demographic factors of interest.** Note, because of how the NC census reports data,
511　the gender and age breakdowns of the 6-county demographics includes only individuals over the age of 4 (including
512　those over age 99), but the race/ethnicity breakdown includes individuals of all ages. Additionally, the 65-99 age
513　category is actually age 65+ for the 6-county demographics.

Table 2. Cohort prevalence estimates									
	Positivity			**BHM prevalence estimates**					
	4/19-6/13	6/14-8/08	8/09-10/03	4/19-6/13		6/14-8/08		8/09-10/03	
				Estimate	95% CI	Estimate	95% CI	Estimate	95% CI
Overall	5·3	10·5	10·8	2·9	(1·7, 4·3)	8·8	(7·1, 10·6)	9·1	(7·2, 11·1)
Age									
5-17	3·1	9·8	9·3	1·4	(0·3, 3·3)	8·1	(3·9, 13·4)	7·6	(3·5, 13·0)
18-49	6·0	12·6	10·5	3·6	(2·2, 5·4)	11·1	(8·6, 13·8)	8·7	(6·2, 11·5)
50-64	5·9	10·4	13·0	3·7	(1·9, 5·8)	8·7	(6·3, 11·3)	11·5	(8·5, 14·7)
65-99	4·3	9·0	9·6	1·5	(0·2, 3·4)	7·1	(5·0, 9·4)	7·7	(5·2, 10·4)
Sex									
Female	4·5	10·3	10·7	2·1	(1·0, 3·5)	8·5	(6·6, 10·6)	8·9	(6·8, 11·3)
Male	6·3	10·7	10·9	3·9	(2·3, 5·8)	9·2	(7·1, 11·3)	9·2	(6·9, 11·8)
Race/Ethnicity									
NL White	3·7	7·5	8·3	1·4	(0·5, 2·7)	5·4	(3·7, 7·3)	6·3	(4·3, 8·4)
NL Black	5·6	12·0	12·8	2·6	(0·6, 5·0)	10·4	(7·5, 13·4)	11·4	(8·2, 14·8)
NL Other	5·7	10·3	11·3	2·0	(0·1, 5·9)	8·5	(3·9, 13·9)	9·3	(4·5, 14·9)
Latinx	15·9	31·9	24·0	14·8	(10·4, 19·6)	33·2	(26·8, 40·0)	23·9	(17·5, 31·1)
In/out patient									
Outpatient	4·3	9·0	9·1	2·0	(1·0, 3·3)	7·1	(5·4, 9·0)	7·1	(5·1, 9·2)
Inpatient	7·7	14·6	14·4	5·0	(2·9, 7·4)	13·3	(10·5, 16·2)	13·3	(10·3, 16·4)
Payor									
Private	5·2	9·0	8·9	2·9	(1·5, 4·6)	7·3	(5·3, 9·6)	7·1	(4·7, 9·6)
Public	5·0	9·8	10·7	2·5	(1·2, 4·2)	7·9	(5·9, 9·9)	8·9	(6·8, 11·2)
Self-Pay	4·0	18·9	17·1	1·3	(0·2, 3·5)	18·3	(13·1, 23·8)	16·3	(10·4, 23·1)
Other/ Unknown	22·2	30·8	43·5	21·1	(11·8, 31·7)	31·2	(19·4, 44·5)	40·4	(22·4, 60·6)

532
533
534 **Table 2. Cohort prevalence estimates**. Raw seropositivity (%) and posterior mean seroprevalence estimates (%)
535 from BHM with 95% credible intervals (lower bound, upper bound). NL, Non-Latinx.

Table 3. Conditional odds ratios of being SARS-CoV-2 seropositive over the study period.

	4/19-6/13		6/14-8/08		8/09-10/03		4/19-10/03 (overall)	
	Estimate	95% CI	Estimate	95% CI	Estimate	95% CI	Estimate	95% CI
Gender								
Female	—	—	—	—	—	—	—	—
Male	**2·05**	**(1·08, 4·25)**	1·10	(0·80, 1·51)	0·91	(0·64, 1·29)	1·27	(0·98, 1·69)
Race/Ethnicity								
NL White	—	—	—	—	—	—	—	—
NL Black	1·66	(0·53, 4·28)	**1·94**	**(1·31, 2·92)**	**1·82**	**(1·20, 2·79)**	**1·80**	**(1·19, 2·65)**
NL Other	1·26	(0·12, 5·74)	1·58	(0·74, 3·19)	1·81	(0·87, 3·57)	1·54	(0·66, 2·84)
Latinx	**14·53**	**(6·47, 36·72)**	**7·43**	**(4·70, 11·97)**	**4·34**	**(2·61, 7·41)**	**7·77**	**(5·20, 12·10)**
Age								
5-17	—	—	—	—	—	—	—	—
18-49	3·09	(0·99, 11·43)	1·38	(0·68, 3·05)	0·89	(0·42, 2·03)	1·56	(0·92, 2·77)
50-64	**3·62**	**(1·13, 13·56)**	1·34	(0·64, 2·99)	1·56	(0·76, 3·54)	**1·96**	**(1·15, 3·55)**
65-99	1·62	(0·28, 6·90)	1·49	(0·71, 3·34)	1·13	(0·52, 2·64)	1·40	(0·71, 2·61)
In/out patient								
Outpatient	—	—	—	—	—	—	—	—
Inpatient	**2·50**	**(1·31, 5·10)**	**1·91**	**(1·38, 2·68)**	**1·92**	**(1·34, 2·80)**	**2·09**	**(1·59, 2·85)**
Payor								
Private	—	—	—	—	—	—	—	—
Public	0·85	(0·41, 1·73)	0·89	(0·58, 1·34)	1·16	(0·74, 1·85)	0·96	(0·70, 1·30)
Self-Pay	**0·18**	**(0·03, 0·64)**	**1·78**	**(1·07, 2·93)**	**1·94**	**(1·03, 3·63)**	0·85	(0·45, 1·41)
Other/ Unknown	**3·08**	**(1·15, 8·23)**	**2·73**	**(1·25, 5·98)**	**6·60**	**(2·29, 18·71)**	**3·81**	**(2·23, 6·54)**

556
557
558 **Table 3. Conditional odds ratios of being SARS-CoV-2 seropositive over the study period.** Data is broken down
559 into three two-month long periods in central North Carolina. Odds ratios of seropositivity calculated from the BHM
560 with 95% credible intervals (lower bound, upper bound) are reported where the baseline groups for comparison are
561 female, Non-Latinx white, age 5-17, outpatient, and private insurance. Odds ratios that do not overlap a value of one
562 are bolded.
563

A. Samples by county

N in sample
- 1,000 to 2,000
- 500 to 999
- 250 to 499
- 1 to 249
- No samples

B. Rate of sampling (samples by county population)

per 1,000
- approx. 26
- 5 to 10
- 1 to 5
- <1
- No samples

● UNC Hospital

N=9,504 (120 sampled individuals reside outside of North Carolina)
Sampling period: April 12, 2020 to October 3, 2020