

1 **Clustering Lifestyle Risk Behaviors among Vietnamese Adolescents and Roles**  
2 **of School: A Bayesian Multilevel Analysis of Global School-Based Student**  
3 **Health Survey 2019**

4 **Authors:**

5 Khuong Quynh Long<sup>1</sup>, Hoang Thi Ngoc-Anh<sup>1</sup>, Nguyen Hong Phuong<sup>2</sup>, Tran Thi Tuyet-Hanh<sup>1</sup>,  
6 Kidong Park<sup>3</sup>, Momoe Takeuchi<sup>3</sup>, Nguyen Tuan Lam<sup>3</sup>, Pham Thi Quynh Nga<sup>3</sup>, Le Phuong-Anh<sup>3</sup>,  
7 Le Van Tuan<sup>4</sup>, Tran Quoc Bao<sup>5</sup>, Ong Phuc Thinh<sup>1</sup>, Nguyen Van Huy<sup>6,7</sup>, Vu Thi Hoang Lan<sup>1</sup>,  
8 Hoang Van Minh<sup>1</sup>

9 <sup>1</sup>Hanoi University of Public Health, Hanoi, 100000, Vietnam

10 <sup>2</sup>Poverty, Health and Nutrition Division, International Food Policy Research Institute,  
11 Washington, DC 20006

12 <sup>3</sup>World Health Organization, Country Office for Vietnam, Hanoi, 100000, Vietnam

13 <sup>4</sup>Vietnam Ministry of Education and Training, Hanoi, 100000, Vietnam

14 <sup>5</sup>General Department of Preventive Medicine, Ministry of Health, Hanoi, 100000, Vietnam

15 <sup>6</sup>Health Innovation and Transformation Centre, Federation University, Victoria, 3353, Australia

16 <sup>7</sup>Graduate School of Public Health, St. Luke's International University, Tokyo, 104-0044, Japan

17

18 **Corresponding author:**

19 Hoang Thi Ngoc-Anh

20 Address: 1A Duc Thang Road, Duc Thang Ward, North Tu Liem District, Hanoi, 100000,  
21 Vietnam

22 Email address: [hnta@huph.edu.vn](mailto:hnta@huph.edu.vn)

23 Mobile: +84 86 577 0967

## 24 **Declarations**

25 **Declaration of Interests:** The authors declare that they have no conflict of interest.

26 **Consent to participate:** Written informed consent was obtained from all participants'  
27 parents/guardians before the study.

28 **Data sharing statement:** The datasets analyzed for the current study are not publicly available  
29 but are available upon reasonable request.

## 30 **Authors' contributions**

31 Conceptualization: KQL

32 Literature review: KQL, HTNA, and OPT

33 Data curation: KQL, TTTH, and NTL

34 Formal Analysis: KQL

35 Results interpretation: KQL, HTNA, and NHP

36 Methodology: KQL and NHP

37 Funding acquisition: HVM

38 Project administration: TTTH and HVM

39 Supervision: NTL, PTQN, MT, and KP

40 Visualization: KQL

41 Writing – original draft: KQL, HTNA, and NHP

42 Writing – review & editing: TTTH, KP, MT, NTL, PTQN, LPA, LVT, TQB, OPT, NVH,  
43 VTHL, and HVM

44 All authors read and approved the final manuscript

45 **Acknowledgments**

46 The authors would like to thank all students who participated in the study as well as the  
47 individuals and institutions that made this study possible: Departments of Education and  
48 Training from selected provinces, principals, and teachers from 81 schools who helped us to  
49 prepare for the data collection. Associate Prof. Tran Dac Phu and Dr. Truong Dinh Bac from the  
50 General Department of Preventive Medicine – Ministry of Health; Mr. Nguyen Thanh De and  
51 Mr. Le Manh Hung from the Ministry of Education and Training; Ms. Leanne Riley from World  
52 Health Organization; Ms. Veronica Lea, Ms. Curtis Blanton and Mr. Timothy McManus from  
53 the US CDC; and Mr. Cao Huu Quang from Hanoi University of Public Health.

54 **Abstract**

55 **Background**

56 Adolescence is a vulnerable period for many lifestyle risk behaviors. In this study, we aimed to  
57 1) examine a clustering pattern of lifestyle risk behaviors; 2) investigate roles of the school  
58 health promotion programs on this pattern among adolescents in Vietnam.

59 **Methods**

60 We analyzed data of 7,541 adolescents aged 13–17 years from the 2019 nationally representative  
61 Global School-based Student Health Survey, conducted in 20 provinces and cities in Vietnam.  
62 We applied the latent class analysis to identify groups of clustering and used Bayesian 2-level  
63 logistic regressions to evaluate the correlation of school health promotion programs on these  
64 clusters. We reassessed the school effect size by incorporating different informative priors to the  
65 Bayesian models.

66 **Findings**

67 The most frequent lifestyle risk behavior among Vietnamese adolescents was physical inactivity,  
68 followed by unhealthy diet, and sedentary behavior. Most of students had a cluster of at least two  
69 risk factors and nearly a half with at least three risk factors. Latent class analysis detected 23%  
70 males and 18% females being at higher risk of lifestyle behaviors. Consistent through different  
71 priors, high quality of health promotion programs associated with lower the odds of lifestyle risk  
72 behaviors (highest quality schools vs. lowest quality schools; males: Odds ratio (OR) = 0.67,  
73 95% Highest Density Interval (HDI): 0.46 – 0.93; females: OR = 0.69, 95% HDI: 0.47 – 0.98).

74 **Interpretation**

75 Our findings demonstrated the clustering of specific lifestyle risk behaviors among Vietnamese  
76 in-school adolescents. School-based interventions separated for males and females might reduce  
77 multiple health risk behaviors in adolescence.

#### 78 **Funding**

79 The 2019 Global School-based Student Health Survey was conducted with financial support  
80 from the World Health Organization. The authors received no funding for the data analysis, data  
81 interpretation, manuscript writing, authorship, and/or publication of this article.

82 **Keywords:** *Risk behaviors clustering, Adolescents, GSHS, Vietnam, Latent class analysis,*  
83 *Bayesian*

## 84 **Introduction**

85 Non-communicable diseases (NCDs) are responsible for almost 70% of all deaths worldwide [1]  
86 and becoming more common among youth nowadays [2]. Health risk behaviors are activities  
87 that increase the risk of diseases or injuries if doing with enough frequency or intensity [3].  
88 These behaviors begin in early life that affects health both at that time and later years [4].  
89 According to WHO, two-thirds of premature deaths in adults are associated with childhood risk  
90 behaviors, such as 81% of youth aged 11–17 years were physical inactivity and 11.7% were  
91 heavy drinking [5]. Most NCDs share predisposing risk factors, for example, people suffered  
92 from diabetes and cancer as concurrent exposure to unhealthy diets, physical inactivity, and  
93 harmful use of tobacco and/or alcohol [5]. These factors are unlikely to isolate, but, instead,  
94 typically cluster and interact to exponentially elevate the risks of NCDs [6]. Findings in more  
95 than 300,000 adolescents from 89 countries showed that 82.4% of them exposed to several  
96 NCDs risk factors, including unhealthy diet and physical inactivity; or unhealthy diet and  
97 cigarette smoking [7].

98 Adolescence is a critical period for developing and forming a healthy lifestyle [8], and also a  
99 vulnerable period for several unhealthy behaviors which might continue into maturity [9].  
100 Adolescents tend to be involved in more than one problem due to shared linkages of such  
101 behaviors [10-13]. School going adolescents spent at least six hours a day at school on their  
102 social, physical, and intellectual development. The school setting becomes an ideal place for  
103 targeted lifestyle programs that equipped healthy behaviors for adolescents before their transition  
104 into adulthood [14]. Some previous studies suggested that school health programs can reduce the  
105 prevalence of health risk behaviors among youth [15]. Although school environments can affect  
106 student health [16], important evidence gaps about the roles of schools in alleviating health risk

107 factors in adolescents remain. Understanding this gap is essential to improve adolescent health  
108 and reduce subsequent disease burden in adulthood.

109 In Vietnam, a surged prevalence of NCDs in recent years is associated with the country's  
110 remarkable economic growth and lifestyle changes [17]. In 2016, NCDs contributed to 73% of  
111 total deaths due to dietary risks, tobacco smoke, alcohol use, and physical inactivity [18]. The  
112 Vietnam national NCDs strategy 2015-2025 focuses on preventing NCDs among adolescents  
113 aimed to reduce smoking prevalence below 3-6%; overweight and obesity prevalence below 10%  
114 in this population [19].

115 Existing literature about the clustering of lifestyle risk behaviors in low- and middle-income  
116 countries (LMICs) is currently limited. However, most of the prior studies used descriptive  
117 techniques to analyze risk behaviors individually using arbitrary cut-off points [20] or  
118 determining high-risk groups via the ratio of observed-to-expected prevalence [21, 22]. Also,  
119 limited evidence has focused on the roles of school health promotion programs. In Vietnam, to  
120 the best of our knowledge, no study published on clustering of risk behaviors to date. In this  
121 paper, we aimed to: 1) describe a clustering pattern of six lifestyle risk behaviors (smoking,  
122 drinking, physical inactivity, sedentary behaviors, low fruit/vegetable intake, and unhealthy diet)  
123 using latent class analysis (LCA) - an analytical technique to find underlying patterns of such  
124 behaviors; 2) to investigate the roles of the school health promotion programs on these behavior  
125 patterns using Bayesian multilevel models with different informative priors.

## 126 **Methods**

### 127 **Study population and survey design**

128 The 2019 Global School-based Student Health Survey (GSHS) is a population-based survey of  
129 school-going adolescents aged 13-17 years, which has been conducted over 101 countries [23],

130 providing data on different aspects of adolescent behaviors and protective factors to help  
131 countries develop suitable adolescent health programs and policies [23]. In this study, we used  
132 the nationally representative sample of Vietnam, which was conducted from May to December  
133 2019 across 20 provinces and cities in Vietnam.

#### 134 **Questionnaire development**

135 The 2019 Vietnam GSHS employed a set of global self-administered questions that adapted to  
136 the local socio-cultural context. A panel including four language and content experts was  
137 established to validate the content. First, the experts translated forward the original English  
138 questionnaire into Vietnamese, then backward translation from Vietnamese into English. The  
139 original English version was compared to backward translation to check consistency. Face  
140 validity was assessed from a pilot study through pre-testing the translated questionnaire in 120  
141 students in one secondary school and one high school in Hanoi city, Vietnam. The validated  
142 questionnaire has two main components: socio-demographics (age, gender, living status, and  
143 place of residence) and health behaviors (alcohol use, tobacco use, dietary behavior, self-  
144 perceived body mass index (BMI) status, illicit drug use, mental health, physical activity, sexual  
145 behaviors, and violence and unintentional injuries).

#### 146 **Sample size and sampling procedures**

147 We used a two-stage cluster sampling method to recruit adolescents. At the first stage, we chose  
148 schools based on probability proportional to size method, then following by the selection of  
149 classes using the simple random sampling technique at the second stage. Two classes for each  
150 secondary school (grades 8 and 9) and three classes for each high school (grades 10, 11, and 12)  
151 were selected. All students who were 13–17 years old and had Vietnamese citizenship in selected  
152 classes were eligible to participate. The sample included 7,796 students in 210 classes from 81

153 schools. The school response rate of 96.4% and the student response rate of 97.0% made up the  
154 overall response rate of 93.5%.

155 Before data collection, school administrators assisted to distribute written assent forms to the  
156 parents/guardians in selected classes. Parents/guardians were asked to return the forms regardless  
157 of their agreement. During data collection days, we only recruited students with parental  
158 permission to participate into the study. Trained researchers briefed and guided on the self-  
159 administered questionnaire, then students marked their responses on a separate computer scan-  
160 able answer sheet. The time to complete the questionnaire was maximum 30 minutes. All  
161 completed anonymous sheets were sealed in envelopes to ensure confidentiality.

## 162 **Variables**

### 163 *Lifestyle risk behaviors*

#### 164 *Smoking*

165 We defined the current smoking status by asking students the number of days that they used any  
166 tobacco products in the past 30 days. The variable response was dichotomized into “0 days” and  
167 “at least 1 day”.

#### 168 *Alcohol consumption*

169 We assessed the current drinking by a question on the number of days that students drank at least  
170 one standard drink of alcoholic beverage (~14gram of pure alcohol) in the past 30 days. The  
171 variable response was categorized as “0 days” and “at least 1 day”.

#### 172 *Physical inactivity*

173 Physical activity is defined as any body movement generated by the contraction of skeletal  
174 muscles that raises energy expenditure above resting metabolic rate [24].

175 We asked students a question on the number of days that they had at least 60 minutes of  
176 moderate-to-vigorous-intensity physical activity in the past seven days. We defined the physical  
177 inactivity if respondents did not meet the WHO recommendation for physical activity [25]. Thus  
178 students were categorized as “physical inactivity” if they did not report doing at least an average  
179 of 60 minutes per day of moderate-to-vigorous intensity physical activities in seven days a week.

#### 180 *Sedentary behavior*

181 Sedentary behaviors are defined as any waking behaviors that consumed an energy expenditure  
182 equal to or below 1.5 metabolic equivalents. Screen time and sitting time are usually the two  
183 main indicators used to quantify the time devoted to sedentary behaviors [26, 27].

184 To assess the sedentariness, we asked students a question on the hour spent on sitting and  
185 watching television, playing computer games, using social media, or doing other sitting  
186 activities, on an average day in the past seven days. We classified respondents as having  
187 sedentary behavior if they spent on more than two hours a day doing those activities [28].

#### 188 *Low fruit/vegetable intake*

189 We asked students questions on the average daily frequency intake of fruits and vegetables in the  
190 past 30 days. We defined the low fruit/vegetable intake if they did not consume both fruits and  
191 vegetables at least two times per day [29].

#### 192 *Unhealthy diet*

193 We defined unhealthy diet students if they drank carbonated soft drinks at least one time per day  
194 during the 30 days before the survey or ate fast-food at least one day during the 7 days before the  
195 survey [29].

#### 196 *School health promotion programs*

197 We asked students whether they received training for five soft-skills (Yes/no), including (1) the  
198 benefits of eating more fruits and vegetables, (2) signs of depression and suicidal behaviors, (3)  
199 problems associated with drinking alcohol, (4) problems associated with using drugs, and (5)  
200 benefits of physical activity. We then aggregated these variables at the school level and  
201 constructed the proxy measurement of school health promotion programs quality by using PCA  
202 to compute a composite index. This composite index was categorized into tertiles [30], with  
203 higher tertiles indicating better school program quality.

#### 204 *Other covariates*

205 The other covariates include demographic characteristics (age, place of residence, living with  
206 mother/father), body mass index–Z score, parental monitoring, peer-relationship (number of  
207 close friends), mental health (loneliness and worrying), and truancy. The details on variable  
208 definitions are provided in **Supplemental materials S2**.

#### 209 **Data analysis**

210 After removing missing values on covariates (3.3%), which was assumed to be missing at  
211 random, the complete case sample analyzed in this paper was 7,541 (3,495 males and 4,046  
212 females).

#### 213 *Sampling weights calculation*

214 We calculated the sampling weights reflecting the likelihood of sampling each student and to  
215 reduce bias by compensating for different patterns of nonresponse. The sampling weights were  
216 given by

$$217 \quad W = W1 * W2 * f1 * f2 * f3$$

218 In which, W1 is the inverse of the probability of selecting the school; W2 is the inverse of the  
219 probability of selecting the classroom within the school; f1 is a school level nonresponse

220 adjustment factor; f2 is a student-level nonresponse adjustment factor; f3 is a post-stratification  
221 adjustment factor calculated by rural/urban and grade.

### 222 *Descriptive analysis*

223 We used frequencies and percentages to summarize categorical variables, mean and standard  
224 deviation (SD) to describe continuous variables. We adjusted all estimations for the complex  
225 survey design, including sampling weights, clustering, and stratification. To describe the  
226 combinations of six lifestyle risk behaviors, we used the UpSet diagrams [31], which visualize  
227 complex intersections of a lifestyle risk behaviors matrix where the rows represent different sets  
228 of combinations and the columns represent the number of students who had these combinations.

### 229 *Lifestyle risk behaviors clustering and related factors*

230 We developed an analysis framework (**Figure 1**) that includes two steps to analyze the clustering  
231 and related factors.

232 **(Figure 1 is about here)**

233 In the first step, we used LCA to identify homogeneous unobservable subgroups of lifestyle risk  
234 behaviors. LCA is a statistical method for identifying unmeasured class membership among  
235 subjects using categorical or continuous observed variables [32]. To explore the potential  
236 number of latent classes, we tested the LCA with a different number of classes (i.e., from two to  
237 five classes). We determined the final number of classes based on the interpretability of the class  
238 memberships [32]. The details in LCA results were provided in the **Supplemental materials S3**.

239 In the second step, we fitted a series of Bayesian 2-level random intercept logistic regressions  
240 with students at level-1, and school at level-2 to evaluate the effects of factors in school and  
241 student levels on latent class memberships obtained from LCA in the first step. Multilevel  
242 modeling was used to account for the nature of hierarchical data i.e., students nested in schools.

243 We applied the vague priors (i.e., flat prior for model parameters). Model 1 was a null model  
244 with no independent variable, Model 2 included the school-level factors ( $X_{1j}$ ), and Model 3  
245 controlled for both school-level and student-level covariates ( $X_{2ij}$ ).

246 
$$y_{ij} \sim \text{Binomial}(n_{ij}, \pi_{ij})$$

247 
$$\text{Model 1: } \text{logit}(\pi_{ij}) = \beta_0 + \mu_{0j}$$

248 
$$\text{Model 2: } \text{logit}(\pi_{ij}) = \beta_0 + \beta_1 * X'_{1j} + \mu_{0j}$$

249 
$$\text{Model 3: } \text{logit}(\pi_{ij}) = \beta_0 + \beta_1 * X'_{1j} + \beta_2 * X'_{2ij} + \mu_{0j}$$

250 The priors for model parameters were specified as:

251 
$$\text{Prior } \beta_0 \sim \text{Student}_3(0, 10)$$

252 
$$\text{Prior } \beta_1 \text{ and prior } \beta_2 \sim \text{flat}$$

253 
$$\text{Prior } \mu_{0j} \sim \text{Student}_3(0, 10)$$

254 Where  $\mu_{0j}$  is the school-specific residual that presents the deviate of each school from the median  
255 log odds of high risk of lifestyle behaviors. For each model, we calculated the variance partition  
256 coefficient (VPC) and the proportion of variance explained by the added factors (i.e., %  
257 explained). Since logistic regression has a variance of 3.29 [33], the VPC is defined as  $VPC =$   
258  $\sigma^2_{\mu_0} / (\sigma^2_{\mu_0} + 3.29)$ . To obtain % explained, we subtracted the variance of the simpler model to the  
259 model with more terms and then converted to percentages. The school-level and student-level  
260 covariates included in the model were determined by the conceptual framework of effects of  
261 school health promotion programs on lifestyle risk behaviors that was based on the conceptual  
262 framework developed by Akseer et al [34]. The conceptual framework is shown in  
263 **Supplemental materials S4.**

264 We presented the median of posterior distributions as odds ratio (OR) with 95% highest density  
265 interval (95% HDI). We also calculated the probability of the posterior distribution of school

266 effect size (i.e., ORs) less than the cut-offs of 1, 0.9, and 0.8 as well as Bayes factors (BF). The  
267 BF is the ratio of the likelihood of a specific hypothesis to the opposite hypothesis [35].  
268 Therefore, in this case, the larger value of the BF indicates the stronger evidence of the  
269 hypothesis that the school effect size was lower than the cut-offs.

### 270 *Sensitivity analysis*

271 We conducted sensitivity analyses to test whether the school effects remain consistent across  
272 different priors. We constructed two priors to reflect two degrees of belief that the school health  
273 promotion programs had either no impact (*equivocal prior*) or positive impact (*optimistic prior*)  
274 on lifestyle risk behaviors of students. We assumed that prior distributions of school effects  
275 were the normal distribution (i.e.,  $\beta_l \sim \text{Normal}(\mu, \sigma^2)$ ).

276 Equivocal prior:  $\beta_l \sim \text{Normal}(0, 0.207^2)$

277 Optimistic prior:  $\beta_l \sim \text{Normal}(-0.693, 0.207^2)$

278 Details of sensitivity analyses were provided in **Supplemental materials S4**.

279 We fitted the Bayesian multilevel models using Hamiltonian Monte Carlo (HMC) algorithm.  
280 With the expected effect sample sizes were greater than 10000, we generated four HMC chains  
281 in parallel, the iterations of 8000, the burn-in period of 1000, and the thinning of 2. The  
282 convergence of HMC chains was diagnosed by trace plot and the Gelman-Rubin coefficient [36].  
283 We used Stata v16 SE (StataCorp, College Station, TX, USA) to clean the data and conduct  
284 descriptive analysis; R (version 4.0.0, R Foundation for Statistical Computing, Vienna, Austria)  
285 to conduct the LCA using package *poLCA* [37] and the Bayesian multilevel models using  
286 package *brms* that interfaces with Stan [36].

### 287 **Ethical approval**

288 All procedures performed in this study followed the ethical standards of the Institution Review  
289 Board of Hanoi University of Public Health (IRB decision No. 421/2019/YTCC-HD3, dated:  
290 06/08/2019).

## 291 **Results**

### 292 *Participant characteristics*

293 **Table 1** describes the characteristics of the study participants. Our weighted sample represented  
294 a population of 5,400,584 adolescents. More than half of students were females (54.1%) and  
295 nearly two thirds of them lived in rural areas. Most of them lived with their mother or father and  
296 had more than two close friends. The proportion of students felt lonely and anxious in the past 12  
297 months was 11.7% and 4.8% in males and 12.8% and 7.2% in females, respectively. The  
298 percentages of students truanted in the past 30 days were 17.1% and 13.0% among males and  
299 females.

300 More than 70% of students were taught about the benefits of eating more fruits and vegetables,  
301 problems associated with drinking alcohol, drug consequences, and benefits of physical activity,  
302 but only about 20% of them received training about signs of depression and suicidal behaviors.  
303 Less than half of students achieved three such modules together (45.4% in males and 45.9% in  
304 females). Results from PCA analysis showed that more than 40% of students studied in high-  
305 quality schools.

306 **(Table 1 is about here)**

### 307 *Distribution of lifestyle risk behaviors*

308 **Figure 2** shows the profile of lifestyle risk behaviors among in-school adolescents in Vietnam.  
309 The physical inactivity was the most frequent lifestyle risk behavior, followed by unhealthy diet  
310 and sedentary behavior. The percentage of smoking and drinking were higher among males than

311 females (4.4% vs. 1.0%, and 24.7% vs. 20.0%, respectively). In contrast, females were more  
312 inactive than males, with a high prevalence of physical inactivity (90.1% vs. 77.9%) and  
313 sedentary behavior (47.8% vs. 37.3%).

314 Almost all students had at least one risk factor (96.8% in males and 98.5% in females). Many  
315 students had a combination of two (34.2% in males and 34.6% in females) and three factors  
316 (27.3% in males and 31.5% in females). Only <1% students had none of the six risk factors.  
317 Males and females equally distributed in each category of the number of risk factors.

318 **(Figure 2 is about here)**

319 **Figure 3** describes the frequent combinations formed by six lifestyle risk behaviors. The most  
320 frequent cluster in both sexes was physical inactivity and unhealthy diet, with the proportion of  
321 15.7% and 16.7% in males and females, respectively. The second most frequency was physical  
322 inactivity, sedentary behavior, and unhealthy diet in females, while physical inactivity in males.  
323 The combination of four risk factors physical inactivity, sedentary behavior, low fruit/vegetable  
324 intake, and unhealthy diet was the third most common combination.

325 **(Figure 3 is about here)**

### 326 *Cluster of lifestyle risk behaviors*

327 Based on the criteria of interpretability from LCA analysis, we decided to choose two latent  
328 classes as the outcome for the clustering of lifestyle risk behaviors modeling. We named the two  
329 classes based on the distribution of each risk behavior in each class (i.e., higher and lower risk  
330 class of lifestyle behaviors). **Table 2** presents the distribution of lifestyle risk behaviors among  
331 participants by two class memberships. We detected 23.1% of males and 18.0% of females were  
332 at higher risk cluster of lifestyle behaviors. Details on LCA results were in the **Supplemental**  
333 **material S3**.

334 (Table 2 is about here)

335

336 *Role of the school health promotion programs on high-risk cluster of lifestyle behaviors*

337 **Table 3** shows factors related to a high level of lifestyle risk behaviors in Vietnamese male and  
338 female adolescents. After controlling for student-level covariates, students in high-quality  
339 schools (3rd tertile) were less likely to have high level of lifestyle risk behaviors than those in  
340 low-quality schools (1st tertile) (Males: OR = 0.67, 95% HDI: 0.46 – 0.93; Females: OR = 0.69,  
341 95% HDI: 0.47 – 0.98).

342 In the base model without independent variable (Model 1), most variation of lifestyle risk  
343 behaviors clustering attributed to the student level; the school-level accounted for 12.4% and  
344 13.3% of the variation among males and females, respectively. The inclusion of school-level  
345 factor explained ~12% the between-school variation of lifestyle risk behaviors clustering, while  
346 student-level covariates explained further 42.9% and 36.0% of the variation.

347 (Table 3 are about here)

348 *Sensitivity analysis*

349 **Figure 4** and **Supplemental materials S4** provide the sensitivity analysis of the effects of school  
350 health promotion programs on students' behaviors. These effects were different due to different  
351 priors, however, yielded the same conclusion: high-quality schools were associated with lower  
352 odds of high level of lifestyle risk behaviors in both males and females. Among males, the  
353 proportion of posterior distribution less than the cutoff of OR = 1 was from 96.8% (BF = 30.3) in  
354 equivocal prior to 100% (BF > 10000) in the optimistic prior. Among females, all results favored  
355 the hypothesis that the proportion of posterior distribution was less than the cutoff of OR = 1,  
356 with a minimum of 94.8% (BF = 18.1) in the equivocal prior. However, in equivocal prior, only

357 56.4% the posterior distribution of the effects of school quality on students' behaviors was below  
358 the cutoff of OR = 0.8 among males; this figure for females was 49.7%.

359 **(Figure 4 is about here)**

### 360 *Other factors related to high-risk cluster of lifestyle behaviors*

361 In both sexes, parental monitoring is associated with lower odds of lifestyle risk behaviors. In  
362 contrast, older students, those who felt lonely, or did truancy were more likely to have a higher  
363 level of lifestyle risk behaviors than those who did not. We also found that worrying was  
364 associated with higher lifestyle risk behaviors among females (**Table 3**).

### 365 **Discussion**

366 This is the first study to investigate the prevalence, clustering pattern of six major lifestyle risk  
367 behaviors of NCDs (smoking, drinking, physical inactivity, sedentary behavior, low  
368 fruit/vegetable intake, and unhealthy diet) and its determinants among school-going adolescents  
369 aged 13-17 years in the nationwide scope in Vietnam. We found that lifestyle risk behaviors  
370 were common in Vietnamese adolescents. Among all NCDs' risk factors, physical inactivity  
371 constituted the highest proportion in both males and females, followed by unhealthy diet and  
372 sedentary behavior intake. Most of students had a cluster of at least two risk factors and nearly a  
373 half with at least three risk factors. These factors tend to be clustered with the common patterns  
374 among physical inactivity and unhealthy diet in both sexes and sedentary behavior in females.  
375 The school health promotion programs quality is associated with lower the odds of lifestyle risk  
376 behaviors. Our study provides important empirical evidence to guide health promotion,  
377 education, and interventional programs of these lifestyle risk behaviors.

378 Previous studies have demonstrated the health risk behaviors among adolescents, however, we  
379 cautioned our comparisons due to variations in investigating different risk factors, definitions

380 and cut-off points; and differing in targeted populations. Compared to a study in 2538 Malaysian  
381 school-going adolescents that employed the same definitions for smoking, alcohol use, sedentary  
382 behavior, and low fruit/vegetable intake, Vietnamese ones had a higher prevalence of alcohol use  
383 (24.7% in Vietnamese males and 20.0% in Vietnamese females vs. 6.6% in Malaysian males and  
384 3.5% in Malaysian females) but lower levels of smoking, sedentary behavior, and low  
385 fruit/vegetable intake [22]. However, compared to another study conducted among 3990  
386 Brazilian adolescents, Vietnamese adolescents had a similar prevalence of drinking alcohol  
387 (~25%) but lower in smoking (4.4% in males and 1% in females in the present study vs 5.7%)  
388 [21].

389 Clustering multiple lifestyle risk behaviors are prevalent in Vietnamese adolescents. The  
390 prevalence of simultaneous occurrences of at least two risk behaviors was lower than that  
391 reported among Malaysian adolescents (~60% vs. 83%) [22], Canadian children and adolescents  
392 (64%) [38] and Brazilian adolescents (53.8% in males and 71.1% in females) [21]. In  
393 concordance with previous literature, an physical inactivity, sedentary behavior, and unhealthy  
394 diet were prone to cluster either alone or together with other risk behaviors in both sexes [39,  
395 40]. Physical inactivity, sedentary behavior, and unhealthy diet are three common obesogenic  
396 characteristics. The mechanism of clustering these obesogenic behaviors among adolescents is  
397 complex [41]. Prolonged physical inactivity and sedentary behavior are associated with an  
398 unhealthy diet by increasing consuming junk foods while doing sedentary activities such as  
399 sitting, watching television, or using a computer. This disrupts energy balance by increasing time  
400 for recreational physical activity that in turn can promote fat accumulation [42]. Physical  
401 inactivity and sedentary behavior combined with an unhealthy diet is negatively associated with  
402 health and wellbeing among adolescents and has been recognized as important lifestyle risk

403 behaviors [43]. The clustering of health risk behaviors follow Jessor's problem-behavior theory  
404 that adolescents engaged in one problem behavior tend to be involved in other problems due to  
405 the shared linkages of such behaviors in the social ecology [10]. We recommend that joint  
406 interventions on multiple risk behaviors can be more effective than individual approaches to  
407 tackle these synergistic clustering effects.

408 We observed more profound clustering effects of lifestyle risk behaviors in Vietnamese females  
409 than in males, and such findings corroborated current evidence [21, 22, 38]. Although generally  
410 known as lower risk-takers [44], adolescent females who engaged in one high-risk behavior tend  
411 to be more involved in others than their male counterparts. Furthermore, they have higher levels  
412 of perceived stress [45], a tendency to ruminate, and feelings of helplessness than males [46] that  
413 could predispose them to engage in dysfunctional coping measures [47]. Previous evidence  
414 indicated that adolescent females were more likely to be influenced by psychosocial motives [45,  
415 46] and therefore more likely to engage in unhealthy risk behaviors [47]. As such, in-depth  
416 investigations into the actual causal factors could probably shed some light on the significant  
417 determinants of the clustering of risk behaviors among Vietnamese females. This finding  
418 implicated the separated design of public health interventions for males and females are needed.

419 Our evidence supports the idea that the school health promotion programs are important for  
420 promoting and supporting healthy lifestyles among adolescents [48-50]. Previous studies  
421 indicated that school-based interventions are effective and feasible to improve healthy dietary  
422 habits, physical activity levels, and weight control [51]. Schools also reach a wide range of  
423 children over a considerable amount of time. Therefore, enhancing the school health promotion  
424 programs could be a prominent way to improve children's health and well-being [52, 53]. In  
425 Vietnam, education on preventing lifestyle risk behaviors is not prerequisites in school; they are

426 considered as elective modules; therefore, the quality, as well as content, might be not sufficient.

427 It is noticed that despite of the high prevalence of studying an individual module for preventing

428 lifestyle risk behavior, only less than half of students completed three or more such modules.

429 This lack of joint knowledge on preventing lifestyle risk behaviors might lead to the

430 pervasiveness of lifestyle risk behaviors in our sample. We highlight a need for the required

431 courses that not only training on the harms of each risk behavior separately but also emphasize

432 the connection of these behaviors [54].

433 Regarding sensitivity analysis, we used the Bayesian approach to analyze the effect of school

434 health promotion programs quality on students' behaviors among different scenarios of belief.

435 All the scenarios, even in the equivocal view which is conservative to our belief, yield the same

436 conclusion, it is more certain and confident to claim the robustness of findings. Another

437 advantage of using Bayesian approach is its ability to measure the effect size. In our study, we

438 found the strong evidence that high tertile of school health promotion programs quality had

439 positively affect students' behaviors (i.e., ORs < 1) compared to low tertile of quality, however,

440 this effect became uncertain when compared to the cutoff of 0.8, (i.e., only 56.4% and 49.7% the

441 posterior distribution was below the cutoff of OR = 0.8 among males and females in the

442 equivocal prior, respectively). Together, our findings suggest that the effect size of school health

443 promotion programs quality on students' behaviors probably ranges from 0.8 to 1.

444 Strengths of our study included using robust statistical technique to investigate clusters of NCDs

445 risk factors and its determinants among adolescents using large nationally representative sample.

446 We used latent class analysis that was able to identify the underlying pattern of lifestyle risk

447 behaviors co-occurrence and Bayesian regression models with different informative priors to

448 investigate the relationship between these cluster and school health promotion programs. This is

449 the first study to investigate the current status of lifestyle risk behaviors in Vietnam. Most current  
450 evidence investigating clusters of risk factors was done on adult populations or in high-income  
451 countries; not many studies conducted in lower-middle income countries. Despite its innovative  
452 approach, the present analysis has certain limitations. First, information on lifestyle risk  
453 behaviors was self-reported, which may lead to under or overestimation. Second, due to the  
454 availability of the data, the cutoff point for low fruit/vegetable intake cannot follow WHO  
455 recommendation. Third, we are unable to collect various aspects of both the school and student's  
456 levels, such as parental education, income, social-economic status, which maybe potential  
457 predictors of clustering risk behaviors. Fourth, due to the cross-sectional nature of the data, we  
458 could not confirm the causal inferences. Fifth, although most adolescents in Vietnam are in  
459 school [55], our study focused only in-school adolescents, which may not represent for those  
460 who are not in school.

## 461 **Conclusion**

462 Our findings demonstrated the clustering of specific combinations of lifestyle risk behaviors  
463 among Vietnamese in-school adolescents. School-based interventions separated for males and  
464 females might reduce multiple health risk behaviors in adolescence. A deeper understanding of  
465 clustered patterns may lead to developing new and comprehensive interventions to prevent the  
466 burden of NCDs in Vietnamese adolescents.

## 467 **References**

- 468 1. World Health Organization. Noncommunicable diseases: WHO;  
469 [https://www.who.int/health-topics/noncommunicable-diseases#tab=tab\\_1](https://www.who.int/health-topics/noncommunicable-diseases#tab=tab_1). Accessed 25 Nov  
470 2020.
- 471 2. Baker R, Taylor E, Essafi S, Jarvis JD, Odok C. Engaging young people in the prevention  
472 of noncommunicable diseases. *Bull World Health Organ.* 2016;94(7):484.
- 473 3. Steptoe A. Health Behavior and Stress\*. In: Fink G, editor. *Encyclopedia of Stress*  
474 (Second Edition). New York: Academic Press; 2007. p. 262-6.
- 475 4. El Achhab Y, El Ammari A, El Kazdough H, Najdi A, Berraho M, Tachfouti N, et al.  
476 Health risk behaviours amongst school adolescents: protocol for a mixed methods study. *BMC*  
477 *public health.* 2016;16(1):1209.
- 478 5. World Health Organization. WHO Global Coordination Mechanism on the Prevention  
479 and Control of NCDs: NCD and Youth: WHO; [https://www.who.int/global-coordination-](https://www.who.int/global-coordination-mechanism/ncd-themes/ncd-and-youth/en)  
480 [mechanism/ncd-themes/ncd-and-youth/en](https://www.who.int/global-coordination-mechanism/ncd-themes/ncd-and-youth/en). Accessed 25 Nov 2020.
- 481 6. Arena R, Guazzi M, Lianov L, Whitsel L, Berra K, Lavie CJ, et al. Healthy lifestyle  
482 interventions to combat noncommunicable disease-a novel nonhierarchical connectivity model  
483 for key stakeholders: a policy statement from the American Heart Association, European Society  
484 of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and  
485 American College of Preventive Medicine. *European heart journal.* 2015;36(31):2097-109.
- 486 7. Uddin R, Lee EY, Khan SR, Tremblay MS, Khan A. Clustering of lifestyle risk factors  
487 for non-communicable diseases in 304,779 adolescents from 89 countries: A global perspective.  
488 *Preventive medicine.* 2020;131:105955.

- 489 8. Lee RL, Loke AY, Wu CS, Ho AP. The lifestyle behaviours and psychosocial well-being  
490 of primary school students in Hong Kong. *Journal of clinical nursing*. 2010;19(9-10):1462-72.
- 491 9. Ness AR, Maynard M, Frankel S, Smith GD, Frobisher C, Leary SD, et al. Diet in  
492 childhood and adult cardiovascular and all cause mortality: the Boyd Orr cohort. *Heart*.  
493 2005;91(7):894-8.
- 494 10. Jessor R. Risk behavior in adolescence: A psychosocial framework for understanding and  
495 action. *Journal of Adolescent Health*. 1991;12(8):597-605.
- 496 11. Nascente FM, Jardim TV, Peixoto MD, Carneiro CS, Mendonça KL, Póvoa TI, et al.  
497 Sedentary lifestyle and its associated factors among adolescents from public and private schools  
498 of a Brazilian state capital. *BMC public health*. 2016;16(1):1177.
- 499 12. Oyeyemi AL, Ishaku CM, Oyekola J, Wakawa HD, Lawan A, Yakubu S, et al. Patterns  
500 and Associated Factors of Physical Activity among Adolescents in Nigeria. *PLoS One*.  
501 2016;11(2):e0150142-e.
- 502 13. Urrutia-Pereira M, Oliano VJ, Aranda CS, Mallol J, Solé D. Prevalence and factors  
503 associated with smoking among adolescents. *Jornal de pediatria*. 2017;93(3):230-7.
- 504 14. DeBar LL, Ritenbaugh C, Aickin M, Orwoll E, Elliot D, Dickerson J, et al. Youth: a  
505 health plan-based lifestyle intervention increases bone mineral density in adolescent girls.  
506 *Archives of pediatrics & adolescent medicine*. 2006;160(12):1269-76.
- 507 15. Bonell C, Wells H, Harden A, Jamal F, Fletcher A, Thomas J, et al. The effects on  
508 student health of interventions modifying the school environment: systematic review. *Journal of*  
509 *Epidemiology and Community Health*. 2013;67(8):677.

- 510 16. Bonell C, Parry W, Wells H, Jamal F, Fletcher A, Harden A, et al. The effects of the  
511 school environment on student health: a systematic review of multi-level studies. *Health & place*.  
512 2013;21:180-91.
- 513 17. Nguyen TT, Hoang MV. Non-communicable diseases, food and nutrition in Vietnam  
514 from 1975 to 2015: the burden and national response. *Asia Pacific journal of clinical nutrition*.  
515 2018;27(1):19-28.
- 516 18. World Health Organization. United Nations Interagency Task Force on the prevention  
517 and control of noncommunicable diseases. Joint mission, Vietnam. Geneva: WHO Press; 2016.
- 518 19. Ministry of Health of Vietnam. National strategy for the prevention and control of  
519 communicable diseases, period 2015-2025. Hanoi, Vietnam 2015.
- 520 20. Bener A, Ghuloum S, Abou-Saleh MT. Prevalence, symptom patterns and comorbidity of  
521 anxiety and depressive disorders in primary care in Qatar. *Social psychiatry and psychiatric*  
522 *epidemiology*. 2012;47(3):439-46.
- 523 21. Dumith SC, Muniz LC, Tassitano RM, Hallal PC, Menezes AM. Clustering of risk  
524 factors for chronic diseases among adolescents from Southern Brazil. *Preventive medicine*.  
525 2012;54(6):393-6.
- 526 22. Teh CH, Teh MW, Lim KH, Kee CC, Sumarni MG, Heng PP, et al. Clustering of  
527 lifestyle risk behaviours and its determinants among school-going adolescents in a middle-  
528 income country: a cross-sectional study. *BMC public health*. 2019;19(1):1177.
- 529 23. World Health Organization. Global school-based student health survey (GSHS): WHO;  
530 <https://www.who.int/ncds/surveillance/gshs/en>. Accessed 25 Nov 2020.

- 531 24. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical  
532 fitness: definitions and distinctions for health-related research. Public health reports  
533 (Washington, DC : 1974). 1985;100(2):126-31.
- 534 25. World Health Organization. Physical activity 2020. [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/physical-activity)  
535 [sheets/detail/physical-activity](https://www.who.int/news-room/fact-sheets/detail/physical-activity). Accessed 25 Nov 2020.
- 536 26. Gibbs BB, Hergenroeder AL, Katzmarzyk PT, Lee IM, Jakicic JM. Definition,  
537 measurement, and health risks associated with sedentary behavior. Medicine and science in  
538 sports and exercise. 2015;47(6):1295-300.
- 539 27. Thivel D, Tremblay A, Genin PM, Panahi S, Rivière D, Duclos M. Physical Activity,  
540 Inactivity, and Sedentary Behaviors: Definitions and Implications in Occupational Health. Front  
541 Public Health. 2018;6:288-.
- 542 28. Chaput J-P, Willumsen J, Bull F, Chou R, Ekelund U, Firth J, et al. 2020 WHO  
543 guidelines on physical activity and sedentary behaviour for children and adolescents aged 5–  
544 17□years: summary of the evidence. International Journal of Behavioral Nutrition and Physical  
545 Activity. 2020;17(1):141.
- 546 29. Shayo FK. Co-occurrence of risk factors for non-communicable diseases among in-  
547 school adolescents in Tanzania: an example of a low-income setting of sub-Saharan Africa for  
548 adolescence health policy actions. BMC public health. 2019;19(1):972.
- 549 30. Vyas S, Kumaranayake L. Constructing socio-economic status indices: how to use  
550 principal components analysis. Health policy and planning. 2006;21(6):459-68.
- 551 31. Lex A, Gehlenborg N, Strobel H, Vuillemot R, Pfister H. UpSet: Visualization of  
552 Intersecting Sets. IEEE Trans Vis Comput Graph. 2014;20(12):1983-92.

- 553 32. Hagenaars JA, McCutcheon AL. Applied Latent Class Analysis. Cambridge: Cambridge  
554 University Press; 2002.
- 555 33. Harvey G. Multilevel Statistical Models, 4th Edition. West Sussex: John Wiley & Sons,  
556 Ld; 2010.
- 557 34. Akseer N, Mehta S, Wigle J, Chera R, Brickman ZJ, Al-Gashm S, et al. Non-  
558 communicable diseases among adolescents: current status, determinants, interventions and  
559 policies. BMC public health. 2020;20(1):1908.
- 560 35. Goodman SN. Toward Evidence-Based Medical Statistics. 2: The Bayes Factor. Annals  
561 of Internal Medicine. 1999;130(12):1005-13.
- 562 36. Bürkner P-C. brms: An R Package for Bayesian Multilevel Models Using Stan. 2017.  
563 2017;80(1):J Journal of Statistical Software.
- 564 37. Linzer DA, Lewis JB. polCA: An R Package for Polytomous Variable Latent Class  
565 Analysis. Journal of Statistical Software; Vol 1, Issue 10 (2011). 2011.
- 566 38. Alamian A, Paradis G. Clustering of chronic disease behavioral risk factors in Canadian  
567 children and adolescents. Preventive medicine. 2009;48(5):493-9.
- 568 39. Leech RM, McNaughton SA, Timperio A. The clustering of diet, physical activity and  
569 sedentary behavior in children and adolescents: a review. Int J Behav Nutr Phys Act. 2014;11:4-.
- 570 40. Matias TS, Silva KS, Silva JAd, Mello GTd, Salmon J. Clustering of diet, physical  
571 activity and sedentary behavior among Brazilian adolescents in the national school - based health  
572 survey (PeNSE 2015). BMC public health. 2018;18(1):1283.
- 573 41. Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause  
574 mortality: a systematic review and meta-analysis. Preventive medicine. 2012;55(3):163-70.

- 575 42. Uddin R, Khan A. Sedentary behaviour is associated with overweight and obesity among  
576 adolescents: evidence from a population-based study. *Acta Paediatrica*. 2019;108(8):1545-6.
- 577 43. Carson V, Hunter S, Kuzik N, Gray CE, Poitras VJ, Chaput JP, et al. Systematic review  
578 of sedentary behaviour and health indicators in school-aged children and youth: an update.  
579 *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et*  
580 *metabolisme*. 2016;41(6 Suppl 3):S240-65.
- 581 44. Elliott MR, Shope JT, Raghunathan TE, Waller PF. Gender differences among young  
582 drivers in the association between high-risk driving and substance use/environmental influences.  
583 *Journal of studies on alcohol*. 2006;67(2):252-60.
- 584 45. Nolen-Hoeksema S. Sex differences in unipolar depression: evidence and theory.  
585 *Psychological bulletin*. 1987;101(2):259-82.
- 586 46. Nolen-Hoeksema S, Girgus JS. The emergence of gender differences in depression  
587 during adolescence. *Psychological bulletin*. 1994;115(3):424-43.
- 588 47. Dumont M, Provost MA. Resilience in Adolescents: Protective Role of Social Support,  
589 Coping Strategies, Self-Esteem, and Social Activities on Experience of Stress and Depression.  
590 *Journal of Youth and Adolescence*. 1999;28(3):343-63.
- 591 48. Choudhry S, McClinton-Powell L, Solomon M, Davis D, Lipton R, Darukhanavala A, et  
592 al. Power-up: a collaborative after-school program to prevent obesity in African American  
593 children. *Progress in community health partnerships : research, education, and action*.  
594 2011;5(4):363-73.
- 595 49. Foster GD, Sherman S, Borradaile KE, Grundy KM, Vander Veur SS, Nachmani J, et al.  
596 A policy-based school intervention to prevent overweight and obesity. *Pediatrics*.  
597 2008;121(4):e794-802.

- 598 50. Lobstein T, Baur L, Uauy R. Obesity in children and young people: a crisis in public  
599 health. *Obesity Reviews*. 2004;5(s1):4-85.
- 600 51. Fung C, Kuhle S, Lu C, Purcell M, Schwartz M, Storey K, et al. From "best practice" to  
601 "next practice": the effectiveness of school-based health promotion in improving healthy eating  
602 and physical activity and preventing childhood obesity. *Int J Behav Nutr Phys Act*. 2012;9:27.
- 603 52. Faught EL, Ekwaru JP, Gleddie D, Storey KE, Asbridge M, Veugelers PJ. The combined  
604 impact of diet, physical activity, sleep and screen time on academic achievement: a prospective  
605 study of elementary school students in Nova Scotia, Canada. *Int J Behav Nutr Phys Act*.  
606 2017;14(1):29.
- 607 53. Veugelers PJ, Fitzgerald AL. Effectiveness of school programs in preventing childhood  
608 obesity: a multilevel comparison. *Am J Public Health*. 2005;95(3):432-5.
- 609 54. Langford R, Bonell C, Jones H, Poulidou T, Murphy S, Waters E, et al. The World Health  
610 Organization's Health Promoting Schools framework: a Cochrane systematic review and meta-  
611 analysis. *BMC public health*. 2015;15(1):130.
- 612 55. Ministry of Domestic Affairs of Vietnam, UNFPA. National report on Vietnames youth.  
613 Hanoi, Vietnam2015.
- 614

615 **Table 1: Participants' characteristics**

Characteristics	Males		Females	
	n <sup>a</sup>	% <sup>b</sup>	n <sup>a</sup>	% <sup>b</sup>
<b>N weighted = 5,400,584</b>	3495	45.9	4046	54.1
<b>Student-level variables</b>				
<b>Age</b>				
13	426	15.5	569	17.3
14	667	28.6	830	32.2
15	795	21.7	828	18.1
16	744	16.3	884	15.9
17	863	18.0	935	16.5
<b>BMI-Z score, mean (SD)</b>		-0.4 (1.3)		-0.5 (1.0)
<b>Place of residence</b>				
Rural	1717	62.7	2041	64.2
Urban	1778	37.3	2005	35.8
<b>Living with mother/father</b>				
No	583	17.2	669	17.5
Yes	2909	82.8	3376	82.5
<b>Parental monitoring, mean (SD)</b>		5.7 (2.3)		5.4 (2.2)
<b>Number of close friends</b>				
0	347	8.4	417	9.0
1	386	10.3	665	15.2
2	447	12.9	699	17.6
≥3	2269	68.3	2218	58.2
<b>Loneliness</b>				
No	3026	88.3	3438	87.2
Yes	468	11.7	608	12.8
<b>Anxiety</b>				
No	3295	95.2	3700	92.8
Yes	194	4.8	344	7.2
<b>Truancy</b>				
No	2782	82.9	3418	87.0
Yes	614	17.1	544	13.0
<b>School health promotion programs</b>				
<b>Percentage of students taught about, mean (SD)</b>				
Benefits of eating more fruits and vegetables		82.3 (10.1)		82.9 (9.6)
Signs of depression and suicidal behaviors		20.8 (8.4)		21.4 (8.1)
Problems associated with drinking alcohol		70.9 (9.7)		71.5 (9.2)
Problems associated with using drugs		84.4 (6.8)		85.2 (6.7)
Benefits of physical activity		81.2 (11.7)		81.8 (11.1)
<b>Percentage of students taught &gt; 3 soft skills, mean (SD)</b>		45.4 (11.1)		45.9 (10.3)
<b>School quality proxy</b>				
1 <sup>st</sup> tertile	1232	25.4	1295	21.4
2 <sup>nd</sup> tertile	1204	34.4	1385	35.7
3 <sup>rd</sup> tertile	1059	40.2	1366	42.9

616 <sup>a</sup>Unweighted frequency, <sup>b</sup>Weighted percentage

617 **Table 2: Distribution of lifestyle risk behaviors among in-school adolescents in Vietnam,**  
 618 **by two class memberships**

Lifestyle health risk behaviors	Males		Females	
	Lower risk cluster (n = 2579, 76.9%)	Higher risk cluster (n = 916, 23.1%)	Lower risk cluster (n = 3250, 82.0%)	Higher risk cluster (n = 796, 18.0%)
Smoking, %	0.0	19.1	0.0	5.7
Alcohol consumption, %	4.1	93.5	2.8	98.3
Physical inactivity, %	78.1	77.2	90.1	90.1
Sedentary behavior, %	31.7	55.8	43.5	67.1
Low fruit/vegetable intake, %	30.6	34.9	29.1	35.3
Unhealthy diet, %	61.8	81.8	63.9	83.1
<b>Number of health risk behaviors, %</b>				
0	4.2	0.0	1.8	0.0
1	23.1	0.1	16.6	0.0
2	41.6	9.5	41.2	4.8
3	24.6	36.4	31.2	32.9
4	6.5	37.8	9.3	41.8
5	0.0	14.6	0.0	19.1
6	0.0	1.6	0.0	1.4

619 **Table 3: Bayesian multivariable models of factors related to high-risk cluster of lifestyle behaviors among in-school**  
 620 **adolescents in Vietnam**

	Model 1		Model 2		Model 3	
	Males OR (95% HDI)	Females OR (95% HDI)	Males OR (95% HDI)	Females OR (95% HDI)	Males OR (95% HDI)	Females OR (95% HDI)
<b>Fixed part</b>						
<b>Intercept</b>	0.30 (0.25–0.35)	0.20 (0.17–0.24)	0.35 (0.26–0.47)	0.23 (0.17–0.30)	0.24 (0.14–0.43)	0.13 (0.08–0.23)
<b>School quality proxy (Ref: 1<sup>st</sup> tertile)</b>						
2nd tertile			1.08 (0.73–1.62)	1.18 (0.77–1.78)	0.98 (0.70–1.38)	1.10 (0.77–1.59)
3rd tertile			<b>0.58 (0.39–0.88)</b>	<b>0.64 (0.42–0.94)</b>	<b>0.67 (0.46–0.93)</b>	<b>0.69 (0.47–0.98)</b>
<b>Student-level covariates</b>						
<b>Age (Ref: 13)</b>						
14					1.08 (0.75–1.57)	<b>1.54 (1.08–2.22)</b>
15					1.34 (0.93–1.95)	1.48 (0.99–2.22)
16					<b>2.10 (1.41–3.06)</b>	<b>2.36 (1.58–3.55)</b>
17					<b>3.40 (2.32–5.00)</b>	<b>2.24 (1.51–3.40)</b>
<b>BMI-Z score</b>					1.03 (0.96–1.10)	0.94 (0.86–1.03)
<b>Place of residence (Ref: Rural)</b>						
Urban					1.18 (0.90–1.58)	1.30 (0.96–1.76)
<b>Living with mother/father (Ref: No)</b>						
Yes					<b>0.78 (0.63–0.98)</b>	0.85 (0.68–1.05)
<b>Parental monitoring</b>					<b>0.93 (0.90–0.97)</b>	<b>0.93 (0.89–0.97)</b>
<b>Number of close friends (Ref: 0)</b>						
1					1.15 (0.78–1.61)	1.02 (0.74–1.42)
2					1.00 (0.69–1.40)	1.22 (0.90–1.71)
≥3					1.15 (0.86–1.53)	1.17 (0.89–1.57)
<b>Loneliness (Ref: No)</b>						
Yes					<b>1.34 (1.04–1.70)</b>	<b>1.53 (1.21–1.92)</b>
<b>Worrying (Ref: No)</b>						
Yes					1.20 (0.84–1.72)	<b>1.68 (1.27–2.21)</b>
<b>Truancy (Ref: No)</b>						
Yes					<b>2.23 (1.82–2.74)</b>	<b>2.01 (1.50–2.49)</b>
<b>Random part</b>						
Variance estimate	0.46 (0.29–0.72)	0.50 (0.29–0.72)	0.40 (0.25–0.64)	0.43 (0.27–0.71)	0.23 (0.12–0.40)	0.28 (0.15–0.48)
VPC (%) <sup>§</sup>	12.4	13.3	11.0	11.8	6.6	7.9
% Explained <sup>†</sup>	--	--	12.5	12.2	42.9	36.0

621 HDI: Highest density interval; OR: Odds ratio

622 Model 1: A null 2-level random effects model, with students at level-1, and school at level-2; Model 2: Model 1 + main predictors (i.e., School quality proxy); Model 3: Model 2 +

623 all student-level covariates; <sup>§</sup>VPC calculated as:  $[\sigma_{\mu 0}^2 / (\sigma_{\mu 0}^2 + 3.29)] * 100$ ; <sup>†</sup>% Explained calculated as:  $[(\sigma_{\text{Model N}}^2 - \sigma_{\text{Model N} + 1}^2) / \sigma_{\text{Model N}}^2] * 100$

624 **Figure 1: Analysis framework**

625 *Note: LCA: Latent Class Analysis; PCA: Principal Component Analysis.*

626

627 **Figure 2: Distribution of lifestyle risk behaviors among in-school adolescents in Vietnam**

628 *Note: A: Percentage of lifestyle risk behaviors among males and females; B: Distribution of*  
629 *lifestyle risk behavior co-occurrence among males and females.*

630

631 **Figure 3: UpSet diagrams for combinations of lifestyle risk behaviors**

632 *Note: The combinations that have proportion less than 1% are not shown.*

633

634 **Figure 4: Prior and posterior distributions the effects of school health promotion programs**  
635 **on high-risk cluster of lifestyle behaviors among in-school adolescents in Vietnam**

636 *Note: The red dash lines represent the three priors: Vague prior:  $\sim N(0, 1002)$ ; Equivocal prior:*  
637  *$\sim N(0, 0.2072)$ ; Optimistic prior:  $\sim N(-0.693, 0.2072)$ . In each scenario, the posterior*  
638 *distributions represent the the effect of school promotion programs quality on lifestyle risk*  
639 *behavior clustering (i.e., the ORs of school promotion programs quality tertile 3 and tertile 2 vs.*  
640 *tertile 1 – reference group). The probability of the posterior distributions less than the cut-offs of*  
641 *1, 0.9, and 0.8 (represented by three vertical lines) are shaded in a lighter gray color.*

642 *All models were adjusted for age, body mass index Z-score, place of residence, living with*  
643 *mother/father, parental monitoring, number of close friend, loneliness, worrying, and truancy.*