Retrospective feasibility study of *Mycobacterium tuberculosis* modified

lipoprotein as a potential biomarker for TB detection in children

Xinxin Yang,[a,b] Matthew F. Wipperman,[a,c] Sharon Nachman,[d] Nicole S. Sampson[a,#]

[a]Department of Chemistry, Stony Brook University, Stony Brook, NY 11794-3400

[b]Chronus Pharmaceuticals, Inc., 25 Health Sciences Drive, Stony Brook, NY 11790

[c]Current address: Clinical and Translational Science Center, Weill Cornell Medicine,

New York, NY 10065

[d]Department of Pediatrics, Renaissance School of Medicine, Stony Brook University,

Stony Brook, NY 11794-3400

Running title: Retrospective Mtb lipoprotein biomarker in children

[#]Corresponding author: nicole.sampson@stonybrook.edu

Abstract

Background: Current TB diagnostic methods available have been developed for adults

and development efforts have neglected the differences in disease and sampling that

occur between adults and children. Diagnostic challenges are even greater in HIV co-

infected children and infants.

Methods and results: We established a sandwich ELISA assay to detect

Mycobacterium tuberculosis modified lipoprotein (TLP) ex vivo in plasma. The study

population contains plasma samples from 21 patients with active TB and 24 control

samples with no TB, collected in the International Maternal Pediatric Adolescent AIDS

Clinical Trails (IMPAACT) P1041 study. Retrospective analysis was performed and the

result demonstrate that TLP level is associated with TB disease.

Conclusions: Plasma levels of TLP associate with active TB disease in HIV positive

subjects and can be used as an indicator for TB detection in children.

Keywords

Childhood TB, HIV co-infection, diagnosis, sandwich ELISA, lipoprotein, low-density

lipoprotein, apoB, biomarker

BACKGROUND

Childhood TB is estimated to contribute 12% of the disease burden, with approximately 10 million cases in 2019 worldwide [1]. Existing gold-standard diagnostic culture tools fail to confirm TB in most children, who typically have low bacterial counts and cannot produce sputum like adults. Diagnostic challenges are great in HIV co-infected children and infants [2], where the clinical presentation of pulmonary TB may be non-specific [3], acute [4, 5], and chest radiographs may be atypical [2]. Although rates of bacteriological confirmation appear similar in HIV-infected and uninfected children [6], diagnostic delay and HIV-related immune pathology contribute to higher rates of TB disease progression with increased disease severity, morbidity, and mortality [7-10]. Infants, young children, and HIV-infected children are at increased risk of developing TB following infection and of disseminated or severe disease, including TB meningitis [11]. Because TB in children is typically paucibacillary, even if a sputum sample or gastric aspirate is obtained, the utility of PCR based diagnostic methods like GeneXpert is limited. More sensitive and child-friendly diagnostic tools are urgently needed to diagnose TB in children [12].

Mycobacterium tuberculosis (*Mtb*) is an intracellular pathogen that is capable of surviving and replicating within macrophages, the frontline of the innate host defense, and has evolved multiple mechanisms to escape these immune cells [13]. One of the immune evasion strategies of *Mtb* is the deregulation of lipid metabolism, leading to the formation of foamy macrophages (FM), a hallmark of granulomata in tuberculosis lesions [14]. However, only pathogenic mycobacterial strains including *M. tuberculosis* [15], *M. avium*, *M. abscessus, or M. bovis* , but not non-pathogenic mycobacterial

strains, like *M. smegmatis*, induce foam cell formation upon infection [16, 17]. Foam

cells are associated with chronic inflammation in many metabolic diseases and certain

cancers besides infectious diseases [18, 19]. For example, foamy macrophages are

critical to the initial formation, development, and instability of atherosclerotic plaques

and are therefore therapeutic targets in atherosclerosis. Foam cell formation is induced

primarily by malondialdehyde-modified low density lipoprotein (MDA-LDL) and not by

native or extensively oxidized LDL in atherosclerosis [20]. Elevated plasma levels of

MDA-LDL in patients are associated with acute coronary syndromes and are used as

diagnostic tools clinically [21].

In this study we identified human lipoprotein modification specific to exposure to

Mtb. The pathogen modified host lipoprotein, TLP, is detectable ex vivo in plasma using

a sandwich ELISA assay. We further evaluated the association between the presence of

TLP and TB disease status in children.

METHODS

Preparation of malondialdehyde-conjugated LDL (MDA-LDL) [20] and Mtb-

modified lipoprotein (TLP). HepG2 human liver cells (ATCC HB-8065) were grown to

80% confluence in HepG2 growth media (DMEM, 10% fetal bovine serum, 20 mM L–

glutamine, 100 U/mL penicillin, 100 µg/mL streptomycin, and 10 mM HEPES). Cells

were grown for 4-5 days and culture supernatants were then harvested and

concentrated. LDL particles were separated by density gradient ultracentrifugation and

desalted by ultrafiltration through a 100-kDa molecular weight filter. MDA-LDL was

prepared by incubation 2 mg protein/mL of LDL with 2 µM acrolein at 37 °C for 24 h

under nitrogen atmosphere. *Mycobacterium tuberculosis* was cultured in Middlebrook

7H9 (broth) supplemented with 0.2% glycerol, 0.5% BSA, 0.08% NaCl, 0.05% (v/v)

tyloxapol to OD~0.7. LDL was added and the culture was incubated at 37 °C for 7 days.

TLP was isolated from the culture supernatant, concentrated and washed with PBS.

Agarose gel electrophoresis and western blot analysis. Agarose gel

electrophoresis was performed in 0.06 M barbital buffer (pH 8.6). The gel was 0.8% and

stained with Sudan Red 7B. Lipoproteins were separated on a 6% sodium dodecyl

sulfate polyacrylamide gel electrophoresis (SDS-PAGE), proteins were transferred to a

PVDF membrane and immunoblotted with specific antibodies. The primary antibodies

used for western blot were monoclonal anti-LDL (MDA oxidized) antibody (Abcam) and

polyclonal anti-apoB Ab (H-300, Santa Cruz). Membranes were treated with anti-mouse,

or anti-rabbit IgG HRP conjugates as secondary antibodies.

THP-1 macrophage preparation and treatment. THP-1 cells (ATCC TIB-202)

were maintained in RPMI-1640 medium containing heat-inactivated 10% fetal bovine

serum (FBS), 0.05 mM 2-mercaptoethanol, 100 U/mL penicillin and 100 mg/mL

streptomycin (RPMI complete medium). The cells were plated into 6-well plates

containing coverslips (1.5×10^6 /well), treated with phorbol 12-myristate 13-acetate

(PMA, 150 nM) in RPMI complete media. Media was replaced with fresh RPMI

complete media at 48 hours after plating. At 72 hours after media change, THP-1

macrophages were treated with LDL, MDA-LDL and TLP at 200 µg/mL in PBS for 24

hours. All cells were incubated in a humid atmosphere at 37°C with 95% air and 5%

CO_2.

Lipid body staining and immunostaining. Following lipoprotein treatment,

macrophages were washed with PBS and fixed with 4% paraformaldehyde at room

temperature for 30 minutes. Cells were stained with Oil Red O solution for 20 minutes at

room temperature. The slides were then counterstained with haematoxylin and

observed under an inverted microscope (Zeiss Axiovert 200M). Percent foam cell

formation was quantified by counting stained versus total cells in 10 fields.

Generation of anti-TLP antibodies [22, 23]. Monoclonal antibodies against

TLP were identified utilizing Human Combinatorial Antibody Library (HuCAL, Bio-Rad

AbD Serotec GmbH). The HuCAL phage display library was depleted of antibodies that

recognize intact LDL. The depleted library was panned for three rounds of binding,

elution, and amplification to isolate antibodies specific for TLP. Binding of anti-TLP

antibodies to control antigens including BSA, HSA, N1-CD33-His6 and intact LDL, was

checked by indirect ELISA assays, with anti-TLP antibodies as primary antibodies and

an anti-Fab-AP conjugate (Bio-Rad) as a secondary antibody.

Sandwich ELISA assay. Briefly, 96 well plates were coated overnight at 4°C

with 5 µg/mL capture antibody (AbD28580, amino acid sequence shown in Table S2) in

phosphate buffered saline (PBS). Plates were washed with PBS containing 0.05%

Tween-20 (PBST) 3 times, blocked with 3% BSA in PBST for 1 hour at room

temperature. Then serial dilutions of standards and test samples in HISPEC assay

diluent (Bio-Rad) were loaded and allowed to react for 2 hours at room temperature.

Plates were then washed 3 times with PBST and treated with 2 µg/mL of HRP

conjugated detection antibody (AbD28582) in HISPEC assay diluent for 1 hour at room

temperature. After washing 6 times with PBST, the plates were developed with

QuantaBlu fluorogenic peroxidase substrate kit (Thermo Scientific) for 30 min at room

temperature. Fluorescence was recorded (ex. 320 ± 25nm, em. 430 ± 35nm).

Subjects and samples. Retrospective analysis was performed on a total of 45

plasma samples from International Maternal Pediatric Adolescent AIDS Clinical Trails

(IMPAACT) P1041 [24, 25]. P1041 was a Phase II/ III, randomized, double-blind,

placebo-controlled clinical trial to evaluate the efficacy of isoniazid prophylaxis on TB

disease and latent *Mtb* infection free survival in HIV-infected and HIV-exposed, but

uninfected infants up to 192 weeks of follow up [25, 26]. Samples include 24 with noTB

and 21 with TBDIS, all HIV infected. Patients without *Mtb* infection were categorized as

noTB; patients with a positive tuberculin skin test but lacking any clinical, radiographic

or laboratory evidence of disease caused by *Mtb* were categorized as latent TB

infection (LTBI); and patients presenting clinical, radiographic or laboratory evidence of

disease caused by *Mtb* were categorized as TBDIS. Plasma samples were collected,

frozen and thawed before use.

Statistical Analysis. Categorical variables were compared using t-student test,

whenever appropriate. Non-parametric tests (Mann-Whitney) were used for non-

normally distributed variables. Assay accuracy, including 95% confidence intervals, was

assessed using sensitivity, specificity, predictive values and area under the ROC in the

TB and non-TB groups. Statistical calculations were performed with GraphPad

Prism® Software.

RESULTS

Mtb modified lipoprotein (TLP) is different from MDA-LDL. We purified and

treated native LDL with *Mycobacterium tuberculosis* in vitro to obtain *Mt*b modified

lipoprotein (TLP). Decreased mobility of TLP in agarose electrophoresis indicates that

TLP is more positively charged than native LDL (Figure 1A). In contrast, MDA-LDL, one

of the end products of lipid peroxidation, is more negatively charged than native LDL

[20]. The increased size of TLP compared to native LDL was also confirmed by dynamic

light scattering analysis (Table S1). Furthermore, TLP is recognized by anti-

apolipoprotein B (apoB) antibodies but not anti-MDA-LDL antibodies, indicating TLP

does not contain a malondialdehyde-modified apoB derivative (Figure 1B). We

demonstrate that LDL is altered in the presence of *Mtb*, and the resulting TLP is distinct

from the typical atherosclerotic species malondialdehyde-modified LDL (MDA-LDL)

present in human blood [27].

A   B

Figure 1. TLP is different from MDA-LDL. (A) an agarose gel electrophoresis showing

TLP is larger in size and more positively charged than native LDL. (B) Immunoblots

showing TLP is recognized by anti- apolipoprotein B (apoB) antibodies (top) but not

anti- MDA-LDL antibodies (bottom). Images of the entire gel and blots are in Figure S1.

Mtb modified LDL induces lipid body accumulation in macrophages. The

induction of foamy macrophages have been reported as hallmarks in many pathologies

associated with chronic proinflammatory stimuli including atherosclerosis [15]. It has

also been reported that *Mtb* lipids induce the formation of giant multinuclear

macrophages [28]. We therefore used foamy macrophage formation as a tool to assess

the bioactive components that trigger a tissue response similar to granuloma formation

in TB. Lipid bodies were stained with Oil red O. The buffer control showed no

accumulation of lipid bodies and the cells remained rounded (Figure 2A). LDL treated

macrophage showed some lipid accumulation, with a commensurate increase in cell

size (Figure 2B). In contrast, MDA-LDL, commonly associated with atherosclerosis,

treated macrophages exhibited more extensive accumulation of lipid bodies than the

LDL treated macrophages. These macrophage phenotypes are consistent with previous

reports of in vitro foamy macrophage formation [20, 29]. Most importantly, TLP treated

macrophages had extensive lipid body accumulation per cell compared to macrophages

treated with other LDLs (Figure 2 and S2B). Moreover, the TLP-treated macrophage

morphology was distinct from other samples. The majority of the TLP-treated

macrophages became multinucleated (Figure 2D and S2A) and developed long

filopodia consistent with cell migration or exocytosis. Thus, we hypothesize that the

modified LDL is detectable in vivo to serve as a biomarker for TB disease, and an

antibody specific for recognition of TLP is an *Mtb*-specific indicator of infection and

progression to (active) TB disease.

Figure 2. TLP induces lipid body accumulation in macrophages. Microscopic images of Oil Red O-hematoxylin stained THP-1 macrophage culture treated with PBS control (A), LDL (B), MDA-LDL (C) and TLP (D) at 200 µg/mL for 2 days.

196 ***Anti-TLP monoclonal antibodies were generated and a TLP Sandwich ELISA assay was established.*** We generated 13 anti-TLP monoclonal antibodies (Fad-A-FH, bivalent Fab-bacterial alkaline phosphatase fusion antibody followed by FLAG® and His6-tag) from the Human Combinatorial Antibody Library (HuCAL, Bio-Rad AbD Serotec GmbH) using in vitro selection and counter selection [22, 23]. HuCAL is a phage display library containing highly specific, fully human monoclonal antibodies, with DNA sequence encoded for each antibody fragment. The HuCAL phage display library was depleted of antibodies that recognize LDL (Figure S3). The depleted library was panned for three rounds of binding, elution, and amplification to isolate antibodies specific for TLP. Among the antibody hits selected, 13 antibodies were identified as unique by DNA sequencing. These antibodies were produced heterologously for further development. The cross-reactivity with native LDL, human serum albumin, bovine serum albumin, and His$_6$ affinity tag was determined in an indirect ELISA format. Eight antibodies showed a specific signal for TLP at least 4-fold above native LDL background (Figure S3). We tested pairwise

combinations of antibodies to identify antibody pairs that have non-overlapping epitopes

suitable for use in a sandwich ELISA (Figure S4A). 5 sandwich pairs were identified with

good signal over background and the assay conditions were optimized for the best pair,

AbD28582 and AbD28580 (amino acid sequence shown in Table S2). We established

the linear response of the sandwich ELISA (Figure S4B) to antigen spiked into pediatric

plasma. TLP is specifically detected at >=4ppm in the presence of plasma LDL, which is

around 10^6 ng/mL [30].

Levels of TLP are increased in patients with active TB compared to no TB

controls. We assayed stored plasma samples from the HIV IMPAACT P1041 trial for

the presence of TLP. The median plasma levels of TLP was 494 ng/mL in control

subjects, and 2.7 fold higher in active TB subjects (Figure 3A). The difference between

control subjects and active TB subjects is significant (p<0.001). The receiver operating

characteristic (ROC) analysis describes the relationship between the sensitivity and

specificity at any cut-off values. An ROC curve of TLP concentration for active TB

versus control subjects is presented in Figure 3B. The area under the ROC curve (AUC)

is 0.86 (95% CI=0.75-0.97, p<0.0001). An optimal cutoff of 1064 ng/mL rendered 71%

sensitivity and 88% specificity. Alternatively, a cutoff of 1230 ng/mL provided 57%

sensitivity and of 96% specificity.

A   B

Figure 3. Retrospective analysis of P1041 HIV positive plasma samples. (A)

Plasma levels of TLP in HIV positive subjects clinically diagnosed probably having

active stage tuberculosis disease (TB) and not having TB (noTB). Individual patients are

depicted as dots with group mean. The dashed line represents the detection limit of the

assay. *** $p < 0.001$. (B) ROC curve of TLP concentration for active TB versus control

subjects of HIV positive patients. AUC, area under curve. The dashed line represents

the no-discrimination line from the left bottom to the top right corners.

We also monitored apoB levels, i.e., LDL levels, of all the samples using Human

apoB ELISA kits (ab190806, Abcam) and found there was no linear correlation between

TLP and apoB levels (Figure S6A), suggesting that plasma LDL fluctuation does not

interfere with TLP detection. There is a weak correlation between TLP levels and time

before or after diagnosis (Figure S6B). The samples collected before diagnosis may

represent disease progression from no infection or infection but without clinical signs of

disease to outward manifestation of disease. Samples collected after diagnosis may

represent a mixed outcome of disease progression and anti- tubercular treatment. We

do not have additional sample sets to demonstrate if anti-tubercular treatment affects

TLP levels. However, the discrimination of TLP levels between control subjects and

active TB subjects (post diagnosis with treatment less than 40 weeks) was improved

with the exclusion of samples collected after 40 weeks of treatment (p<0.0001), which

may represent patients clear of TB (Figure S6). ROC (Receiver operating characteristic)

analysis provided an AUC (area under the ROC curve) of 0.88 (95% CI=0.76-0.99,

p<0.0001). An optimal cutoff of 632 ng/mL gave 67% sensitivity and 93% specificity.

The results provide further evidence that TLP levels are associated with active TB

disease. Overall, the results clearly highlight the potential of TLP as a biomarker in the

diagnosis of active TB in HIV positive children.

DISCUSSION AND CONCLUSIONS

The wide spectrum of disease observed in children, and the non-specific signs and

symptoms especially in young and HIV+ children, contribute to diagnostic delay and

missed opportunities to detect TB, which in turn, favors TB disease progression and poor

treatment outcomes. There is urgent need to search for a "Rapid Non-sputum-based

Biomarker Test for Tuberculosis Detection" [12] .Most of the biomarkers currently studied

are host-derived biomarkers including metabolites and cytokines,[31, 32] the levels of which

can be affected by many confounding factors. On the other hand, *Mtb* products can be

detected directly in blood, sputum or urine, and are increasingly being used for

diagnosis.[33] We have characterized a unique biomolecule, *Mtb* modified lipoprotein,

which results from a combination of host and *Mtb* pathogen activity. Functionally, TLP

increases the flux of lipid into activated macrophages, thereby stimulating their conversion

into foamy multi-nucleated macrophages. The TLP-stimulated macrophages accumulate

lipid bodies but have a distinct phenotype from MDA-LDL-stimulated macrophages

(Figure 2). We reason that detection of this biomolecule in patient plasma may withstand

the heterogeneity associated with variations derived from host and be pathogen or

disease specific. Thus, TLP detection may be used as TB diagnosis in young children.

The lipoprotein modification can take place in macrophages in contact with *M.*

tuberculosis. There are also reports demonstrating that extracellular vesicles carrying

bacterial components are released from infected macrophages and circulated beyond

the site of infection [34], where host lipoproteins could be modified via contact with the

extracellular vesicles. In addition, the intracellular bacilli are able to avoid killing by

escaping phagosome-lysosome fusion. In certain individuals, especially children below

5 years of age and immunosuppressed subjects, the extracellular bacilli may disperse to

distant metastatic sites via lymphatics and the bloodstream [35], where host lipoproteins

may be modified by direct contact with *M. tuberculosis*. Our data demonstrate that the

modified lipoprotein in the present study is distinct from other commonly recognized

disease associated LDLs, which are typically negatively charged and smaller in particle

size compared to native LDL [36].

The present study established a rapid, non-invasive, and robust sandwich ELISA

assay for the detection of a novel biomolecule TLP in plasma. We demonstrate that

plasma levels of TLP are significantly distinguishable in HIV positive subjects with active

TB disease compared to no TB controls (Figure 3). The significant elevation of plasma

levels of TLP in HIV positive subjects with active TB disease suggests the detection of

TLP associate with TB active disease in HIV positive subjects. The AUC value of 0.86

evidenced the diagnostic value of TLP for TB detection in child. The target product profiles

(TPP) adopted by the World Health Organization (WHO) in 2015 for a "Rapid Non-

sputum-based Biomarker Test for Tuberculosis Detection" is a diagnostic sensitivity ≥ 66%

for microbiologically confirmed pediatric TB and 98% specificity for a pediatric test [12].

The TPP proposed sensitivity is similar to the sensitivity of the Xpert MTB/RIF assay in

microbiologically confirmed samples. Our assay renders 57% sensitivity and 96%

specificity, which has the potential to meet the WHO TPP for detection of active TB in

children, given the highly heterogenous population of P1041 samples, with respect to the

bacteriological and clinical status.

Samples used in our study were obtained from IMPAACT P1041 clinical trial. The

accuracy of TB diagnosis for the P1041 trial samples is not known due to the absence of

good pediatric TB diagnostics. In addition, microbiological confirmation of TB disease was

not obtained for any of these patients in the P1041 trial. TB disease subjects used in the

present study are patients infected with TB presenting clinical symptoms of disease. The

ELISA assays were performed retrospectively and the variance in storage conditions

might affect the antigen integrity and therefore the accuracy of the assay. We have

insufficient samples to include other clinical characteristics of the patients into data

analysis. The clinical characteristics include but not limited to BCG administration,

immunosuppression status like CD4 counts, or clinical manifestation of tuberculosis

(pulmonary and extra-pulmonary TB). The consideration of those characteristics may help

define the limitation of our TLP assay or render improved accuracy with the combination

of other clinical factors.

We focused on HIV positive subjects in the current study. TB diagnosis is particularly

difficult among HIV co-infected individuals who may have atypical, nonspecific clinical

presentation, high rates of smear negative disease and high rates of extrapulmonary TB

[37]. P1041 samples were obtained from TB endemic settings, therefore patients were

typically being diagnosed at more advanced stages than those living in the US. Therefore,

the heterogenous stages of TB disease may contribute to the wide dispersity of TLP levels

in TB disease subjects (Figure 3A), which is common among TB diagnostic assays. In

addition, the duration of anti-tubercular treatment was not consistent among patients in

the present study. Change of LDL modification is a dynamic process and may have the

potential to monitor treatment outcome. Although not statistically significant, the slight

decrease in the TLP level post diagnosis/ anti-tubercular treatment of 40 weeks may be

related to a reduction in microbial load (Figures S6A and S6B). Further studies are

required to elucidate the effect of different disease status and/or treatment status on TLP

levels in plasma, and to extend our preliminary results to other populations such as HIV

negative subjects and adults.

It is a growing notion in the field that a single biomarker will not be sufficient for

distinguishing TB status in different patient groups, and multiple biomarkers may be

used to increase sensitivity and specificity. Further gains in clinical sensitivity and

specificity may be obtained by combining the current assay with other biomarkers or

diagnostic tests.

LIST OF ABBREVIATIONS

TB, tuberculosis; *Mtb, Mycobacterium tuberculosis*; ELISA, enzyme-linked

immunosorbent assay; FM, foamy macrophages; MDA-LDL, malondialdehyde-modified

low density lipoprotein; TLP, *Mycobacterium tuberculosis* modified lipoprotein; ROC,

receiver operating characteristic; AUC, area under the ROC curve.

DECLARATIONS

Ethics approval and consent to participate

The NIH-IMPAACT Scientific Leadership Group (SLG) reviewed and approved the

use of P1041 trial samples under NWCS127. The Wits Health Consortium,

University of Witwatersrand, South Africa provided deidentified samples to Stony

Brook University for which informed consent had been obtained from a parent

and/or legal guardian. The Stony Brook University CORIHS determined the

experiments in this manuscript were not human subjects research under either

the Common Rule or FDA Regulations.

Consent for publication

All authors have approved the manuscript for submission.

Availability of data and materials

The datasets used and/or analysed during the current study are available from

the corresponding author on reasonable request.

Competing interests

NSS is President and co-owner of Chronus Pharmaceuticals, Inc. XY was an

employee of both Chronus Pharmaceuticals, Inc and Stony Brook University. All

intellectual property is currently owned by Stony Brook University. SN and MFM

have no conflict of interest.

Funding

Research reported in this publication was supported by the National Heart, Lung,

and Blood Institute of the National Institutes of Health under Award Number

U01HL127522, and National Institute of Allergy and Infectious Diseases (NIAID)

under award number R41AI136071 to Chronus Pharmaceuticals, Inc with a

subcontract to Stony Brook University (NSS) and the National Center for

Advancing Translational Sciences under award number TL1TR002386 (MFW).

Overall support for the International Maternal Pediatric Adolescent AIDS Clinical

Trials (IMPAACT) Network was provided by the National Institute of Allergy and

Infectious Diseases (NIAID) of the National Institutes of Health (NIH) under Award

Numbers UM1AI068632 (IMPAACT LOC), UM1AI068616 (IMPAACT SDMC) and

UM1AI106716 (IMPAACT LC), with co-funding from the Eunice Kennedy Shriver

National Institute of Child Health and Human Development (NICHD) and the

National Institute of Mental Health (NIMH). The content is solely the

responsibility of the authors and does not necessarily represent the official views

of the NIH.

Authors' contributions

SN, MFW, XY, NSS conceived the experiments

SN, MFW, obtained sample resource

XY performed the experiments

XY, NSS wrote the main manuscript text

XY prepared all figures

All authors reviewed the manuscript

Acknowledgements

We thank Guannan Chen for help with initial LDL purification experiments.

REFERENCES

1.   WHO. 2020. Global tuberculosis report

https://apps.who.int/iris/bitstream/handle/10665/336069/9789240013131-eng.pdf

Accessed 16 March 2021.

2.   Cotton MF, Schaaf HS, Hesseling AC, Madhi SA: **HIV and childhood tuberculosis: the way**

forward. *The international journal of tuberculosis and lung disease : the official journal of the*

International Union against Tuberculosis and Lung Disease 2004, **8**(5):675-682.

3.   Marais BJ, Gie RP, Obihara CC, Hesseling AC, Schaaf HS, Beyers N: **Well defined symptoms**

are of value in the diagnosis of childhood pulmonary tuberculosis. *Archives of disease in*

childhood 2005, **90**(11):1162-1165.

4.   Moore DP, Klugman KP, Madhi SA: **Role of Streptococcus pneumoniae in hospitalization**

for acute community-acquired pneumonia associated with culture-confirmed

Mycobacterium tuberculosis **in children: a pneumococcal conjugate vaccine probe**

study. *The Pediatric infectious disease journal* 2010, **29**(12):1099-1004.

5.   Zar HJ, Tannenbaum E, Apolles P, Roux P, Hanslo D, Hussey G: **Sputum induction for the**

diagnosis of pulmonary tuberculosis in infants and young children in an urban setting

in South Africa. *Archives of disease in childhood* 2000, **82**(4):305-308.

6.   Walters E, Cotton MF, Rabie H, Schaaf HS, Walters LO, Marais BJ: **Clinical presentation and**

outcome of tuberculosis in human immunodeficiency virus infected children on anti-

retroviral therapy. *BMC pediatrics* 2008, **8**:1.

7.   Jeena PM, Mitha T, Bamber S, Wesley A, Coutsoudis A, Coovadia HM: **Effects of the human**

immunodeficiency virus on tuberculosis in children. *Tubercle and lung disease : the*

official journal of the International Union against Tuberculosis and Lung Disease 1996,

77(5):437-443.

8.   Schaaf HS, Gie RP, Beyers N, Smuts N, Donald PR: **Tuberculosis in infants less than 3**

months of age. *Archives of disease in childhood* 1993, **69**(3):371-374.

9.   Jeena PM, Pillay P, Pillay T, Coovadia HM: **Impact of HIV-1 co-infection on presentation**

and hospital-related mortality in children with culture proven pulmonary

tuberculosis in Durban, South Africa. *The international journal of tuberculosis and lung*

disease : the official journal of the International Union against Tuberculosis and Lung Disease

2002, **6**(8):672-678.

10.   Madhi SA, Huebner RE, Doedens L, Aduc T, Wesley D, Cooper PA: **HIV-1 co-infection in**

children hospitalised with tuberculosis in South Africa. *The international journal of*

tuberculosis and lung disease : the official journal of the International Union against

Tuberculosis and Lung Disease 2000, **4**(5):448-454.

11.   Marais BJ, Gie RP, Schaaf HS, Hesseling AC, Obihara CC, Starke JJ, Enarson DA, Donald PR,

Beyers N: **The natural history of childhood intra-thoracic tuberculosis: a critical review**

of literature from the pre-chemotherapy era. *The international journal of tuberculosis and*

lung disease : the official journal of the International Union against Tuberculosis and Lung

Disease 2004, **8**(4):392-402.

12. Denkinger CM, Kik SV, Cirillo DM, Casenghi M, Shinnick T, Weyer K, Gilpin C, Boehme CC,

Schito M, Kimerling M *et al*: **Defining the needs for next generation assays for**

tuberculosis. *J Infect Dis* 2015, **211 Suppl 2**:S29-38.

13. Abdalla AE, Ejaz H, Mahjoob MO, Alameen AAM, Abosalif KOA, Elamir MYM, Mousa MA:

Intelligent Mechanisms of Macrophage Apoptosis Subversion by Mycobacterium.

Pathogens 2020, **9**(3).

14. Genoula M, Marin Franco JL, Dupont M, Kviatcovsky D, Milillo A, Schierloh P, Morana EJ, Poggi

S, Palmero D, Mata-Espinosa D *et al*: **Formation of Foamy Macrophages by Tuberculous**

Pleural Effusions Is Triggered by the Interleukin-10/Signal Transducer and Activator

of Transcription 3 Axis through ACAT Upregulation. *Front Immunol* 2018, **9**:459.

15. Russell DG, Cardona PJ, Kim MJ, Allain S, Altare F: **Foamy macrophages and the progression**

of the human tuberculosis granuloma. *Nat Immunol* 2009, **10**(9):943-948.

16. Viljoen A, Blaise M, de Chastellier C, Kremer L: **MAB_3551c encodes the primary**

triacylglycerol synthase involved in lipid accumulation in Mycobacterium abscessus.

Mol Microbiol 2016, **102**(4):611-627.

17. Caire-Brandli I, Papadopoulos A, Malaga W, Marais D, Canaan S, Thilo L, de Chastellier C:

Reversible lipid accumulation and associated division arrest of Mycobacterium avium

in lipoprotein-induced foamy macrophages may resemble key events during latency

and reactivation of tuberculosis. *Infect Immun* 2014, **82**(2):476-490.

18. Guerrini V, Gennaro ML: **Foam Cells: One Size Doesn't Fit All**. *Trends Immunol* 2019,

40(12):1163-1179.

19.   Agarwal P, Combes TW, Shojaee-Moradie F, Fielding B, Gordon S, Mizrahi V, Martinez FO:

449   **Foam Cells Control Mycobacterium tuberculosis Infection**. *Front Microbiol* 2020, **11**:1394.

450   20.   Watanabe K, Nakazato Y, Saiki R, Igarashi K, Kitada M, Ishii I: **Acrolein-conjugated low-**

451   **density lipoprotein induces macrophage foam cell formation**. *Atherosclerosis* 2013,

452   **227**(1):51-57.

453   21.   Holvoet P, Vanhaecke J, Janssens S, Van de Werf F, Collen D: **Oxidized LDL and**

454   **malondialdehyde-modified LDL in patients with acute coronary syndromes and stable**

455   **coronary artery disease.** *Circulation* 1998, **98**(15):1487-1494.

456   22.   Knappik A, Ge L, Honegger A, Pack P, Fischer M, Wellnhofer G, Hoess A, Wolle J, Pluckthun A,

457   Virnekas B: **Fully synthetic human combinatorial antibody libraries (HuCAL) based on**

458   **modular consensus frameworks and CDRs randomized with trinucleotides**. *J Mol Biol*

459   2000, **296**(1):57-86.

460   23.   Prassler J, Thiel S, Pracht C, Polzer A, Peters S, Bauer M, Norenberg S, Stark Y, Kolln J, Popp A

461   *et al*: **HuCAL PLATINUM, a synthetic Fab library optimized for sequence diversity and**

462   **superior performance in mammalian expression systems**. *J Mol Biol* 2011, **413**(1):261-

463   278.

464   24.   Madhi SA, Nachman S, Violari A, Kim S, Cotton MF, Bobat R, Jean-Philippe P, McSherry G,

465   Mitchell C, Team PS: **Primary isoniazid prophylaxis against tuberculosis in HIV-exposed**

466   **children**. *The New England journal of medicine* 2011, **365**(1):21-31.

467   25.   **A randomized, double blind, placebo controlled trial to determine the efficacy of**

468   **isoniazid (INH) in preventing tuberculosis disease and latent tuberculosis infection**

469   **among infants with perinatal exposure to HIV.**

470   [https://impaactnetwork.org/DocFiles/P1041/P1041V2_11Jul07.pdf]

471   26.   Gupta A, Montepiedra G, Gupte A, Zeldow B, Jubulis J, Detrick B, Violari A, Madhi S, Bobat R,

472   Cotton M *et al*: **Low Vitamin-D Levels Combined with PKP3-SIGIRR-TMEM16J Host**

Variants Is Associated with Tuberculosis and Death in HIV-Infected and -Exposed
Infants. *PloS one* 2016, **11**(2):e0148649.

27.   Yoshida M, Higashi K, Kobayashi E, Saeki N, Wakui K, Kusaka T, Takizawa H, Kashiwado K,
Suzuki N, Fukuda K *et al*: **Correlation between images of silent brain infarction, carotid**
atherosclerosis and white matter hyperintensity, and plasma levels of acrolein, IL-6
and CRP. *Atherosclerosis* 2010, **211**(2):475-479.

28.   Puissegur MP, Lay G, Gilleron M, Botella L, Nigou J, Marrakchi H, Mari B, Duteyrat JL, Guerardel
Y, Kremer L *et al*: **Mycobacterial lipomannan induces granuloma macrophage fusion via**
a TLR2-dependent, ADAM9- and beta1 integrin-mediated pathway. *J Immunol* 2007,
178(5):3161-3169.

29.   Nezami N, Ghorbanihaghjo A, Rashtchizadeh N, Argani H, Tafrishinejad A, Ghorashi S,
Hajhosseini B: **Atherogenic changes of low-density lipoprotein susceptibility to**
oxidation, and antioxidant enzymes in pulmonary tuberculosis. *Atherosclerosis* 2011,
217(1):268-273.

30.   Sacks FM, Moye LA, Davis BR, Cole TG, Rouleau JL, Nash DT, Pfeffer MA, Braunwald E:
Relationship between plasma LDL concentrations during treatment with pravastatin
and recurrent coronary events in the cholesterol and recurrent events trial. *Circulation*
1998, **97**(15):1446-1452.

31.   Weiner J, Kaufmann SH: **Recent advances towards tuberculosis control: vaccines and**
biomarkers. *Journal of internal medicine* 2014.

32.   Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A: **Immunological biomarkers of**
tuberculosis. *Nature reviews Immunology* 2011, **11**(5):343-354.

33.   Goletti D, Lee MR, Wang JY, Walter N, Ottenhoff THM: **Update on tuberculosis biomarkers:**
from correlates of risk, to correlates of active disease and of cure from disease.
Respirology 2018, **23**(5):455-466.

34.   Athman JJ, Wang Y, McDonald DJ, Boom WH, Harding CV, Wearsch PA: **Bacterial membrane**
499   **vesicles mediate the release of Mycobacterium tuberculosis lipoglycans and**
500   **lipoproteins from infected macrophages**. *J Immunol* 2015, **195**(3):1044-1053.

501   35.   Grosset J: **Mycobacterium tuberculosis in the extracellular compartment: an**
502   **underestimated adversary**. *Antimicrob Agents Chemother* 2003, **47**(3):833-836.

503   36.   Orekhov AN: **LDL and foam cell formation as the basis of atherogenesis**. *Curr Opin Lipidol*
504   2018, **29**(4):279-284.

505   37.   Scott L, da Silva P, Boehme CC, Stevens W, Gilpin CM: **Diagnosis of opportunistic infections:**
506   **HIV co-infections - tuberculosis**. *Curr Opin HIV AIDS* 2017, **12**(2):129-138.