Long Title: Long non-coding RNAs (lncRNAs) NEAT1 and MALAT1 are differentially expressed in severe

COVID-19 patients: An integrated single cell analysis

Short Title: LncRNAs NEAT1 and MALAT1 differentiate inflammation in severe COVID-19 patients

Authors: Kai Huang BS[1,2], Catherine Wang MS[3], Christen Vagts MD[1], Vanitha Raguveer BS[3], Patricia W.

Finn MD[1,2,4*], David L. Perkins MD PhD[2,5,6]

[1]Division of Pulmonary, Critical Care, Sleep, and Allergy, Department of Medicine, University of Illinois at

Chicago, Chicago IL, 60612 United States

[2]Department of Bioengineering, University of Illinois at Chicago, Chicago IL, 60612 United States

[3]College of Medicine, University of Illinois at Chicago, Chicago IL, 60612 United States

[4] Department of Microbiology and Immunology, University of Illinois at Chicago, Chicago IL, 60612

United States

[5]Division of Nephrology, Department of Medicine, University of Illinois at Chicago, Chicago IL, 60612

United States

[6]Department of Surgery, University of Illinois at Chicago, Chicago IL, 60612 United States

*Corresponding author: pwfinn@uic.edu

Author Contributions:
Conceptualization, K.H.; Methodology, K.H.; Software, K.H.; Investigation, K.H, C.W., and V.R.; Writing –
Original Draft, K.H., C.W.; Writing Review & Editing, K.H., C.W., V.R., C.V., D.L.P., and P.W.F;
Visualization, K.H.; Supervision, C.V., D.L.P., and P.W.F.; Funding Acquisition, K.H., D.L.P., and P.W.F.
24

Abstract:

Hyperactive and damaging inflammation is a hallmark of severe rather than mild COVID-19 syndrome.

To uncover key inflammatory differentiators between severe and mild COVID-19 disease, we applied an

unbiased single-cell transcriptomic analysis. We integrated a bronchoalveolar lavage (BAL) dataset with

a peripheral blood mononuclear cell dataset (PBMC) and analyzed the combined cell population,

focusing on genes associated with disease severity. Distinct cell populations were detected in both BAL

and PBMC where the immunomodulatory long non-coding RNAs (lncRNAs) NEAT1 and MALAT1 were

highly differentially expressed between mild and severe patients. The detection of other severity

associated genes involved in cellular stress response and apoptosis regulation suggests that the pro-

inflammatory functions of these lncRNAs may foster cell stress and damage. The lncRNAs NEAT1 and

MALAT1 are potential components of immune dysregulation in COVID-19 that may provide targets for

severity related diagnostic measures or therapy.

Introduction:

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) pandemic continues around the world,(1,2) but the underlying pathophysiology of coronavirus disease 2019 (COVID-19) is ill-defined. Symptoms and progression of COVID-19 vary widely(3) as some patients may be asymptomatic while others exhibit disease with varying severity.(4,5) Common symptoms include fever, cough, and fatigue, which generally appear 2 to 14 days after exposure,(6,7) while rarer symptoms include dyspnea, headache/dizziness, nausea, diarrhea, and hemoptysis.(8) Severe cases of COVID-19 are distinguished by strong inflammatory responses that can lead to multiorgan damage and death.(9) The mechanisms that separate mild and severe disease remain poorly understood.

After viral exposure, the inflammatory response to COVID-19 commences with signaling cascades that lead to secretion of type I interferons, cytokines and chemokines.(10) This initial exposure also activates inflammasomes, multimeric protein complexes that play an important role in triggering inflammation and the subsequent initiation of an adaptive immune response.(11–13) The Nod-like receptor pyrin domain-containing 3 (NLRP3) inflammasome is a major cause of cytokine storm associated the with clinical manifestations of severe COVID-19 disease.(14) Furthermore, coronavirus viroporin proteins activate the NLRP3 inflammasome which regulates the secretion of IL-1β and IL-18.(15) Pyroptosis, a programmed cell death pathway that leads to immune cell depletion, is also regulated by activation of the NLRP3 inflammasome and is an important mechanism of viral pathogenesis in both SARS-CoV-2 and SARS-CoV.(16–18) These studies suggest that investigation of inflammasome regulation may elucidate understanding of COVID-19 disease pathophysiology.

Single-cell studies of COVID-19 patients have found dysregulated immune compartments in the respiratory tract as well as peripheral blood.(19–24) However, it is challenging to directly compare results across studies in different tissues due to differences in cell cluster identification between physiological compartments. We postulated that simultaneous analysis of severe versus mild COVID-19

patients across respiratory and peripheral immune compartments using integrated clustering would

uncover overall effectors of immune dysregulation in the COVID-19 immune response. To achieve this

goal, we integrated single-cell datasets, one from bronchoalveolar lavage (BAL) and one from peripheral

blood mononuclear cells (PBMC),(19,20) in order to examine disease transcriptomics across severities as

well as between local and peripheral cellular environments. We utilized an unbiased analytical strategy

that was agnostic to specific gene functions and focused on genes with severity dependent expression

across different cell types. Taken together, we uncovered genes contributing to the dysregulated COVID-

19 immune response prominent in severe relative to mild disease. Moreover, we identified cell types

where these inflammatory regulators manifest in local and peripheral compartments.

Methods:

Dataset preprocessing and integration

78   We selected publicly available single-cell datasets with patient severity metrics and ample

sequencing depth to pass our quality filters for integration into our combined dataset. The raw count

matrices for BAL cells and PBMC cells were downloaded from the NCBI Gene Expression Omnibus

(accession number GSE145926) and the COVID-19 Cell Atlas (https:/www.covid19cellatlas.org/#wilk20),

respectively. Patients who were mechanically ventilated or had $PaO_2/FiO_2 \leq 300$ mmHg indicating

hypoxemia consistent with acute respiratory distress syndrome (ARDS)(25) were designated as severe

patients while all others were considered to have mild disease (**Table S1**). The BAL dataset contained

three healthy controls, while the PBMC dataset contained six (**Table S2**). Both datasets were

preprocessed using the R program Seurat.(26) Briefly, cells were filtered to only include cells with

unique molecular identifier (UMI) counts greater than 1000, gene count between 200 and 6000, and less

than 10% of genes mapping to mitochondrial genes. The function SCTransform from the Seurat package

was applied to each dataset separately to regress out technical variability as well as the percentage of

mitochondrial gene expression.(27) Transformed BAL and PBMC datasets were integrated with 3000

integration features and 50 integration anchors as recommended in Seurat.(28) We found that the "M3"

mild patient sample from the BAL dataset contained only 369 total cells, while every other patient

sample for BAL or PBMC had at least 1200 cells. The M3 sample was removed before differential

expression analysis to avoid skewing results due to extremely low cell counts.

Clustering and Identification

97   The integrated dataset was dimension reduced using principal component analysis (PCA) and

clustered with a resolution set to 0.5 and including the top 30 principle components. The clustering was

visualized using uniform manifold approximation and projection (UMAP).(29) The raw integrated dataset

was normalized by applying SCTransform to the full integrated dataset. This normalized count matrix is

utilized for all subsequent analysis. Marker genes for each cluster were computed using the

FindAllMarkers function with the Model-based Analysis of Single-cell Transcriptomics algorithm (MAST)

for differential expression with UMI count as a latent variable.(30,31) Cluster markers were then

inspected and labeled according to known cell markers(Supplemental file 1).(32,33) A second round of

clustering with a resolution of 1 was then conducted to further classify subtypes of identified cells.

Clusters with fewer than 300 cells were reassigned to larger clusters using Seurat integration label

transfer. Cluster identities were scored and verified using a signature matrix generated from flow

cytometry sorted RNA-seq data of immune cells.(34) Plots of cell clusters and key cell type markers were

generated using Seurat's plotting functions.

Cell Proportions

Cell types were tallied for each sample, and the percentage abundance of each cell type was

calculated. Cell proportions for healthy controls, mild patients, and severe patients were compared

using a two-sided pairwise test of equal proportion with false discovery rate (FDR) p-value adjustment.

The resulting proportions were plotted using the ggplot2 R package.(35)

Differential Expression

For each cell type, differentially expressed genes (DEGs) were calculated separately for BAL and

PBMC cells using MAST with UMI count as a latent variable. To support MAST differential gene

expression analysis between three sample groups, the Seurat built-in differential expression function

FindMarkers was modified to generating iterations of the hurdle model corresponding to each set of

two compared conditions. Genes from each cell type that were differentially expressed across all three

comparisons between healthy controls, mild, and severe patients were tallied. For BAL, only genes with

FDR adjusted p-values less than 1e-7 across all comparisons were considered (17.4% of all DEGs in BAL),

while PBMC DEGs were considered if all FDR adjusted p-values across comparisons for that gene were

less than 0.05 (16.4% of all DEGs in PBMCs). The difference in p-value threshold was used to filter a

similar proportion of genes from BAL and PBMC data due to the smaller number of DEGs detected in

PBMCs (200-400 in PBMCs, 1000+ in BAL). These highly differentially expressed genes were further

filtered by removing genes that were not differentially expressed across all conditions in at least 5

different cell types (1/3 of our total cell types). The lists of PBMC and BAL highly differentially expressed

genes were then combined, removing duplicates.

Some of the genes identified were highly cell type specific. These genes also had the highest

residual variance. Since these genes represent intrinsic cell type differences rather than biologically

interesting differentially expressed genes, they were removed. To determine this filter threshold, the

residual variance of the top 100 variable genes were plotted in decreasing order to determine the

"elbow" point where the variance stops decreasing at a rapid rate.(**Figure S1**) This resulted in the 21

most variable genes being removed. We termed the 50 remaining differentially expressed genes

recurrent differentially expressed genes (rDEGs) since they were found in multiple cell types and showed

differential expression between patients and healthy controls as well as between severities. The rDEG

expression data was exported to Monocle 3 to generate modules for gene ontology (GO) enrichment

analysis. We generated 4 modules from the 50 rDEGs using the find_gene_modules function in Monocle

3 with 30 principle components and a resolution of 0.8.(36–38) Differential module expression was

calculated using ANOVA using aggregated module expression levels and processed with a tukey posthoc

test. Module gene ontology enrichment was computed topGO with default settings.(39) Module and

gene level plots were generated using the R packages ggplot2, ComplexHeatmap, and Circlize.(35,40,41)

Validation

We compared rDEG trends from our analysis with two additional COVID-19 datasets, one with

nasopharyngeal data to compare against the local inflammatory environment of our BAL data, and one

in PBMCs to compare against the peripheral environment in our PBMC data.(22,23) Cell types from our

analysis were transferred onto the validation datasets using Seurat to identify the validation cells for

comparison. Each validation dataset was filtered and preprocessed separately using the same

parameters as the main dataset. After preprocessing and label transfer, DEGs were generated

independently for each validation set for our transferred cell types. We compared the rDEGs we focused

on in the manuscript with DEG results from each validation set, noting whether the same DEG was

detected and whether the direction of change was the same.

Results:

Integrated PBMC and BAL analysis identified 26 clusters consolidated into 15 cell types.

After quality filtering, we recovered 100,739 single cell transcriptomes. From these, we

recovered 26 cell clusters from Seurat. (**Figure S2**). Since we identified cell types from the integrated

dataset containing both PBMC and BAL cells, we were able to examine how each of our cell clusters

behaves across both physiological compartments. The clusters did not aggregate based on sample type

or patient condition (**Figure S2**), indicating successful integration clustering of the two datasets.

From the 26 clusters, 11 were identified as monocyte/macrophage (Mo/Ma) clusters. Since our

clusters contain both monocytes from the PBMC sample as well as macrophages from the BAL, we

designated them as MoMa clusters. Six of the MoMa clusters showed classically M1 associated

transcriptomes with increased expression of VCAN, FCN1 and CD14 expression.(42,43) These clusters

also expressed other pro-inflammatory factors such as S100A8, CCL2, CCL3, CCL7, and CCL8.(44,45)

Three other MoMa clusters showed M2 polarization with increased FN1 expression along with

decreased VCAN and FCN1 expression. These M2 MoMa clusters also expressed Th2 associated

inflammatory factors such as MRC1 and CCL18.(45) All MoMa clusters expressed FCGR3A (CD16a).(42)

Two additional clusters were labeled as intermediate MoMa because they did not show distinct

transcriptomes corresponding to either M1 or M2 groups. One intermediate MoMa cluster

overexpressed MALAT1, while the other overexpressed metallothionein proteins including MT1F and

MT1G.

We also identified two clusters of CD4+ T cells (CD4 and IL7R), one cluster of T regulatory cells

(IL2RA and LAG3),(46,47) and three clusters of CD8+ T cells (CD8A). Two of the CD8+ T cell clusters were

labeled as CD8+ memory cells due to their high CCL5 and GZMH expression.(48) Other identified

immune cell clusters include natural killer (NK) cells (SPON2 and NCAM1), neutrophils (NAMPT),(34)

naïve B cells (MS4A1), plasmablasts (IGJ and MZB1), plasmacytoid dendritic cells (IRF8 and PLD4),(49,50)

and myeloid dendritic cells (CD1C)and LGALS2).(34,51) In addition to immune cell types, we also found

two epithelial clusters. One contained a mixture of epithelial and granulocyte markers including KRT19

and SLPI(34,52) while the other also contained the additional markers PPIL6 and CFAP300 for

pneumocytes and ciliary cells, respectively.(53,54)

The 26 clusters were consolidated into 15 cell types (**Figure S2)** to streamline further analysis by

combining clusters that are not distinguishable when examining their canonical marker expression

levels. This consolidation also prevents cell groups with many clusters such as the M1 MoMa group from

overshadowing those with fewer clusters in our subsequent differential gene expression ranking

analysis. Cells within each cluster were compared against their original identifications from both their

respective dataset, and most clusters identifications were consistent. The one exception was the

intermediate MoMa type, which was predominantly composed of macrophages from BAL, but also

contained a mixture of monocytes, CD4+ and CD8+ T cell identifications from PBMCs (**Table S3**).

Proinflammatory cell types are enriched in severe COVID-19 patients.

Cell type proportions within BAL and PBMC sample groups showed distinct differences between

patients and healthy controls as well as between mild and severe patients (**Figure 1**). In BAL samples of

patients with severe disease, proinflammatory cells such as M1 MoMa and neutrophils showed

increased abundance ($p <<< .01$, **Table S4**), while immunoregulatory cell types including M2, intermediate

MoMa, and Tregs were less abundant ($p <<< .01$). BAL samples of patients with mild disease showed

decreased abundance of M1 MoMa and neutrophils and increased Tregs and CD8+ memory T cells

compared to healthy controls and severe patients.

Fig 1: Severe patients show increased proportions of proinflammatory cell types. A. Overall average abundance of each major

cell type for all cells. **B.** Per patient abundance of all major cell types for all cells. **C.** Per cohort (bronchoaveolar lavage (BAL) and

peripheral blood mononuclear cells (PBMC)) and per condition (healthy, mild, severe) abundance for each cell type. Conditions

that are significant versus their respective controls are labeled with a triangle ($p < .05$). Conditions that are significant between

severe and both mild as well as healthy controls are labeled with a star ($p < .05$). Conditions that are significant between severe

and mild, but not between severe and healthy controls are labeled with a diamond ($p < .05$).

In PBMCs, the trends in M1 MoMa and neutrophils are reversed. Tregs and CD8+ memory T cells

are less abundant in PBMCs of mild patients. These opposing patterns may illustrate heavy recruitment

of the cell types abundant in BAL, resulting in depletion in the PBMCs that results in an increase in

relative abundance of non-recruited cells in PBMCs. Mild patients also showed an increase of

intermediate MoMa in PBMCs, reinforcing the pattern of relative increases in abundance of

immunoregulatory cell types in mild patients in both BAL and PBMC compartments.

Recurrent DEG (rDEG) modules highlight key pathways in COVID-19 immune response.

We identified an average of 1158 DEGs per cell type for BAL samples, and 260 DEGs per cell type

for PBMC samples.(**Table S5,S6**) After filtering, we identified 50 rDEGs across our 15 cell types that

formed 4 distinct modules (**Figure 2**). Module 1 showed significant GO enrichment for developmental

processes (p<.05) but did not show differential expression between conditions. Module 2 showed

significance for viral defense and Type I interferon GO terms. Most genes in this module were interferon

induced genes including the first three IFIT family genes, ISG15, CXCL10, and MX1. This module was

significantly overexpressed in BAL of all patients versus healthy controls (p<.01).

Fig 2: rDEGs grouped into four distinct modules with immune regulation enriched GO terms. A. Heatmap of modules

generated from the recurrent differentially expressed geens (rDEGs). Triangles indicate significance (p<.05) versus healthy

control within the sample cohort; diamonds indicate significance between severe and mild patients (p<.05). **B.** Module

membership for each module. Modules 1 and 4 contain a mixture of metabolic and immune response related genes. Module 2

contains genes related to interferon activated viral defense. Module 3 contains other inflammatory regulation genes and stress

response genes (generated using the Circlize R package). **C.** Per module GO term enrichment showing the top enriched terms

for each module and their respective p-values with the red line indicating -log10(0.05). The first three modules contain

inflammation related terms in their most enriched terms, while module 4 only contains metabolism related terms.

Module 3 was enriched for macromolecule synthesis and cellular processes. This module

includes the immunomodulatory lncRNAs NEAT1 and MALAT1.(55,56) It also includes MTRNR2L12, an

anti-apoptotic lncRNA, and NFKBIA which is an NF-κB inhibitor. The module was significantly

underexpressed in BAL of mild patients versus healthy controls, and it was overexpressed in BAL of

severe patients versus mild patients. Module 4 had significant terms related to negative regulation of

metabolic processes. This module included the NUPR1 stress response gene, and CSTB which is an

inhibitor of cathepsins like CTSL and CTSB that are involved in COVID-19 viral entry.(57) Module 4 was

significantly underexpressed in BAL of severe patients versus healthy controls.

Stress response, apoptosis, and viral entry related genes show severity dependent expression.

We analyzed individual rDEGs in each of our five most abundant cell types: M1 MoMa, M2

MoMa, CD4+ T cells, NK cells, and CD8+ memory T cells (**Figure 3**). This analysis confirmed previous

reports of downregulation of HLA genes(20,58) such as HLA-DRA and HLA-DRB5 in COVID-19 patients,

with severe patients showing the most downregulation. We also saw upregulation of interferon related

genes including MX1, and IFIT1-3. This increase was greatest in mild patients, correlating with previous

findings of immune exhaustion (**Table S7**).(59,60) Further examination showed additional severity

dependent patterns of differential expression of transcripts related to the stress response, cell death,

and viral entry in cell types involved in the viral immune response.

Fig 3: rDEG expression in most abundant cell types highlights differential immune regulation between mild and severe

patients in both BAL and PBMC cohorts. A. Heatmaps visualizing rDEGs within each of the top five most abundant cell types in

our dataset (generated using the ComplexHeatmap R package). For each cell type, the full rDEG list was filtered via the same p-

values (p<10e-7 for BAL, p<.05 for PBMC) and only rDEGs that are differentially expressed below these thresholds for either BAL

or PBMC are included in the plot. Expression levels are normalized separately for each cohort. The first sidebar indicates which

cohort the particular gene passed the rDEG threshold for, while the second sidebar indicates the ratio of expression of the

particular gene between BAL and PBMC with green (positive values) indicating higher expression levels detected in BAL. **B.**

Visualization of select rDEGs representing pathways outside of the main interferon activated gene group that are relevant to

disease. These genes are visualized separately for each cohort and condition using the sample UMAP projection of cell types

from Figure 1. Each gene shows cell type, cohort, and condition specific differences in localization across the dataset.

265   The NF-κB inhibitor NFKBIA was upregulated in all five most abundant cell types within the BAL

266  of severe patients compared to healthy controls and mild groups. In PBMCs of severe patients, NFKBIA

267  was downregulated compared to healthy controls and mild patients except in CD8+ memory T cells. This

268  pattern of localized overexpression in BAL may indicate increased NFKBIA activity in response to local

269  hyperactivity of NF-κB. Furthermore, the stress response gene NUPR1, whose downregulation leads to

270  cell death, was downregulated in M1 and M2 MoMa in the BAL of severe patients and upregulated in

271  mild patients, indicating a pro-apoptotic shift in severe patient MoMa clusters. NUPR1 was

272  downregulated in BAL of both mild and severe patients for NK cells, CD4+ T cells, and CD8+ Memory T

273  cells.

274   Mild and severe patients also had variable expression of two anti-apoptotic genes, the BCL2

275  inhibitor BCL2A1 and the lncRNA MTRNR2L12. BCL2A1 was significantly upregulated in BAL of severe

276  patients over healthy controls and mild groups for M1 and M2 MoMa, NK cells, and CD4+ T cells. Mild

277  patients showed downregulation of BCL2A1 versus healthy controls in NK and CD4+ T cells. Additionally,

278  MTRNR2L12 was upregulated in BAL of both mild and severe patients in M1 and M2 MoMa, NK cells,

279  CD4+ T cells and CD8+ Memory T cells. The upregulation of these anti-apoptotic genes shows a

280  defensive response to apoptotic cell stresses, particularly in BAL.

281   CTSL, which is a critical protein in the viral entry pathway for COVID-19, was upregulated in BAL

282  of severe patients in M1 and M2 MoMa in mild patients and healthy controls. This suggests a faster viral

283  entry pathway in severe patients, which may contribute to the formation of a hyperinflammatory

284  response. In BAL of NK, CD4+ T cells, and CD8+ Memory T cells, CTSL was downregulated in mild patients

285  and upregulated in severe patients. CTSB, also implicated in viral entry, showed similar patterns.

287  *NEAT1 and MALAT1 are differential regulators of inflammation in severe COVID-19.*

The pro-inflammatory lncRNA NEAT1 passed our rDEG threshold in BAL samples for nine

different cell types, more than any other gene in our analysis. These cell types include M1, M2 and

intermediate MoMa, NK cells, CD4+ T cells, CD8+ memory T cells, naïve B cells, myeloid dendritic cells,

and epithelium/basal cells (**Figure 4**). NEAT1 is localized to the site of infection and inflammation since it

is not differentially expressed in PBMCs. Additionally, among rDEGs, it has one of the highest averages in

log2-fold change between severe and mild patients (**Figure 4**). NEAT1 is overexpressed in BAL of severe

patients and underexpressed in mild patients. The epithelial/basal cell group is the exception where mild

groups also show NEAT1 overexpression over healthy controls, but expression is still significantly higher

in severe patients versus mild patients.

Another immunomodulatory lncRNA, MALAT1, was the second most frequent rDEG in PBMCs. It

passed our rDEG threshold in 6 cell types (tied with ISG15) and 3 cell types in BAL. In BAL derived M1

and M2 MoMa, MALAT1 was underexpressed in mild patients compared to both healthy controls and

severe patients. In CD4+ T cells, MALAT1 shows consistent overexpression in mild patients and

underexpression in severe patients. In PBMCs, MALAT1 was underexpressed in severe patients versus

both healthy controls and mild patients in M1, M2 and intermediate MoMa, NK cells, plasmablasts, and

epithelial/basal cells.

Validation of rDEG expression patterns.

We projected the cell type classifications in our analysis via Seurat's label transfer feature to two

other COVID-19 datasets, one with nasopharyngeal samples(22) and one with PBMC samples(23).

Comparison of cell labels from the original datasets versus our transferred labels shows general

agreement among the cell IDs, allowing us to use our cell type labels for direct comparison of rDEG

patterns in the validation data (**Figure S3&S4**). We compared rDEG expression patterns of the genes

analyzed in the previous two sections in the same five cell types. Among each gene's statistically

significant changes in our analysis between healthy, mild, and severe cases, 36.3% were significant in

our validation cohorts. However, more than two thirds of the non-significant findings in validation were

from comparisons with healthy control in the nasopharyngeal dataset. This is likely due to the small

number of cells recovered from these controls after filtering. Only 1148 cells were recovered from

control samples after filtering compared to 35715 and 25546 for moderate and severe cases

respectively. Among the findings that were significant in both our data and in validation, we found that

79% were significant in the same direction. Notably, NEAT1 showed 100% agreement in validation of

BAL, while MALAT1 showed 100% agreement in validation of PBMCs (**Table S8,S9 & S10**).

Discussion:

Our analysis of BAL and PBMC single cell data in COVID-19 patients has elucidated key

differences between mild and severe disease. We were able to combine cells from both PBMC and BAL

in an integrated analysis. Although our intermediate MoMa group had a mixed group of PBMC cells, our

overall identifications were consistent across both datasets. Furthermore, the cells in the intermediate

MoMa group consisted of cells with weak expression of a wide range of canonical markers. These cells

may be intermediate immune cells from different lineages that share a similar transcriptomic profile. By

conducting analysis simultaneously on cells from the local infection site in the lung as well as the

peripheral immune system, we contrast how the disease manifests and interacts across both

compartments. We have identified differentially expressed genes that vary with severity, are highly

differentially expressed across multiple cell types, and represent key functions related to the

hyperinflammatory disease state. NEAT1 was the most widely differentially expressed gene across cell

types within BAL; it also exhibited a high log-fold change that correlated with disease severity. The

ubiquity of NEAT1, its specific localization to BAL cells, and pro-inflammatory functions suggests that it

may be a key mediator of the inflammation seen in severe COVID-19. NEAT1 is a well characterized

activator of the NLRP3 inflammasome, as well as NLRC4 and AIM2 inflammasomes, which in turn

amplify the inflammatory response.(55) However, an overactive immune response contributes to lasting

tissue damage in severe COVID-19 disease. Intense inflammation through activation of the NLRP3

inflammasome can also lead to pyroptosis, driven by the upregulation of NEAT1.(55,61) These highly

inflammatory and damaging effects of NEAT1 illustrate how overexpression in severe patients might

lead to the inflammatory tissue damage seen in severe COVID-19.

353   MALAT1 also exerts various immunological effects including the mediation of NLRP3

inflammasome activation.(62,63) MALAT1 has been linked to M1-like activity in macrophages,

promoting inflammation.(64) Our finding that MALAT1 is overexpressed in BAL MoMa of severe versus

mild patients suggests that it might be involved in precipitating a shift towards M1 macrophages that

exacerbates inflammation. This is further supported by our findings that severe patients show expansion

of M1 macrophages and decrease of M2 and intermediate macrophages in BAL, while mild patients

show decrease of M1 macrophages. Furthermore, MALAT1 was overexpressed in CD4+ T cells of mild

patients. This is also reflective of MALAT1's protective role in T cells. Loss of MALAT1 expression has

been shown to push T cells towards the inflammatory Th1 and Th17 phenotype while also decreasing

Treg differentiation.(65) This function matches our observed increase in abundance of Tregs in mild

patients. Thus, the upregulation of MALAT1 we see in mild patients may be contributing to the more

subdued immune response observed in these patients.

365   The severity dependent differential expression of other genes in our analysis provides further

evidence of increased cellular stress reflective of a NEAT1 and MALAT1 enhanced hyperinflammatory

state. NF-κB is induced in COVID-19 infection.(66) Although we did not detect differential activity of NF-

κB directly, we found upregulation of its inhibitor NFKBIA in BAL of severe patients which suggests a

feedback response to strong NF-κB activity. NFKBIA's downregulation in PBMCs of severe patients may

be due to localization of cells expressing NFKBIA to the site of infection in attempts to regulate the

hyperactive inflammatory state.(67) The upregulation of BCL2A1 and MTRNR2L12 is also indicative of

extensive cellular stress.(68,69) While MTRNR2L12 is upregulated in both mild and severe disease,

BCL2A1 is upregulated exclusively in severe disease. The increased activity of these anti-apoptotic genes,

particularly in BAL of severe patients, shows additional evidence of the cellular stress induced by

infection and inflammation. These genes may be responding to pyroptosis pathways triggered by

inflammasome activation via NEAT1 and MALAT1. Further evidence of inflammatory cell damage is seen

in the downregulation of NUPR1 in BAL of M1 and M2 macrophages of severe patients with

upregulation in mild patients. Downregulation of this stress response gene has been shown to cause

mitochondrial dysfunction and ROS production that can lead to cell death.(70) Lastly, our observation

that CTSL, a protein crucial for COVID-19 viral entry is upregulated across multiple cell types in severe

patients provides a potential initial mechanism for the induction of the NEAT1 and MALAT1 mediated

inflammatory state through increased efficiency of viral entry.(57)

Limitations in our study include the small sample size, variable clinical presentation and

treatment. Additionally, time from presentation to sample collection varied across patients. The

stratification of patients as severe or mild may also introduce unknown factors due to patient variability

in presentation and classification. Although our validation shows promising reproduction of expression

patterns, additional studies with more subjects and stringent recruiting and sample collection would

further elucidate these findings.

We have demonstrated a clear ensemble of differential gene activity associated with severe

disease in COVID-19 infection that revolves around the lncRNAs NEAT1 and MALAT1. Their specific

activity changes in severe patients coupled with inflammasome promoting functions, suggest important

roles in the COVID-19 hyperinflammatory process. These findings indicate that NEAT1 and MALAT1 may

be candidates for treatment targeting or biological marker exploration.

References

1. Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. RESEARCH High Contagiousness

and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. Emerg Infect Dis

[Internet]. 2020 Jul 1 [cited 2020 Oct 14];26(7):1470–7. Available from:

/pmc/articles/PMC7323562/?report=abstract

2. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates

the rapid dissemination of novel coronavirus (SARS-CoV-2). Science (80-) [Internet]. 2020 May 1

[cited 2020 Oct 14];368(6490):489–93. Available from: http://science.sciencemag.org/

3. García LF. Immune Response, Inflammation, and the Clinical Spectrum of COVID-19 [Internet].

Vol. 11, Frontiers in Immunology. Frontiers Media S.A.; 2020 [cited 2020 Oct 14]. Available from:

https://pubmed.ncbi.nlm.nih.gov/32612615/

4. Vetter P, Eckerle I, Kaiser L. Covid-19: A puzzle with many missing pieces [Internet]. Vol. 368, The

BMJ. BMJ Publishing Group; 2020 [cited 2020 Oct 14]. Available from:

https://twitter.com/WHO/status/

5. Wang F, Qu M, Zhou X, Zhao K, Lai C, Tang Q, et al. The timeline and risk factors of clinical

progression of COVID-19 in Shenzhen, China. J Transl Med [Internet]. 2020 Jul 3 [cited 2020 Oct

14];18(1):270. Available from: https://translational-

medicine.biomedcentral.com/articles/10.1186/s12967-020-02423-8

6. Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus Disease 2019

in China. N Engl J Med [Internet]. 2020 Apr 30 [cited 2020 Oct 14];382(18):1708–20. Available

from: http://www.nejm.org/doi/10.1056/NEJMoa2002032

7. Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, et al. Clinical Characteristics of Covid-19

in New York City. N Engl J Med [Internet]. 2020 Jun 11 [cited 2020 Oct 14];382(24):2372–4.

Available from: http://www.nejm.org/doi/10.1056/NEJMc2010419

8.   Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LFP. The trinity of COVID-19: immunity, inflammation

and intervention [Internet]. Vol. 20, Nature Reviews Immunology. Nature Research; 2020 [cited

2020 Oct 14]. p. 363–74. Available from: /pmc/articles/PMC7187672/?report=abstract

9.   Zaim S, Chong JH, Sankaranarayanan V, Harky A. COVID-19 and Multiorgan Response [Internet].

Vol. 45, Current Problems in Cardiology. Mosby Inc.; 2020 [cited 2020 Oct 14]. Available from:

/pmc/articles/PMC7187881/?report=abstract

10.   van den Berg DF, te Velde AA. Severe COVID-19: NLRP3 Inflammasome Dysregulated. Front

Immunol [Internet]. 2020 Jun 26 [cited 2020 Oct 14];11:1580. Available from:

https://www.frontiersin.org/article/10.3389/fimmu.2020.01580/full

11.   Evavold CL, Kagan JC. How Inflammasomes Inform Adaptive Immunity [Internet]. Vol. 430,

Journal of Molecular Biology. Academic Press; 2018 [cited 2020 Oct 14]. p. 217–37. Available

from: /pmc/articles/PMC5766381/?report=abstract

12.   Li G, Fan Y, Lai Y, Han T, Li Z, Zhou P, et al. Coronavirus infections and immune responses

[Internet]. Vol. 92, Journal of Medical Virology. John Wiley and Sons Inc.; 2020 [cited 2020 Oct

14]. p. 424–32. Available from: https://pubmed.ncbi.nlm.nih.gov/31981224/

13.   Guo H, Callaway JB, Ting JPY. Inflammasomes: Mechanism of action, role in disease, and

therapeutics [Internet]. Vol. 21, Nature Medicine. Nature Publishing Group; 2015 [cited 2020 Oct

14]. p. 677–87. Available from: /pmc/articles/PMC4519035/?report=abstract

14.   Freeman TL, Swartz TH. Targeting the NLRP3 Inflammasome in Severe COVID-19 [Internet]. Vol.

11, Frontiers in Immunology. Frontiers Media S.A.; 2020 [cited 2020 Sep 25]. p. 1518. Available

from: www.frontiersin.org

15.   Farag NS, Breitinger U, Breitinger HG, El Azizi MA. Viroporins and inflammasomes: A key to

understand virus-induced inflammation [Internet]. Vol. 122, International Journal of Biochemistry

and Cell Biology. Elsevier Ltd; 2020 [cited 2021 Mar 2]. p. 105738. Available from:

/pmc/articles/PMC7102644/

16.   Yap JKY, Moriyama M, Iwasaki A. Inflammasomes and Pyroptosis as Therapeutic Targets for

COVID-19. J Immunol [Internet]. 2020 Jul 15 [cited 2020 Oct 14];205(2):307–12. Available from:

http://www.jimmunol.org/content/205/2/307

17.   Yue Y, Nabar NR, Shi CS, Kamenyeva O, Xiao X, Hwang IY, et al. SARS-Coronavirus Open Reading

Frame-3a drives multimodal necrotic cell death. Cell Death Dis [Internet]. 2018 Sep 1 [cited 2020

Oct 14];9(9):1–15. Available from: https://www.nature.com/articles/s41419-018-0917-y

18.   Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: Host cell death and inflammation [Internet]. Vol.

7, Nature Reviews Microbiology. NIH Public Access; 2009 [cited 2020 Sep 25]. p. 99–109.

Available from: /pmc/articles/PMC2910423/?report=abstract

19.   Liao M, Liu Y, Yuan J, Wen Y, Xu G, Zhao J, et al. Single-cell landscape of bronchoalveolar immune

cells in patients with COVID-19. Nat Med [Internet]. 2020 May 12 [cited 2020 May 15];1–3.

Available from: http://www.nature.com/articles/s41591-020-0901-9

20.   Wilk AJ, Rustagi A, Zhao NQ⬚. A single-cell atlas of the peripheral immune response in patients

with severe COVID-19. Nat Med [Internet]. 2020; Available from:

https://doi.org/10.1038/s41591-020-0944-y

21.   Zhang JY, Wang XM, Xing X, Xu Z, Zhang C, Song JW, et al. Single-cell landscape of immunological

responses in patients with COVID-19. Nat Immunol [Internet]. 2020 Sep 1 [cited 2020 Sep

25];21(9):1107–18. Available from: https://doi.org/10.1038/s41590-020-0762-x

22.   Chua RL, Lukassen S, Trump S, Hennig BP, Wendisch D, Pott F, et al. COVID-19 severity correlates

with airway epithelium–immune cell interactions identified by single-cell analysis. Nat Biotechnol

[Internet]. 2020 Aug 1 [cited 2020 Sep 25];38(8):970–9. Available from:

https://doi.org/10.1038/s41587-020-0602-4

23.   Schulte-Schrepping J, Reusch N, Paclik D, Baßler K, Schlickeiser S, Zhang B, et al. Severe COVID-19

Is Marked by a Dysregulated Myeloid Cell Compartment. Cell [Internet]. 2020 Sep 17 [cited 2020

Sep 25];182(6):1419. Available from: /pmc/articles/PMC7405822/?report=abstract

24.   Lee JS, Park S, Jeong HW, Ahn JY, Choi SJ, Lee H, et al. Immunophenotyping of covid-19 and

influenza highlights the role of type i interferons in development of severe covid-19. Sci Immunol

[Internet]. 2020 Jul 1 [cited 2020 Oct 14];5(49):1554. Available from:

http://immunology.sciencemag.org/

25.   Diaz J V, Baller A, Banerjee A, Bertagnolio S, Bonet M, Bosman A, et al. Clinical management of

COVID-19: interim guidance. 2020;WHO/2019-nCoV/clinical/2020.5. Available from:

https://www.who.int/publications/i/item/clinical-management-of-covid-19

26.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al. Comprehensive

Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-1902.e21.

27.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data using

regularized negative binomial regression. Genome Biol [Internet]. 2019 Dec 23 [cited 2020 Jun

8];20(1):296. Available from:

https://genomebiology.biomedcentral.com/articles/10.1186/s13059-019-1874-1

28.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data

across different conditions, technologies, and species. Nat Biotechnol [Internet]. 2018 Apr 2

[cited 2018 Jul 25];36(5):411–20. Available from:

http://www.nature.com/doifinder/10.1038/nbt.4096

29.   McInnes L, Healy J, Saul N, Großberger L. UMAP: Uniform Manifold Approximation and

Projection. J Open Source Softw [Internet]. 2018 Sep 2 [cited 2020 Oct 14];3(29):861. Available

from: http://joss.theoj.org/papers/10.21105/joss.00861

30.   Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, et al. MAST: A flexible statistical

framework for assessing transcriptional changes and characterizing heterogeneity in single-cell

RNA sequencing data. Genome Biol [Internet]. 2015 Dec 10 [cited 2020 Jun 8];16(1):278.

Available from: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-015-0844-5

31.   Soneson C, Robinson MD. Bias, robustness and scalability in single-cell differential expression

analysis. Nat Publ Gr. 2018;15.

32.   Zhang X, Lan Y, Xu J, Quan F, Zhao E, Deng C, et al. CellMarker: A manually curated resource of

cell markers in human and mouse. Nucleic Acids Res. 2019;47(D1):D721–8.

33.   Franzén O, Gan LM, Björkegren JLM. PanglaoDB: A web server for exploration of mouse and

human single-cell RNA sequencing data. Database [Internet]. 2019 Jan 1 [cited 2020 Oct

14];2019(1):46. Available from:

https://academic.oup.com/database/article/doi/10.1093/database/baz046/5427041

34.   Monaco G, Lee B, Xu W, Mustafah S, Hwang YY, Carré C, et al. RNA-Seq Signatures Normalized by

mRNA Abundance Allow Absolute Deconvolution of Human Immune Cell Types. Cell Rep

[Internet]. 2019 Feb 5 [cited 2019 Apr 4];26(6):1627-1640.e7. Available from:

https://www.sciencedirect.com/science/article/pii/S2211124719300592

35.   Wickham H. ggplot2: Elegant Graphics for Data Analysis [Internet]. Cham: Springer International

Publishing; 2016 [cited 2020 Oct 14]. (Use R!). Available from:

http://link.springer.com/10.1007/978-3-319-24277-4

36.   Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell transcriptional

landscape of mammalian organogenesis. Nature [Internet]. 2019 Feb 20 [cited 2019 Mar

13];566(7745):496–502. Available from: http://www.nature.com/articles/s41586-019-0969-x

37.   Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, et al. Reversed graph embedding resolves

complex single-cell trajectories. Nat Methods [Internet]. 2017;14(10):979–82. Available from:

http://dx.doi.org/10.1038/nmeth.4402

38.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and regulators

of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol

[Internet]. 2014 Mar 23 [cited 2020 Oct 14];32(4):381–6. Available from:

https://www.nature.com/articles/nbt.2859

39.   Alexa A, Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology. Bioconductor. 2020;R

package.

40.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics [Internet]. 2016 Sep 15 [cited 2020 Oct

16];32(18):2847–9. Available from: https://pubmed.ncbi.nlm.nih.gov/27207943/

41.   Gu Z, Gu L, Eils R, Schlesner M, Brors B. Circlize implements and enhances circular visualization in

R. Bioinformatics [Internet]. 2014 Apr 2 [cited 2020 Oct 16];30(19):2811–2. Available from:

https://pubmed.ncbi.nlm.nih.gov/24930139/

42.   Kapellos TS, Bonaguro L, Gemünd I, Reusch N, Saglam A, Hinkley ER, et al. Human monocyte

subsets and phenotypes in major chronic inflammatory diseases [Internet]. Vol. 10, Frontiers in

Immunology. Frontiers Media S.A.; 2019 [cited 2020 Aug 19]. p. 2035. Available from:

www.frontiersin.org

43.   Chang MY, Kang I, Gale M, Manicone AM, Kinsella MG, Braun KR, et al. Versican is produced by

trif- and type I interferon-dependent signaling in macrophages and contributes to fine control of

innate immunity in lungs. Am J Physiol - Lung Cell Mol Physiol [Internet]. 2017 Dec 1 [cited 2020

Aug 19];313(6):L1069–86. Available from: /pmc/articles/PMC5814701/?report=abstract

44.   Wang S, Song R, Wang Z, Jing Z, Wang S, Ma J. S100A8/A9 in inflammation [Internet]. Vol. 9,

Frontiers in Immunology. Frontiers Media S.A.; 2018 [cited 2020 Aug 19]. p. 1298. Available from:

www.frontiersin.org

45.   Palomino DC arolin. T, Marti LC avalheir. Chemokines and immunity [Internet]. Vol. 13, Einstein

(São Paulo, Brazil). Instituto de Ensino e Pesquisa Albert Einstein; 2015 [cited 2020 Aug 19]. p.

469–73. Available from: /pmc/articles/PMC4943798/?report=abstract

46.   Mohr A, Malhotra R, Mayer G, Gorochov G, Miyara M. Human FOXP3+ T regulatory cell

heterogeneity [Internet]. Vol. 7, Clinical and Translational Immunology. Wiley-Blackwell; 2018

[cited 2020 Oct 14]. Available from: /pmc/articles/PMC5822410/?report=abstract

47.   Okamura T, Fujio K, Shibuya M, Sumitomo S, Shoda H, Sakaguchi S, et al. CD4+CD25-LAG3+

regulatory T cells controlled by the transcription factor Egr-2. Proc Natl Acad Sci U S A [Internet].

2009 Aug 18 [cited 2020 Oct 14];106(33):13974–9. Available from:

www.pnas.org/cgi/content/full/

48.   Weng NP, Araki Y, Subedi K. The molecular basis of the memory T cell response: Differential gene

expression and its epigenetic regulation [Internet]. Vol. 12, Nature Reviews Immunology. NIH

Public Access; 2012 [cited 2020 Oct 14]. p. 306–15. Available from:

/pmc/articles/PMC4686144/?report=abstract

49.   Sichien D, Scott CL, Martens L, Vanderkerken M, Van Gassen S, Plantinga M, et al. IRF8

Transcription Factor Controls Survival and Function of Terminally Differentiated Conventional and

Plasmacytoid Dendritic Cells, Respectively. Immunity. 2016 Sep 20;45(3):626–40.

50.   Gavin AL, Huang D, Huber C, Mårtensson A, Tardif V, Skog PD, et al. PLD3 and PLD4 are single-

stranded acid exonucleases that regulate endosomal nucleic-acid sensing. Nat Immunol

[Internet]. 2018 Sep 1 [cited 2020 Oct 15];19(9):942–53. Available from:

/pmc/articles/PMC6105523/?report=abstract

51. Collin M, Bigley V. Human dendritic cell subsets: an update [Internet]. Vol. 154, Immunology.

Blackwell Publishing Ltd; 2018 [cited 2020 Oct 15]. p. 3–20. Available from:

/pmc/articles/PMC5904714/?report=abstract

52. Camper N, Glasgow AMA, Osbourn M, Quinn DJ, Small DM, McLean DT, et al. A secretory

leukocyte protease inhibitor variant with improved activity against lung infection. Mucosal

Immunol [Internet]. 2016 May 1 [cited 2020 Oct 15];9(3):669–76. Available from:

www.nature.com/mi

53. Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Tissue-based

map of the human proteome. Science (80-) [Internet]. 2015 Jan 23 [cited 2020 Oct

15];347(6220):1260419–1260419. Available from:

https://www.sciencemag.org/lookup/doi/10.1126/science.1260419

54. Zietkiewicz E, Bukowy-Bieryllo Z, Rabiasz A, Daca-Roszak P, Wojda A, Voelkel K, et al. CFAP300:

Mutations in slavic patients with primary ciliary dyskinesia and a role in ciliary dynein arms

trafficking. Am J Respir Cell Mol Biol [Internet]. 2019 Oct 1 [cited 2020 Oct 15];61(4):400–49.

Available from: https://www.atsjournals.org/doi/10.1165/rcmb.2018-0260OC

55. Zhang P, Cao L, Zhou R, Yang X, Wu M. The lncRNA Neat1 promotes activation of inflammasomes

in macrophages. Nat Commun [Internet]. 2019 Dec 1 [cited 2020 Sep 20];10(1):1–17. Available

from: https://doi.org/10.1038/s41467-019-09482-6

56. Amodio N, Raimondi L, Juli G, Stamato MA, Caracciolo D, Tagliaferri P, et al. MALAT1: A druggable

long non-coding RNA for targeted anti-cancer approaches [Internet]. Vol. 11, Journal of

Hematology and Oncology. BioMed Central Ltd.; 2018 [cited 2020 Oct 14]. p. 1–19. Available

from: https://doi.org/10.1186/s13045-018-0606-4

57.   Bittmann S, Weissenstein A, Villalon G, Moschuring-Alieva E, Luchter E. Simultaneous Treatment

of COVID-19 With Serine Protease Inhibitor Camostat and/or Cathepsin L Inhibitor? J Clin Med

Res [Internet]. 2020 [cited 2020 Oct 14];12(5):320–2. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7239585/

58.   Zmijewski JW, Pittet JF. Human Leukocyte Antigen-DR Deficiency and Immunosuppression-

Related End-Organ Failure in SARS-CoV2 Infection [Internet]. Anesthesia and Analgesia.

Lippincott Williams and Wilkins; 2020 [cited 2020 Oct 14]. p. 989–92. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7386673/

59.   Zheng M, Gao Y, Wang G, Song G, Liu S, Sun D, et al. Functional exhaustion of antiviral

lymphocytes in COVID-19 patients [Internet]. Vol. 17, Cellular and Molecular Immunology.

Springer Nature; 2020 [cited 2020 Oct 14]. p. 533–5. Available from:

https://doi.org/10.1038/s41423-020-0402-2

60.   Mathew D, Giles JR, Baxter AE, Oldridge DA, Greenplate AR, Wu JE, et al. Deep immune profiling

of COVID-19 patients reveals distinct immunotypes with therapeutic implications. Science (80-)

[Internet]. 2020 Sep 4 [cited 2020 Oct 14];369(6508). Available from:

https://doi.org/10.1126/science.abc8511

61.   Zhan J-F, Huang H-W, Huang C, Hu L-L, Xu W-W. Long Non-Coding RNA NEAT1 Regulates

Pyroptosis in Diabetic Nephropathy via Mediating the miR-34c/NLRP3 Axis. Kidney Blood Press

Res [Internet]. 2020 [cited 2020 Oct 15];45(4):589–602. Available from:

https://www.karger.com/Article/FullText/508372

62.   Menon MP, Hua K-F. The Long Non-coding RNAs: Paramount Regulators of the NLRP3

Inflammasome. Front Immunol [Internet]. 2020 Sep 25 [cited 2020 Oct 15];11:569524. Available

from: www.frontiersin.org

63.   Yu S yang, Dong B, Tang L, Zhou S hua. LncRNA MALAT1 sponges miR-133 to promote NLRP3

inflammasome expression in ischemia-reperfusion injured heart [Internet]. Vol. 254,

International Journal of Cardiology. Elsevier Ireland Ltd; 2018 [cited 2020 Oct 15]. p. 50. Available

from: https://pubmed.ncbi.nlm.nih.gov/29407129/

64.   Cui H, Banerjee S, Guo S, Xie N, Ge J, Jiang D, et al. Long noncoding RNA Malat1 regulates

differential activation of macrophages and response to lung injury. JCI insight [Internet]. 2019

Feb 21 [cited 2020 Oct 15];4(4). Available from: /pmc/articles/PMC6478413/?report=abstract

65.   Masoumi F, Ghorbani S, Talebi F, Branton WG, Rajaei S, Power C, et al. Malat1 long noncoding

RNA regulates inflammation and leukocyte differentiation in experimental autoimmune

encephalomyelitis. J Neuroimmunol [Internet]. 2019 Mar 15 [cited 2020 Sep 20];328:50–9.

Available from: https://pubmed.ncbi.nlm.nih.gov/30583215/

66.   Hirano T, Murakami M. COVID-19: A New Virus, but a Familiar Receptor and Cytokine Release

Syndrome. Immunity. 2020 May 19;52(5):731–3.

67.   Ali S, Hirschfeld AF, Mayer ML, Fortuno ES, Corbett N, Kaplan M, et al. Functional Genetic

Variation in NFKBIA and Susceptibility to Childhood Asthma, Bronchiolitis, and

Bronchopulmonary Dysplasia . J Immunol [Internet]. 2013 Apr 15 [cited 2020 Oct

15];190(8):3949–58. Available from: http://www.jimmunol.org/content/190/8/3949

68.   Vogler M. BCL2A1: The underdog in the BCL2 family [Internet]. Vol. 19, Cell Death and

Differentiation. Nature Publishing Group; 2012 [cited 2020 Oct 15]. p. 67–74. Available from:

www.nature.com/cdd

69.   Du C, Xie H, Zang R, Shen Z, Li H, Chen P, et al. Apoptotic neuron-secreted HN12 inhibits cell

apoptosis in Hirschsprung's disease. Int J Nanomedicine [Internet]. 2016 Nov 7 [cited 2020 Oct

15];11:5871–81. Available from: /pmc/articles/PMC5106231/?report=abstract

70.   Santofimia-Castaño P, Lan W, Bintz J, Gayet O, Carrier A, Lomberk G, et al. Inactivation of NUPR1

promotes cell death by coupling ER-stress responses with necrosis. Sci Rep [Internet]. 2018 Dec 1

[cited 2020 Oct 15];8(1). Available from: /pmc/articles/PMC6242935/?report=abstract

Supplementary Figures

S1 Figure: Filtering of genes with extremely high residual variance. A. A plot of mean expression levels

versus residual variance for all genes detected in dataset. Genes with the highest residual variance are

nearly all immunoglobulin and ribosome genes. **B.** Plot of 100 genes with highest residual variance in

descending order. The rate of variance decrease stabilizes after the 21^{st} gene. **C.** Table of top 21 genes

filtered out of downstream analysis due to very high residual variance.

S2 Figure: Unbiased clustering of combined BAL and PBMC data reveals common cell types between

cohorts and sample conditions. A/B. UMAP plots showing the distribution of all cells in both cohorts

within defined cell types. Plot A shows the 15 major cell groups we identified with labels over the cluster

centers. Plot B shows the subclusters making up those groups. The major groups are abbreviated as

prefixes with a letter suffix indicating subgroup. M1, M2 and Int are the macrophage/monocytes. NK is

natural killer cells, E/G is epithelial cells and granulocytes, and E/P/C is epithelia, pneumocyte, and ciliary

cells. **C.** Select markers for major cell groups plotted on the same UMAP projection. This illustrates the

specificity of these markers for different regions of the plot corresponding to our respective cell types.

D. Dot plot visualization of top markers utilized for identification of each cell cluster. The size of each dot

indicates the percentage of cells with detectable expression of each gene, with color indicating

expression level. **E.** Splitting the UMAP by cohort and by severity, with "H" indicating healthy control,

"M" indicating mild disease, and "S" indicating severe disease illustrates that cell clusters do not

organize according to sample type or patient condition, indicating successful integration of the datasets.

S3 Figure: Transferred cell identities correspond to original cell identities from the nasopharyngeal

validation set. Dot plot shows cells from the nasopharyngeal validation dataset, identified by their

transferred labels on the Y axis and their original labels on the X axis. Color of each dot indicates the

numeric log frequency of cells that fit each corresponding set of labels. The size of each dot represents

the percentage of each transferred cell types which is represented by each original cell type. From left to

right, the full cell type names from the nasopharyngeal dataset are: non-resident macrophage (nrMa),

monocyte-derived macrophage (MoD-Ma), monocyte-derived dendritic cell (moDC), resident

macrophage (rMa), cytotoxic T cell (CTL), regulatory T cell (Treg), natural killer T cell (NKT), proliferating

natural killer T cell (NKT-p), natural killer cells (NK), neutrophils (Neu), B cells, plasmacytoid dendritic cell

(pDC), basal cell, ciliated cell, differentiating ciliated cell, FOXN4+ epithelial cell, ionocyte, interferon

responsive cell (IRC), mast cell (MC), epithelial outliers, secretory, differentiating secretory, squamous,

and unknown epithelial.

S4 Figure: Transferred cell identities correspond to original cell identities from the PBMC validation

set. Dot plot shows cells from the PBMC validation set, identified by their transferred labels on the Y axis

and their original labels on the X axis. Color of each dot indicates the numeric log frequency of cells that

fit each corresponding set of labels. The size of each dot represents the percentage of each transferred

cell types which is represented by each original cell type. This dataset contained several cell types where

the same label was applied to more than one subcluster, resulting in numeric suffixes for similar cell

types. mDCs correspond to myeloid dendritic cells and pDCs correspond to plasmacytoid dendritic cells.

Supplementary Excel Tables

S1 Table: Key characteristics of patients within each dataset. Patient information from the BAL and

PBMC cohorts used in this analysis. Patients who were intubated or had PaO2/FiO2 ⬚≤⬚300 mmHg were

classified as severe. In BAL, patient "Mild 3" had only 369 cells recovered after filtering and was only

used for initial clustering. Most patients in the BAL cohort. Exact ages were not available in the PBMC

cohort. The first patient in this cohort was sampled twice, once while classified as a mild patient, and

once after their symptoms worsened and required mechanical ventilation. Several patients in the PBMC

cohort received azithromycin, which can have immunomodulatory effects before sample collection.

S2 Table: Demographic characteristics of healthy subjects. All healthy controls used from both the BAL

and PBMC cohorts are listed.

S3 Table: Cell identities from original datasets versus new combined identification. For BAL and

PBMCs, each cell's original cell identification and new identification are tabulated for the 26 clusters in

the subgroup sheets, and the 15 consolidated cell groups in the coarse sheets.

S4 Table: P-value tables for cell proportion comparisons across each cell type. Conditions compared

are listed in the first two columns, and FDR adjusted p-values are listed in the third column. Each sheet is

labeled by cell type.

S5 and S6 Tables: DEGs for BAL and PBMC samples respectively, separated by cell type, with raw p-

values as well as FDR adjusted p-values. Each sheet is labeled by cell type. Column headings include

indicators for which conditions are being compared where applicable. The conditions are numbered:

"1=healthy control", "2=mild COVID-19 patient", "3=severe COVID-19 patient". For example, the prefix

"g2_1" indicates the comparison of mild patient expression levels minus healthy control expression

levels. Log-fold change is reported relative to the natural log. Columns labeled "pct.1", "pct.2", or

"pct.3" indicate the percentage of cells in the condition corresponding to that number with detectable

expression of a particular gene.

S7 Table: rDEGs with their expression levels in each cell type where each rDEG's adjusted p-values

passed the p-value filter as defined in our Methods section. Column headings include indicators for

which conditions are being compared where applicable. The conditions are numbered: "1=healthy

control", "2=mild COVID-19 patient", "3=severe COVID-19 patient". For example, the prefix "g2_1"

indicates the comparison of mild patient expression levels minus healthy control expression levels. Log-

fold change is reported relative to the natural log. Columns labeled "pct.1", "pct.2", or "pct.3" indicate

the percentage of cells in the condition corresponding to that number with detectable expression of a

particular gene. The "celltype" and "sample" columns indicate which cell type and which sample

condition the rDEG passed filter in.

S8 Table: Comparisons between differentially expressed rDEGs discussed in our results shows strong

agreement in validation datasets. A/B. Tables representing the tally of differential expression results of

our discussed rDEGs which agreed or disagreed between analysis and validation groups based on the

direction of detected differential expression. Tables are split by BAL/nasopharyngeal and PBMC groups.

The first three columns correspond to cases where a comparison is not available due to a lack of

differential expression detected in the original analysis (na.orig), the validation set (na.val), or both

(na.all). The top half of each table reports the results for severe vs mild cases only (SvsM) while the

bottom half reports results for all three comparisons: healthy vs mild, healthy vs severe, and severe vs

mild.

S9 and S10 Tables: Per gene level validation tables for BAL and PBMC groups respectively. Each sheet

name corresponds to the cell type presented. Row names indicate the gene being compared, and

column names indicate the cohorts being compared: healthy (H), mild COVID (M), severe COVID (S).

When differential expression is detected in both the original analysis and validation, the column is labled

agree if the change occurred in the same direction and disagree if it is opposite. Other labels indicate

where a comparison is not available due to a lack of differential expression detected in the original

analysis (na.orig), the validation set (na.val), or both (na.all).

A

B

C

A

B