

How unequal vaccine distribution promotes the evolution of vaccine escape

Philip J Gerrish^{1,2,3,*}, Fernando Saldaña⁴, Benjamin Galeota-Sprung⁵, Alexandre Colato⁶,
Erika E Rodriguez⁷, Jorge X Velasco Hernández⁴

1. Department of Biology, University of New Mexico, Albuquerque NM 87131 USA
2. Theoretical Biology & Biophysics, Los Alamos National Lab, Los Alamos NM 87545 USA
3. Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, México
4. Instituto de Matemáticas, Universidad Nacional Autónoma de México, Juriquilla, México
5. Department of Biology, University of Pennsylvania, Philadelphia PA 19104 USA
6. Departamento Ciências da Natureza, Matemática e Educação, Universidade Federal de São Carlos, Araras SP, Brazil
7. Instituto de Ciencias Básicas e Ingeniería (ICBI), Universidad Autónoma del Estado de Hidalgo, Pachuca, México

Classification: Evolution, Epidemiology, Applied Mathematics

Keywords: Evolutionary epidemiology, Viral evolution, SARS-CoV-2, Antigenic mutation

* To whom correspondence should be addressed:

Philip Gerrish
167 Castetter Hall, MSC03 2020
1 University of New Mexico
Albuquerque, NM 87131-0001

email: pgerrish@unm.edu
phone: +1 505 695 2770

Abstract

In the context of the ongoing SARS-CoV-2 pandemic, health officials warn that vaccines must be uniformly distributed within and among countries if we are to quell the pandemic. Yet there has been little critical assessment of the underlying reasons for this warning. Here, we explicitly show why vaccine equity is necessary. We begin by drawing an analogy to studies showing how disparities in drug concentration within a single host can promote the evolution of drug resistance, and we then proceed to mathematical modeling and simulation of vaccine escape evolution in structured host populations. Perhaps counter-intuitively, we find that vaccine escape mutants are less likely to come from vaccinated regions where there is strong selection pressure for vaccine escape and more likely to come from a neighboring unvaccinated region where there is no selection for escape. Unvaccinated geographic regions thus provide evolutionary reservoirs from which vaccine escape mutants can arise and infect neighboring vaccinated regions, causing new local epidemics within those regions and beyond. Our findings have timely implications for vaccine rollout strategies and public health policy.

In the face of the ongoing SARS-CoV-2 pandemic, researchers and companies around the world have vacillated between competition and cooperation in the race for a vaccine that could restore some societal and economic normalcy. As of 16 March 2021, there are 82 vaccines in clinical development and another 182 vaccines in pre-clinical stages. These very hopeful advances happened in record time: the vaccine from the University of Oxford and the AstraZeneca vaccine took approximately 10 months to be ready for use.

Vaccine rollout and distribution are still just beginning at the time of writing this paper. In the western hemisphere, the three largest countries – the US, Brazil, and Mexico – have vaccinated 12%, 2%, and 1% of their populations, respectively [25]. Significantly, the distribution of the vaccine to date has been highly non-uniform among and within these three countries and around the globe.

As vaccines are being distributed, the emergence of vaccine escape presents a major concern [6, 12, 14, 16, 22, 31–33]. Vaccine escape is the appearance and spread of viral variants that can infect and cause illness in vaccinated hosts. These variants can achieve this because they have acquired mutations allowing them to escape detection by antibodies created in response to vaccination. Researchers around the globe continue to identify emerging variants of SARS-CoV-2 that are increasingly refractory to vaccine-induced antibodies. These strains carry mutations in the receptor-binding domain (RBD) and the N-terminal domain (NTD) of the spike surface glycoprotein targeted by vaccines [6, 31, 32]. Put differently, vaccine escape in the ongoing pandemic is, to some degree, already occurring.

The threat of vaccine escape in the past has depended largely on the particular virus. Polio and measles

vaccines are two stellar examples of highly effective vaccines to which the respective viruses have not evolved escape strains. Flu vaccines, on the other hand, are notoriously “leaky”, with rampant vaccine escape emerging every flu season and creating the need for a new flu vaccine every year. Vaccine escape has effectively prevented the development of an HIV vaccine because mutants able to “escape” any conceivable vaccine target preexist in circulating virus. Where the many different SARS-CoV-2 vaccines stand in this wide spectrum of vaccine-escape susceptibility is still a matter of debate, but increasingly the evidence indicates escape is a real threat [6, 12, 14, 16, 22, 31–33]. The E484K mutation in the backgrounds of UK variant B.1.1.7 or South African variant B.1.135 are two particularly worrisome variants [14, 32]. There is even some concern, and evidence, that new variants may be able to evade natural immunity to SARS-CoV-2 in previously-infected hosts through “immune escape” [9, 14, 31]; this does not bode well for prospects of lasting vaccine-induced immunity [1, 20]. Finally, a recent study [8] reveals that closely-related endemic human coronavirus 229E displays evidence of “antigenic drift” – the same process of rapid antigenic evolution that occurs in Influenza.

Vaccine escape can be viewed as analogous to drug resistance. In the evolution of drug resistance, it is well-established that “privileged sites” in an infected host in which the administered drug is somehow restricted due to physiological constraints (e.g, the blood-brain barrier [19]), can play a very key role [3, 17, 19]. In such sites, the population size of the infectious agent can remain large because there is essentially no drug to suppress it. In these large sub-populations, mutants that are resistant to the drug can increase in frequency without selective constraints for or against. When resistant mutants from such privileged sites migrate back into sites with unrestricted drug concentrations, these “unprivileged sites” quickly succumb to fixation of resistant mutants. The same phenomenon is well-documented in biofilms [7, 17, 28, 30], wherein regions of a biofilm that are shielded from antibiotics provide reservoirs in which resistance mutations evolve neutrally and can subsequently migrate to unshielded regions, rendering the antibiotic ineffective in these regions and eventually in the entire biofilm. The same principle even applies to cancer: disparities in drug concentration can promote the evolution of drug resistance [11].

The lessons learned from drug resistance point to two key factors that could facilitate the evolution of vaccine escape, namely, population size and population structure [5, 18]. Population size is important because the overwhelming majority of mutations occur during replication. Smaller populations mean fewer replications which means reduced opportunity for mutations to arise; this simple principle is the basis for health officials’ repeated pleas for continued social distancing and facemask usage (in the context of the current SARS-CoV-2 pandemic) despite the existence of vaccines. Population structure is important because local epidemics can

vary in size and vaccine coverage and thus harbor the vaccine equivalent of privileged sites, mentioned above, albeit at the host-population and not within-host level. A recent study [13] looks at the effects of different kinds of population compartmentalization on the risk of vaccine escape, namely, age and vulnerability.

A further lesson from drug resistance studies derives from the observation that variability in drug distribution can have more of an impact on the evolution of resistance than overall drug concentration [5, 18]. Extrapolating to vaccine escape, this would indicate that geographic variability in vaccine distribution can pose a bigger threat of vaccine escape than factors such as public distrust and fake news that reduce vaccine participation throughout the population.

To assess claims that vaccine equity is essential, and to validate our verbal extrapolations from drug resistance evolution, we employ mathematical models and simulations of vaccine escape evolution. In our basic model, there are just two local epidemics in geographically neighboring regions or “patches”. One patch has access to a vaccine, the other does not. We study how the unvaccinated patch affects the probability of vaccine escape in the vaccinated patch.

We can assume that an escape mutant will always have a selective advantage in a vaccinated population (SM), simply because there is a larger number of hosts it can infect (susceptible and vaccinated hosts) than the wildtype (infects susceptible hosts only). Thus, we do not need to explicitly model the transmission of and dynamics of escape mutants after they have emerged; we can focus simply on the timing of emergence of the first escape mutant. To this end, we simply model the accumulation of escape mutations from wildtype and focus on the timing of the first infection event in which a new host is infected with an escape mutant, which we will call an “escape-infection” event.

In vaccinated Patch 1, there will be strong selection for vaccine escape but limited opportunity for escape mutations to arise simply because of the reduced number of unvaccinated susceptible hosts. In unvaccinated Patch 2, escape mutations have no selective advantage, but there is a larger number of unvaccinated susceptible hosts.

Our model is described by the following equations:

$$\begin{aligned} \dot{S}_j &= - \sum_k^n \beta_{kj} I_k S_j - \phi_j S_j, & \dot{I}_j &= \sum_k^n \beta_{jk} I_j S_k - (\gamma + U) I_j, \\ \dot{V}_j &= \phi_j S_j, & \dot{E}_j &= U I_j - \gamma E_j, & \dot{R}_j &= \gamma (I_j + E_j), \end{aligned} \tag{1}$$

where S_j , V_j , I_j , E_j , and R_j are the fraction of the population that are susceptible, vaccinated, infected,

infected with escape mutant, and recovered, respectively, in Patch j ; β_{ij} is the transmission rate from Patch i to Patch j (β_{jj} is the transmission rate within Patch j); ϕ_j is vaccination rate in Patch j ; γ is recovery rate; U is a composite per-host mutation rate from wildtype virus to escape mutant virus (see discussion below); n is number of patches; dots indicate time derivatives.

We note the absence of a contagion term in the equation for E_j . This term is not needed for our purposes because our focus is only on the first escape-infection event – a discrete event. Furthermore, this term can lead to erroneous results because ours is a continuous model: a contagion term would allow for transmission to fractions of individual hosts that can erroneously amplify the vaccine escape mutant prior to the first escape-infection event.

Here, we assume there are only two patches, $n = 2$ and $j \in [1, 2]$. Our more complex models and detailed simulations are described in the Supplementary Materials (SM).

We define random variable T_{ij} as the time of the first infection event in which a new host in Patch j is infected by an escape mutant that arose in Patch i . Such infection events occur with rate $r_{ij}(t) = \beta_{ij}E_i(t)(S_j(t) + \sigma V_j(t))$, where σ allows for varying levels of escape reflecting the observed spectrum of partial immunity against different variants ranging from no escape $\sigma = 0$ to full escape $\sigma = 1$.

For now, we will assume intra-patch transmission rates are equal, $\beta_{jj} = \beta$, and inter-patch transmission rates are equal, $\beta_{ij|_{i \neq j}} = \beta_x$. We let $\beta_x = \lambda\beta$ and we assume $\lambda \ll 1$ to reflect the fact that inter-patch transmission will typically be much less frequent than intra-patch transmission. We define random variable T_f as the time at which the last infected individual recovers.

The three quantities of interest are: 1) $p = \mathbb{P}\{T_{11} > T_{21} \mid T_{12} < T_f \vee T_{11} < T_f\}$, the probability that vaccine escape in Patch 1 comes not from Patch 1 but from neighboring unvaccinated Patch 2, conditioned on vaccine escape emerging in Patch 1 from one of the two patches, 2) $f = \mathbb{P}\{T_{21} < T_f \vee T_{11} < T_f\} / \mathbb{P}\{T_{11} < T_f\}$, the factor by which the probability of vaccine escape in Patch 1 is increased by having neighboring unvaccinated Patch 2, and 3) $\varepsilon = \mathbb{P}\{T_{11} < T_f \vee T_{12} < T_f \vee T_{21} < T_f \vee T_{22} < T_f\}$, the total probability of vaccine escape in the two patches as a function of vaccine distribution between the two patches. These quantities are rather immediate functions of the $r_{ij}(t)$; they are derived in the SM and plotted in Figs 1 and 2.

A striking feature of Fig 1 is how large the effect of an unvaccinated neighboring region can be: the probability of vaccine escape can be orders of magnitude higher with an unvaccinated neighboring region than without it. At the time of writing this paper, an area of the world that approximates our model in a rather extreme

corner of parameter space – in the context of the ongoing SARS-CoV-2 pandemic – is Israel and its neighbors: Israel presently has the highest vaccine coverage in the world [24] while its neighbors have among the lowest. Our findings would recommend vigilance for vaccine escape in this and many other areas of the world that have significant disparity in vaccine distribution both between and within countries. Figure 2 explicitly shows the effect of vaccine disparity between the two patches. Equal vaccination between the two patches gives the lowest probability of vaccine escape. Curiously, for medium to high reproductive numbers and low mutation rates, moderate disparities in vaccination can promote the emergence of vaccine escape more strongly than extreme (all or nothing) disparities.

Our parameter U is a composite parameter: it is the rate at which the transmission chain among hosts finally leads to one host infecting another host with a vaccine escape mutant, which we refer to above as an escape-infection event. As such, this parameter incorporates the mutation rate of the virus as well as any effect on within-host fitness it may have: a decrease (increase) in within-host fitness will effectively decrease (increase) U . Within-host fitness of SARS-CoV-2 should not be affected by humoral immunity of the host because transmissibility of SARS-CoV-2 peaks around the time of onset of symptoms [15], whereas a robust antibody response is not mounted until roughly ten days after the onset of symptoms [15, 29]. Thus any effect that escape mutations have on within-host fitness will not be antigenic in nature and will thus expose any pleiotropic fitness effects of vaccine escape. In this light, U may be viewed as primarily encapsulating two factors: viral mutation rate, and any pleiotropic fitness effect vaccine escape may have.

The parameters of our model most readily affected by public policy are λ , β and ϕ_j : λ can be reduced, for example by closing borders or otherwise limiting the movement of people, β can be reduced by facemasks and social distancing, and ϕ_j can be made more uniform by equitable vaccine distribution. As vaccines become increasingly available, restrictions on movement and contact are slowly being lifted in parallel, increasing both λ and β . The relaxation of these restrictions may have adverse effects if done too quickly [2, 26, 27], and may further increase the probability of vaccine escape.

Our findings provide a solid theoretical basis to support arguments for vaccine equity. Geographic regions of the human population that are not being vaccinated can serve as evolutionary reservoirs from which vaccine escape mutations may arise and give rise to renewed and unfettered spread of SARS-CoV-2. It may be that vaccine escape is inevitable and that SARS-CoV-2 will eventually become endemic [4, 23], in which case vaccine updating and exploration of new antigenic targets [10] will become the norm. Or we may have a window of opportunity now to prevent that outcome, in which case present vaccine rollout strategies could

make the difference. Vaccine updating, optimizing deployment of the many different available vaccines [21] and, as we have shown, vaccine equity, are key ingredients of these strategies.

Acknowledgements

The authors thank Matthew Fricke, the Center for Advanced Research Computing, and the Biology Department at the University of New Mexico for critical infrastructure and computational support. P.G. and A.C. received financial support from the USA/Brazil Fulbright scholar program. J.V. and F.S. received financial support from grants DGAPA-PAPIIT UNAM IV100220 and DGAPA-PAPIIT IN115720 UNAM.

References

- [1] S. S. Abdool Karim and T. de Oliveira. “New SARS-CoV-2 Variants - Clinical, Public Health, and Vaccine Implications”. *N. Engl. J. Med.* (Mar. 2021).
- [2] M. A. Acuña-Zegarra, M. Santana-Cibrian, and J. X. Velasco-Hernandez. “Modeling behavioral change and COVID-19 containment in Mexico: A trade-off between lockdown and compliance”. *Math. Biosci.* 325 (July 2020), p. 108370.
- [3] J. S. Ayres and D. S. Schneider. “Tolerance of infections”. *Annu. Rev. Immunol.* 30 (Jan. 2012), pp. 271–294.
- [4] L. Berman and M. Riedel. “Coping with COVID”. *Vaccine* 39 (2021), pp. 777–778.
- [5] F. Blanquart, S. Lehtinen, M. Lipsitch, and C. Fraser. “The evolution of antibiotic resistance in a structured host population”. *J. R. Soc. Interface* 15 (June 2018).
- [6] D. A. Collier, A. De Marco, I. A. T. M. Ferreira, et al. “SARS-CoV-2 B.1.1.7 sensitivity to mRNA vaccine-elicited, convalescent and monoclonal antibodies”. *medRxiv* (Feb. 2021).
- [7] J. W. Costerton, P. S. Stewart, and E. P. Greenberg. “Bacterial biofilms: a common cause of persistent infections”. *Science* 284 (May 1999), pp. 1318–1322.
- [8] R. Eguia, K. H. D. Crawford, T. Stevens-Ayers, et al. “A human coronavirus evolves antigenically to escape antibody immunity”. *bioRxiv.org* (Mar. 2021).
- [9] N. R. Faria, T. A. Mellan, C. Whittaker, et al. “Genomics and epidemiology of a novel SARS-CoV-2 lineage in Manaus, Brazil”. *medRxiv* (Mar. 2021).
- [10] A. Fernández. “Toward the next-generation COVID-19 vaccines that circumvent antigenic drift while defusing viral infection”. *ACS Pharmacol. Transl. Sci.* (Mar. 2021).
- [11] F. Fu, M. A. Nowak, and S. Bonhoeffer. “Spatial heterogeneity in drug concentrations can facilitate the emergence of resistance to cancer therapy”. *PLoS Comput. Biol.* 11 (Mar. 2015), e1004142.
- [12] W. F. Garcia-Beltran, E. C. Lam, K. St. Denis, et al. “Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral immunity”. *Cell* (Mar. 2021).
- [13] J. R. Gog, E. M. Hill, L. Danon, and R. Thompson. “Vaccine escape in a heterogeneous population: insights for SARS-CoV-2 from a simple model”. *medRxiv* (Mar. 2021).

- [14] A. J. Greaney, A. N. Loes, K. H. D. Crawford, et al. “Comprehensive mapping of mutations to the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human serum antibodies”. *bioRxiv* (Jan. 2021), p. 2020.12.31.425021.
- [15] X. He, E. H. Y. Lau, P. Wu, et al. “Temporal dynamics in viral shedding and transmissibility of COVID-19”. *Nat. Med.* 26 (May 2020), pp. 672–675.
- [16] M. Hoffmann, P. Arora, R. Groß, et al. “SARS-CoV-2 variants B.1.351 and B.1.1.248: Escape from therapeutic antibodies and antibodies induced by infection and vaccination”. *bioRxiv* (Feb. 2021), p. 2021.02.11.430787.
- [17] T. B. Kepler and A. S. Perelson. “Drug concentration heterogeneity facilitates the evolution of drug resistance”. *Proc. Natl. Acad. Sci. U. S. A.* 95 (Sept. 1998), pp. 11514–11519.
- [18] M. S. Krieger, C. E. Denison, T. L. Anderson, et al. “Population structure across scales facilitates coexistence and spatial heterogeneity of antibiotic-resistant infections”. *PLoS Comput. Biol.* 16 (July 2020), e1008010.
- [19] A. Licht and G. Alter. “A Drug-Free Zone—Lymph Nodes as a Safe Haven for HIV”. *Cell Host Microbe* 19 (Mar. 2016), pp. 275–276.
- [20] K. D. McCormick, J. L. Jacobs, and J. W. Mellors. “The emerging plasticity of SARS-CoV-2”. *Science* 371 (Mar. 2021), pp. 1306–1308.
- [21] J. P. Moore. “Approaches for Optimal Use of Different COVID-19 Vaccines: Issues of Viral Variants and Vaccine Efficacy”. *JAMA* (Mar. 2021), E1–2.
- [22] J. P. Moore and P. A. Offit. “SARS-CoV-2 Vaccines and the Growing Threat of Viral Variants”. *JAMA* 325 (Mar. 2021), pp. 821–822.
- [23] N. Phillips. “The coronavirus is here to stay - here’s what that means”. *Nature* 590 (Feb. 2021), pp. 382–384.
- [24] B. Rosen, R. Waitzberg, and A. Israeli. “Israel’s rapid rollout of vaccinations for COVID-19”. *Isr. J. Health Policy Res.* 10 (Jan. 2021), p. 6.
- [25] M. Roser, H. Ritchie, E. Ortiz-Ospina, and J. Hasell. *Coronavirus Pandemic (COVID-19)*. <https://ourworldindata.org/coronavirus/country/sao-tome-and-principe>. May 2020.
- [26] F. Saldaña and J. X. Velasco-Hernández. “The trade-off between mobility and vaccination for COVID-19 control: a metapopulation modeling approach”. *bioRxiv* (Mar. 2021).

- [27] M. Santana-Cibrian, M. A. Acuña-Zegarra, and J. X. Velasco-Hernandez. “Lifting mobility restrictions and the effect of superspreading events on the short-term dynamics of COVID-19”. *Math. Biosci. Eng.* 17 (2020), pp. 6240–6258.
- [28] P. S. Stewart. “Mechanisms of antibiotic resistance in bacterial biofilms”. *Int. J. Med. Microbiol.* 292 (July 2002), pp. 107–113.
- [29] K. K.-W. To, O. T.-Y. Tsang, W.-S. Leung, et al. “Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study”. *Lancet Infect. Dis.* 20 (May 2020), pp. 565–574.
- [30] J. G. Tyerman, J. M. Ponciano, P. Joyce, et al. “The evolution of antibiotic susceptibility and resistance during the formation of *Escherichia coli* biofilms in the absence of antibiotics”. *BMC Evol. Biol.* 13 (Jan. 2013), p. 22.
- [31] D. Van Egeren, A. Novokhodko, M. Stoddard, et al. “Risk of evolutionary escape from neutralizing antibodies targeting SARS-CoV-2 spike protein”. *bioRxiv* (Nov. 2020).
- [32] P. Wang, M. S. Nair, L. Liu, et al. “Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7”. *Nature* (Mar. 2021).
- [33] J. Wise. “Covid-19: The E484K mutation and the risks it poses”. *BMJ* 372 (Feb. 2021), n359.




Fig 1 | Representative plots of: p , the probability that vaccine escape emerging in vaccinated Patch 1 comes not from Patch 1 but from Patch 2 (*left column*), and f , the factor by which the probability of vaccine escape emerging in Patch 1 is increased as a consequence of having unvaccinated neighboring Patch 2 (*right column*). Horizontal axes indicate Patch 1 vaccination rate, ϕ_1 , and $\phi_2 = 0$. Parameters not specified in the plots are: $\lambda = 0.02$, $\gamma = 0.1$, $\beta = \gamma R_0$, and $N = 10^5$. Expressions for p and f are specified in the main text and derived in the SM. For these plots, we have assumed that vaccination begins at the time the epidemic begins. Departures from this assumption as well as exploration of parameter space are in the SM.




Fig 2 | Effect of vaccine disparity on probability of vaccine escape. Vaccine escape factor, ε , is plotted as a function the fraction of vaccine that goes to Patch 2. Vaccine escape factor is defined as vaccine escape probability divided by the escape probability when all vaccine goes to one of the two patches (maximum disparity). Parameters are: $\lambda = 0.05$, $\gamma = 0.1$, $\beta = \gamma R_0$, $N = 10^5$ and *left column*: $V(0) = 0$, $\phi_1 + \phi_2 = 0.05$; *middle column*: $V(0) = 0.2$, $\phi_1 + \phi_2 = 0.02$; *right column*: $V(0) = 0.6$, $\phi_1 + \phi_2 = 0.1$.