Thyroid dysfunction diagnosis from routine laboratory tests based on machine learning

Min Hu[1#], Ph.D., Chikashi Asami[1#], BS., Hiroshi Iwakura[2], M.D., Ph.D., Yasuyo Nakajima[3],

M.D., Ph.D., Ryousuke Sema[1], MBA., Tsuyoshi Kikuchi[1], MBA., Koji Sakamaki[5], M.S.,

Takumi Kudo[4], M.D., Ph.D., Masanobu Yamada[3], M.D., Ph.D., Takashi Akamizu[4], M.D., Ph.D.,

Yasubumi Sakakibara[1,6], Ph.D.

[1]Cosmic Corporation Co., Ltd., Tokyo, Japan

[2]Wakayama Medical University Hospital, Wakayama, Japan

[3]Gunma University Hospital, Gunma, Japan

[4]Kuma Hospital, Hyogo, Japan

[5]Hidaka Hospital, Gunma, Japan

[6]Keio University, Tokyo, Japan

[#]These authors contributed equally to this work.

Running title

AI-based system for thyroid dysfunction diagnosis

Keywords: misdiagnosed thyroid dysfunction; hyperthyroidism; hypothyroidism; machine

learning

Abstract

Approximately 2.4 million patients need treatment for thyroid disease, including Graves' disease

and Hashimoto's disease, in Japan. However, only 450,000 of them are receiving treatment, and

many patients with thyroid dysfunction remain largely overlooked. In this retrospective study,

we aimed to screen patients with hyperthyroidism and hypothyroidism who would greatly benefit

from prompt medical treatment, and examined routine laboratory finding data and machine

learning algorithms to investigate whether such accurate and robust screening is possible to

prevent overlooking and misdiagnosing thyroid dysfunction. We succeeded in developing a

machine learning method to construct the classification model for detecting hyperthyroidism and

hypothyroidism in patients using 11 routine laboratory tests. We collected electronic health

record and medical checkup data from four hospitals in Japan. As a result of cross-validation and

external evaluation, we achieved a high classification accuracy for the hyperthyroidism and

hypothyroidism models.

Introduction

Thyroid dysfunction is a leading endocrine disorder with major health implications, including an

increased risk of heart disease and hypercholesterolemia. One of the greatest challenges in

thyroid dysfunction treatment is to prevent overlooking and misdiagnosing these diseases.

Thyroid hormone excess and deficiency are frequently misunderstood and are too often

overlooked and misdiagnosed (1). For hyperthyroidism, the diagnosis may be delayed or missed

because some symptoms can be easily attributed to other conditions such as stress (2), and often

mistaken for cardiac disease or gastrointestinal malignancies. Hypothyroidism can present with

nonspecific constitutional and neuropsychiatric complaints (3), and patients with hypothyroidism

are often misdiagnosed as dementia, cardiac disease, liver disease, or hyperlipidemia, and hence

not given the proper treatment (4). The American Association of Clinical Endocrinologists has

estimated that in the United States, approximately 4.78% of the population has misdiagnosed

thyroid dysfunction (5). Another study argues that it can be calculated that approximately 15

million adults have unrecognized thyroid disease (6). In Japan, it is estimated that approximately

2.4 million patients need treatment for thyroid disease (7). However, only approximately 450,000

of them are receiving treatment. Thus, patients with thyroid dysfunction are frequently

overlooked and misdiagnosed (6,7).

Hyperthyroidism is the condition that occurs due to excessive production of thyroid

hormones. The first step to diagnose hyperthyroidism is to measure free thyroxine (FT4) and free

triiodothyronine (FT3) thyroid hormones and thyroid-stimulating hormone (TSH) (6). In

contrast, hypothyroidism is a condition in which serum thyroid hormones decrease. Typical

diseases of hypothyroidism include Hashimoto's disease and are diagnosed by anti-thyroid

antibody tests such as anti-thyroid peroxidase antibody (TPO) and anti-thyroglobulin antibody

(TgAb) (5). Despite their clinical significance, thyroid function tests and anti-thyroid antibody

tests were not included in the Japanese national health checkups.

As popular and effective approaches to predictive analytics, machine learning is highly

regarded due to their success in diagnosis, prediction, and choice of treatment. Recently, an

emerging technique in the field of medical informatics has employed machine learning to

accurately derive insights from medical records to support clinical screening and predict

misdiagnosed disease (8). For instance, there is a study that emphasized the superiority of

machine learning technology for predicting cardiovascular risk from routine clinical data (9). In

another study, the incidence of myocardial infarction or cerebral infarction was predicted using

the results of health checkup (10). Numerous studies have also attempted to assess the efficacy of

detecting misdiagnosed diseases, including thyroid dysfunction (11-17). Aoki et al. (16,17)

found that there were strong, multiple correlations between the set of routine clinical parameters

and FT4 in patients with both overt hyperthyroidism and overt hypothyroidism. These studies

used pattern recognition methods such as neural networks and predicted the likelihood of thyroid

dysfunction from a set of routine clinical tests.

Despite such great efforts, there are still several concerns on the machine learning

application to diagnosis of disease. Those includes the issues of data cleansing, missing value

completion, dysfunction labeling criterion , integration of multiple hospital datasets, validation

and interpretation of machine learning model. In this study, we developed an explainable

artificial intelligence diagnosis support system using machine learning algorithms to identify

thyroid dysfunction with routine clinical data to improve medical screening and prevent

overlooking and misdiagnosing thyroid dysfunction. Our study addresses those concerns on the

machine learning application and provides some possible solutions.

We devised two criteria for dysfunction labeling of data: thyroid test criterion and

prescription criterion. Thyroid test criterion, which includes the thyroid function tests TSH and

FT4, can be used to clearly model the overt and subclinical thyroid dysfunction. However, both

TSH and FT4 tests are required so that the number of available data tends to be smaller. More

data are available through prescription criterion based on the presence or absence of doctor's

prescriptions, though it can lead to a problem of confounding overt and subclinical thyroid

dysfunction with euthyroidism. Second, we integrated data from four hospitals including

electronic medical record of Wakayama Medical University hospital, Gunma University hospital,

and Kuma hospital, and annual medical checkup data of Hidaka hospital. Among the four

hospitals, a machine learning model was trained and evaluated via cross-validation by combining

patient data of Wakayama Medical University hospital and Gunma University hospital with

medical checkup data of healthy individuals in Hidaka hospital. Furthermore, electronic medical

record data of Kuma hospital was used as the external evaluation for the trained models. Third,

we examined four typical machine learning algorithms for the structured data: gradient boosting

decision tree, support vector machines and neural networks used in related studies, as well as

logistic regression, which is a common tool in medical studies. Fourth, in terms of the input

feature used in machine learning models, features including AST (aspartate aminotransferase),

ALT (alanine aminotransferase), γ-GTP, total cholesterol, hemoglobin, red blood cell count

(RBC), creatinine, and sex were selected from the health checkup test list specific in Japan.

alkaline phosphatase (ALP), uric acid (UA), and UA to serum creatinine (S-Cr) ratio were

further added, and hence totally 11 features were used. To further verify the performance of

models depending on the set of input features, we trained and evaluated models in the case

limited to five routine tests including AST, ALT, γ-GTP, total cholesterol, and sex. Finally, all

24 laboratory findings available in this study were also applied and validated.

Methods

Data source

In the present study, we acquired laboratory finding datasets from different clinical university

medical institutions in Japan, including Wakayama Medical University Hospital, Gunma

University Hospital, Hidaka Hospital, and Kuma Hospital. The anonymized electronic medical

records include age, sex, diagnosis codes for insurance billing, prescribed drugs, and biochemical

test results. The institutional ethical review boards of the three institutions at which the study was

conducted gave their approval.

A sample of 176,727 subjects in total were included in our study, aged between 13 and 88

and from different regions in Japan between 2004 and 2019, as illustrated in Table 1. Among the

four institutions, Wakayama Medical University hospital and Gunma University hospital are

hospitals affiliated with a medical college, Hidaka hospital is a regional medical care support

hospital, and Kuma hospital is a hospital specialized on thyroid diseases. The data of the

176,727 subjects consisted of doctor evaluations, prescriptions, clinical examinations, and

laboratory findings. The doctor evaluations addressed medical history, medication use, and

differential diagnosis, among other topics. If a subject was prescribed medication, the name and

dose of the prescription were recorded. The examinations involved anthropometric

measurements and laboratory tests, among others. The institutional ethical review boards of the

three institutions at which the study was conducted gave their approval (Approval Number of

Wakayama Medical University Hospital: 2301, Hidaka Hospital: 257, Gunma University

Hospital : HS2018-245)

<center>Table 1. Summary of the data from each institution</center>

Institution	Wakayama Medical University	Gunma University	Hidaka Hospital	Kuma Hospital
Number of prescriptions	8,249,286	34,561,268	23,450	61,590
Number of patients	14,249	27,133	10,482	124,863
Average age	60.9	51.7	47.7	50.3
Male/female ratio	1.03 (5,888/5,723)	0.53 (8,143/15,296)	1.82 (15,125/8,325)	0.21
Data period	2010-2018	2004-2019	2004-2007	2007-2020

The K-nearest neighbor (KNN) algorithm was used to predict and complement the

missing values, with k set to 3 in the data filling process. A previous study (11) has reported

KNN to substantially increase the number of applicable subjects. Compared with missing value

deletion, it is easily applied, performs well for nonparametric datasets and provides a larger

sample size. Furthermore, since the age and sex distributions were different among the

institutions, as shown in Table 1, we also conducted random under sampling to fix the gaps in

these differences. From this dataset, the model was constructed using the thyroid patient data

from Wakayama Medical University and Gunma University and the data of control groups from

Hidaka hospital, and was evaluated using cross-validation. To validate on external data, the

model was also evaluated on the dataset of Kuma hospital.

141
Construction of machine learning model

As shown in Table 2, four verification items were devised in this study to improve the

performance of our machine learning model. The criteria of data labeling and the combination of

multiple institutions were evaluated at first. Then four different machine learning algorithms and

three sets of input features were evaluated to achieve the best performance of our thyroid

dysfunction classification models.

148   Table 2. List of verification items

No.	Verification item	Option			
1	Training data labeling	Thyroid function test criterion	Prescription criterion		
2	Institution combination (for patient data and control group data)	Institution combination 1 (Inst. comb. 1)	Institution combination 2 (Inst. comb. 2)	Institution combination 3 (Inst. comb. 3)	External
3	Machine learning algorithm	GBDT	SVM	Logistic regression	ANN
4	Input features	Feature set 1	Feature set 2	Feature set 3	

149
150

Data labeling criterion

According to the guidelines of Japanese Society of Laboratory Medicine for the diagnosis of

hyperthyroidism and hypothyroidism, if the disorder is suspected from the clinical findings, first

the thyroid function test (TSH and FT4 measurement) is conducted, from which the disorder is

classified into three categories, hyperthyroidism, hypothyroidism, and euthyroidism (5).

Therefore, we devised and compared the performance of two data labeling criteria.

157   We firstly devised the labeling criterion by using the result of the thyroid function test as

a reference (hereinafter referred to as the "thyroid function test criterion"). Specifically, in the

dataset of Wakayama Medical University, FT4 and TSH were measured with the ECLusys kits.

TSH < 0.5 and FT4 > 1.7 was defined as overt hyperthyroidism, TSH < 0.5 and $0.9 \leq FT4 \leq 1.7$

as subclinical hyperthyroidism, TSH > 5.0 and FT4<0.9 as overt hypothyroidism, and TSH > 5.0

and $0.9 \leq FT4 \leq 1.7$ as subclinical hypothyroidism (TSH unit: μIU/mL; FT4 unit: ng/dL). In the

dataset of Gunma University, in which FT4 and TSH were measured with the Architect kit, TSH

< 0.35 and FT4 > 1.48 was defined as overt hyperthyroidism, TSH < 0.35 and $0.7 \leq FT4 \leq 1.48$

as subclinical hyperthyroidism, TSH > 4.94 and FT4 < 0.7 as overt hypothyroidism, and TSH >

4.94 and $0.7 \leq FT4 \leq 1.48$ as subclinical hypothyroidism. In this study, overt and subclinical

hyperthyroid patients are collectively referred to as the hyperthyroidism group, and overt and

subclinical hypothyroid patients are collectively referred to as hypothyroidism group.

Data for the control group were extracted from the third institution, Hidaka hospital,

which consisted of the test results from regular medical examinations. We extracted

comprehensive medical examination data for subject who did not have any symptoms suggesting

thyroid dysfunction or abnormal values in the laboratory tests of thyroid-stimulating hormone

(TSH) and serum free T4 (fT4) The normal ranges were set to 0.34-3.88 μIU/mL for TSH and

0.95-1.74 ng/dL for fT4. Random under sampling was conducted for the control group in such a

way that the sample size of the control group was equivalent to the size of the hyperthyroidism

and hypothyroidism groups. The thyroid function test criterion required both TSH and FT4 test

results, but a smaller number of patient records tended to have both of these levels. Therefore, as

an alternative solution, we devised another criterion of labeling the training data according to the

presence of prescription (hereinafter referred to as the "prescription criterion") for thyroid

disorder. Specifically, the procedure of prescription criterion satisfies the following conditions:

(a) it includes patient records with standard prescribed medications for thyroid dysfunction

(including thiamazole, propylthiouracil, and potassium iodide for the hyperthyroidism group, and

levothyroxin and thyronamine for the hypothyroidism group) obtained on the patient's first

visits, (b) the patient is not diagnosed with thyroid nodules, (c) patient records contain laboratory

findings obtained within four weeks after the patient's first prescription, and (d) exclude records

with missing values of more than half of our selected features. Since the age distributions were

different among the institutions, as shown in Table 1, we also conducted data under sampling to

fix the gaps in these differences.

In machine learning, a control group is generally used as negative label. Since

hyperthyroidism and hypothyroidism are thyroid dysfunction, both often express similar

symptoms and effects on some routine laboratory findings (e.g. Hb is decreased in both

hyperthyroidism and hypothyroidism patients). Therefore, we consider the confounding of

hyperthyroidism and hyperthyroidism as "crosstalk" and refined the labeling criteria in such a

way that the negative label is set as both the healthy subjects of the control group and the patients

of the opposite type of thyroid dysfunction. For instance, in the data labeling process of the

hyperthyroidism classification model, hyperthyroidism group was set as positive label whereas

both healthy subjects of the control group and hypothyroidism patients were set as negative

label.

Integrating multiple hospital datasets

The demographics were different among the three institutions from different districts. To

investigate the effect of integrating three hospital datasets, we explored three combinations of the

datasets to increase the generalization ability of our models. Specifically, three options on

datasets, namely, thyroid dysfunction group data from both Wakayama Medical University and

Gunma University and control group data from Hidaka hospital (referred to as Inst. comb. 1),

thyroid dysfunction group data from Wakayama Medical University and control group data from

Hidaka hospital (referred to as Inst. comb. 2), and thyroid dysfunction group data from Gunma

University and control group data from Hidaka hospital (referred to as Inst. comb. 3), were set to

train and evaluate the models.

Machine learning algorithms

Four representative machine learning algorithms were applied and evaluated of the performance

on thyroid dysfunction classification:

214   Gradient boosting decision tree (GBDT), as proposed by Friedman (18), produces a

prediction model in the form of an ensemble of weak prediction models, typically decision trees.

It is based on a machine learning technique that consists of an "ensemble" family of algorithms,

creates multiple models (called weak learners), and combines them to increase the prediction

accuracy. The main idea of this technique is to build a set of decision trees and use them to

classify a new case. Each decision tree is generated using randomly selected variable subsets

from all feature variables and a randomly selected subset of data combined by bootstrapping

(19). In this study, we employed the most accurate algorithm, called CATBoost (20), in the

GBDT family.

223   The artificial neural network (ANN) is a well-established classification technique that is

widely used in pattern recognition studies. In general, an ANN consists of 3 layers: an input layer

that receives information, a hidden layer that processes information, and an output layer that

calculates the results (21). In the present study, a standard feed-forward ANN was applied due to

its relative simplicity and stability.

Support vector machine (SVM) is a supervised machine learning technique that is widely

used in pattern recognition and classification problems (22). In the approach of this method, each

data sample is a vector whose dimensions are equal to the number of features to be considered,

and the SVM creates a hyperplane that separates samples into two categories. The induced

hyperplane is constructed to maximize its distance from the samples of both classes. This

algorithm achieves high classification performance by using special nonlinear functions called

kernels to transform the input space into a multidimensional space (22). In this study, the radial

basis function kernel is used.

Logistic regression is a statistical classifier that provides the probability for predicting the

labeled class of categorical type by using a number of attributes. Logistic regression is frequently

used to examine the risk relationship between disease and exposure, with the ability to test for

statistical interaction and control for multi-variable confiding (23). It is a linear model and used

as the baseline model for the performance comparison,

241
Explanatory features (variables) for machine learning

Features from a subject's record were designed to sufficiently explain factors that were related to

thyroid dysfunction. We used 11 variables as explanatory variables in this study as the first

experimented set of features (referred to as Feature set 1) in this study, of which eight tests are

tests measured in routine health checkup: sex, AST, ALT, γ-GTP, total cholesterol, Hemoglobin

(Hb), RBC, and creatinine (S-Cr). In addition, since ALP, UA, and S-Cr ratio are reported to be

highly relevant to thyroid dysfunction (24, 25), these were added to the above items. We also

included UA/S-Cr ratio in this study considering that the reduction of S-Cr has been reported in

hyperthyroidism, while UA has not been confirmed to fluctuate with thyroid dysfunction. To

discriminate hyperthyroidism with renal dysfunction, which usually leads to the rise of both S-Cr

and UA, we introduced UA/S-Cr ratio as one of the features to improve the classification

performance. 11 tests (Feature set 1) in total were used as features to train machine learning

models in this study.

With an aim to quantify the necessity of each of the 11 tests mentioned above, the

performance of five items (referred to as Feature set 2) out of the 11 tests was checked. Feature

set 2 excluded three items, Hb, S-Cr, and RBC, which are the tests measured only at the doctor's

discretion.

Model validation

Cross-validation was applied to evaluate the performance of our machine learning method in

classifying patients. The evaluation was conducted by extracting 9/10 training data and 1/10 test

data by conducting 10-fold cross-validation. This was repeated 10 times to extract the training

and test data uniformly, and the average and standard deviation of each evaluation score of each

time were calculated. During the model training and test process, we avoided including the same

subject to both training dataset and test dataset. The following measures were used for the

performance evaluation criteria: area under the receiver operating characteristic curve (AUROC),

area under the precision-recall curve (AUPRC), sensitivity is defined by TP/(TP+FN), and

specificity is defined by TN/(TN+FP), where TP is the number of true positives, TN is the

number of true negatives, FP is the number of false positives, FN is the number of false

negatives. Note that the cutoff value for classifying as positive or negative is determined by

Youden index (26). Finally, the AUROC performance difference between models was verified as

statistically significant by the Wilcoxon signed-rank test.

In addition, the data of Kuma hospital were employed as an external validation. The

model was constructed using the hyperthyroidism group and the hypothyroidism group of

Wakayama Medical University and Gunma University and the control group of Hidaka hospital

as the training data. The model was evaluated using the hyperthyroidism group and

hypothyroidism group of Kuma hospital and the control group of Hidaka hospital (referred to as

External).

Classification of subclinical thyroid dysfunction

In the guideline of Japan Thyroid Association (27), subclinical hypothyroidism is defined as

when FT4 is within the normal limit but the TSH measured is higher than normal .On the other

hand subclinical hyperthyroidism is defined as when FT4 is within normal limit and TSH is

lower than normal. Compared to the overt thyroid dysfunction where both TSH and FT4 are out

of the standard ranges, it is difficult to classify subclinical thyroid dysfunction. This study

evaluated the classification performance of the machine learning model by using subclinical

standards in the thyroid function test criterion labeling method. We further extended the feature

set in the attempt of improving model performance and selected 24 tests (referred to as Feature

set 3), which was the all the laboratory tests available in this study [1].

Feature importance

---

[1] Feature set 3 includes sex, AST, ALT, γ-GTP, Total cholesterol, RBC, hemoglobin, uric acid, S-Cr, uric acid/S-Cr ratio, ALP, albumin-globulin ratio, albumin, blood urea nitrogen, C-reactive protein, hematocrit, lactate dehydrogenase, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, mean corpuscular volume, platelet count, total bilirubin, total protein, white blood count.

To further understand how each feature contributes to the classification of patients in our model,

we introduced feature importance. Feature importance represents the factor by which the model

error is increased compared to the original model error. In the decision tree–based machine

learning algorithms, including GBDT, impurities and the features at which the node is split are

recorded for all the nodes when the decision tree learning is finished, and the decision tree

calculates the features importance using this information (19).

Results

Model validation

Table 3 is a summary of the performance results f of the machine learning model constructed in

this study. As the result of 10-fold cross-validation, as shown in No. I of Table 3, the best

classification model for overt hyperthyroidism achieved an accuracy of AUROC = 92.4%,

sensitivity = 83.3%, and specificity = 90.9%. The best classification model for overt

hypothyroidism achieved an accuracy of AUROC = 90.5%, sensitivity = 84.4%, and specificity

= 86.4%. In the external evaluation, as shown in No. IX of Table 3, the classification model for

overt hyperthyroidism achieved an accuracy of AUROC = 96.3%, and the classification model

for overt hypothyroidism achieved an accuracy of AUROC = 92.9%. As shown in No. XI of

Table 3, the classification model for subclinical hyperthyroidism achieved an accuracy of

AUROC = 73.8%, and the classification model for subclinical hypothyroidism achieved an

accuracy of AUROC = 75.2%.

313   The result of comparing different labeling criteria is shown in No. I and II of Table 3.

314   When the prescription criterion was applied as the labeling criterion, the accuracy of the

315   hyperthyroidism classification model achieved AUROC = 88.2%, and that of the hypothyroidism

316   classification model achieved AUROC = 82.4%. On the other hand, as shown in No. I, when the

317 thyroid function test criterion was used, the accuracy of the hyperthyroidism classification model

318 achieved AUROC = 92.4%, and that of the hypothyroidism classification model achieved

319 AUROC = 90.5%. The model trained on the thyroid function test criterion data achieved a

320 superior performance, which was statistically significant by the Wilcoxon test at p-value 0.05.

321 The result of comparing models built on different institution combinations is shown in No. I, III,

322 and IV of Table 3, as the highest performance was obtained when institution combination 1 was

323 used as training set, and the accuracy of the hyperthyroidism classification model achieved

324 AUROC = 92.4%, while and that of the hypothyroidism classification model achieved AUROC

325 = 90.5%.

326   Among the four machine learning algorithms used in this study, including GBDT, SVM,

327 logistic regression, and ANN, the highest performance was obtained when the GBDT method

328 was applied as shown in No. I, V, VI, and VII of Table 3. The accuracy of the hyperthyroidism

329 classification model achieved AUROC = 92.4%, while that of the hypothyroidism classification

330 model achieved AUROC = 90.5%, which were statistically significant at p-value 0.05 by the

331 Wilcoxon test. After comparing the performance of different feature sets, as shown in I and VIII

332 of Table 3, when the feature set 3 was applied, the accuracy of the hyperthyroidism classification

333 model was reduced to AUROC = 87.4%, and the performance of the hypothyroidism

334 classification model was reduced to AUROC = 85.5%, which shows significant differences by

335 the Wilcoxon test at p-value 0.05.

Table 3. Results of validation on different models

No.	I	II	III	IV	V	VI	VII	VIII	IX	X	XI
Training — Data labeling	Thyroid function test criterion	Prescription criterion	Thyroid function test criterion								
Training — Institution combination	Inst. comb. 1	Inst. comb. 1	Inst. comb. 2	Inst. comb. 3	Inst. comb. 1				Inst. comb. 1		
Training — Machine learning algorithm	GBDT		GBDT		SVM	Logistics regression	ANN	GBDT			
Training — Input features	Feature set 1							Feature set 2	Feature set 1		Feature set 3
Validation — Institution combination	Inst. comb. 1								External	Inst. comb. 1	
Validation — Over/Subclinical	Overt									Subclinical	
Validation — Labeling criteria	Thyroid function test criterion										
Hyperthyroidism AUROC	92.4±3.2%	88.2±3.2%	91.4±3.0%	90.7±4.1%	85.7±5.2%	86.1±4.6%	86.7±4.0%	87.4±4.2%	96.3±0.3%	73.0±5.7%	73.8±6.3%
Hyperthyroidism AUPRC	88.5±5.2%	82.8±4.9%	87.4±3.3%	85.7±6.9%	80.7±5.5%	79.8±4.5%	81.5±4.5%	80.0±7.3%	99.1±0.1%	56.6±8.3%	57.6±7.9%
Hyperthyroidism Sensitivity	83.3±9.0%	77.5±8.5%	81.9±9.2%	88.4±7.8%	70.8±11.9%	76.4±11.4%	73.5±9.3%	82.3±6.0%	87.7±1.8%	71.7±12.4%	78.7±11.2%
Hyperthyroidism Specificity	90.9±6.8%	86.4±6.2%	89.8±5.0%	80.5±12.5%	91.4±4.6%	86.0±5.8%	87.8±5.6%	83.9±7.7%	93.5±1.1%	66.6±14.0%	61.6±16.3%
Hypothyroidism AUROC	90.5±3.6%	82.4±5.3%	85.9±4.9%	86.5±3.5%	77.9±4.4%	81.4±5.7%	79.8±5.3%	85.5±3.1%	92.9±0.7%	69.1±4.1%	75.2±3.3%
Hypothyroidism AUPRC	84.1±5.7%	71.3±9.9%	77.1±7.5%	79.0±5.8%	69.5±6.8%	71.3±6.8%	70.3±7.3%	78.2±4.9%	75.7±2.7%	51.3±4.2%	59.9±7.1%
Hypothyroidism Sensitivity	84.4±9.6%	77.4±8.9%	83.6±4.6%	80.1±10.1%	66.9±14.2%	79.5±8.8%	73.5±8.8%	76.0±7.2%	87.1±3.3%	68.2±15.8%	77.7±10.4%
Hypothyroidism Specificity	86.4±5.2%	77.0±11.7%	78.3±8.8%	81.0±9.2%	78.4±14.9%	73.1±10.7%	77.1±8.4%	82.9±9.0%	84.2±3.7%	64.3±16.4%	64.3±13.6%

The mean and standard deviation for the 10 folds are shown in each score.

The model with the best performance was evaluated using the external dataset for Kuma

Hospital, as shown in No. IX of Table 3. High classification performance was achieved using the

external data: AUROC = 96.3%, sensitivity = 87.7%, and specificity = 93.5% for the

hyperthyroidism classification model and AUROC = 92.9%, sensitivity = 75.7%, and specificity

= 87.1% for the hypothyroidism classification model. No. X and XI of Table 3 show that using

feature set 3 improved the classification performance: for subclinical thyroid dysfunction,

AUROC = 73.8%, sensitivity = 78.7%, and specificity =61.6%; for hypothyroidism, AUROC =

75.2%, sensitivity = 59.9%, and specificity = 77.7%. In particular, the significance of the

hypothyroidism classification models was statistically confirmed by the Wilcoxon test at p-value

0.05.

Feature Importance

The features importance of each model was examined using the feature set 1. The left picture of

Figure 1 shows the features importance of the overt hyperthyroidism classification model and the

overt hypothyroidism classification model. The three most important features in the overt

hyperthyroidism model were ALP, UA/S-Cr ratio, and total cholesterol. The three most

important features in the overt hypothyroidism model were AST, total cholesterol, and RBC. On

the other hand, the right picture of Figure 1 shows the features importance of the subclinical

hyperthyroidism classification model and the subclinical hypothyroidism classification model.

The three most important features in the subclinical hyperthyroidism model were ALP, UA/S-Cr

ratio, and S-Cr, and the three most important features in the subclinical hypothyroidism model

were total cholesterol, AST, and RBC. For both overt and subclinical disease, ALP and S-Cr

were the top related features in the hyperthyroidism classification model, and total cholesterol,

AST, and RBC were the top features in the hypothyroidism classification model.

Furthermore, the features importance in the subclinical hyperthyroidism and subclinical

hypothyroidism classification models using the feature set 3 was conducted. As shown in Figure

2, ALP and the UA/S-Cr ratio were among the three most important features in the subclinical

hyperthyroidism classification model when the feature set 1 was used, as well as when the

feature set 3 was used. If the five most important features were considered, MCV and MCH, two

features added to the feature set 3, were included. These findings suggest that these two features

are also likely to be effective in hyperthyroidism classification. On the other hand, as shown on

right side of Figure 2, a difference was seen in the subclinical hypothyroidism classification

model when the feature set 1 was used vs. when the feature set 3 was used. The three most

important features in the model that used the feature set 1 were total cholesterol, AST, and RBC,

whereas the three most important features in the model that used the feature set 3 were total

protein, total cholesterol, AST, and the UA/S-Cr ratio. These findings suggest that total protein is

likely to be effective in classifying subclinical hypothyroidism.

Figure 1.   Comparison of feature importance between overt and subclinical thyroid dysfunction classification models

Figure 2.   Comparison of feature importance between models built on the Feature set 1 and Feature set 3

Discussion
Feature importance

The correlation of routine laboratory tests such as ALP, S-Cr, UA, and RBC, etc. with thyroid

dysfunction has been pointed out in many previous studies. According to studies on the

relationship between thyroid dysfunction and liver function (28, 29), a correlation was confirmed

between the increase in ALP and hyperthyroidism, as the ALP value was significantly higher

when bone metabolism increases in Graves's disease, which is a typical disorder of

hyperthyroidism. Sönmez (30) examined data from 433 patients and reported that S-Cr in the

hyperthyroidism group was significantly lower than in the euthyroid group. TSH and S-Cr were

also reported to have a significantly negative correlation with overt hypothyroidism (31).

Dorgalaleh (32) suggested that thyroid dysfunction directly affects most of the blood values,

including RBC, and health professionals must pay attention to such effects. The correlation

between hypothyroidism and hyperuricemia has also been confirmed by in multiple studies (33,

34).

Comparison with related studies

Several previous studies revealed promising results from the use of machine learning approaches

for predicting thyroid dysfunction (16, 17).

Similar to the present study, Aoki's (17) study used pattern recognition methods such as

neural networks to predict the likelihood of thyroid dysfunction from a set of routine test

parameters such as ALP, S-Cr, and TC. Their results suggested that most patients with overt

thyroid dysfunction could be screened by using a set of routine clinical data without measuring

thyroid hormone levels. The correct rate of 91.3% was reported in the hyperthyroidism

classification model, and the correct rate of 90.0% was reported in the hypothyroidism

classification model. Their results suggested that there is a high correlation between a set of

routine laboratory tests and thyroid dysfunction. However, the model verification of these studies

used the leave-one-out method instead of cross-validation and used the correct rate as the

indicator instead of AUROC. Thus, the model evaluation was considered insufficient.

414   Unlike the present study, one drawback of these previous studies is that those have not

considered crosstalk in the data labeling process. For hyperthyroidism classification in this study,

the hyperthyroidism group was used as a positive label, and both the control and hypothyroidism

groups were negatively labeled. For the hypothyroidism classification in this study, the

hypothyroidism group was used as a positive label, whereas both the control and

hyperthyroidism groups were negatively labeled (referred to as "crosstalk on"). On the other

hand, related studies (16, 17) performed classification by setting thyroid dysfunction patients

(with hyperthyroidism or hypothyroidism) as positive label and only control group as negative

label (referred to as "crosstalk off"). Therefore, we evaluated the performance of the models with

similar settings as these studies. As shown in A-1 column of Table 4, when only control group

was labeled negative in both the training data and validation data, a high classification

performance of AUROC = 94.9% and AUROC = 91.3% was achieved in the classification of

overt hyperthyroidism and overt hypothyroidism, respectively. However, as shown in A-2

column of Table 4, when both control group and hypothyroidism group were labeled negative in

the validation data of overt hyperthyroidism and when both control group and hyperthyroidism

group were labeled negative in the validation data of overt hypothyroidism, the classification

performance was reduced to AUROC = 78.5% and AUROC = 68.1%, respectively. The

classification performance dropped significantly in the models in which crosstalk was not

considered during the negative labeling process.

434   Table 4. Evaluation result obtained without considering crosstalk

No.		A-1	A-2
Training	Thyroid function test criterion	Overt + subclinical	
	Negative label setting	Crosstalk off	
Validation	Thyroid function test criterion	Overt	
	Negative label setting	Crosstalk off	Crosstalk on
Hyperthyroidism	AUROC	94.9±2.4%	78.5±3.1%
Hypothyroidism	AUROC	91.3±4.0%	68.1±3.1%

The mean and standard deviation for the 10 folds are shown in AUROC scores.

Limitations

In the current study, subjects under medication may be included in the data extraction process of

this study. Though we extracted only the laboratory tests at each subject's first visit to avoid

including the influence of thyroid dysfunction treatment, some subjects might be already on

medication before being referred to the hospitals in our study. These subjects on medications

may have an unexpected impact on the models we built in this study.

Another limitation of this study is that the hypothyroidism classification models exhibited

lower performance than the hyperthyroidism classification models. This result is attributed to

differences in the respective serum hormones and underlying molecular mechanisms (35). The

various nonspecific symptoms of hypothyroidism may not manifest simultaneously, resulting its

subclinical rate larger than that of hyperthyroidism. In addition, patients with hypothyroidism

such as Hashimoto's thyroiditis are dependent upon long-term levothyroxine treatment, which

may affect the manifestation of routine laboratory findings.

Furthermore, in the external evaluation of this study, the subclinical classification model

showed lower overall results than the overt classification models. Among subclinical thyroid

dysfunctions, the cause of subclinical hypothyroidism is associated with chronic thyroiditis

(Hashimoto's disease), of which approximately 60-80% of cases are related thyroid

autoantibodies (36). On the other hand, the causes of subclinical thyrotoxicosis are classified into

extrinsic overdose of thyroid hormone drugs, and endogenous hyperthyroidism such as Graves'

disease (37). Most of the subclinical thyroid dysfunctions such as subclinical thyrotoxicosis and

subclinical hypothyroidism have no subjective symptoms and are usually considered to be

transient (38, 39). Performance may have been limited due to the fact that symptoms of

subclinical thyroid dysfunction are usually minor compared to overt thyroid dysfunction, and the

phenotype of subclinical thyroid dysfunction may not be reflected in the results of routine

laboratory examination.

Conclusion

This study evaluated the screening method to discriminate hyperthyroidism and hypothyroidism

from the electronic medical records or routine laboratory finding data from health checkups

using a machine learning method with an aim to prevent missed diagnosis of thyroid

dysfunction. This is a versatile new screening method that was successfully developed from a

machine learning model construction method to discriminate patients with hyperthyroidism and

hypothyroidism using 11 features. High accuracy was achieved in the discrimination of evident

hyperthyroidism or hypothyroidism, although the discrimination accuracy of subclinical

hyperthyroidism or hypothyroidism was not satisfactory, these alerts can be useful for non-

specialists for thyroid diseases.

It is expected that the quality of life of patients will improve by applying the model

developed in this study. If thyroid dysfunction is screened using our method in healthcare

facilities, including hospitals and health checkup facilities, prompt and accurate diagnostic

support can be provided from only routine laboratory tests.

References

(1)   Garmendia Madariaga A, Santos Palacios S, Guillén-Grima F, et al. The incidence and

481   prevalence of thyroid dysfunction in Europe: A meta-analysis. J Clin Endocrinol Metab.

482   2014;99:923-931.

483   (2)   Cooper DS. Hyperthyroidism. Lancet. 2003;362:459-468.

484   (3)   Roberts CG, Ladenson PW. Hypothyroidism. Lancet. 2004;363:793-803.

485   (4)   Garber JR, Cobin RH, Gharib H, et al. Clinical practice guidelines for hypothyroidism in

486   adults: Cosponsored by the American Association of Clinical Endocrinologists and the

487   American Thyroid Association. Endocr Pract. 2012;18:988-1028.

488   (5)   Cooper DS, Ridgway EC. Thoughts on prevention of thyroid disease in the United States.

489   Thyroid. 2002;12:925-929.

490   (6)   Hamada N. The frequency of thyroid diseases that should not be overlooked in general

491   outpatient settings. Jap Med J. 1995;3740:22.

492   (7)   Japanese Ministry of Health Patient Survey Database. Available at www.mhlw.go.jp.

493   Accessed February 17, 2020.

494   (8)   Ngiam KY, Khor IW. Big data and machine learning algorithms for health-care delivery.

495   Lancet Oncol. 2019;20:e262-e273.

496   (9)   Weng S.F., et al. Can machine-learning improve cardiovascular risk prediction using

497   routine clinical data?. PloS one, 2017, 12.4: e0174944.

498   (10)   Yatsuya, Hiroshi, et al. Development of a Risk Equation for the Incidence of Coronary

499   Artery Disease and Ischemic Stroke for Middle-Aged Japanese–Japan Public Health

500   Center-Based Prospective Study. Circulation Journal 80.6 (2016): 1386-1395.

501   (11)   Chung JW, Kim WJ, Choi SB, et al. Screening for pre-diabetes using support vector

502   machine model. Presented at the 36th Annual International Conference of the IEEE

503   Engineering in Medicine and Biology Society. Chicago, IL, August 26-30, 2014.

504   (12)   Soguero-Ruiz C, Mora-Jiménez I, Rojo-Alvarez JL, et al. Feature selection using Kernel

505   component analysis for early detection of anastomosis leakage. Presented at the 2nd

506   International Workshop on Pattern Recognition for Healthcare Analytics, Stockholm,

507   Sweden, August 24, 2014.

508   (13)   Kawakami J, Hoshi K, Sato W, et al. Screening of the patient with hyperthyroidism using

509   routine test data. J Tohoku Pharm Univ. 2005;52:141-148.

510   (14)   Hoshi K, Kawakami J, Sato W, et al. Assisting the diagnosis of thyroid diseases with

511   Bayesian-type and SOM-type neural networks making use of routine test data. Chem

512   Pharm Bull (Tokyo). 2006;54:1162-1169.

513   (15)   Sato W, Hoshi K, Kawakami J, et al. Assisting the diagnosis of Graves' hyperthyroidism

514   with Bayesian-type and SOM-type neural networks by making use of a set of three routine

515   tests and their correlation with free T4. Biomed Pharmacother. 2010;64:7-15.

516   (16)   Aoki S, Hoshi K, Kawakami J, et al. Assisting the diagnosis of Graves' hyperthyroidism

517   with pattern recognition methods and a set of three routine tests parameters, and their

518   correlations with free T4 levels: Extension to male patients. Biomed Pharmacother.

519   2011;65:95-104.

(17) Aoki S, Hoshi K, Kawakami J, et al. Assisting the diagnosis of overt hypothyroidism with
pattern recognition methods, making use of a set of routine tests, and their multiple
correlation with total T4. Biomed Pharmacother. 2012;66:195-205.

(18) Liang W, Luo, S, et al. Predicting hard rock pillar stability using GBDT, XGBoost, and
LightGBM algorithms. Mathematics 8.5 (2020): 765.

(19) Friedman JH. Greedy function approximation: A gradient boosting machine. Ann Stat.
2001;29:1189-1232.

(20) Prokhorenkova L, Gusev G, Vorobev A, et al. CatBoost: unbiased boosting with
categorical features. In: Advances in neural information processing systems. 2018. p. 6638-
6648.

(21) Bishop CM, Nasrabadi N. Pattern recognition and machine learning. Pattern Recognit.
2006;4:738.

(22) Cortes C, Vapnik V. Support-vector networks. Machine learning. 1995;20(3):273–97.

(23) Hosmer DW, Lemeshow S, and Sturdivant RX. Applied logistic regression. John Wiley &
Sons. 2013; Vol. 398.

(24) Hoshi K, Kawakami J, Sato W, Sato K, Sugawara A, Saito Y, et al. Assisting the diagnosis
of thyroid diseases with Bayesian-type and SOM-type neural networks making use of
routine test data. Chem Pharm Bull 2006;54:1162–9.

(25) Sato W, Hoshi K, Kawakami J, Sato K, Sugawara A, Saito Y, Yoshida K. Assisting the
diagnosis of Graves' hyperthyroidism with Bayesian-type and SOM-type neural networks
by making use of a set of three routine tests and their correlation with free T4. Biomed
Pharmacother. 2010 Jan;64(1):7-15.

(26) Youden, W. J. Index for rating diagnostic tests. Cancer. 1950 Jan;3(1):32-5.

(27) Japan Thyroid Association Guidelines 2013:

http://www.japanthyroid.jp/doctor/guideline/japanese.html ［2021．1．19］ Japanese

(28) Cooper DS, Kaplan MM, Ridgway EC, Maloof F, Daniels GH. Alkaline phosphatase

isoenzyme patterns in hyperthyroidism. Ann Intern Med. 1979;90(2):164-168.

(29) Malik R, Hodgson H. The relationship between the thyroid gland and the liver. QJM: An

International Journal of Medicine. 2002;559-569.

(30) Sönmez E, Bulur O, Ertugrul DT, Sahin K, Beyan E, Dal K. Hyperthyroidism influences

renal function. Endocrine. 2019 Jul;65(1):144-148.

(31) Saini, Vandana, et al. Correlation of creatinine with TSH levels in overt hypothyroidism—

A requirement for monitoring of renal function in hypothyroid patients?. Clinical

biochemistry 45.3 (2012): 212-214.

(32) Dorgalaleh A, Mahmoodi M, Varmaghani B, et al. Effect of thyroid dysfunctions on blood

cell count and red blood cell indice. Iran J Pediatr Hematol Oncol. 2013;3(2):73-77.

(33) Kuhlback B Creatine and creatinine metabolism in thyrotoxicosis and hypothyroidism: a

clinical study. Acta Med Scand Suppl. 1957;331:1–70.

(34) Erickson AR, Enzenauer RJ, Nordstrom DM et al. The prevalence of hypothyroidism in

gout. Am J Med. 1994;97:231–234.

(35) Okamura K, Nakashima T, Ueda K, et al. Thyroid disorders in the general population of

Hisayama Japan, with special reference to prevalence and sex differences. International

Journal of Epidemiology. 1987; 16(4), 545–549.

(36) Cooper DS,Biondi B : Subclinical thyroid disease. Lancet 2012 ; 379 : 1142-1154

(37) Biondi B, Cooper DS : The clinical significance of subclinical thyroid dysfunction. Endocr

Rev 2008 ; 29 :76-131

(38) Fatourechi V. Subclinical hypothyroidism: an update for primary care physicians. Mayo

Clinic Proceedings. Vol. 84. No. 1. Elsevier, 2009.

(39) Biondi B, et al. Endogenous subclinical hyperthyroidism affects quality of life and

cardiac morphology and function in young and middle-aged patients. The Journal of

Clinical Endocrinology & Metabolism, 2000, 85.12: 4701-4705.

List of Abbreviations

TSH: thyroid-stimulating hormone

FT3: free triiodothyronine

FT4: free triiodothyronine

GBDT: gradient boosting decision tree

ANN: artificial neural network

SVM: support vector machine

AUROC: area under the receiver operating characteristic curve

AUPRC: area under the precision-recall curve

ALP :alkaline phosphatase

UA : uric acid

S-Cr: serum creatinine

AST : glutamic aspartate transaminase

ALT: alanine aminotransferase

RBC : red blood cell count

Declarations

Data Availability

The data that support the findings of this study are available from Japan Thyroid Association but

restrictions apply to the availability of these data, which were used under license for the current

study, and so are not publicly available. Data are however available from the authors upon

reasonable request and with permission of Japan Thyroid Association.

Correspondence address

Min Hu

Koishikawa 2-7-3 Tomisaka Building, Bunkyo-ku, Tokyo, Japan

E-mail: m.hu@cosmic-jpn.co.jp

Yasubumi Sakakibara

3-14-1 Hiyoshi, Kohoku-ku, Yokohama, 223-8522, Japan

E-mail: yasu@bio.keio.ac.jp

Competing Interests

YS is a paid scientific advisory board of Cosmic corporation co., ltd. The other authors declare

no competing financial interests.

Funding statement

This research received no specific grant from any funding agency in the public, commercial, or

not-for-profit sectors.

Author Contributions

MH, CA: implemented the software, analyzed the data, and co-wrote the paper. MY, TA:

supervised the research, and collected the medical data. HI, YN, KS, TK: aided in the feature

sets designing and interpreting the results of the models, as well as collected the medical data.

RS, TM: contributed to the design of the research, and to the writing of the manuscript. YS:

designed and supervised the research, analyzed the data, and co-wrote the paper. All authors read

and approved the final manuscript.

Acknowledgements

This study would not have been possible without the exceptional support of Dr. Akira Miyauchi,

who shared insightful comments on this project and provided the opportunity to validate the

models on external dataset and improved this study in innumerable ways. Dr. Masako Akuzawa

and Dr. Yoshitaka Ando facilitated this project to accessing the dataset of control group in

Hidaka Hospital, which significantly improved the generalization performance of the thyroid

dysfunction classification models built in this project.