1

Title:

Reasons for non-attendance to cervical cancer screening and acceptability of HPV

self-sampling among Bruneian women: A cross-sectional study.

Authors:

Liling Chaw[1*], Shirley HF Lee[1], Nurul Iffah Hazwani Ja'afar[1], Edwin Lim[2], Roslin

Sharbawi[3]

Affiliations:

1) PAPRSB Institute of Health Sciences, Gadong, Brunei Darussalam.

2) Histopathology Department, RIPAS Hospital, Ministry of Health, Brunei

Darussalam

3) Community Maternal Health Service, Department of Health Services, Ministry of

Health, Bandar Seri Begawan, Brunei Darussalam

* Corresponding Author:

Email: liling.chaw@ubd.edu.bn (LC)

Running title:

23   Reasons for cervical cancer screening non-attendance and HPV self-sampling acceptability in

24   Brunei

27   **Number of words:** 410 words (abstract), 4272 words (manuscript and tables, excluding

28   references)

Abstract

Objective: Uptake for cervical cancer screening remains well below the 80% target as recommended by Brunei's National Cervical Cancer Prevention and Control plan. We conducted a pilot study to determine the reasons for non-attendance and explore their acceptance of human papillomavirus (HPV) self-sampling as an alternative to the Pap test.

Methods: A cross-sectional study was conducted at a primary healthcare center in Brunei, from January to December 2019. We recruited screening non-attendees, defined as women who were eligible for Pap test but who either never, or did not have one within the recommended screening interval of 3 years. This recruitment was done conveniently among women attending outpatient care and/or child health services at the primary healthcare center. Participants were first asked to complete a self-administered paper-based questionnaire on their reasons for screening non-attendance, and then invited for HPV self-sampling. Among those who agreed to participate in HPV self-sampling, they were asked to complete a second questionnaire on the self-sampling procedure and their samples were tested for high-risk HPV (hr-HPV). Results were analyzed using descriptive and inferential statistics.

Result: We enrolled 174 screening non-attendees, out of which 97 (55.7%) also participated in HPV self-sampling. The main reasons for not attending Pap test screening were fear of bad results (16.1%, n = 28); embarrassment (14.9%, n = 26) and lack of time due to home

commitments (10.3%, n = 18). When compared to those who agreed to participate in HPV

self-sampling, those who declined were significantly older (p = 0.002) and less likely to agree

that they are susceptible to cervical cancer (p = 0.023). They preferred to receive Pap test-

related information from healthcare workers (59.0%, n = 155), social messaging platforms

(28.7%, n = 51) and social media (26.4%, n = 47). HPV self-sampling kits were positively

received among the 97 participants, where > 90% agreed on its ease and convenience. Nine

(9.3%) tested positive for hr-HPV, out of which eight were non-16/18 HPV genotypes.

Conclusion: Our findings suggest that promoting awareness on cervical cancer, clarifying

any misconceptions of Pap test results, and highlighting that the disease is preventable and

that early detection through screening can facilitate successful treatment would help

increase screening uptake among Bruneian non-attendees. Response to HPV self-sampling

was highly positive, suggesting the possibility of implementing this strategy in the local

setting. Our high detection of non-16/18 HPV genotypes suggest high prevalence of other

hr-HPV genotypes in Brunei. Larger studies should be conducted to further validate our

findings.

Keywords: Cervical cancer screening, HPV self-sampling, women's health, Brunei

Introduction

Cervical cancer is highly preventable but still remains one of the most common cancers among women worldwide. Globally, an estimated 604,000 women were diagnosed with cervical cancer, and 342,000 women died from the disease in 2020 [1]. Cervical cancer screening has drastically reduced the incidence of invasive cervical cancer in countries that have implemented such screening programs [2], which traditionally involves the use of the Papanicolaou (Pap) test.

Brunei Darussalam (population 459,500) is a small Southeast Asian country with a predominant Muslim population and a crude birth rate of 15.3 per 1,000 population [3]. Within this region, it has one of the highest age-standardized incidence rate (ASR) for cervical cancer: 20.6 in Brunei when compared to 10.5 and 7.7 in Malaysia and Singapore, respectively [4]. Brunei has initiated an organized cervical screening program since 2009, where married or ever married Bruneian women between 20 and 65 years old were invited to attend cervical cancer screening through periodic mail invitations. Pap test is the only screening test used in the country, and currently, liquid-based cytology is being used since 2012. Women with any positive cytology result, defined as with atypical squamous cells of undetermined significance (ASC-US) or worse, are followed up with colposcopy-guided cervical biopsy to diagnose cervical intra-epithelial neoplasia (CIN). The latter refers to premalignant lesions that are mainly caused by infection with certain types of HPV [5], and can be categorized into any one of three stages (CIN1, CIN2, or CIN3) depending on the degree of dysplasia. If untreated, either CIN2 or CIN3 (collectively referred to as CIN2+) can progress to cervical cancer.

Despite this screening service being offered free of charge, the national screening coverage

rate remains low at 44% in 2018 (unpublished data). Reasons for screening non-attendance

can vary across settings [6,7], but they can be broadly categorized into two groups: practical

and organizational barriers (such as forgot to schedule an appointment, work and childcare

commitments) [6], and emotional barriers (such as feeling healthy, lack of time, discomfort

associated with gynecologic examination embarrassment, fear of smear test, previous

negative experiences and dissatisfaction with their general practitioner) [6,8–10]. As women

who do not attend screening are at increased risk of developing cervical cancer [6], it is thus

important to first understand why women chose not to attend screening in the local

context.

Human papillomavirus (HPV) DNA detection has been recommended by the World Health

Organization as the primary screening test for cervical cancer as specific high-risk HPV

subtypes (hr-HPV) are known to be a causative agent [1]. In particular, the use of HPV self-

sampling kits was suggested to increase screening uptake particularly among screening non-

attendees [11,12], due to its ease of access (where kits could be mailed to women's homes)

and also flexibility for women to perform the test by themselves. Previous studies have

shown HPV self-sampling to be highly acceptable among screening non-attendees [13,14].

In addition, HPV self-sampling results exhibit similar sensitivity and specificity compared to

those from samples taken by trained professionals [15]. Repeated HPV self-sampling and

testing were shown to increase screening uptake [11,12,16], and also resulted in at least

two-fold higher detection rate of CIN2+ when compared to the Pap test [17,18]. In the

United Kingdom and Australia, early detection through an organized screening program

using HPV testing as the primary screening test was shown to reduce cervical cancer

morbidity and mortality [19,20]. .

With Brunei's relatively high incidence of cervical cancer and low screening uptake, we

conducted a pilot study to explore the reasons behind non-attendance and to assess the

acceptability of HPV self-sampling as a possible alternative to the Pap test among non-

attendees. Study findings could be used to strategize ways to improve screening uptake and

provide preliminary evidence towards implementing HPV testing as the primary screening

test for cervical cancer in Brunei.

Methods
138

139　## Study design and data collection
140

A cross-sectional survey was conducted at the Jubli Perak Sengkurong Health Center

(JPSHC), from January to December 2019. JPSHC is a government-funded primary healthcare

center located at Brunei-Muara District, where the majority (69.3%) of the country's

population resides. This center provides primary health care services to Mukim Sengkurong,

a sub-district with about 32 000 people from various socioeconomic backgrounds.

Eligible women attending either the outpatient or child health clinic at JPHSC were

conveniently recruited by triage nurses. We defined screening non-attendees as married or

ever married women between 20 and 65 years old and have never undergone cervical

cancer screening, or did not have one within the recommended screening interval of 3

years. We excluded women who could not comprehend Brunei-Malay or English language,

were pregnant, have had total hysterectomy, or with a history of malignancies.

We implemented a two-stage recruitment procedure. In the first stage, participants were

first recruited to complete a self-administered questionnaire on the reasons for screening

non-attendance (Q1) onsite. Q1 consists of 19 questions on the participant's socio-

demographics, reasons for not getting Pap test, attitude and knowledge on cervical cancer,

and lastly, preferred sources to acquire information about Pap test. From a prepared list of

16 possible reasons for not attending the screening program, participants were also asked

to select one "Major" reason (defined as the main reason) and one or more "Minor" reason

(defined as other reasons for not attending the screening program). Responses for

questions on their attitude and knowledge on cervical cancer were recorded using a five-

point Likert scale, ranging from "Strongly Agree" to "Strongly Disagree".

In the second stage, all participants were given an envelope containing information on HPV

self-sampling and an instruction leaflet on the procedure, after completing Q1. Within the

following two weeks, they were contacted via telephone by a trained nurse, and those who

gave verbal consent were given an appointment at JPSHC. On the day, the nurse first

explained the procedure using an instructional video and answered any questions.

Participants were then given a self-sampling kit and asked to perform the procedure in the

clinic. After completion, they were asked to complete the second self-administered

questionnaire (Q2) on their acceptability of the self-sampling procedure. Q2 consists of 12

five-point Likert scale questions (ranging from "Strongly Agree" to "Strongly Disagree") on

their experiences and opinions of the self-sampling kit.

Two separate written consents were requested from participants: one for completing Q1

and another for performing self-sampling and completing Q2.

178 Questionnaires used
179

The two questionnaires used (Q1 and Q2) were bilingual self-administered paper-based

questionnaires in the two languages commonly used in Brunei (namely, Brunei-Malay and

English language). Both questionnaires were adapted from similar studies [21,22]. They

were first translated to Brunei-Malay language by native speakers, and then back-translated

to check for inconsistencies in comprehension. Both questionnaires were also pre-tested on

six eligible women to assess if the questions could be easily understood. Responses from

pre-testing were not included in the analysis.

190 Self-sample handling and laboratory testing
191

Swabs collected from the self-sampling kits were sent to an overseas laboratory (at BNH

hospital, Thailand) for hr-HPV testing. The self-sampling device used was the Evalyn® brush

from Rovers Medical Devices. Upon receipt at the testing laboratory, the dry Evalyn® brush

was suspended in SurePath medium from which a sample was obtained for the

identification of the presence of hr-HPV using the cobas HPV test (Roche, USA). Cobas HPV

tests are automated qualitative *in vitro* tests for the detection of HPV DNA in patient

specimens. The tests utilize amplification of target DNA by polymerase chain reaction (PCR)

and nucleic acid hybridization for the detection of 14 hr-HPV types (namely genotypes 16,

18, and 12 pooled hr-HPV genotypes 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66 and 68) in a

single analysis. Results obtained from this test can be categorized into four groups:

Negative, HPV-16 positive, HPV-18 positive, and positive for non-16/18 HPV genotypes.

204   Clinical management of hr-HPV positive participants
205

We adopted the cytology triage strategy for HPV self-sampling participants [23]. Participants

with hr-HPV positive results were invited for an immediate clinic-based cytology triage test.

Those found to have negative cytology results were invited for a repeat cytology triage test

after six months. Those with second negative results were returned to routine cervical

screening recall. Participants with any positive cytology result (defined as with ASC-US or

worse) were referred for colposcopic examination.

213   Statistical analysis
214

Descriptive statistics was conducted to characterize the socio-demographic characteristics

of the study population, their reasons for not attending screening, their attitudes and

preference for information access on such screening, as well as responses from Q2. Where

appropriate, Mann-Whitney and/or Fisher's exact tests were used to assess significant

differences in sociodemographic characteristics, reasons for not attending screening, and

attitudes between women who agreed and women declined to join HPV self-sampling.

Count responses for questions with the five-point Likert scale were categorized into three

categories (agree, neutral, and disagree), and those with missing values were classified as

neutral. Statistical analysis was conducted using R ver. 3.6 [24]. Ethics approval was

obtained from the Medical and Health Research and Ethics Committee (MHREC), Ministry of

Health, Brunei Darussalam [Reference no. MHREC/MOH/2018/9(2)].

Results

A total of 174 eligible women were enrolled in this study from January to December 2019,

out of which 97 (55.7%) also participated in HPV self-sampling (Fig. 1). Their median age was

45 years, ranging between 23 and 65 years (Table 1). The participants were mainly of Malay

ethnicity (92.5%, n = 161), married (90.2%, n = 157), and had ≥ 3 births (60.3%, n = 105).

About half of the participants had their last Pap test performed between 4 and 10 years ago

(52.3%, n = 91), and have never received the HPV vaccine (54.6%, n = 95). There were

significant differences between those who agreed and declined to participate in HPV self-

sampling: those who declined were significantly older (p = 0.002) and more likely to have

their last Pap test performed > 10 years ago (p = 0.031).

Fig 1. Flowchart of study participants at JPHSC, Brunei (Jan – Dec 2019).

Table 1. Sociodemographic characteristics of the study population, including comparison

between groups that agreed and declined to join HPV self-sampling. IQR = Interquartile

range.

Characteristics		Total study population (n = 174) n (%)	Joined self-sampling (n = 97) n (%)	Declined self-sampling (n = 77) n (%)	p-value
Median age in years (IQR)		45.0 (15.25)	41.0 (17)	49.0 (14.5)	**0.002***
Age group (in years)	20 – 24	3 (1.7)	3 (100)	0 (0.0)	**0.018***
	25 – 29	20 (11.5)	15 (75.0)	5 (25.0)	
	30 – 34	14 (8.1)	10 (71.4)	4 (28.6)	
	35 – 39	28 (16.1)	18 (64.3)	10 (35.7)	
	40 – 44	18 (10.3)	11 (61.1)	7 (38.9)	
	45 – 49	27 (15.5)	12 (44.4)	15 (55.5)	
	50 – 54	35 (20.1)	15 (42.9)	20 (57.1)	
	55 – 59	14 (8.1)	10 (71.4)	4 (28.6)	
	> 60	13 (7.5)	3 (23.1)	10 (76.9)	
	Missing	2 (1.1)	0 (0.0)	2 (100)	
Race	Malay	161 (92.5)	90 (55.9)	71 (44.1)	0.752
	Chinese	6 (3.5)	4 (66.7)	2 (33.3)	
	Other	7 (4.0)	3 (42.9)	4 (57.1)	
Education level	Primary school	16 (9.2)	10 (62.5)	6 (37.5)	0.396
	Secondary school	96 (55.2)	49 (51.0)	47 (49.0)	
	College / University	57 (32.7)	35 (61.4)	22 (38.6)	
	Missing	5 (2.9)	3 (60.0)	2 (40.0)	
Marital status	Married	157 (90.2)	91 (58.0)	66 (42.0)	0.15
	Divorced	8 (4.6)	2 (25.0)	6 (75.0)	
	Widowed	9 (5.2)	4 (44.4)	5 (55.6)	
Occupation	Housewife	64 (36.8)	39 (60.9)	25 (39.1)	0.087
	Government employee	67 (38.5)	41 (61.2)	26 (38.8)	
	Private employee	31 (17.8)	13 (41.9)	18 (58.1)	
	Retired	9 (5.2)	3 (33.3)	6 (66.7)	
	Unemployed	1 (0.6)	1 (100)	0 (0.0)	
	Other	2 (1.1)	0 (0.0)	2 (100)	
Monthly household income	< $500	27 (15.5)	15 (55.6)	12 (44.4)	0.498
	$500 < $999	27 (15.5)	12 (44.4)	15 (55.6)	
	$1000-$1999	40 (23.0)	22 (55.0)	18 (45.0)	
	$2000-$2999	19 (10.9)	14 (73.7)	5 (26.3)	
	$3000-$5000	24 (13.8)	15 (62.5)	9 (37.5)	

	>$5000	3 (1.7)	2 (66.7)	1 (33.3)	
	Missing	34 (19.5)	17 (50.0)	17 (50.0)	
Number of births	0	27 (15.5)	17 (63.0)	10 (37.0)	0.103
	1	19 (10.9)	14 (73.7)	5 (26.3)	
	2	22 (12.1)	14 (66.7)	7 (33.3)	
	3 or more	105 (60.3)	51 (48.6)	54 (51.4)	
	Missing	2 (1.2)	1 (50.0)	1 (50.0)	
Last Pap test done	Never	41 (23.6)	29 (70.7)	12 (29.3)	**0.031***
	4-10 years	91 (52.3)	48 (52.7)	43 (47.3)	
	> 10 years	36 (20.7)	15 (41.7)	21 (58.3)	
	Missing	6 (3.4)	5 (83.3)	1 (16.7)	
HPV vaccination status	Unvaccinated	95 (54.6)	50 (52.6)	45 (47.4)	0.483
	Fully vaccinated	27 (15.5)	16 (59.3)	11 (40.7)	
	Partly vaccinated	41 (23.6)	26 (63.4)	15 (36.6)	
	Missing	11 (6.3)	5 (45.5)	6 (54.5)	

Fig. 2 and S1 Table show the responses for the major and minor reasons for not attending

cervical cancer screening. The top three major reasons reported were that they were "afraid

of getting a bad result" (16.1%, n = 28), "feeling embarrassed being examined by a doctor or

nurse" (14.9%, n = 26), and "I can't find the time as I'm too busy at home" (10.3%, n = 18).

The top three minor reasons were "feeling embarrassed being examined by a doctor or

nurse" (20.7%, n = 36), "I can't find time as I'm too busy at work" (20.7%, n = 36), and

"afraid of getting a bad result" (20.1%, n = 35).

Fig. 2. Responses on their major and minor reasons for not attending cervical cancer

screening, among non-attendees at JPSHC (Jan – Dec 2019). The x-axis indicates the

percentage, and the number next to each bar indicates the actual number of responses.

When comparing top 10 major reasons for not attending screening between those who

agreed and declined HPV self-sampling (Table 2), those whose major reason was "feeling

embarrassed being examined by a doctor or nurse" were significantly more likely to join

self-sampling (p = 0.020). Also, those whose major reason was "afraid of getting a bad

result" were significantly more likely to decline self-sampling (p = 0.034). Among those who

were employed (n = 98), about a quarter (23.5%, n = 23) reported work-related reasons as

their major reason for not attending screening ("I can't find the time as I'm too busy at

work" and "Difficult to get permission from employer").

Table 2. The top ten major reasons of not attending cervical cancer screening at JPSHC,

Brunei (Jan – Dec 2019), with comparison between those who agreed and declined HPV self-

sampling. Responses from 146 participants (83.9% of the total study population) were

included.

No.	Top ten major reasons of screening non-attendance	Total n (%)	Joined HPV self-sampling n (%)	Declined HPV self-sampling n (%)	p-value
1	I feel embarrassed being examined by a doctor or nurse	26 (14.9)	20 (76.9)	6 (23.1)	**0.020***
2	I am scared of pain because of previous bad experience(s)	16 (9.2)	8 (50.0)	8 (50.0)	0.825
3	I am scared of getting a bad result	28 (16.1)	10 (35.7)	18 (64.3)	**0.034***
4	I can't find the time as I'm too busy at home	18 (10.3)	12 (66.7)	6 (33.3)	0.463
5	I can't find the time as I'm too busy at work	16 (9.2)	10 (62.5)	6 (37.5)	0.759
6	Difficult to get permission from employer	8 (4.6)	5 (62.5)	3 (37.5)	1
7	I have never heard of a Pap Test	7 (4.0)	4 (57.1)	3 (42.9)	1
8	I have forgotten about it	6 (3.4)	5 (83.3)	1 (16.7)	0.229
9	I did not receive any invitation	6 (3.4)	5 (83.3)	1 (16.7)	0.229
10	Others	15 (8.6)	7 (46.7)	8 (53.3)	0.492

While the responses vary when asked about their health and susceptibility to disease, most

agreed on the benefits of undergoing Pap test (92.0%, n = 160), and that cervical cancer is a

severe and potentially lethal disease(82.8%, n = 144; Table 3 and S2 Table). Those who

agreed that they are more susceptible to develop cervical cancer were significantly more

likely to join HPV self-sampling (p = 0.023).

Table 3. Attitudes towards cervical cancer screening among non-attendees at JPSHC, Brunei

(Jan–Dec 2019), between those who agreed and declined HPV self-sampling.

No.	Attitude questions		Total study population (n = 174) n (%)	Joined HPV self-sampling (n = 97) n (%)	Declined HPV self-sampling (n = 77) n (%)	*p*-value
1	I believe I am healthy and free of any diseases	Agree	62 (35.6)	34 (54.8)	28 (45.2)	0.984
		Neutral/ Disagree	112 (64.3)	63 (56.3)	49 (43.7)	
2	Having Pap test taken is beneficial for my health	Agree	160 (92.0)	91 (56.9)	69 (43.1)	0.464
		Neutral/ Disagree	14 (8.0)	6 (42.9)	8 (57.1)	
3	Like any women, I am susceptible to develop cervical cancer	Agree	110 (63.2)	69 (62.7)	41 (37.3)	**0.023***
		Neutral/ Disagree	64 (36.8)	28 (43.8)	36 (56.2)	
4	Cervical cancer can be severe and may lead to death	Agree	144 (82.8)	79 (54.9)	65 (45.1)	0.754
		Neutral/ Disagree	30 (17.2)	18 (60.0)	12 (40.0)	

Most participants would like to obtain more information about cervical cancer screening

(85.6%, n = 149). The top preferred information sources were healthcare workers (59.8%, n

= 104), social messaging platforms (28.7%, n = 50) and social media (26.4%, n = 46; S1 Fig).

Among those who participated in self-sampling (55.7%, n = 97), their responses on Q2

(Table 4) were mostly positive. A majority agreed that the instructions were clear (94.8%, n

= 92), that it was easy to perform the swab (93.8%, n = 91), and that it was more convenient

than the Pap test (91.7%, n = 89). They also reported their confidence in correctly getting

the sample (92.8%, n = 90), would prefer to use this method next time (94.8%, n = 92), and

would recommend this method to other women (93.8%, n = 91). Notably, 54.6% (n = 53) still

prefer a proper Pap test for their subsequent check-up.

Table 4. Acceptability of HPV self-sampling among non-attendees who participated in HPV

self-sampling at JPSHC, Brunei (Jan–Dec 2019). Responses from all Q2 respondents (n = 97)

were included.

No.	Self-sampling Questions	Agree n (%)	Neutral n (%)	Disagree n (%)
1	I thought the instructions were clear	92 (94.8)	0 (0.0)	5 (5.2)
2	It was easy to do the swab	91 (93.8)	1 (1.0)	5 (5.2)
3	Taking the sample with the swab was painful	12 (12.4)	8 (8.2)	77 (79.4)
4	Taking the sample was uncomfortable to do	11 (11.4)	4 (4.1)	82 (84.5)
5	I felt embarrassed doing the self-sampling	9 (9.3)	1 (1.0)	87 (89.7)
6	It was convenient to do without having to undergo a Pap Test	89 (91.7)	3 (3.1)	5 (5.2)
7	I am confident I did it correctly	90 (92.8)	4 (4.1)	3 (3.1)
8	I want to use this method next time	92 (94.8)	2 (2.1)	3 (3.1)
9	I prefer to do this at home	64 (66.0)	15 (15.5)	18 (18.5)
10	I would recommend this method to other women	91 (93.8)	3 (3.1)	3 (3.1)
11	I trust that the result of this self-sampling will be accurate	74 (76.3)	19 (19.6)	4 (4.1)
12	I would like to attend for a proper Pap test in clinic next time	53 (54.6)	23 (23.7)	21 (21.7)

Among the 97 samples taken, nine (9.3%) tested positive for hr-HPV: one was positive for

HPV 16 and eight were positive for non-HPV 16/18 HPV genotype. The HPV 16 positive case

was found to have high-grade ASC-US (ASCUS-H) on the initial follow up smear, but found

negative after subsequent follow up cervical biopsy. Among the other 8 non-HPV 16/18 HPV

genotype positive cases: 2 had negative follow up smears, 4 were reported to have ASC-US

on their initial follow up smears, but had subsequent negative follow up smears, 1 was

reported to have low-grade squamous intraepithelial lesion (LSIL) on her initial follow up

smear, and had a subsequent negative follow up smear, and 1 was found to have CIN 3 with

glandular involvement on the follow up smear and cervical biopsy, and has received

treatment with a cone biopsy where excision of CIN 3 was confirmed. No significant

differences were observed when comparing the sociodemographic characteristics between

those who tested hr-HPV positive and those who tested hr-HPV negative (S3 Table).

Discussion

Our study findings highlight three important points to consider for improving cervical cancer

screening uptake and detection. First, our findings suggest that it is necessary to provide

accurate information among Bruneian women on cervical cancer, the importance of

screening and addressing any misconceptions about the Pap test. Important facts to relay

include the slow development from pre-cancerous changes to cervical cancer, that pre-

cancerous changes are highly treatable, and that screening will help in early detection and

thus facilitate successful treatment.

In our study, emotional barriers (fear of unfavorable test results and embarrassment) were

the most common major reasons of screening non-attendance. Also, about two-thirds of

our participants cited "I am scared of getting a bad result" as their main barrier also

declined to take part in HPV self-sampling, possibly due to relating abnormal Pap test results

to cervical cancer diagnoses [25]. Although such barriers may play a large role at the onset

of screening program, its role diminishes over time with increasing education on the

benefits of screening [26]. Educational interventions could also benefit the small group of

women who cited menopause, cessation of child-bearing and having had HPV vaccination as

reasons for not attending screening (Fig. 2 and S1 Table).

In our study, we observed those who agree that they are susceptible to cervical cancer were

significantly more likely to participate in HPV self-sampling. This suggests that perceived

susceptibility could be an important factor for self-sampling participation, whereby those

who do not perceive themselves as susceptible were less likely to engage in preventative

behaviors [27–29]. Perceived susceptibility can be increased through education to improve

their beliefs on the importance of screening [30]. We suggest that such information could

be more effectively disseminated as simple health messages endorsed by Brunei's Ministry

of Health via website and social media platforms.

Second, we observed high acceptability of HPV self-sampling among our participants. Meta-

analyses have indicated strong acceptance of and preference for self-sampling over clinician

sampling [31], mainly due to logistical reasons [32]. More than half of our participants who

joined HPV self-sampling cited embarrassment or lack of time due to home and/or work

commitments as their major reason of not attending Pap test screening. This suggest that

providing flexibility to accommodate women's screening method preference [33], such as

the option of self-sampling [34], could improve screening uptake.

Thirdly, most of the detected hr-HPV genotypes in our study were non-16/18, with only one

out of nine participants tested positive for HPV 16. Although our sample size is small, this

result suggests that it may not be accurate to assume HPV 16 or 18 as common hr-HPV

genotypes in Brunei, even though this is true in the global context [35]. Other studies have

detected a significant percentage of non-16/18 hr-HPV genotypes, suggesting the presence

of region-specific heterogeneity in the HPV genotype distribution [35–39]. Also, variation in

HPV distribution among different ethnic groups has been reported in an American study

[40]. Our finding has potential implication on Brunei's national school-based HPV

vaccination program [41] which currently provides vaccines which do not confer protection

against non-16/18 genotypes. Larger population-based studies to understand the

distribution of HPV genotypes among Bruneian women will be crucial to determine the

efficacy or impact of the current vaccines.

One notable point for the local context relates to the presence of cultural barriers. Being a

predominantly Islamic society, religious and cultural modesty could be a contributing factor

for embarrassment among Muslim women [42]. Also, having premarital sex is ataboo in

Brunei and is generally not openly discussed [43]. This could prevent any unmarried but

sexually active women from participating in the screening program. It should be emphasized

that only married or ever married females were included in this study; included because it is

part of the eligibility criteria for the national cervical cancer screening program in Brunei.

A major limitation for this study is that non-attendees from only one health center were

recruited, thus our findings are not representative of the adult female population in Brunei.

Secondly, our findings should also be interpreted with caution due to the small sample size

and the non-probability sampling approach used. This study was initially conceived as a pilot

study due to resource and logistics limitations. There were two reasons for choosing JPSHC

as our study site: It is the third largest primary government healthcare center in the country,

and that it serves a sub-district with a sizable percentage of residents in the middle- to low-

income groups. However, even at this pilot stage, our study findings could encourage

stakeholders to conduct similar and larger studies, using random sampling approach. Lastly,

as this study relied on the self-reported history of previous Pap test attendance, we might

have missed recruiting those who may have forgotten their last Pap test date.

In conclusion, our findings indicate the need to further promote knowledge on cervical

cancer, the benefits of screening and clarifying any misconceptions of Pap test results.

Reasons of cervical cancer screening non-attendance were mainly related to emotional and

logistical factors. As we found high acceptance towards HPV self-sampling, this could be

adopted as an alternative for women who refrain from Pap test. Our high detection of non-

16/18 HPV genotypes suggest high prevalence of other hr-HPV genotypes in Brunei. Future

larger studies involving more Bruneian women should be done to verify our results. Follow-

up studies should also be conducted to consider HPV testing as the suitable method for

cervical cancer screening.

Acknowledgements

396
397   The authors would like to thank the JPSHC staff who have assisted in participant recruitment

398   and questionnaire distribution.

References

1. World Health Organization. WHO guideline for screening and treatment of cervical pre-cancer lesions for cervical cancer prevention (2nd edition). 2021. Available: https://www.who.int/publications/i/item/9789240030824

2. Gustafsson L, Pontén J, Zack M, Adami H-O. International incidence rates of invasive cervical cancer after introduction of cytological screening. Cancer Causes Control. 1997;8: 755–763. doi:10.1023/A:1018435522475

3. Ministry of Health Brunei Darussalam. Health Information Booklet 2017 (Second Revision). 2019 [cited 11 Feb 2020]. Available: http://moh.gov.bn/SitePages/Health Information Booklet.aspx

4. Arbyn M, Weiderpass E, Bruni L, de Sanjosé S, Saraiya M, Ferlay J, et al. Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. Lancet Glob Heal. 2020;8: e191–e203. doi:10.1016/S2214-109X(19)30482-6

5. WHO. Guidelines for screening and treatment of precancerous lesions for cervical cancer prevention. WHO Guidel. 2013.

6. Bosgraaf RP, Ketelaars PJW, Verhoef VMJ, Massuger LFAG, Meijer CJLM, Melchers WJG, et al. Reasons for non-attendance to cervical screening and preferences for HPV self-sampling in Dutch women. Prev Med (Baltim). 2014. doi:10.1016/j.ypmed.2014.04.011

7. Virtanen A, Nieminen P, Niironen M, Luostarinen T, Anttila A. Self-sampling experiences among non-attendees to cervical screening. Gynecol Oncol. 2014. doi:10.1016/j.ygyno.2014.09.019

8. Knops-Dullens T, de Vries N, de Vries H. Reasons for non-attendance in cervical

cancer screening programmes: an application of the Integrated Model for

Behavioural Change. Eur J Cancer Prev. 2007;16. Available:

https://journals.lww.com/eurjcancerprev/Fulltext/2007/10000/Reasons_for_non_att

endance_in_cervical_cancer.9.aspx

9.   Waller J, Bartoszek M, Marlow L, Wardle J. Barriers to cervical cancer screening

attendance in England: a population-based survey. J Med Screen. 2009;16: 199–204.

doi:10.1258/jms.2009.009073

10.   Oscarsson MG, Benzein EG, Wijma BE. Reasons for non-attendance at cervical

screening as reported by non-attendees in Sweden. J Psychosom Obstet Gynecol.

2008;29: 23–31. doi:10.1080/01674820701504619

11.   Yeh PT, Kennedy CE, Vuyst H de, Narasimhan M. Self-sampling for human

papillomavirus (HPV) testing: a systematic review and meta-analysis. BMJ Glob Heal.

2019;4: e001351. doi:10.1136/BMJGH-2018-001351

12.   CS R, DR W, D G. Self-collected HPV testing improves participation in cervical cancer

screening: a systematic review and meta-analysis. Can J Public Health. 2013;104.

doi:10.1007/BF03405681

13.   Maza M, Melendez M, Masch R, Alfaro K, Chacon A, Gonzalez E, et al. Acceptability of

self-sampling and human papillomavirus testing among non-attenders of cervical

cancer screening programs in El Salvador. Prev Med (Baltim). 2018.

doi:10.1016/j.ypmed.2018.06.017

14.   Sancho-Garnier H, Tamalet C, Halfon P, Leandri FX, Retraite L Le, Djoufelkit K, et al.

HPV self-sampling or the Pap-smear: A randomized study among cervical screening

nonattenders from lower socioeconomic groups in France. Int J Cancer. 2013.

doi:10.1002/ijc.28283

15.   Chrysostomou AC, Stylianou DC, Constantinidou A, Kostrikis LG. Cervical cancer

screening programs in Europe: The transition towards HPV vaccination and

population-based HPV testing. Viruses. 2018;10. doi:10.3390/v10120729

16.   Arrossi S, Thouyaret L, Herrero R, Campanera A, Magdaleno A, Cuberli M, et al. Effect

of self-collection of HPV DNA offered by community health workers at home visits on

uptake of screening for cervical cancer (the EMA study): A population-based cluster-

randomised trial. Lancet Glob Heal. 2015;3: e85–e94. doi:10.1016/S2214-

109X(14)70354-7

17.   Gustavsson I, Aarnio R, Berggrund M, Hedlund-Lindberg J, Strand A-S, Sanner K, et al.

Randomised study shows that repeated self-sampling and HPV test has more than

two-fold higher detection rate of women with CIN2+ histology than Pap smear

cytology. Br J Cancer. 2018;118: 896–904. doi:10.1038/bjc.2017.485

18.   Arrossi S, Paolino M, Laudi R, Gago J, Campanera A, Marín O, et al. Programmatic

human papillomavirus testing in cervical cancer prevention in the Jujuy

Demonstration Project in Argentina: a population-based, before-and-after

retrospective cohort study. Lancet Glob Heal. 2019. doi:10.1016/S2214-

109X(19)30048-8

19.   Landy R, Pesola F, Castañón A, Sasieni P. Impact of cervical screening on cervical

cancer mortality: Estimation using stage-specific results from a nested case-control

study. Br J Cancer. 2016;115: 1140–1146. doi:10.1038/bjc.2016.290

20.   Canfell K, Sitas F, Beral V. Cervical cancer in Australia and the United Kingdom:

Comparison of screening policy and uptake, and cancer incidence and mortality. Med

J Aust. 2006;185: 482–486. doi:10.5694/j.1326-5377.2006.tb00661.x

21.   Bayu H, Berhe Y, Mulat A, Alemu A. Cervical Cancer Screening Service Uptake and

Associated Factors among Age Eligible Women in Mekelle Zone, Northern Ethiopia,

2015: A Community Based Study Using Health Belief Model. PLoS One. 2016;11:

e0149908. doi:10.1371/journal.pone.0149908

22. Sultana F, Mullins R, English DR, Simpson JA, Drennan KT, Heley S, et al. Women's

experience with home-based self-sampling for human papillomavirus testing. BMC

Cancer. 2015;15: 849. doi:10.1186/s12885-015-1804-x

23. The Hague: Health Council of the Netherlands. Health Council of the Netherlands

Population screening for cervical cancer. 2011;2011/07E. Available:

https://www.gezondheidsraad.nl/sites/default/files/201107E_PopulationSCC_0.pdf

24. R Core Team. R: A language and environment for statistical computing. Vienna,

Austria; 2021. Available: http://www.r-project.org/

25. Fylan F. Screening for cervical cancer: a review of women's attitudes, knowledge, and

behaviour. Br J Gen Pract. 1998;48: 1509–14.

26. Teng FF, Mitchell SM, Sekikubo M, Biryabarema C, Byamugisha JK, Steinberg M, et al.

Understanding the role of embarrassment in gynaecological screening: A qualitative

study from the ASPIRE cervical cancer screening project in Uganda. BMJ Open. 2014.

doi:10.1136/bmjopen-2014-004783

27. Agurto I, Arrossi S, White S, Coffey P, Dzuba I, Bingham A, et al. Involving the

community in cervical cancer prevention programs. Int J Gynaecol Obstet. 2005;89

Suppl 2: S38-45. doi:10.1016/j.ijgo.2005.01.015

28. Shirazi Zadeh Mehraban S, Namdar A, Naghizadeh MM. Assessment of Preventive

Behavior for Cervical Cancer with the Health Belief Model. Asian Pac J Cancer Prev.

2018;19: 2155–2163. doi:10.22034/APJCP.2018.19.8.2155

29. Tapera R, Manyala E, Erick P, Maswabi TM, Tumoyagae T, Letsholo B, et al.

Knowledge and Attitudes towards Cervical Cancer Screening amongst University of

Botswana Female Students. Asian Pac J Cancer Prev. 2017;18: 2445–2450.

doi:10.22034/APJCP.2017.18.9.2445

30. Khademolhosseini F, Noroozi A, Tahmasebi R. The Effect of Health Belief Model-

Based Education through Telegram Instant Messaging Services on Pap smear

performance. Asian Pac J Cancer Prev. 2017;18: 2221–2226.

doi:10.22034/APJCP.2017.18.8.2221

31. Nelson EJ, Maynard BR, Loux T, Fatla J, Gordon R, Arnold LD. The acceptability of self-

sampled screening for HPV DNA: A systematic review and meta-analysis. Sexually

Transmitted Infections. 2017. doi:10.1136/sextrans-2016-052609

32. Lorenzi NPC, Termini L, Longatto Filho A, Tacla M, de Aguiar LM, Beldi MC, et al. Age-

related acceptability of vaginal self-sampling in cervical cancer screening at two

university hospitals: a pilot cross-sectional study. BMC Public Health. 2019.

doi:10.1186/s12889-019-7292-1

33. Bansil P, Wittet S, Lim JL, Winkler JL, Paul P, Jeronimo J. Acceptability of self-collection

sampling for HPV-DNA testing in low-resource settings: a mixed methods approach.

BMC Public Health. 2014;14: 596. doi:10.1186/1471-2458-14-596

34. Jeronimo J, Perkins RB, Scalici J, Pierce JY. Should Self-Sampling Be an Option for

Women in the United States? J Low Genit Tract Dis. 2019;23: 54.

doi:10.1097/LGT.0000000000000453

35. Albawardi A, Quddus MR, Al Awar S, Almarzooqi S. Frequency of rare and multi viral

high-risk HPV types infection in cervical high grade squamous intraepithelial lesions in

a non-native dominant middle eastern country: A polymerase chain reaction-based

pilot study. Diagn Pathol. 2018. doi:10.1186/s13000-018-0716-x

36.　Zeng Z, Yang H, Li Z, He X, Griffith CC, Chen X, et al. Prevalence and Genotype

Distribution of HPV Infection in China: Analysis of 51,345 HPV Genotyping Results

from China's Largest CAP Certified Laboratory. J Cancer. 2016. doi:10.7150/jca.14971

37.　Khoo SP, Bhoo-Pathy N, Yap SH, Anwar Shafii MK, Hairizan Nasir N, Belinson J, et al.

Prevalence and sociodemographic correlates of cervicovaginal human papillomavirus

(HPV) carriage in a cross-sectional, multiethnic, community-based female Asian

population. Sex Transm Infect. 2018. doi:10.1136/sextrans-2017-053320

38.　Peng RR, Li HM, Chang H, Li JH, Wang AL, Chen XS. Prevalence and genotype

distribution of cervical human papillomavirus infection among female sex workers in

Asia: A systematic literature review and meta-analysis. Sexual Health. 2012.

doi:10.1071/SH11066

39.　Al-Awadhi R, Chehadeh W, Jaragh M, Al-Shaheen A, Sharma P, Kapila K. Distribution

of human papillomavirus among women with abnormal cervical cytology in Kuwait.

Diagn Cytopathol. 2013. doi:10.1002/dc.21778

40.　Keller MJ, Burk RD, Massad LS, Eltoum IE, Hessol NA, Anastos K, et al. Racial

differences in human papilloma virus types amongst United States women with HIV

and cervical precancer. AIDS. 2018. doi:10.1097/QAD.0000000000002005

41.　Chaw LL, Lim STW, Md Yussof SR. Human Papillomavirus vaccine coverage among

female students in Brunei Darussalam: results from the first 4 years of the national

school-based vaccination programme. Heliyon. 2019;5.

doi:10.1016/j.heliyon.2019.e02588

42.　Ali S, Skirton H, Clark MT, Donaldson C. Integrative review of cervical cancer screening

in Western Asian and Middle Eastern Arab countries. Nurs Heal Sci. 2017;19: 414–

426. doi:10.1111/nhs.12374

43. Haji Tahamit N. Cultural and Religious Barriers to Setting Up Sex and Relationship

Education in a Muslim Country: A Case Study of Brunei Darussalam. University of

Leeds; 2015 Aug. Available: https://etheses.whiterose.ac.uk/15306/

Supplementary information

S1 Table. Responses on their major and minor reasons for not attending cervical cancer

screening among non-attendees at JPSHC, Brunei (Jan – Dec 2019).

S2 Table. Attitudes towards cervical cancer screening among non-attendees at JPSHC,

Brunei (Jan – Dec 2019). Responses from the total study population (n = 174) were included.

S1 Fig. Preferred sources of information about cervical cancer among non-attendees at

JPSHC, Brunei (Jan–Dec 2019). The x-axis indicates the percentage, and the number next to

each bar indicates the number of responses. Multiple responses were allowed and

responses from the total study population (n = 174) were included.

S3 Table. Socio demographic characteristics and comparison between screening non-

attendees who tested positive and negative for hr-HPV at JPSHC, Brunei (Jan – Dec 2019).

S4 Questionnaire. Questionnaire 1: Reasons why women do not participate in the National

Cervical Cancer Screening Program

S5 Questionnaire. Questionnaire 2: Acceptability of Self Sampling for HPV Testing Among

Non-attendees of Cervical Screening Program in Brunei Darussalam

S6 Dataset. Minimal dataset

Author contributions:

Liling Chaw: Conceptualization, Methodology, Data curation, Formal analysis, Supervision, Writing - original draft, Writing - review & editing

Shirley HF Lee: Visualization, Writing - original draft, Writing - review & editing

Nurul Iffah Hazwani Ja'afar: Investigation, Data curation, Writing - original draft

Edwin Lim: Conceptualization, Methodology, Resources, Funding acquisition, Supervision, Writing - review & editing

Roslin Sharbawi: Conceptualization, Methodology, Resources, Funding acquisition, Investigation, Project administration, Supervision, Writing - original draft, Writing - review & editing

Category	Main reason	Minor reason
I am scared of getting a bad result	28	35
I feel embarrassed being examined by a doctor or nurse	26	36
I can't find the time as I'm too busy at home	18	28
I am scared of pain because of previous bad experience(s)	13	16
I can't find the time as I'm too busy at work	16	36
Difficult to get permission from employer	8	14
I have never heard of a Pap Test	7	8
I did not receive any invitation	0	6
I have forgotten about it	2	6
Not necessary as I don't have menses anymore	4	11
Not necessary as I am not child-bearing anymore	4	14
Not necessary as I am healthy	4	15
Nobody is looking after child(ren) at home	2	13
Nobody to send me to clinic	2	11
Not necessary as I have already had my HPV vaccination	2	2
I don't know what cervical cancer is	1	15
Not necessary as I don't have a partner anymore	0	1
Others	8	4
Missing	9	

Main reason · Minor reason