1

Title:

Reasons for non-attendance to cervical cancer screening and acceptability of HPV self-

sampling among Bruneian women: A cross-sectional study.

Authors:

Liling Chaw[1*], Shirley HF Lee[1], Nurul Iffah Hazwani Ja'afar[1], Edwin Lim[2], Roslin Sharbawi[3]

Affiliations:

1) PAPRSB Institute of Health Sciences, Gadong, Brunei Darussalam.

2) Histopathology Department, RIPAS Hospital, Ministry of Health, Brunei Darussalam

3) Community Maternal Health Service, Department of Health Services, Ministry of Health,
Bandar Seri Begawan, Brunei Darussalam

* Corresponding Author:

Email: liling.chaw@ubd.edu.bn (LC)

Running title:

Reasons for cervical cancer screening non-attendance and HPV self-sampling acceptability in Brunei

Number of words: 327 words (abstract), 3049 words (manuscript, excluding references)

Abstract
28

Objective: Uptake for cervical cancer screening remains well below the recommended rate of 80%.

We conducted a preliminary study to determine the reasons for non-attendance and explore their

acceptance of human papillomavirus (HPV) self-sampling as an alternative to the Pap test.

Methods: A cross-sectional study was conducted at a health center in Brunei, from January to

December 2019. Women who were eligible for Pap test but who either never or did not have one

within the past four years were conveniently recruited. Participants were first asked to complete a

paper-based survey on their reasons for screening non-attendance, and then invited for HPV self-

sampling. Results were analyzed using descriptive and inferential statistics.

Result: We enrolled 174 screening non-attendees, out of which 97 (55.7%) also participated in HPV

self-sampling. The main reasons for not attending Pap test screening were fear of bad results

(16.1%, n = 28); embarrassment (14.9%, n = 26) and lack of time due to home commitments (10.3%,

n = 18). When compared to those who agreed to participate in HPV self-sampling, those who

declined were significantly older (p = 0.002) and less likely to agree that they are susceptible to

cervical cancer (p = 0.023). They preferred to receive Pap test-related information from healthcare

workers (59.0%, n = 155), social messaging platforms (28.7%, n = 51) and social media (26.4%, n =

47). HPV self-sampling kits were positively received, where > 90% agreed on its ease and

convenience. Nine of 97 participants (9.3%) tested positive for hr-HPV, out of which eight were non-

16/18 HPV genotypes.

Conclusion: Our findings suggest that promoting knowledge on cervical cancer, the benefits of

screening and clarifying any misconceptions of Pap test results would help increase screening uptake

among Bruneian non-attendees. Response to HPV self-sampling was highly positive, implying its

receptiveness in the local setting. Our high detection of non-16/18 HPV genotypes suggest high

prevalence of other hr-HPV genotypes in Brunei. Larger studies can be conducted to further validate

our findings.

Keywords: Cervical cancer screening, HPV self-sampling, women's health, Brunei

Introduction

Cervical cancer is highly preventable but still remains one of the most common cancers among women worldwide, accounting for 6.6% of all female cancer deaths in 2018 [1]. Cervical cancer screening has drastically reduced the incidence of invasive cervical cancer in countries that have implemented such screening programs [2], which traditionally involves the use of the standard Papanicolaou (Pap) test. However, as specific high-risk human papillomavirus subtypes (hr-HPV) are known to be a causative agent, the use of HPV self-sampling is becoming either an alternative or a complement to Pap test, with an aim to increase screening uptake rate [3] particularly among screening non-attendees [4].

HPV self-sampling were found to be highly acceptable among screening non-attendees [5,6]. Repeated HPV self-sampling and testing were shown to increase screening uptake [3,4,7], and also resulted in at least two-fold higher detection rate of high-grade cervical intraepithelial neoplasia (CIN2+) when compared to the Pap test [8,9]. In addition, HPV self-sampling results exhibit similar sensitivity and specificity compared to those from samples taken by trained professionals [10]. At the national level, early detection through organized screening was shown to reduce cervical cancer morbidity and mortality in countries such as the United Kingdom and Australia [11,12]. Both countries have further strengthened their national screening programs by using HPV testing as the primary screening test.

Brunei Darussalam (population 459,500) is a small Southeast Asian country with a predominant Muslim population and a crude birth rate of 15.3 per 1,000 population [13]. Within this region, it has one of the highest age-standardized incidence rate (ASR) for cervical cancer: 20.6 in Brunei when compared to 10.5 and 7.7 in Malaysia and Singapore, respectively [14]. Brunei has initiated a national cervical cancer prevention and control program since 2009, which include setting up the

National Pap Test Registry, organizing cervical screening through periodic mail invitations for eligible

women, replacing the Pap test with liquid-based cytology, and initiating HPV vaccination for school-

aged female adolescents [15]. However, despite cervical screening being offered free of charge, the

national screening coverage rate remains low at 44% in 2018 (unpublished data). It is known that

screening non-attendees are at increased risk of developing cervical cancer [16], and that reasons

for screening non-attendance can also vary across settings [17,18].

With Brunei's relatively high incidence of cervical cancer and low screening uptake, we conducted a

preliminary study to explore the reasons behind non-attendance and to assess the acceptability of

HPV self-sampling as a possible alternative to the Pap test among non-attendees. Study findings

could be used to strategize ways to improve screening uptake and provide preliminary evidence

towards implementing HPV testing as the primary screening test for cervical cancer in Brunei.

Methods

Study design and data collection

A cross-sectional survey was conducted at the Jubli Perak Sengkurong Health Center (JPSHC), from

January to December 2019. JPSHC is a government-funded health center located at Brunei-Muara

District, where the majority (69.3%) of the country's population resides. This center provides primary

health care services to Mukim Sengkurong, a sub-district with about 32 000 people from various

socioeconomic backgrounds.

Eligible women attending either the outpatient or child health clinic were conveniently recruited by

triage nurses. The inclusion criteria were married or ever married Bruneian women between 20 and

65 years old, who either never or did not have a Pap test within the last 4 years, and who can

comprehend Brunei-Malay or English language. We excluded women who were pregnant, have had

total hysterectomy, or with a history of malignancies.

Participants who gave written consent to the first part of the study were asked to complete the first

questionnaire (Q1) onsite. They were then given an envelope containing information on the second

part and an instruction leaflet on the HPV self-sampling procedure. Upon second contact about two

weeks later, those who gave verbal consent were given an appointment to JPSHC. On the day, a

trained nurse first explained the procedure using an instructional video and answered any questions.

Participants who gave another written consent were then given a self-sampling kit and asked to

perform the procedure in the clinic. After completion, they were asked to complete the second

questionnaire (Q2).

Questionnaires used
124
125

Two bilingual paper-based questionnaires (Brunei-Malay and English language) were used: one on

the reasons for non-attendance (Q1) and another on their acceptability of the self-sampling

procedure (Q2). Both questionnaires were adapted from similar studies [19,20]. They were first

translated to Brunei-Malay language by native speakers, and then back-translated to check for

inconsistencies in comprehension. Both questionnaires were also pre-tested on six eligible women

to assess if the questions can be easily understood. Responses from pre-testing were not included in

the main analysis.

Q1 consists of 19 questions on the participant's socio-demographics; reasons for not getting Pap

test; attitude and knowledge on cervical cancer; and lastly, preferred sources to acquire information

about Pap test. From a prepared list of 16 options, participants were asked to select one ("Major"

reason) or more reasons (categorized as "Minor" reasons) for not attending the screening program.

Responses for questions on their attitude and knowledge on cervical cancer were assessed using a

five-point Likert scale, ranging from "Strongly Agree" to "Strongly Disagree". Q2 consists of 12 five-

point Likert scale questions (ranging from "Strongly Agree" to "Strongly Disagree") on their

experiences and opinions of the self-sampling kit.

Self-sample handling and laboratory testing
143
144

Swabs collected from the self-sampling kits were sent to an overseas laboratory (at BNH hospital,

Thailand) for hr-HPV testing. The self-sampling device used was the Evalyn® brush from Rovers

Medical Devices. Upon receipt at the testing laboratory, the dry Evalyn® brush was suspended in

SurePath medium from which a sample was obtained for the identification of the presence of hr-

HPV using the cobas HPV test (Roche, USA). Cobas HPV tests are automated qualitative *in vitro* tests

for the detection of HPV DNA in patient specimens. The tests utilize amplification of target DNA by

polymerase chain reaction (PCR) and nucleic acid hybridization for the detection of 14 hr-HPV types

(namely genotypes 16, 18, and 12 pooled hr-HPV genotypes 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66

and 68) in a single analysis. Results obtained from this test can be categorized into four groups:

Negative, HPV-16 positive, HPV-18 positive, and positive for non-16/18 HPV genotypes.

Clinical management of hr-HPV positive participants
156
157

We adopted the cytology triage strategy for HPV self-sampling participants [21]. Participants with hr-

HPV positive results were invited for an immediate clinic-based cytology triage test. Those found to

have negative cytology results were invited for a repeat cytology triage test after six months. Those

with second negative results were returned to routine cervical screening recall. Participants with any

positive cytology result (defined as with Atypical Squamous Cells of Undetermined Significance or

worse) were referred for colposcopic examination.

Statistical analysis

Descriptive statistics was conducted to characterize the socio-demographic characteristics of the study population, their reasons for not attending screening, their attitudes and preference for information access on such screening, as well as responses from Q2. Mann-Whitney and Fisher's exact tests were used to assess significant differences in sociodemographic characteristics between women who agreed and women declined to proceed to the HPV self-sampling part of the study. Responses for questions with the five-point Likert scale were categorized into three categories (agree, neutral, and disagree), and those with missing values were classified as neutral. Statistical analysis was conducted using R ver. 3.6 [22]. Ethics approval was obtained from the Medical and Health Research and Ethics Committee (MHREC), Ministry of Health, Brunei Darussalam [Reference no. MHREC/MOH/2018/9(2)].

Results

A total of 174 eligible women were enrolled into the study from January to December 2019, out of which 97 (55.7%) also participated in HPV self-sampling (Fig. 1). Their median age was 45 years, ranging between 23 and 65 years (Table 1). The participants were mainly of Malay ethnicity (92.5%, n = 161), married (90.2%, n = 157), and had ≥ 3 births (60.3%, n = 105). About half of the participants had their last Pap test performed between 4 and 10 years ago (52.3%, n = 91), and have never received the HPV vaccine (54.6%, n = 95). There were significant differences between those who agreed and declined to participate in HPV self-sampling: those who declined were significantly older (p = 0.002) and more likely to have their last Pap test performed > 10 years ago (p = 0.031).

Fig 1. Flowchart of study participants at JPHSC, Brunei (Jan – Dec 2019).

190   Table 1. Sociodemographic characteristics of the study population, including comparison between

191   groups that agreed and declined to join HPV self-sampling. IQR = Interquartile range.

Characteristics		Total study population (n = 174) n (%)	Joined self-sampling (n = 97) n (%)	Declined self-sampling (n = 77) n (%)	p-value
Median age in years (IQR)		45.0 (15.25)	41.0 (17)	49.0 (14.5)	**0.002***
Age-group (in years)	20 - 24	3 (1.7)	3 (3.1)	0 (0.0)	**0.018***
	25 - 29	20 (11.5)	15 (15.5)	5 (6.5)	
	30 - 34	14 (8.1)	10 (10.3)	4 (5.2)	
	35 - 39	28 (16.1)	18 (18.5)	10 (13.0)	
	40 - 44	18 (10.3)	11 (11.3)	7 (9.0)	
	45 - 49	27 (15.5)	12 (12.4)	15 (19.5)	
	50 - 54	35 (20.1)	15 (15.5)	20 (26.0)	
	55 - 59	14 (8.1)	10 (10.3)	4 (5.2)	
	> 60	13 (7.5)	3 (3.1)	10 (13.0)	
	Missing	2 (1.1)	0 (0.0)	2 (2.6)	
Race	Malay	161 (92.5)	90 (92.8)	71 (92.2)	0.752
	Chinese	6 (3.5)	4 (4.1)	2 (2.6)	
	Other	7 (4.0)	3 (3.1)	4 (5.2)	
Education level	Primary school	16 (9.2)	10 (10.3)	6 (7.8)	0.396
	Secondary school	96 (55.2)	49 (50.5)	47 (61.0)	
	College / University	57 (32.7)	35 (36.1)	22 (28.6)	
	Missing	5 (2.9)	3 (3.1)	2 (2.6)	
Marital status	Married	157 (90.2)	91 (93.8)	66 (85.7)	0.15
	Divorced	8 (4.6)	2 (2.1)	6 (7.8)	
	Widowed	9 (5.2)	4 (4.1)	5 (6.5)	
Occupation	Housewife	64 (36.8)	39 (40.2)	25 (32.4)	0.087
	Government employee	67 (38.5)	41 (42.3)	26 (33.8)	
	Private employee	31 (17.8)	13 (13.4)	18 (23.4)	
	Retired	9 (5.2)	3 (3.1)	6 (7.8)	
	Unemployed	1 (0.6)	1 (1.0)	0 (0.0)	
	Other	2 (1.1)	0 (0.0)	2 (2.6)	
Monthly household income	< $500	27 (15.5)	15 (15.5)	12 (15.6)	0.498
	$500 < $999	27 (15.5)	12 (12.4)	15 (19.5)	
	$1000-$1999	40 (23.0)	22 (22.7)	18 (23.4)	
	$2000-$2999	19 (10.9)	14 (14.3)	5 (6.5)	
	$3000-$5000	24 (13.8)	15 (15.5)	9 (11.7)	
	>$5000	3 (1.7)	2 (2.1)	1 (1.3)	
	Missing	34 (19.5)	17 (17.5)	17 (22.0)	
Number of births	0	27 (15.5)	17 (17.5)	10 (13.0)	0.103
	1	19 (10.9)	14 (14.4)	5 (6.5)	

	2	22 (12.1)	14 (14.4)	7 (9.1)	
	3 or more	105 (60.3)	51 (52.6)	54 (70.1)	
	Missing	2 (1.2)	1 (1.1)	1 (1.3)	
Last Pap test done	Never	41 (23.6)	29 (29.9)	12 (15.6)	**0.031***
	4-10 years	91 (52.3)	48 (49.5)	43 (55.8)	
	> 10 years	36 (20.7)	15 (15.5)	21 (27.3)	
	Missing	6 (3.4)	5 (5.1)	1 (1.3)	
Completed HPV vaccination	Never	95 (54.6)	50 (51.6)	45 (58.4)	0.483
	Yes	27 (15.5)	16 (16.5)	11 (14.3)	
	No	41 (23.6)	26 (26.8)	15 (19.5)	
	Missing	11 (6.3)	5 (5.1)	6 (7.8)	

Fig. 2 and S1 Table show the responses for the major and minor reasons for not attending cervical cancer screening. The top three major reasons reported were that they were "afraid of getting a bad result" (16.1%, n = 28), "feeling embarrassed being examined by a doctor or nurse" (14.9%, n = 26), and "I can't find the time as I'm too busy at home" (10.3%, n = 18). The top three minor reasons were "feeling embarrassed being examined by a doctor or nurse" (20.7%, n = 36), "I can't find time as I'm too busy at work" (20.7%, n = 36), and "afraid of getting a bad result" (20.1%, n = 35).

Fig. 2. Responses on their major and minor reasons for not attending cervical cancer screening, among non-attendees at JPSHC (Jan – Dec 2019). The x-axis indicates the percentage, and the number next to each bar indicates the actual number of responses.

When comparing top 10 major reasons for not attending screening between those who agreed and declined HPV self-sampling (Table 2), those whose major reason was "feeling embarrassed being examined by a doctor or nurse" were significantly more likely to join self-sampling (p = 0.020). Also, those whose major reason was "afraid of getting a bad result" were significantly more likely to decline self-sampling (p = 0.034). Among those who were employed (n = 98), about a quarter (23.5%, n = 23) reported work-related reasons as their major reason for not attending screening ("I can't find the time as I'm too busy at work" and "Difficult to get permission from employer").

212   Table 2. The top ten major reasons of not attending cervical cancer screening at JPSHC, Brunei (Jan –

213   Dec 2019), with comparison between those who agreed and declined HPV self-sampling. Responses

214   from 146 participants (83.9% of the total study population) were included.

No.	Top ten major reasons of screening non-attendance	Total n (%)	Joined HPV self-sampling n (%)	Declined HPV self-sampling n (%)	*p*-value
1	I feel embarrassed being examined by a doctor or nurse	26 (14.9)	20 (76.9)	6 (23.1)	**0.020***
2	I am scared of pain because of previous bad experience(s)	16 (9.2)	8 (50.0)	8 (50.0)	0.825
3	I am scared of getting a bad result	28 (16.1)	10 (35.7)	18 (64.3)	**0.034***
4	I can't find the time as I'm too busy at home	18 (10.3)	12 (66.7)	6 (33.3)	0.463
5	I can't find the time as I'm too busy at work	16 (9.2)	10 (62.5)	6 (37.5)	0.759
6	Difficult to get permission from employer	8 (4.6)	5 (62.5)	3 (37.5)	1
7	I have never heard of a Pap Test	7 (4.0)	4 (57.1)	3 (42.9)	1
8	I have forgotten about it	6 (3.4)	5 (83.3)	1 (16.7)	0.229
9	I did not receive any invitation	6 (3.4)	5 (83.3)	1 (16.7)	0.229
10	Others	15 (8.6)	7 (46.7)	8 (53.3)	0.492

While the responses vary when asked about their health and susceptibility to disease, most agreed

on the benefits of undergoing Pap test (92.0%, n = 160), and on the severity of cervical cancer

(82.8%, n = 144; Table 3 and S2 Table). Those who agreed that they are more susceptible to develop

cervical cancer were significantly more likely to join HPV self-sampling (*p* = 0.023).

Table 3. Attitudes towards cervical cancer screening among non-attendees at JPSHC, Brunei (Jan–

Dec 2019), between those who agreed and declined HPV self-sampling.

No.	Attitude questions		Total study population (n = 174) n (%)	Joined HPV self-sampling (n = 97) n (%)	Declined HPV self-sampling (n = 77) n (%)	*p*-value
1	I believe I am healthy and free of any diseases	Agree	62 (35.6)	34 (54.8)	28 (45.2)	0.984
		Neutral/ Disagree	112 (64.3)	63 (56.3)	49 (43.7)	
2	Having Pap test taken is beneficial for my health	Agree	160 (92.0)	91 (56.9)	69 (43.1)	0.464
		Neutral/ Disagree	14 (8.0)	6 (42.9)	8 (57.1)	
3	Like any women, I am susceptible to develop cervical cancer	Agree	110 (63.2)	69 (62.7)	41 (37.3)	**0.023***
		Neutral/ Disagree	64 (36.8)	28 (43.8)	36 (56.2)	
4	Cervical cancer can be severe and may lead to death	Agree	144 (82.8)	79 (54.9)	65 (45.1)	0.754
		Neutral/ Disagree	30 (17.2)	18 (60.0)	12 (40.0)	

Most participants would like to obtain more information about cervical cancer screening (85.6%, n =

149). The top preferred information sources were healthcare workers (59.8%, n = 104), social

messaging platforms (28.7%, n = 50) and social media (26.4%, n = 46; S1 Fig).

Among those who participated in self-sampling (55.7%, n = 97), their responses on Q2 (Table 4) were

generally positive and encouraging. A majority agreed that the instructions were clear (94.8%, n =

92), that it was easy to perform the swab (93.8%, n = 91), and that it was more convenient than the

Pap test (91.7%, n = 89). They also reported their confidence in correctly getting the sample (92.8%,

n = 90), would prefer to use this method next time (94.8%, n = 92), and would recommend this

method to other women (93.8%, n = 91). Notably, 54.6% (n = 53) still prefer a proper Pap test for

their subsequent check-up.

Table 4. Acceptability of HPV self-sampling among non-attendees who participated in HPV self-

sampling at JPSHC, Brunei (Jan–Dec 2019). Responses from all Q2 respondents (n = 97) were

included.

No.	Self-sampling Questions	Agree n (%)	Neutral n (%)	Disagree n (%)
1	I thought the instructions were clear	92 (94.8)	0 (0.0)	5 (5.2)
2	It was easy to do the swab	91 (93.8)	1 (1.0)	5 (5.2)
3	Taking the sample with the swab was painful	12 (12.4)	8 (8.2)	77 (79.4)
4	Taking the sample was uncomfortable to do	11 (11.4)	4 (4.1)	82 (84.5)
5	I felt embarrassed doing the self-sampling	9 (9.3)	1 (1.0)	87 (89.7)
6	It was convenient to do without having to undergo a Pap Test	89 (91.7)	3 (3.1)	5 (5.2)
7	I am confident I did it correctly	90 (92.8)	4 (4.1)	3 (3.1)
8	I want to use this method next time	92 (94.8)	2 (2.1)	3 (3.1)
9	I prefer to do this at home	64 (66.0)	15 (15.5)	18 (18.5)
10	I would recommend this method to other women	91 (93.8)	3 (3.1)	3 (3.1)
11	I trust that the result of this self-sampling will be accurate	74 (76.3)	19 (19.6)	4 (4.1)
12	I would like to attend for a proper Pap test in clinic next time	53 (54.6)	23 (23.7)	21 (21.7)

Among the 97 samples taken, nine (9.3%) tested positive for hr-HPV: one was positive for HPV 16

and eight were positive for non-HPV 16/18 HPV genotype. Upon follow-up cytology, only one

participant had a normal cytological test result, whilst another participant with non-16/18 HPV

genotype was found to have CIN 3 with glandular involvement following a confirmatory biopsy. No

significant differences were observed when comparing the sociodemographic characteristics

between those who tested hr-HPV positive and those who tested hr-HPV negative (S3 Table).

Discussion

This preliminary cross-sectional study explored the reasons for cervical screening non-attendance

among Bruneian women, and whether HPV self-sampling is a viable alternative for the non-

attendees. Our findings highlight three important points to consider for improving screening uptake

and cervical cancer detection.

First, our findings suggest that it is necessary to provide accurate information among Bruneian

women on cervical cancer, the importance of screening and addressing any misconceptions about

the Pap test. Important facts to relay include the slow development from pre-cancerous changes to

cervical cancer, that pre-cancerous changes are highly treatable, and that screening will help in early

detection and thus facilitate successful treatment.

In our study, emotional barriers (fear of unfavorable test results and embarrassment) were the most

common major reasons of screening non-attendance. Also, about two-thirds of our participants

cited "I am scared of getting a bad result" as their main barrier also declined to take part in HPV self-

sampling, possibly due to relating abnormal Pap test results to cervical cancer diagnoses [23].

Although such barriers may play a large role at the onset of screening program, its role diminishes

over time with increasing education on the benefits of screening [24]. Educational interventions

could also benefit the small group of women who cited menopause, cessation of child-bearing and

having had HPV vaccination as reasons for not attending screening.

In our study, we observed those who agree that they are susceptible to cervical cancer were

significantly more likely to participate in HPV self-sampling. This suggests that perceived

susceptibility could be an important factor for self-sampling participation, whereby those who do

not perceive themselves as susceptible were less likely to engage in preventative behaviors [25–27].

Perceived susceptibility can be increased through education to improve their beliefs on the

importance of screening [28]. We suggest that such information could be more effectively

disseminated as simple health messages endorsed by Brunei's Ministry of Health via website and

social media platforms.

Second, we observed high acceptability of HPV self-sampling among our participants. Meta-analyses

have indicated strong acceptance of and preference for self-sampling over clinician sampling [29],

mainly due to logistical reasons [30]. More than half of our participants who joined HPV self-

sampling cited embarrassment or lack of time due to home and/or work commitments as their

major reason of not attending Pap test screening. This suggest that providing flexibility to

accommodate women's screening method preference [31], such as the option of self-sampling [32],

could improve screening uptake.

Thirdly, most of the detected hr-HPV genotypes in our study were non-16/18, with only one out of

nine participants tested positive for HPV 16. Although our sample size is small, this result suggests

that it may not be accurate to assume HPV 16 or 18 as common hr-HPV genotypes in Brunei, even

though this is true in the global context [33]. Other studies have detected a significant percentage of

non-16/18 hr-HPV genotypes, suggesting the presence of region-specific heterogeneity in the HPV

genotype distribution [33–37]. Also, variation in HPV distribution among different ethnic groups has

been reported in an American study [38]. Our finding has potential implication on Brunei's national

school-based HPV vaccination program which currently provides vaccines which do not confer

protection against non-16/18 genotypes. Larger population-based studies to understand the

distribution of HPV genotypes among Bruneian women will be crucial to determine the efficacy or

impact of the current vaccines.

One notable point for the local context relates to the presence of cultural barriers. Being a

predominantly Islamic society, religious and cultural modesty could be a contributing factor for

embarrassment among Muslim women [39]. Also, having premarital sex is a cultural taboo in Brunei

and is generally not openly discussed [40]. This could prevent any unmarried but sexually active

women from participating in the screening program. It should be emphasized that only married or

ever married females were included in this study; included because it is part of the eligibility criteria

for the national cervical cancer screening program in Brunei.

A major limitation for this study is that non-attendees from only one health center were recruited,

thus our findings are not representative of the adult female population in Brunei. Secondly, our

findings should also be interpreted with caution due to the small sample size and the non-probability

sampling approach used. This study was initially conceived as a preliminary study due to resource

and logistics limitations. There were two reasons for choosing JPSHC as our study site: It is the third

largest primary government healthcare center in the country, and that it serves a sub-district with a

sizable percentage of residents in the middle- to low-income groups. It is hoped that even at this

preliminary stage, our study findings could encourage stakeholders to conduct similar and larger

studies, using random sampling approach. Lastly, as this study relied on the self-reported history of

previous Pap test attendance, we might have missed recruiting those who may have forgotten their

last Pap test date.

In conclusion, our findings indicate the need to further promote knowledge on cervical cancer, the

benefits of screening and clarifying any misconceptions of Pap test results. Reasons of cervical

cancer screening non-attendance were mainly related to emotional and logistical factors. As we

found high acceptance towards HPV self-sampling, this could be adopted as an alternative for

women who refrain from Pap test. Our high detection of non-16/18 HPV genotypes suggest high

prevalence of other hr-HPV genotypes in Brunei. Future larger studies involving more Bruneian

women should be done to verify our results. Follow-up studies should also be conducted to consider

HPV testing as the suitable method for cervical cancer screening.

Acknowledgements

The authors would like to thank the JPSHC staff who have assisted in participant recruitment and

questionnaire distribution.

References

1. World Health Organization. Cervical cancer. [cited 8 Sep 2020]. Available: https://www.who.int/health-topics/cervical-cancer#tab=tab_1

2. Gustafsson L, Pontén J, Zack M, Adami H-O. International incidence rates of invasive cervical cancer after introduction of cytological screening. Cancer Causes Control. 1997;8: 755–763. doi:10.1023/A:1018435522475

3. Yeh PT, Kennedy CE, Vuyst H de, Narasimhan M. Self-sampling for human papillomavirus (HPV) testing: a systematic review and meta-analysis. BMJ Glob Heal. 2019;4: e001351. doi:10.1136/BMJGH-2018-001351

4. CS R, DR W, D G. Self-collected HPV testing improves participation in cervical cancer screening: a systematic review and meta-analysis. Can J Public Health. 2013;104. doi:10.1007/BF03405681

5. Maza M, Melendez M, Masch R, Alfaro K, Chacon A, Gonzalez E, et al. Acceptability of self-sampling and human papillomavirus testing among non-attenders of cervical cancer screening programs in El Salvador. Prev Med (Baltim). 2018. doi:10.1016/j.ypmed.2018.06.017

6. Sancho-Garnier H, Tamalet C, Halfon P, Leandri FX, Retraite L Le, Djoufelkit K, et al. HPV self-sampling or the Pap-smear: A randomized study among cervical screening nonattenders from lower socioeconomic groups in France. Int J Cancer. 2013. doi:10.1002/ijc.28283

7. Arrossi S, Thouyaret L, Herrero R, Campanera A, Magdaleno A, Cuberli M, et al. Effect of self-collection of HPV DNA offered by community health workers at home visits on uptake of screening for cervical cancer (the EMA study): A population-based cluster-randomised trial. Lancet Glob Heal. 2015;3: e85–e94. doi:10.1016/S2214-109X(14)70354-7

8. Gustavsson I, Aarnio R, Berggrund M, Hedlund-Lindberg J, Strand A-S, Sanner K, et al. Randomised study shows that repeated self-sampling and HPV test has more than two-fold

higher detection rate of women with CIN2+ histology than Pap smear cytology. Br J Cancer.

2018;118: 896–904. doi:10.1038/bjc.2017.485

9.   Arrossi S, Paolino M, Laudi R, Gago J, Campanera A, Marín O, et al. Programmatic human

papillomavirus testing in cervical cancer prevention in the Jujuy Demonstration Project in

Argentina: a population-based, before-and-after retrospective cohort study. Lancet Glob

Heal. 2019. doi:10.1016/S2214-109X(19)30048-8

10.   Chrysostomou AC, Stylianou DC, Constantinidou A, Kostrikis LG. Cervical cancer screening

programs in Europe: The transition towards HPV vaccination and population-based HPV

testing. Viruses. 2018;10. doi:10.3390/v10120729

11.   Landy R, Pesola F, Castañón A, Sasieni P. Impact of cervical screening on cervical cancer

mortality: Estimation using stage-specific results from a nested case-control study. Br J

Cancer. 2016;115: 1140–1146. doi:10.1038/bjc.2016.290

12.   Canfell K, Sitas F, Beral V. Cervical cancer in Australia and the United Kingdom: Comparison of

screening policy and uptake, and cancer incidence and mortality. Med J Aust. 2006;185: 482–

486. doi:10.5694/j.1326-5377.2006.tb00661.x

13.   Ministry of Health Brunei Darussalam. Health Information Booklet 2017 (Second Revision).

2019 [cited 11 Feb 2020]. Available: http://moh.gov.bn/SitePages/Health Information

Booklet.aspx

14.   Arbyn M, Weiderpass E, Bruni L, de Sanjosé S, Saraiya M, Ferlay J, et al. Estimates of

incidence and mortality of cervical cancer in 2018: a worldwide analysis. Lancet Glob Heal.

2020;8: e191–e203. doi:10.1016/S2214-109X(19)30482-6

15.   Chaw LL, Lim STW, Md Yussof SR. Human Papillomavirus vaccine coverage among female

students in Brunei Darussalam: results from the first 4 years of the national school-based

vaccination programme. Heliyon. 2019;5. doi:10.1016/j.heliyon.2019.e02588

16.   Peto PJ, Gilham PC, Fletcher O, Matthews FE. The cervical cancer epidemic that screening has

prevented in the UK. Lancet. 2004;364: 249–256. doi:10.1016/S0140-6736(04)16674-9

17.   Bosgraaf RP, Ketelaars PJW, Verhoef VMJ, Massuger LFAG, Meijer CJLM, Melchers WJG, et al.

Reasons for non-attendance to cervical screening and preferences for HPV self-sampling in

Dutch women. Prev Med (Baltim). 2014. doi:10.1016/j.ypmed.2014.04.011

18.   , Nieminen P, Niironen M, Luostarinen T, Anttila A. Self-sampling experiences among non-

attendees to cervical screening. Gynecol Oncol. 2014. doi:10.1016/j.ygyno.2014.09.019

19.   Bayu H, Berhe Y, Mulat A, Alemu A. Cervical Cancer Screening Service Uptake and Associated

Factors among Age Eligible Women in Mekelle Zone, Northern Ethiopia, 2015: A Community

Based Study Using Health Belief Model. PLoS One. 2016;11: e0149908.

doi:10.1371/journal.pone.0149908

20.   Sultana F, Mullins R, English DR, Simpson JA, Drennan KT, Heley S, et al. Women's experience

with home-based self-sampling for human papillomavirus testing. BMC Cancer. 2015;15: 849.

doi:10.1186/s12885-015-1804-x

21.   The Hague: Health Council of the Netherlands. Health Council of the Netherlands Population

screening for cervical cancer. 2011;2011/07E. Available:

https://www.gezondheidsraad.nl/sites/default/files/201107E_PopulationSCC_0.pdf

22.   R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria; 2020. Available: http://www.r-project.org/

23.   Fylan F. Screening for cervical cancer: a review of women's attitudes, knowledge, and

behaviour. Br J Gen Pract. 1998;48: 1509–14.

24.   Teng FF, Mitchell SM, Sekikubo M, Biryabarema C, Byamugisha JK, Steinberg M, et al.

Understanding the role of embarrassment in gynaecological screening: A qualitative study

from the ASPIRE cervical cancer screening project in Uganda. BMJ Open. 2014.

doi:10.1136/bmjopen-2014-004783

25.   Agurto I, Arrossi S, White S, Coffey P, Dzuba I, Bingham A, et al. Involving the community in

cervical cancer prevention programs. Int J Gynaecol Obstet. 2005;89 Suppl 2: S38-45.

doi:10.1016/j.ijgo.2005.01.015

26.   Shirazi Zadeh Mehraban S, Namdar A, Naghizadeh MM. Assessment of Preventive Behavior

for Cervical Cancer with the Health Belief Model. Asian Pac J Cancer Prev. 2018;19: 2155–

2163. doi:10.22034/APJCP.2018.19.8.2155

27.   Tapera R, Manyala E, Erick P, Maswabi TM, Tumoyagae T, Letsholo B, et al. Knowledge and

Attitudes towards Cervical Cancer Screening amongst University of Botswana Female

Students. Asian Pac J Cancer Prev. 2017;18: 2445–2450. doi:10.22034/APJCP.2017.18.9.2445

28.   Khademolhosseini F, Noroozi A, Tahmasebi R. The Effect of Health Belief Model-Based

Education through Telegram Instant Messaging Services on Pap smear performance. Asian

Pac J Cancer Prev. 2017;18: 2221–2226. doi:10.22034/APJCP.2017.18.8.2221

29.   Nelson EJ, Maynard BR, Loux T, Fatla J, Gordon R, Arnold LD. The acceptability of self-sampled

screening for HPV DNA: A systematic review and meta-analysis. Sexually Transmitted

Infections. 2017. doi:10.1136/sextrans-2016-052609

30.   Lorenzi NPC, Termini L, Longatto Filho A, Tacla M, de Aguiar LM, Beldi MC, et al. Age-related

acceptability of vaginal self-sampling in cervical cancer screening at two university hospitals:

a pilot cross-sectional study. BMC Public Health. 2019. doi:10.1186/s12889-019-7292-1

31.   Bansil P, Wittet S, Lim JL, Winkler JL, Paul P, Jeronimo J. Acceptability of self-collection

sampling for HPV-DNA testing in low-resource settings: a mixed methods approach. BMC

Public Health. 2014;14: 596. doi:10.1186/1471-2458-14-596

32.   Jeronimo J, Perkins RB, Scalici J, Pierce JY. Should Self-Sampling Be an Option for Women in

the United States? J Low Genit Tract Dis. 2019;23: 54. doi:10.1097/LGT.0000000000000453

33.   Albawardi A, Quddus MR, Al Awar S, Almarzooqi S. Frequency of rare and multi viral high-risk

HPV types infection in cervical high grade squamous intraepithelial lesions in a non-native

dominant middle eastern country: A polymerase chain reaction-based pilot study. Diagn

Pathol. 2018. doi:10.1186/s13000-018-0716-x

34.   Zeng Z, Yang H, Li Z, He X, Griffith CC, Chen X, et al. Prevalence and Genotype Distribution of

HPV Infection in China: Analysis of 51,345 HPV Genotyping Results from China's Largest CAP

Certified Laboratory. J Cancer. 2016. doi:10.7150/jca.14971

35.   Khoo SP, Bhoo-Pathy N, Yap SH, Anwar Shafii MK, Hairizan Nasir N, Belinson J, et al.

Prevalence and sociodemographic correlates of cervicovaginal human papillomavirus (HPV)

carriage in a cross-sectional, multiethnic, community-based female Asian population. Sex

Transm Infect. 2018. doi:10.1136/sextrans-2017-053320

36.   Peng RR, Li HM, Chang H, Li JH, Wang AL, Chen XS. Prevalence and genotype distribution of

cervical human papillomavirus infection among female sex workers in Asia: A systematic

literature review and meta-analysis. Sexual Health. 2012. doi:10.1071/SH11066

37.   Al-Awadhi R, Chehadeh W, Jaragh M, Al-Shaheen A, Sharma P, Kapila K. Distribution of

human papillomavirus among women with abnormal cervical cytology in Kuwait. Diagn

Cytopathol. 2013. doi:10.1002/dc.21778

38.   Keller MJ, Burk RD, Massad LS, Eltoum IE, Hessol NA, Anastos K, et al. Racial differences in

human papilloma virus types amongst United States women with HIV and cervical precancer.

AIDS. 2018. doi:10.1097/QAD.0000000000002005

39.   Ali S, Skirton H, Clark MT, Donaldson C. Integrative review of cervical cancer screening in

Western Asian and Middle Eastern Arab countries. Nurs Heal Sci. 2017;19: 414–426.

doi:10.1111/nhs.12374

40.   Haji Tahamit N. Cultural and Religious Barriers to Setting Up Sex and Relationship Education

in a Muslim Country: A Case Study of Brunei Darussalam. University of Leeds; 2015 Aug.

Available: https://etheses.whiterose.ac.uk/15306/

Supplementary information

S1 Table. Responses on their major and minor reasons for not attending cervical cancer screening

among non-attendees at JPSHC, Brunei (Jan – Dec 2019).

S2 Table. Attitudes towards cervical cancer screening among non-attendees at JPSHC, Brunei (Jan –

Dec 2019). Responses from the total study population (n = 174) were included.

S1 Fig. Preferred sources of information about cervical cancer among non-attendees at JPSHC,

Brunei (Jan–Dec 2019). The x-axis indicates the percentage, and the number next to each bar

indicates the number of responses. Multiple responses were allowed and responses from the total

study population (n = 174) were included.

S3 Table. Socio demographic characteristics and comparison between screening non-attendees who

tested positive and negative for hr-HPV at JPSHC, Brunei (Jan – Dec 2019).

S4 Questionnaire. Questionnaire 1: Reasons why women do not participate in the National Cervical

Cancer Screening Program

S5 Questionnaire. Questionnaire 2: Acceptability of Self Sampling for HPV Testing Among Non-

attendees of Cervical Screening Program in Brunei Darussalam

S6 Dataset. Minimal dataset

497
498
Author contributions:

Liling Chaw: Conceptualization, Methodology, Data curation, Formal analysis, Supervision, Writing -

original draft, Writing - review & editing

Shirley HF Lee: Visualization, Writing - original draft, Writing - review & editing

Nurul Iffah Hazwani Ja'afar: Investigation, Data curation, Writing - original draft

Edwin Lim: Conceptualization, Methodology, Resources, Funding acquisition, Supervision, Writing -

review & editing

Roslin Sharbawi: Conceptualization, Methodology, Resources, Funding acquisition, Investigation,

Project administration, Supervision, Writing - original draft, Writing - review & editing

Category	Main reason	Minor reason
I am scared of getting a bad result	28	35
I feel embarrassed being examined by a doctor or nurse	26	36
I can't find the time as I'm too busy at home	18	28
I am scared of pain because of previous bad experience(s)	13	16
I can't find the time as I'm too busy at work	16	36
Difficult to get permission from employer	8	14
I have never heard of a Pap Test	7	8
I did not receive any invitation	0	6
I have forgotten about it	2	6
Not necessary as I don't have menses anymore	4	11
Not necessary as I am not child-bearing anymore	4	14
Not necessary as I am healthy	4	15
Nobody is looking after child(ren) at home	2	13
Nobody to send me to clinic	2	11
Not necessary as I have already had my HPV vaccination	2	2
I don't know what cervical cancer is	1	15
Not necessary as I don't have a partner anymore	0	1
Others	8	4
Missing	9	