# Neutralizing activity of Sputnik V vaccine sera against SARS-CoV-2 variants

Satoshi Ikegame[1,#], Mohammed N. A. Siddiquey[2,#], Chuan-Tien Hung[1], Griffin Haas[1], Luca Brambilla[1], Kasopefoluwa Y. Oguntuyo[1], Shreyas Kowdle[1], Ariel Esteban Vilardo[3], Alexis Edelstein[3], Claudia Perandones[3,†], Jeremy P. Kamil[2,†], and Benhur Lee[1,†,*]

**Affiliations.**

1. Department of Microbiology at the Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

2. Department of Microbiology and Immunology, Louisiana State University Health Shreveport, Shreveport, LA 71103, USA.

3. National Administration of Laboratories and Health Institutes of Argentina (ANLIS) Dr. Carlos G. Malbrán, Buenos Aires, Argentina

# These authors contributed equally to the study.

† Senior authors

* Correspondence to: Benhur Lee, benhur.lee@mssm.edu

**Competing interests:** B.L. and K.Y.O. are named inventors on a patent filed by the Icahn School of Medicine for some of the materials used in this work. J.P.K. is a consultant for BioNTech (advisory panel on coronavirus variants).

**ABSTRACT.**

The novel pandemic betacoronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has infected at least 120 million people since its identification as the cause of a December 2019 viral pneumonia outbreak in Wuhan, China. Despite the unprecedented pace of vaccine development, with six vaccines already in use worldwide, the emergence of SARS-CoV-2 'variants of concern' (VOC) across diverse geographic locales suggests herd immunity may fail to eliminate the virus. All three officially designated VOC carry Spike (S) polymorphisms thought to enable escape from neutralizing antibodies elicited during initial waves of the pandemic. Here, we characterize the biological consequences of the ensemble of S mutations present in VOC lineages B.1.1.7 (501Y.V1) and B.1.351 (501Y.V2). Using a replication-competent EGFP-reporter vesicular stomatitis virus (VSV) system, rcVSV-CoV2-S, which encodes S from SARS coronavirus 2 in place of VSV-G, and coupled with a clonal HEK-293T ACE2 TMPRSS2 cell line optimized for highly efficient S-mediated infection, we determined that only 1 out of 12 serum samples from a cohort of recipients of the Gamaleya Sputnik V Ad26 / Ad5 vaccine showed effective neutralization ($IC_{90}$) of rcVSV-CoV2-S: B.1.351 at full serum strength. The same set of sera efficiently neutralized S from B.1.1.7 and showed only moderately reduced activity against S carrying the E484K substitution alone. Taken together, our data suggest that control of some emergent SARS-CoV-2 variants may benefit from updated vaccines.

## INTRODUCTION.

In the 15 months since its emergence in late 2019 [1], SARS-CoV-2 has caused over 131 million confirmed COVID-19 cases worldwide, leading to at least 2.85 million deaths [2]. SARS-CoV-2 is closely related to two other zoonotic betacoronaviruses, MERS-CoV and SARS-CoV, that also cause life-threatening respiratory infections [3].

This global health emergency has spurred the development of COVID-19 preventive vaccines at an unprecedented pace. Six are already authorized for human use across the globe [4-9]. These vaccines focus on the SARS-CoV-2 spike protein (S), due to its critical roles in cell entry. Indeed, the presence of serum neutralizing antibodies directed at S correlate strongly with protection against COVID-19 [10,11]. Although these six vaccines are efficacious, the recent emergence of novel SARS-CoV-2 variants has reignited concerns that the pandemic may not be so easily brought under control.

In December 2020, the United Kingdom reported the sudden emergence of a novel SARS-CoV-2 lineage, termed B.1.1.7 (501Y.V1, VOC 202012/01), which was designated as the first SARS-CoV-2 variant of concern (VOC) . The lineage had rapidly increased in prevalence since first being detected in November 2020 [12]. Its genome showed an unusually high number of non-synonymous substitutions and deletions, including eight in the S gene, suggesting a substantial degree of host adaptation that may have occurred during prolonged infection of an immunocompromised person [13]. The B.1.1.7 lineage has now been shown to exhibit enhanced transmissibility [14] as well as an increased case fatality rate [15,16].

Soon afterwards, two additional SARS-CoV-2 VOC, B.1.351 and P.1, were reported from S. Africa and Brazil, respectively, which each showed substantial escape from neutralizing antibodies elicited by first wave pandemic viruses, leading to documented cases of re-infection [17-19]. The S genes of B.1.351 and P.1 viruses each carry a number of mutations, but include three in the receptor binding domain (RBD) that are particularly notable, the S: N501Y substitution, found in B.1.1.7, alongside polymorphisms at positions 417 and 484, K417N/T and E484K. S: E484K had already been identified in multiple independent laboratories to confer escape from convalescent sera and monoclonal antibodies [20-22]. As expected, the P.1 and B.1.351 variants escape or resist neutralization by first wave convalescent sera, as well as antibodies elicited by COVID-19 vaccines [23-27].

Although the P.1 and B.1.351 lineages are dominant in Brazil and S. Africa, unlike B.1.1.7 they have not increased greatly in number in the United States since originally being detected here. In contrast, the E484K polymorphism is recurrently emergent, and is found in a number of other lineages that are increasing in the U.S. and other countries. For example, a B.1.526 sub-lineage carrying E484K in recent weeks has expanded more rapidly than B.1.1.7 [28,29], which may be indicative of the ability of S: E484K variants to penetrate herd immunity. The P.2 lineage, originally

detected in Rio de Janeiro, carries only the E484K mutation in the RBD and has spread
to other parts of South America, including Argentina [30].

The six COVID-19 vaccines currently in use around the world employ different
strategies, and do not all incorporate the two proline substitutions that "lock" S into the
pre-fusion conformer.  Vaccines that do not utilize pre-fusion "locked" S are expected
to produce lower levels of neutralizing antibodies, and hence may be less efficacious
against infection, even if they do protect against severe COVID-19[31,32].  Indeed, a two-
dose regimen of the AstraZeneca ChAdOx1 based vaccine, which does not use a
"locked" S, did not protect against mild-to-moderate COVID-19 in S. Africa, where
93% of COVID-19 cases in trial participants were caused by the B.1.351 variant [33]. Like
the AstraZeneca ChAdOx1 vaccine, the Sputnik V vaccine (Gam-COVID-Vac) is based
on adenovirus vectored expression of a native S sequence, rather than a pre-fusion
"locked" S [34]. Although the Sputnik V vaccine has a reported vaccine efficacy of 91.6%
in the interim analysis of Phase 3 trials held in Russia between Sept 7 and Nov 24,
2020, none of the VOC mentioned above nor independent lineages containing the
E484K mutation were prevalent in Russia during this time period. Since the Sputnik
vaccine is now in use not only in Russia, but also in countries like Argentina, Mexico,
and Hungary, where some of the VOC and emerging lineages bearing the E484K
mutation are more widespread, it is critical to assess the neutralizing activity of Sputnik
vaccine elicited antibody responses against these cognate VOC and mutant spikes.

This study characterizes the neutralization activity of sera from a dozen Sputnik
V vaccine recipients in Argentina. Our work was spurred by Argentina's nascent
genomic surveillance efforts, which detected multiple independent lineages with S:
E484K (B.1.1.318 and P.2) and/or S: N501Y substitutions (B.1.1.7 and P.1) in common,
just as Argentina had started rolling out its vaccination campaign, which commenced
on Dec 29, 2020. Here, we generated isogenic replication-competent vesicular
stomatitis virus bearing the prevailing wild-type (WT=D614G) SARS-CoV-2 S (rcVSV-
CoV2-S), or the B.1.1.7, B.1.351 or E484K mutant S and used them in a robust virus
neutralization assay. Our results show that Sputnik V vaccine sera effectively
neutralized S: WT and S: B.1.1.7. viruses, albeit with highly variable titers. The same
sera, however, exhibited moderate and markedly reduced neutralization titers,
respectively, against S: E484K and S: B.1.351. Analyses of dose response curves
indicate that S: B.1.351 exhibits resistance to neutralizing sera in a manner that is
qualitatively different from the E484K mutant. Taken together, our data argue that
surveillance of the neutralizing activity elicited by vaccine sera will be necessary on an
ongoing basis. Viral neutralization assays can indicate which SARS-CoV-2 variants are
likely capable of transmission in the face of vaccine elicited immunity, and whether
updated vaccines will be needed to control their emergence and spread.

## RESULTS.

### Robust reverse genetics for generating replication-competent VSV expressing SARS-CoV-2 Spike proteins.

Several groups have now generated replication-competent VSV expressing SARS-CoV-2 spike in place of VSV-G (rcVSV-CoV2-S)[35–37,38]. These rcVSV-CoV2-S can be used in BSL-2 compatible virus neutralization assays (VNAs), which correlate very well with VNAs using live SARS-CoV-2 (Spearman's r > 0.9 across multiple studies). rcVSV-CoV2-S has been assessed as a candidate vaccine [37,39], and used in forward genetics experiments to generate antibody escape mutants or perform comprehensive epitope mapping studies [40,20,38]. Indeed, the now concerning E484K mutation, present in many variants of concern (VOC), was identified as an antibody escape mutation using rcVSV-CoV-2-S [20,38].

However, many groups passage their rcVSV-CoV-2-S extensively in Vero cells after the initial rescue, either to generate higher titer stocks and/or to remove confounding components such as the vaccinia virus expressing T7-polymerase and/or transfected VSV-G, both of which were deemed necessary for efficient rescue [38]. Serial passage of rcVSV-CoV-2-S in Vero cells invariably leads to mutations in the S1/S2 furin cleavage site, as well as truncations in the cytoplasmic tail of the S protein [39]. The latter promotes S incorporation into VSV without compromising the conformational integrity of the ectodomain, whereas the former is problematic when assessing the neutralization sensitivity and structure-function phenotype of Spike VOC with multiple mutations that likely have complex epistatic interactions.

To generate rcVSV-CoV2-S containing different variants or mutants on demand, without the need for extensive passaging, we developed a robust reverse genetics system and VNA which leverages the cell lines we previously developed for a standardized SARS-CoV-2 VNA that correlates well with live virus neutralization [41]. Salient improvements include the addition of a hammerhead ribozyme immediately upstream of the 3' leader sequence which cleaves *in cis* to give the exact 3' termini, the use of a codon-optimized T7-polymerase which alleviates the use of vaccinia-driven T7-polymerase, and a highly permissive and transfectable 293T-ACE2+TMPRSS2 clone (F8-2) [41] (Extended Data Fig S1). A 6-plasmid transfection into F8-2 cells results in GFP+ cells 2-3 days post-transfection (dpt), which turn into foci of syncytia by 4-5 dpt indicating virus replication and cell-to-cell spread (Fig. 1A). Transfer of F8-2 cell supernatant into interferon-defective Vero-TMPRSS2 cells allowed for rapid expansion of low-passage viral stocks that maintain only the engineered Spike mutations. Clarified viral supernatants from Vero-TMPRSS2 cells were aliquoted, sequenced verified, then titered on F8-2 cells to determine the linear range of response (Fig. 1B).

Next, we generated isogenic rcVSV-CoV2-S expressing the B.1.1.7, B.1.351 (Fig. 2A),
or E484K S to evaluate the neutralizing activity of Sputnik V vaccine sera from
Argentina. The relevant Spike substitutions that make up these variants are indicated in
Fig. 2A.  The characteristics of the vaccine recipient cohort (n=12) receiving the two-
dose regimen of the Sputnik vaccine are given in Table 1. At one month post-
completion of the two-dose regimen, the Sputnik V vaccine generated respectable
virus neutralizing titers (VNT) against rcVSV-CoV2-S bearing the WT (D614G) and
B.1.1.7 spike proteins (Fig. 2B). The geometric mean titer (GMT) and 95% CI for WT
(1/$IC_{50}$ GMT 49.4, 23.4 - 105) in our cohort of vaccine recipients was remarkably similar
to that reported in the phase III Sputnik vaccine trial (GMT 44.5, 31.8 - 62.2)[48,49].
However, GMT against B.1.351 and E484K was reduced by a median 6.8- and 2.8-fold,
respectively compared to WT (Fig. 2C).

Sputnik vaccine recipients appeared to generate qualitatively different neutralizing
antibody responses against SARS-CoV-2 that could be segregated into three different
groups (Fig. 3). Group (A) sera showed reasonable VNT against wild-type (WT) and
B.1.1.7 (Fig. 3A and E). However, the Hill slope of their neutralization curves for B.1.351
were extremely shallow (h<0.40), resulting in a low IC50s and maximal neutralization of
50-60% even when extrapolated to full serum strength (Fig. 3E and Fig. 4). In contrast,
Group (B) sera neutralized E484K and B.351 with similar potencies to WT and B.1.1.7,
especially at high serum concentrations (Fig. 3B and E). This group of sera reveals that
qualitatively different neutralizing responses can be generated that effectively neutralize
B.1.351. Group (C) sera generally exhibited effective neutralization of WT, B.1.1.7, and
even E484K at high serum concentrations, but not B.1.351 (Fig. 3C and E). The
decreased potency and shallow Hill Slope result in <90% neutralization of B.1.351 even
at full serum strength (see next section). One serum sample (SP012) exhibited little to
no neutralizing activity against WT, E484K and B.1.351, yet it neutralized B.1.1.7 as
well as Group A-C sera (Fig. 3D-E).  That these three groups exhibit qualitatively
distinct neutralization patterns is further highlighted by the Hill slopes of their
neutralization curves (Fig. 3F). Group A sera not only have the lowest slopes against
B.1.351, but as a group, they have slopes significantly <1 against the other viruses
(median/IQR = 0.4965 / 0.2880 – 1.186). Group B sera mostly have slope values
around 1 (median/IQR = 0.8855 / 0.7865 – 1.065) while Group C sera have the highest
overall slopes (median/IQR = 1.348 / 0.8395 – 1.820) that are significantly >1 (Fig. 3F).

The Hill Slope of the neutralization curves against B.1.351 was significantly different
from WT, B.1.1.7 and E484K (Fig. 4A). As a consequence, the maximal neutralization
attainable when extrapolated to full serum strength was also significantly lower for
B.1.351 compared to the rest (Fig. 4B). Conversely, the steep Hill slope for the E484K
curves resulted in maximal neutralization potencies that were not significantly different
from WT or B.1.1.7 despite significantly lower reciprocal $IC_{50}$ values (compare Fig. 2B

with Fig. 4B). Notably, the maximal percent inhibition was strongly correlated with the

Hill Slope for WT and VOC/mutant spikes across all valid pairs of sample values (Fig.

4C, n=45 pairs), suggesting that antibody co-operativity likely plays a role at high serum

concentrations (see Discussion) [42]. While we acknowledge the limitations of

extrapolating values from nonlinear regression curves, the striking correlation between

slope and maximal percent inhibition attainable at full serum strength supports the

robustness of our nonlinear regression model.

The heterogenous dose-response curves described in Fig. 3-4 is a property of Sputnik

V vaccine elicited responses as soluble RBD-Fc inhibition of WT and VOC S-mediated

entry produced classical dose response curves with Hill slopes close to -1.0 (Fig. 5).

Both B.1.1.7 and B.1.351 were modestly but significantly more resistant to RBD-Fc

inhibition (Fig. 5A-B). This is not surprising as both harbor the N501Y mutation known to

enhance affinity of RBD for ACE2 [43–45]. However, this 1.5 to 2-fold increase in RBD-Fc

$IC_{50}$ for B.1.1.7 and B.1.351, respectively, does not explain the neutralization-resistant

versus sensitive phenotype of B.1.351 versus B.1.1.7 in our virus neutralization assays.

Furthermore, the E484K mutant was more sensitive to RBD-Fc inhibition than B.1.1.7

(Fig. 5C-D), and yet also remained more neutralization-resistant relative to B.1.1.7.

Experimental measurements of both RBD and trimeric spike binding to ACE2 have

revealed that the E484K mutation alone does not confer increase binding affinity for

ACE2 unlike N501Y [44,46]. Our RBD-Fc inhibition studies in the context of virus infection

confirm and extend these results. Our data reinforces the notion that the mechanism

underlying the increased neutralization resistance of E484K containing variants and

mutants do not involve ACE2 binding affinity per se, but rather affects a key

immunodominant epitope targeted by a significant class of human neutralizing

antibodies, variably termed as RBM class II, RBS-B, or Cluster 2 antibodies [47–50].

**DISCUSSION**

A key public health concern related to emergent SARS-CoV-2 variants is that by incrementally accruing mutations that escape neutralizing antibodies, they will penetrate herd immunity and spread to reach unvaccinated individuals, some of whom will be susceptible to severe or fatal disease.

Three of the six COVID-19 vaccines currently in use worldwide, namely Moderna mRNA-1273, BioNTech BNT162b2, and Janssen Ad26.COV2.S, each express S harboring K986P and V987P substitutions (2P) within a loop abutting the central helix of the S2' membrane fusion machinery [51–53]. This modification locks the spike in a prefusion conformation and elicits higher titers of neutralizing antibodies [54,55]. The Janssen vaccine has an additional deletion in the furin cleavage site, while the yet-to-be approved Novavax vaccine contains arrays of stabilized spikes conjugated onto a nanoparticle (Table 2). Of the three vaccines that do not appear to make use of 2P Spike mutants, Gamaleya's Sputnik V and AstraZeneca's AZD1222 are adenovirus-vectored vaccines encoding native S. The third is CoronaVac, a preparation of inactivated SARS-CoV-2 virions. Although all six vaccines are highly efficacious at preventing severe COVID-19 outcomes, they do not all uniformly prevent infection. Moreover, in all cases thus far examined, these first generation vaccines are less effective against variants with certain non-synonymous substitutions in Spike, such as E484K.

The most concerning variants are those with multiple mutations in the receptor binding domain (RBD) that confer both enhanced affinity for the hACE2 receptor and escape from neutralizing antibody responses [17,24,27,33,56,57]. B.1.351 and P.1 have in common three RBD substitutions (K417N/T, E484K and N501Y) whereas B.1.351, P.1 and B.1.1.7 contain the N501Y substitution. Although B.1.1.7 shows enhanced transmissibility and more severe disease outcomes[52], it does not appear to be consistently more resistant to serum neutralizing responses elicited by vaccines or natural infection [58,59]. The same is not true, however, for the B.1.351 variant.

In live virus plaque reduction neutralization assays, sera from AstraZeneca vaccine recipients in South Africa exhibited 4.1 to 32.5-fold reduction in neutralizing activity against B.1.351 [33]. The actual reduction is even more marked because 7 of 12 vaccine recipients who had neutralizing activity against the parental B.1.1 variant, had undetectable neutralization against the B.1.351 strain. Comparator sera from recipients of Moderna and BioNTech mRNA vaccines showed smaller, 6.5 to 8.6-fold reductions in neutralization [60].

As of this writing, there are no peer-reviewed data on the protective efficacy of Sputnik V and CoronaVac against SARS-CoV-2 S variants. Here, we showed that sera from Sputnik vaccine recipients in Argentina had a median 6.1-fold and 2.8-fold reduction in GMT against B.1.351 and the E484K mutant spike, respectively. Even more revealing is

their dose-response curves. When extrapolated to full serum strength, half of the sera
samples failed to achieve an $IC_{80}$ and only 1 out 12 achieved an $IC_{90}$ against B.1.351
(Fig. 4A). Table 2 summarizes peer-reviewed studies that have tested post-vaccination
sera from the major vaccines against the VOC/mutant spikes used in this study. Our
study shows a similar mean reduction in GMT (reciprocal IC50) against E484K and
B.1.351 using 1-month post-Sputnik vaccine sera when compared to other vaccines.
Our sample number is admittedly small but matches the median and modal number
used in other studies to date. Nonetheless, we caution that comparing only the mean
reduction in IC50 can be misleading as an aggregate measure of serum neutralizing
activity. The neutralization curves for B.1.351 in our study are not classically sigmodal
and have significantly shallower slopes than WT, B.1.17 and E484K, which result in $\leq$
90% neutralization for all but one sample when extrapolated to full serum strength. The
possible mechanisms for the varying slope values are discussed below.

E484K is present not only as part of an ensemble of RBD mutations present in B.1.351
and P.1, but in many of the 17 lineages detected from South America that carry it, such
as P.2, E484K is the only RBD substitution (Supplementary Table 1). A more detailed
report covering the genomic surveillance efforts in Argentina that detected the VOC
which spurred our study is currently in preparation (Dr. Claudia Perandones, personal
communication).

While the E484K substitution appears to be a common route of escape from many
RBD-targeting monoclonal antibodies, it is somewhat surprising that a single mutation
can confer a significant degree of neutralization resistance from polyclonal responses.
Nonetheless, our data show that resistance conferred by E484K mutation be overcome
by higher titer antibodies present in undiluted patient sera.  But the neutralization
resistance conferred by the suite of mutations present in B.1.351 appears qualitatively
different. In the majority of cases, the slope of the dose response curve indicates a
failure to neutralize even at full strength. We had previously shown that the dose-
response curve slope is a major predictor of therapeutic potency for HIV broadly
neutralizing antibodies at clinically relevant concentrations [42]. Importantly, the slope
parameter is independent of IC50 but is specifically related to an antibody's epitope
class. Here, we show that defining the neutralization phenotype of a given spike variant
or mutant by both its relative IC50 and slope provides a fuller characterization of serum
neutralizing activity against SARS-CoV-2 and emergent VOC.

The deletion of residue 242-244 in the NTD of the B.1.351 spike appear to cause large-
scale resurfacing of the NTD antigenic surface resulting in greater conformational
heterogeneity[44]. Variable neutralization responses across such a heterogenous virus
population may result in the shallow slopes (<1) seen. Furthermore, three major classes
of neutralizing antibodies (RBS-A, -B, and -C) identified from convalescent patients are
sensitive to either the K417N (RBS-A) or E484K (RBS-B and -C) substitutions present

in B.1.351.  On the other hand, at high serum concentrations, co-operative effects from
low-affinity spike binding antibodies or increased spike occupancy by different classes
of antibodies can result in the steep Hill slope observed. The steep Hill slopes (>1.0)
observed for E484K suggest such co-operative effects might be occurring.

The emergence of variants is fluid situation. B.1.427/1.429, B.1.526, and B.1.617 are
other emergent VOI/VOC that could be tested.  These strains have substitutions in the
RBD (L452R and E484K/Q) and elsewhere in the spike that might confer some degree
of neutralization resistance in our in vitro assays. However, all vaccines are effective
against most variants, and we do not know what degree of resistance in *in vitro* assays
translate to a decrease in the real world efficacy of any given vaccine.

Although we stress that the Gameyla Sputnik V vaccine is likely to retain strong
efficacy at preventing severe COVID-19, even in the case of infection by VOC, our data
reveal a concerning potential of B.1.351, and to a lesser extent, any variant carrying the
E484K substitution (e.g. P.2), to escape the neutralizing antibody responses that this
immunization elicits. Furthermore, we acknowledge that *in vivo* protective efficacy can
be derived from Fc effector functions of antibodies that bind but do not neutralize. In
addition, an adenoviral vectored vaccine should induce potent cell-mediated immunity
against multiple epitopes, which were not measured in our study. Nevertheless, given
the crucial roles neutralizing antibodies play in preventing infection, our results suggest
that updated SARS-CoV-2 vaccines will be necessary to eliminate the virus.

**Materials and Methods**

**Cell lines**

Vero-CCL81 TMPRSS2, HEK 293T-hACE2 (clone 5-7), and 293T-hACE2-TMPRSS2
(clone F8-2) cells were described previously [41], and were maintained in DMEM +
10%FBS.  The HEK 293T-hACE2-TMPRSS2 cells were plated on collagen coated
plates or dishes. BSR-T7 cells [61], which stably express T7-polymerase were maintained
in DMEM with 10% FBS.

**VSV-eGFP-CoV2 spike (Δ21aa) genomic clone and helper plasmids.**

We cloned VSV-eGFP sequence into pEMC vector (pEMC-VSV-eGFP), which includes
an optimized T7 promoter and hammerhead ribozyme just before the 5' end of the viral

genome. The original VSV-eGFP sequence was from pVSV-eGFP, a generous gift of
Dr. John Rose [62].

We generated pEMC-VSV-eGFP-CoV2-S (Genbank Accession: MW816496) as follows:
the VSV-G open reading frame of pEMC-VSV-eGFP was replaced with the SARS-CoV-
2 S, truncated to lack the final 21 amino acids [63]. We introduced a Pac-I restriction
enzyme site just after the open reading frame of S transcriptional unit, such that the S
transcriptional unit is flanked by MluI / PacI sites. SARS-CoV-2 S is from pCAGGS-
CoV-2-S [64], which codes the codon optimized S from the Wuhan Hu-1 isolate (NCBI
ref. seq. NC_045512.2) with a point mutation of D614G, resulting in B.1 lineage. The
B.1.1.7 Spike we used carries the mutations found in GISAID Accession Number
EPI_ISL 668152: del 69-70, del145, N501Y, A570D, D614G, P681H, T716I, S982A, and
D1118H.  The B.1.351 Spike carries the mutations D80A, D215G, del242-244, K417N,
E484K, N501Y, D614G, and A701V (from EPI_ISL_745109). The Spike sequences of
WT, B.1.1.7, B.1.351, and E484K are available at Genbank (Accession Numbers:
MW816497, MW816498, MW816499, and MW816500; please also see Supplemental
Table 2).

Sequences encoding the VSV N, P, M, G, and L proteins were also cloned into pCI
vector to make expression plasmids for virus rescue, resulting in plasmids: pCI-VSV-N,
pCI-VSV-P, pCI-VSV-M, pCI-VSV-G, and pCI-VSV-L. These accessory plasmids were
a kind gift from Dr. Benjamin tenOever.

**Generation of VSV-CoV2 spike from cDNA**

$4 \times 10^5$ 293T-ACE2-TMPRSS2 cells per well were seeded onto collagen-I coated 6 well
plates. The next day, 2000 ng of pEMC-VSV-EGFP-CoV2 spike, 2500 ng of pCAGGS-
T7opt [65], 850 ng of pCI-VSV-N, 400 ng of pCI-VSV-P, 100 ng of pCI-VSV-M, 100 ng of
pCI-VSV-G, 100 ng of pCI-VSV-L were mixed with 4 mL of Plus reagent and 6.6 mL of
Lipofectamine LTX (Invitrogen). 30 min later, transfection mixture was applied to 293T-
hACE2-TMPRSS2 cells in a dropwise fashion. Cells were maintained with medium
replacement every day for 4 to 5 days until GFP positive syncytia appeared. Rescued
viruses were amplified in Vero-CCL81 TMPRSS2 cells [41], then titered and used for the
assay.

**Virus neutralization assay**

5 x 10E4 293T-hACE2-TMPRSS2 cells per well were seeded onto collagen-coated 96
well cluster plates one day prior to use in viral neutralization assays. Virus stocks were
mixed with serially diluted serum for 10 minutes at room temperature, then infected to
cells.  Note:  all sera assayed in this study were previously heat inactivated by 56
degrees for 30 min before use in any viral neutralization studies.  At 10 h post infection,
GFP counts were counted by Celigo imaging cytometer (Nexcelom).  Each assay was

done in triplicate. For calculation of IC50, GFP counts from "no serum" conditions were
set to 100%; GFP counts of each condition (serum treated) were normalized to no
serum control well.  Inhibition curves were generated using Prism 8.4.3 (GraphPad
Software) with 'log (inhibitor) vs normalized response - variable slope' settings.

**Design of RBD-Fc producing Sendai virus**

Sendai virus (SeV) Z strain cDNA sequence (AB855655.1) was generated and cloned
into pRS vector with the addition of eGFP transcriptional unit at the head of SeV
genome. The sequence of F transcriptional unit was from SeV fushimi strain
(KY295909.1) due to the cloning reason. We refer to the pRS-based plasmid coding
this sequence as pRS-SeVZ-GFP-F[fushimi] in this paper. For the introduction of foreign
gene into SeV, we generated additional transcriptional unit for RBD-Fc between P gene
and M gene. RBD-Fc construct was generated as below; codon optimized DNA
sequence of from SARS-CoV-2 spike (MN908947) in pCAGGS a gift of Dr. Florian
Krammer [64]. S amino acids 319 – 541 (corresponding to the RBD domain) sequence
were C-terminally fused to the Fc region of human IgG$_1$ (220 – 449 aa of P0DOX5.2)

**Generation of recombinant Sendai virus from cDNA.**

2x10E5 BSR-T7 cells per well were seeded onto 6-well cluster plates. The next day, 4
µg of pRS-SeVZ-GFP-F[fushimi], 4 µg of pCAGGS-T7opt, 1.44 µg of SeV-N, 0.77 ug of
SeV-P, 0.07 ug of  SeV-L were mixed with 5.5 µl of Plus reagent and 8.9 µl of
Lipofectamine LTX (Invitrogen). 30 min later, transfection mixtures were applied to Bsr-
T7 cells in a dropwise fashion, as described previously [65]. At one day post transfection,
medium was replaced with DMEM + 0.2 µg/ml of TPCK-trypsin (Millipore Sigma,
#T1426), with subsequent medium replacement each day until infection reached 100%
cytopathic effect. Supernatants were stored at -80℃ until use in experiments.

**Titration of viruses.**

For SeV titration, 2 x 10E4 Bsr-T7 cells per well were seeded onto 96-well plates. The
next day, 100 µL of serially diluted virus stock (in DMEM + 10% FBS) were applied to
each well. GFP positive foci were counted at 24 hours post infection using a Celigo
imaging cytometer (Nexcelom, Inc.). Infectivity is presented in infectious units (IU) per
mL.

For VSV-CoV2 titration, 5 x 10E4 293T-hACE2-TMPRSS2 cells per well were seeded
onto a collagen-coated 96 well plate. Serially diluted virus stocks were then applied to

the cells, and GFP positivity was scored at 10 h post infection using a Celigo imaging
cytometer.

**Production of proteins and purification.**

$5\times10E6$ Bsr-T7 cells are seeded in T175cm$^2$-flask one day before infection. Cells were
infected by SeV at MOI of 0.1 for one hour, followed by replacement of medium with
DMEM supplemented with 0.2 mg/mL TPCK-trypsin. Medium was replaced with fresh
0.2 mg/ml TPCK-trypsin containing DMEM each day until infection reached 100%
CPE, at which point medium was exchanged for DMEM lacking TPCK-trypsin. Cells
were incubated for additional 24 h to allow protein production. Supernatants were
centrifuged at 360 $g$ for 5 min, then filtered with 0.1 µm filter (Corning$^®$ 500 mL Vacuum
Filter/Storage Bottle System, 0.1 µm Pore) to remove virions and debris. Supernatant
including RBD-Fc were applied to Protein G Sepharose (Millipore Sigma, #GE17-0618-
01) containing column (5ml Polypropylene Columns ;ThermoFisher, #29922), followed
by wash and elution.

**Human Subjects Research.**

Human subjects research was conducted following the Declaration of Helsinki and
related institutional and local regulations. Studies and serum collection relating to the
Sputnik vaccine at ANLIS Dr. Carlos G. Malbrán (NatIonal Administration Laboratories
and Health Institutes - Carlos G. Malbrán, Argentina) were approved by the Research
Ethics Committee of its Unidad Operativa Centro de Contención Biológica (UOCCB) on
9 Feb 2021.

**Authors contributions**

S.I., C.P., B.H.L., J.P.K, conceived of and supervised the study. C.P., A.E.V. and A.E.
supervised, collected, analyzed, and provided materials relevant to this study.   S.I.
generated VSV-CoV-2 S plasmid and rescued viruses. S.I., G.H., S.K., and M.N.A.S.
were involved in the generation of S mutant viruses. S.I, L.B., M.N.A.S., K.Y.O.
conducted neutralization assays. S.I. and C.T.H. developed the Sendai virus protein
expressing system and purified RBD-Fc protein. S.I., B.H.L., and J.P.K. wrote the
paper with input from C.P. and all co-authors.

**Acknowledgements**

We gratefully acknowledge all submitting authors and collecting authors on whose
work this research is based, and to all researchers, clinicians, and public health

authorities who make SARS-CoV-2 sequence data available in a timely manner via the
GISAID initiative[66,67].

**Funding information**

We acknowledge the following finding. K.Y.O. was supported by Viral-Host
Pathogenesis Training Grant T32 AI07647 and additionally by a NRSA F31 AI154739.
S.I. and C.-T.H. were supported by postdoctoral fellowships from CHOT-SG (Fukuoka
University, Japan) and the Ministry of Science and Technology (MOST, Taiwan),
respectively. B.L. acknowledges flexible funding support from NIH grants AI123449
and AI138921; a grant from the Department of Microbiology, Icahn School of Medicine
at Mount Sinai; and the Ward-Coleman estate, which endowed the Ward-Coleman
Chairs at the ISMMS. J.P.K. was supported by a COVID-19 Fast Grants award from
Emergent Ventures, an initiative of the Mercatus Center at George Mason University,
and by an intramural grant and other funding from the Office of the Vice Chancellor for
Research at LSU Health Sciences Center Shreveport (J.P.K., M.N.A.S.). Processing
costs recovered from multiple users of our standardized SARS-CoV-2 VSV
pseudotyped particles provided additional support (BL). Work at ANLIS-MALBRAN
(A.E.V., A.E., C.P.) was supported by the Ministry of Health (Ministerio de Salud),
Argentina.

**Table 1. Cohort characteristics of Sputnik vaccine recipients from ANLIS MALBRÁN (Buenos Aires, República Argentina).**

| Sera ID | 1st DOSE | 2nd DOSE | Vaccine Status | SEX | AGE |
|---|---|---|---|---|---|
| SP001 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 45-50 |
| SP002 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 40-45 |
| SP003 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 55-60 |
| SP004 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 50-55 |
| SP005 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 35-40 |
| SP006 | Late Dec/2020 | Mid Jan/2021 | (+) | F | 35-40 |
| SP007 | Late Dec/2020 | Mid Jan/2021 | (+) | F | 20-25 |
| SP008 | Late Dec/2020 | Early Feb/2021 | (+) | M | 35-40 |
| SP009 | Late Dec/2020 | Early Feb/2021 | (+) | F | 30-35 |
| SP010 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 30-35 |
| SP011 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 40-45 |
| SP012 | Late Dec/2020 | Mid Jan/2021 | (+) | M | 25-30 |
| | | | | Median Age | **39.5** |
| | | | | Range | **25-56** |
| SP013 | N.A. | N.A. | (-) | F | 45-50 |
| SP014 | N.A. | N.A. | (-) | F | 50-55 |
| SP015 | N.A. | N.A. | (-) | M | 40-45 |

**N.A., Not Applicable**

**Table 2.** Summary of post-vaccine sera evaluated for neutralization potency against the indicated SARS-CoV-2 variants of concern (VOC). Table format adapted and updated from Abdool Karim and de Olivera[60].

| Vaccine | Company | Spike Construct | Neutralization Assay ($IC_{50}$ Fold-reduction vs WT) | | | | Reference PMID |
|---|---|---|---|---|---|---|---|
| | | | B.1.1.7 Variant | P.1 Variant | B.1.351 Variant | # samples tested (n) | |
| Adapted from **Table I** of SS Abdool Karim and T de Oliveira, New SARS-CoV-2 Variants – Clinical, Public Health, and Vaccine Implications. *NEJM*, 24 Mar, 2021, DOI: 10.1056/NEJMc2100362. | | | B.1.1.7 Variant | E484K | B.1.351 Variant | # samples tested (n) | |
| Ad26.COV2.S | Johnson&Johnson | 2P & ΔFurin | ≤2× (n.s.) | NA | ≤5× | 8* | 33909009 |
| BNT162b2 | Pfizer/BioNTech | "2P" | 2× | NA | ≤6.5× | 10 | 33684923 |
| | | | 2x (n.s.) | 6.7× | 35x | 30 | 33743213 |
| | | | 3.3x | NA | NA | 25 | 33743891 |
| | | | NA | NA | 7.9 | | 33730597 |
| mRNA-1273 | Moderna | "2P" | 1.8× | NA | ≤8.6× | 12 | 33684923 |
| | | | (n.s.) | 4.5× | 28x | 35 | 33684923 |
| Pfizer/BioNTech (Pfz) OR Moderna (Mod) | | "2P" | NA | ≤3× | NA | 4^Pfz, 11^Mod | 33567448 |
| NVX-CoV2373 | Novavax | Stabilized | 2× | NA | NA | 28 | 33705729 |
| AZD1222 | AstraZeneca | Native | NA | NA | 4× | 13 | 33725432 |
| | | | 8.9x | NA | NA | 49 | 33798499 |
| | | | 2.1~2.5x | NA | NA | 25 | 33743891 |
| | | | NA | NA | 9x | | 33730597 |
| CoronaVac | Sinovac | Native | NA | NA | NA | N.D. | N.D. |
| BBIBP-CorV | Sinopharm | Native | NA | NA | 1.6× (n.s.) | 12** | 33870240 |
| Current Study: Ikegame et al, Apr, 2021 | | | B.1.1.7 | E484K | B.1.351 | | |
| **Sputnik V** | **Gamaleya** | **Native** | (n.s.) | 2.8x | 6.1x | 12 | 33821288 |

\* Non-human Primate data only

\*\* Preprint only

| | | |
|---|---|---|
| Mean | 19.6 |
| Median | 12.5 |
| Mode | 12.0 |

**SUPPLEMENTAL TABLE 1.** Acknowledgement of S: E484K viruses from South America shared on GISAID.

**SUPPLEMENTAL TABLE 2.** Acknowledgement of B.1.1.7 and B.1.351 viruses used for selection of S variants evaluated in this study.

**Figure 1. Replication-competent VSV bearing wild-type and variant SARS-CoV-2 spike (rcVSV-CoV2-S). (A)** Representative images of *de novo* generation of rcVSV-CoV2-S, carrying an EGFP reporter, in transfected 293T-ACE2+TMPRSS2 (F8-2) cells as described in Extended Data Fig. S1. Single GFP+ cells detected at 2-3 days post-transfection (dpt) form a foci of syncytia by 4 dpt. Images are taken by Celigo imaging cytometer (Nexcelom) and are computational composites from the identical number of fields in each well. White bar is equal to 1 millimeter. **(B)** Entry efficiency of rcVSV-CoV2-S in parental 293T cells, 293T stably expressing ACE2 alone (293T-ACE2) or with TMPRSS2 (293T-ACE2+TMPRSS2). Serial dilutions of virus stocks amplified on Vero-TMPRSS2 cells were used to infect the indicated cell lines in 96-well plates in triplicates. GFP signal was detected and counted by a Celigo imaging cytometer (Nexcelom) 10 hours post-infection. Symbols are individual data points from triplicate infections at the indicated dilutions. Bars represent the average of 3 replicates with error bars indicating standard deviation. A two-way ANOVA was used to compare the differences

between cell lines at any given dilution. Adjusted p values from Tukey's multiple comparisons test are given (ns; not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). **(C)** rcVSV-CoV-2-S containing the prevailing WT (D614G) and VOC (B.1.1.7 and B.1.351) spikes were inoculated into one 6-well each of F8-2 cells (MOI 0.1) and subsequently overlaid with methylcellulose-DMEM to monitor syncytia formation. Representative images of syncytial plaques at 48 hpi are shown. White bar equals 1 millimeter. **(D)** shows the growth of GFP positive area / infectious unit (IU) in the 6 well plate. GFP positive areas were imaged and measured by the Celigo imaging cytometer. IU was checked at 10 hpi in the same well. Bar shows the average of 3 independent experiments with error bar indicating standard deviation. No statistically significant differences were detected between WT and VOC spikes in the size of GFP+ syncytia at any given time point (two-way ANOVA as above, 'ns' not indicated in graph).

| | | | | |
|---|---|---|---|---|
| GMT | 87.1 | 49.4 | 7.9 | 17.0 |
| Lower 95% CI | 49.5 | 23.4 | 3.2 | 8.6 |
| Upper 95% CI | 154 | 105 | 19.5 | 33.8 |

| | | |
|---|---|---|
| Median Fold-Change | 6.1 | 2.8 |
| Lower 95% CI | 2.3 | 1.8 |
| Upper 95% CI | 10.1 | 3.3 |

**Figure 2. Neutralization activity of antibody responses elicited by the Sputnik V vaccine.**
**(A)** Schematic of the Spike substitutions that make up the variants being evaluated in this study. The amino acid positions and corresponding 'Ancestral' sequence of the Wuhan isolate is shown. The prevailing WT sequence now has a D614G substitution. All the variants and mutants have D614G. **(B)** Neutralization activity of individual serum samples against rcVSV-CoV2-S with the WT, variant (B.1.1.7 or B.1.351), or mutant E484K spike proteins. Neutralization is represented by the reciprocal 50% inhibitory dilution factor ($1/IC_{50}$). Sera samples with no appreciable neutralization against a given virus were assigned a defined $1/IC_{50}$ value of 1.0, as values $\leq 1$ are not physiological (Grey shaded area). Dashed line indicates the lowest serum dilution tested ($1/IC50 = 20$). Geometric mean titers (GMT and 95% CI) for the neutralizing activity of all vaccine sera are indicated below each of the viral spike proteins examined . NS; not significant, *; $p<0.05$, $p < 0.01$; ** are adjusted p values from non-parametric one-way ANOVA with Dunn's multiple comparisons test. **(C)** For each serum sample, the fold-change in $IC_{50}$ (reciprocal inhibitory dilution factor) against the indicated variant and mutant spike proteins relative to its $IC_{50}$ against wild-type (WT) spike (set at 1) is plotted. Adjusted p values were calculated as in (B). Medians are represented by the bars and whiskers demarcate the 95% CI. Neutralization dose-response curves were performed in triplicates, and the mean values from each triplicate experiment are shown as the single data points for each sera sample.

**Figure 3. Sputnik vaccine recipients generate qualitatively different neutralizing antibody responses against SARS-CoV-2. (A-C)** Group A (SP001, SP005, SP006, SP012), Group B (SP002, SP007), and Group C (SP003, SP004, SP008, SP010, SP011) represent potentially distinct classes of virus neutralizing activity present in the sera samples analyzed. Full

neutralization curves for all sera tested against all viruses bearing the variant and mutant spike proteins are shown. **(D)** shows a singular example of a serum that only neutralized the B.1.1.7 spike. **(E)** graphs the serum neutralizing titers (SNT = $1/IC_{50}$) and 95% CI that can be extrapolated from the nonlinear regression curves shown for all the sera samples analyzed. Colored filled symbols represent the indicated viruses, open symbols in (E) represent assigned SNT values of 1.0 when no significant neutralization activity could be detected (SP012, B.1.351 and E484K). The dotted black line represents a reciprocal serum dilution of 1.0. The red dashed line and shaded boundaries represent the geometric mean titer and 95% CI, respectively, for B.1.351. **(F)** The Hill slope values for all the neutralization curves are aggregated according to their groups. The different colored symbols in each group represent the indicated virus tested. P values are from a non-parametric Wilcoxon signed rank test  using a theoretical median of 1.0.

**Figure 4. Maximal inhibition and slope help to define the distinct classes of neutralizing sera in Sputnik vaccine recipients.** **(A)** Paired comparison of Hillslopes from the neutralization curves of all samples except for SP012 where no significant neutralization was observed for viruses other than B.1.1.7. NS; not significant, p<0.05, *; p<0.01; **, p < 0.0001; ****, are adjusted p values from non-parametric one-way ANOVA with Dunn's multiple comparisons test, which assumes non-Gaussian distribution of values being analyzed. **(B)** Maximal percent inhibition (MPI) at full serum strength extrapolated from nonlinear regression of log(inhibitor) versus normalized response, variable slope curve. Model used is from PRISM v9.1 where Y= 100/(1+10^((LogIC50-X)*HillSlope))). Log IC50 and Hill slope values were obtained for each curve generated in Fig. 3. MPI = 100-Y, when X= 0 for reciprocal serum dilution of 1 (10^0 =1). Data points for one serum (SP012) against WT, B.1.351 and E484K could not be calculated because there was no best-fit value. The dotted line indicates 90% inhibition. Median (central bar) and interquartile values (whiskers) are indicated. Adjusted p values was calculated as in (A). **(C)** Correlation analysis of MPI versus the Hill Slope parameter for all sera samples tested against all spike proteins. SP012 was excluded for the abovementioned reasons. Non-parametric Spearman r values and 95% confidence interval are shown. X-axis is plotted as an asymptotic cumulative probability scale as x approaches 100% (PRISM v9.1.1) only to resolve the many MPI values >90%.

**Figure 5. Competitive inhibition of rcVSV-CoV2-S entry by soluble RBD-Fc. (A)** Recombinant RBD-Fc was serially titrated with the infection inoculum containing a fixed amount of rcVSV-CoV2-S bearing WT or the indicated VOC spike proteins. 10 hpi, GFP+ cells were quantified by the Celigo image cytometer. Data points are means of six independent replicates with error bars representing S.D. The number of GFP+ cells in the absence of any RBD-Fc was set to 100% and used to normalize the infection response in the presence of increasing amounts of RBD-Fc. Log[inhibitor] versus normalized response variable slope nonlinear regression curves were generated using GraphPad PRISM (v9.1.0). **(B)** The IC50 values from each replicate dose response curve generated for a given virus were grouped. The mean (central bar) and SD (whiskers) for each group are indicated. Adjusted p values (*; p<0.05, **; p<0.01, ****; p<0.0001) from ordinary one-way ANOVA with Dunnett's multiple comparisons test are indicated. **(C)** is a repeat of the experiment done in A with the E484K mutant using a different preparation of recombinant RBD-Fc (see methods). B.1.1.7 serves as the common reference control. **(D)** The IC50 values were calculated and analyzed as in (B).

**Extended Data Figure S1. Robust and efficient generation of an EGFP-reporter replication-competent VSV bearing SARS-CoV-2 spike (rcVSV-CoV2-S). (A)** Schematic of the rcVSV-CoV2-S genomic coding construct and the virus rescue procedure. The maximal T7 promoter (T7prom) followed by a hammer-head ribozyme (HhRbz) and the HDV ribozyme (HDVRbz) plus T7 terminator (T7term) are positioned at the 3' and 5' ends of the viral cDNA, respectively. An EGFP(E) transcriptional unit is placed at the 3' terminus to allow for high level transcription. SARS-CoV-2-S is cloned in place of VSV-G using the indicated restriction sites designed to facilitate easy exchange of spike variant or mutants. (**B**) For virus rescue, highly permissive 293T cells stably expressing human ACE2 and TMPRSS2 (293T-[ACE2+TMPRSS2], F8-2 clone) cells were transfected with the genome coding plasmid, helper plasmids encoding CMV-driven N, P, M, and L genes, and pCAGS encoding codon-optimized T7-RNA polymerase(T7opt). 48-72 hpi, transfected cells turn EGFP+ and start forming syncytia. Supernatant containing rcVSV-CoV2-S are then amplified in Vero-TMPRSS2 cells at the scale shown. The blue arrows at the bottom indicate the timeline for production of each sequence verified stock.

# REFERENCES

1. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497–506 (2020).

2. COVID-19 Map - Johns Hopkins Coronavirus Resource Center. https://coronavirus.jhu.edu/map.html.

3. Boni, M. F. *et al.* Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. *Nat Microbiol* **382**, 1199 (2020).

4. Logunov, D. Y. *et al.* Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. *Lancet* **397**, 671–681 (2021).

5. Baden, L. R. *et al.* Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N. Engl. J. Med.* **384**, 403–416 (2021).

6. Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N. Engl. J. Med.* **383**, 2603–2615 (2020).

7. Zhang, Y. *et al.* Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18--59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *Lancet Infect. Dis.* (2020).

8. Sadoff, J. *et al.* Interim Results of a Phase 1–2a Trial of Ad26.COV2.S Covid-19 Vaccine. *N. Engl. J. Med.* (2021) doi:10.1056/NEJMoa2034201.

9. Folegatti, P. M. *et al.* Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial. *Lancet* **396**, 467–478 (2020).

10. Addetia, A. *et al.* Neutralizing Antibodies Correlate with Protection from SARS-CoV-2 in Humans during a Fishery Vessel Outbreak with a High Attack Rate. *J. Clin. Microbiol.* **58**, (2020).

11. Khoury, D. S. *et al.* What level of neutralising antibody protects from COVID-19? *bioRxiv* (2021) doi:10.1101/2021.03.09.21252641.

12. Andrew Rambaut, Nick Loman, Oliver Pybus, Wendy Barclay, Jeff Barrett, Alesandro Carabelli, Tom Connor, Tom Peacock, David L Robertson, Erik Volz, COVID-19 Genomics Consortium UK (CoG-UK). Preliminary genomic characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations. *Virological.org* https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563 (2020).

13. Choi, B. *et al.* Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *N. Engl. J. Med.* **383**, 2291–2293 (2020).

14. Davies, N. G. *et al.* Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science* (2021) doi:10.1126/science.abg3055.

15. Davies, N. G. *et al.* Increased mortality in community-tested cases of SARS-CoV-2 lineage B.1.1.7. *Nature* (2021) doi:10.1038/s41586-021-03426-1.

16. Grint, D. J. *et al.* Case fatality risk of the SARS-CoV-2 variant of concern B.1.1.7 in England. *bioRxiv* (2021) doi:10.1101/2021.03.04.21252528.

17. Tegally, H. *et al.* Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. *bioRxiv* (2020) doi:10.1101/2020.12.21.20248640.

18. Naveca, F. *et al.* SARS-CoV-2 reinfection by the new Variant of Concern (VOC) P. 1 in Amazonas, Brazil. *virological. org. Preprint available at: https://virological. org/t/sars-cov-2-reinfection-by-thenew-variant-of-concern-voc-p-1-in-a mazonas-brazil/596* (2021).

19. Dejnirattisai, W. *et al.* Antibody evasion by the Brazilian P.1 strain of SARS-CoV-2. *Cold Spring Harbor Laboratory* 2021.03.12.435194 (2021) doi:10.1101/2021.03.12.435194.

20. Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein variants. *Elife* **9**, (2020).

21. Liu, Z. *et al.* Identification of SARS-CoV-2 spike mutations that attenuate monoclonal and serum antibody neutralization. *Cell Host Microbe* **0**, (2021).

22. Greaney, A. J. *et al.* Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human plasma antibodies. *Cell Host Microbe* **29**, 463-476.e6 (2021).

23. Wibmer, C. K. *et al.* SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19 donor plasma. *Nat. Med.* (2021) doi:10.1038/s41591-021-01285-x.

24. Zhou, D. *et al.* Evidence of escape of SARS-CoV-2 variant B.1.351 from natural and vaccine-induced sera. *Cell* (2021) doi:10.1016/j.cell.2021.02.037.

25. Wang, P. *et al.* Increased Resistance of SARS-CoV-2 Variant P. 1 to Antibody Neutralization. *BioRxiv* (2021).

26. Liu, Y. *et al.* Neutralizing Activity of BNT162b2-Elicited Serum. *N. Engl. J. Med.* (2021) doi:10.1056/NEJMc2102017.

27. Garcia-Beltran, W. F. *et al.* Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral immunity. *Cell* (2021) doi:10.1016/j.cell.2021.03.013.

28. West, A. P., Barnes, C. O., Yang, Z. & Bjorkman, P. J. SARS-CoV-2 lineage B.1.526 emerging in the New York region detected by software utility created to query the spike mutational landscape. *Cold Spring Harbor Laboratory* 2021.02.14.431043 (2021) doi:10.1101/2021.02.14.431043.

29. Annavajhala, M. K. *et al.* A novel SARS-CoV-2 variant of concern, B.1.526, identified in New York. *bioRxiv* (2021) doi:10.1101/2021.02.23.21252259.

30. Voloch, C. M. *et al.* Genomic characterization of a novel SARS-CoV-2 lineage from Rio de Janeiro, Brazil. *J. Virol.* (2021) doi:10.1128/JVI.00119-21.

31. Sanders, R. W. & Moore, J. P. Virus vaccines: proteins prefer prolines. *Cell Host Microbe* **29**, 327–333 (2021).

32. Amanat, F. *et al.* Introduction of two prolines and removal of the polybasic cleavage site lead to higher efficacy of a recombinant spike-based SARS-CoV-2 vaccine in the mouse model. *MBio* **12**, (2021).

33. Madhi, S. A. *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N. Engl. J. Med.* (2021) doi:10.1056/NEJMoa2102214.

34. Logunov, D. Y. *et al.* Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia. *Lancet* **396**, 887–897 (2020).

35. Dieterle, M. E. *et al.* A Replication-Competent Vesicular Stomatitis Virus for Studies of SARS-CoV-2 Spike-Mediated Cell Entry and Its Inhibition. *Cell Host Microbe* **28**, 486-496.e6 (2020).

36. Li, H. *et al.* Establishment of replication-competent vesicular stomatitis virus-based recombinant viruses suitable for SARS-CoV-2 entry and neutralization assays. *Emerg. Microbes Infect.* **9**, 2269–2277 (2020).

37. Case, J. B. *et al.* Replication-Competent Vesicular Stomatitis Virus Vaccine Vector Protects against SARS-CoV-2-Mediated Pathogenesis in Mice. *Cell Host Microbe* **28**, 465-474.e4 (2020).

38. Baum, A. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science* **369**, 1014–1018 (2020).

39. Yahalom-Ronen, Y. *et al.* A single dose of recombinant VSV-ΔG-spike vaccine provides protection against SARS-CoV-2 challenge. *Nat. Commun.* **11**, 6402 (2020).

40. Greaney, A. J. *et al.* Complete Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody Recognition. *Cell Host Microbe* **29**, 44-57.e9 (2021).

41. Oguntuyo, K. Y. *et al.* Quantifying absolute neutralization titers against SARS-CoV-2 by a standardized virus neutralization assay allows for cross-cohort comparisons of COVID-19 sera. *MBio* **12**, (2021).

42. Webb, N. E., Montefiori, D. C. & Lee, B. Dose–response curve slope helps predict therapeutic potency and breadth of HIV broadly neutralizing antibodies. *Nat. Commun.* **6**, 1–10 (2015).

43. Collier, D. A. *et al.* Sensitivity of SARS-CoV-2 B.1.1.7 to mRNA vaccine-elicited antibodies. *Nature* **593**, 136–141 (2021).

44. Cai, Y. *et al.* Structural basis for enhanced infectivity and immune evasion of SARS-CoV-2 variants. *bioRxivorg* (2021) doi:10.1101/2021.04.13.439709.

45. Zhu, X. *et al.* Cryo-electron microscopy structures of the N501Y SARS-CoV-2 spike protein in complex with ACE2 and 2 potent neutralizing antibodies. *PLoS Biol.* **19**, e3001237 (2021).

46. Laffeber, C., de Koning, K., Kanaar, R. & Lebbink, J. H. G. Experimental evidence for enhanced receptor binding by rapidly spreading SARS-CoV-2 variants. *bioRxiv* (2021) doi:10.1101/2021.02.22.432357.

47. Finkelstein, M. T. *et al.* Structural analysis of neutralizing epitopes of the SARS-CoV-2 spike to guide therapy and vaccine design strategies. *Viruses* **13**, 134 (2021).

48. Yuan, M. *et al.* Structural and functional ramifications of antigenic drift in recent SARS-CoV-2 variants. *bioRxivorg* (2021) doi:10.1101/2021.02.16.430500.

49. Yin, R. *et al.* Structural and energetic profiling of SARS-CoV-2 antibody recognition and the impact of circulating variants. *bioRxiv* (2021) doi:10.1101/2021.03.21.436311.

50. Gowthaman, R. *et al.* CoV3D: a database of high resolution coronavirus protein structures. *Nucleic Acids Res.* **49**, D282–D287 (2021).

51. Bos, R. *et al.* Ad26 vector-based COVID-19 vaccine encoding a prefusion-stabilized SARS-CoV-2 Spike immunogen induces potent humoral and cellular immune responses. *NPJ Vaccines* **5**, 91 (2020).

52. Walsh, E. E. *et al.* Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *N. Engl. J. Med.* **383**, 2439–2450 (2020).

53. Corbett, K. S. *et al.* SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen preparedness. *Nature* **586**, 567–571 (2020).

54. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260–1263 (2020).

55. Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E7348–E7357 (2017).

56. Cele, S. *et al.* Escape of SARS-CoV-2 501Y. V2 from neutralization by convalescent plasma. *medRxiv* (2021).

57. Hoffmann, M. *et al.* SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing antibodies. *Cell* (2021) doi:10.1016/j.cell.2021.03.036.

58. Muik, A. *et al.* Neutralization of SARS-CoV-2 lineage B.1.1.7 pseudovirus by BNT162b2 vaccine-elicited human sera. *Science* **371**, 1152–1153 (2021).

59. Shen, X. *et al.* SARS-CoV-2 variant B.1.1.7 is susceptible to neutralizing antibodies elicited by ancestral spike vaccines. *Cell Host Microbe* (2021) doi:10.1016/j.chom.2021.03.002.

60. Abdool Karim, S. S. & de Oliveira, T. New SARS-CoV-2 Variants — Clinical, Public Health, and Vaccine Implications. *N. Engl. J. Med.* (2021) doi:10.1056/NEJMc2100362.

61. Buchholz, U. J., Finke, S. & Conzelmann, K. K. … respiratory syncytial virus (BRSV) from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the human RSV leader region acts as a functional …. *J. Virol.* (1999).

62. Ramsburg, E. *et al.* A vesicular stomatitis virus recombinant expressing granulocyte-macrophage colony-stimulating factor induces enhanced T-cell responses and is highly attenuated for replication in animals. *J. Virol.* **79**, 15043–15053 (2005).

63. Case, J. B. *et al.* Neutralizing Antibody and Soluble ACE2 Inhibition of a Replication-Competent VSV-SARS-CoV-2 and a Clinical Isolate of SARS-CoV-2. *Cell Host Microbe* **28**, 475-485.e5 (2020).

64. Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat. Med.* (2020) doi:10.1038/s41591-020-0913-5.

65. Beaty, S. M. *et al.* Efficient and Robust Paramyxoviridae Reverse Genetics Systems. *mSphere* **2**, (2017).

66. Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative contribution to global health. *Glob Chall* **1**, 33–46 (2017).

67. Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill.* **22**, (2017).