Regular article

Neutrophil-mediated Oxidative Stress and Albumin Structural Damage Predict COVID-19-associated Mortality

Mohamed A. Badawy,[1,*] Basma A. Yasseen,[1,*] Riem M. El-Messiery,[2,*] Engy A. Abdel-Rahman,[1,3*] Aya A. Elkhodiry,[1,*] Azza G. Kamel,[1,*] Asmaa M. Shedra,[1] Rehab Hamdy,[1] Mona Zidan,[1] Diaa Al-Raawi,[1] Mahmoud Hammad,[4] Nahla Elsharkawy,[5] Mohamed El Ansary,[6] Ahmed Al-Halfawy,[7] Alaa Elhadad,[4] Ashraf Hatem,[8] Sherif Abouelnaga,[4] and Sameh S. Ali[1,†]

[1] Research Department, Children's Cancer Hospital Egypt 57357, Cairo, Egypt.

[2] Infectious Disease Unit, Internal Medicine Department, Faculty of Medicine, Cairo University, Cairo, Egypt.

[3] Pharmacology Department, Faculty of Medicine, Assuit University, Assuit, Egypt.

[4] Pediatric Oncology Department, National Cancer Institute, Cairo University and Children's Cancer Hospital 57357, Cairo, Egypt.

[5] Clinical pathology department, National Cancer Institute, Cairo University and Children's Cancer Hospital 57357, Cairo, Egypt.

[6] Department of Intensive Care, Faculty of Medicine, Cairo University, Cairo, Egypt.

[7] Department of Pulmonary Medicine, Faculty of Medicine, Cairo University, Cairo, Egypt.

[8] Department of Chest Diseases, Faculty of Medicine, Cairo University, Cairo, Egypt.

Running Title: Damage of Human Serum Albumin and COVID-19 Mortality.

*Equal contributions

[†] Corresponding author: Prof. Sameh S. Ali, Research Department, Children's Cancer Hospital Egypt 57357, Cairo, Egypt, Tel: +202225351500 Ext-7233, sameh.ali@57357.org

Word count for text: 5094; **Abstract word count**: 249

Number of Figures: 5 colored figures; **Number of Tables**: 2 + 1 supplementary table.

Reference count: 45

Abstract

Human serum albumin (HSA) is the frontline antioxidant protein in blood with established anti-inflammatory and anticoagulation functions. Here we report that COVID-19-induced oxidative stress inflicts structural damages to HSA and is linked with mortality outcome in critically ill patients. We recruited 25 patients who were followed up for a median of 12.5 days (1-35 days), among them 14 had died. Analyzing blood samples from patients and healthy individuals (n=10), we provide evidence that neutrophils are major sources of oxidative stress in blood and that hydrogen peroxide is highly accumulated in plasmas of non-survivors. We then analyzed electron paramagnetic resonance (EPR) spectra of spin labelled fatty acids (SLFA) bound with HSA in whole blood of control, survivor, and non-survivor subjects (n=10-11). Non-survivors' HSA showed dramatically reduced protein packing order parameter, faster SLFA correlational rotational time, and greater S/W ratio (strong-binding/weak-binding sites within HSA), all reflecting remarkably fluid protein microenvironments. Stratified at the means, Kaplan–Meier survival analysis indicated that lower values of S/W ratio and accumulated H_2O_2 in plasma significantly predicted in-hospital mortality (S/W<0.16, 80% (9/12) vs. S/W>0.16, 20% (2/10), p=0.008; plasma $[H_2O_2]$>7.1 μM, 83.3% (5/6) vs. 16.7% (1/6), p=0.049). When we combined these two parameters as the ratio (($S/W)/[H_2O_2]$) to derive a risk score, the resultant risk score lower than the mean (< 0.0253) predicted mortality with 100% accuracy (100% (6/6) vs. 0% (0/6), logrank χ^2 = 12.01, p = 5x10^{-4}). The derived parameters may provide a surrogate marker to assess new candidates for COVID-19 treatments targeting HSA replacements.

Keywords: Critically ill COVID-19 patients, COVID-19 Mortality, Neutrophils, Oxidative Stress, Human Serum Albumin Damage, Spin Labels, Electron Paramagnetic Resonance

Introduction

COVID-19 pandemic continues as a global health crisis while the underlying SARS-CoV-2 virus

defies all attempted treatment strategies. While writing this report there have been more than 110

million confirmed cases including around 2.5 million deaths worldwide according to the World Health

Organization Coronavirus Disease Dashboard (https://covid19.who.int/). Although 50% of cases are

reported to be in the 25-64 age group, the percentage of deaths increases dramatically with age, and

approximately 75% of deaths are in those aged 65 years and above (COVID-19 Hospitalization and Death

by Age | CDC). People in the age groups 30-39 years, 40-49 years, and 50-64 years are 4, 10, and 30

times more likely to die from COVID-19 complications compared to 18-29 years' age group.

Nevertheless, molecular and cellular factors contributing to mortality outcome in a homogeneous

cohort of patients are not yet clear. Lack of diagnostic markers that predict mortality in COVID-19

patients impedes current efforts to siege the pandemic. It is thus critical to identify prognostic tests that

can assess the risk of death in critically ill patients to guide clinical protocols and prioritize interventions.

Furthermore, mechanistic clues for determining the underlying molecular factors contributing to the

hypercoagulability, inflammation, and cytokine storm have been so far illusive. It is therefore imperative

to intensify efforts focusing on understanding the molecular pathophysiology of COVID-19 infection and

to identify prognostic markers to guide and prioritize clinical decisions.

Human serum albumin (HSA) is the most abundant constituent of soluble proteins in the

circulatory system. HSA has been suggested and used as a diagnostic and prognostic marker of

numerous diseases and conditions including ischemia, rheumatoid arthritis, cancer, septic shock among

many others. In addition to its numerous physiological and pharmacological functions including the

maintenance of blood/tissue osmotic balance (1), blood pH, metal cation transport and homeostasis (2,

3), nutrients and drug shuttling (4, 5), toxin neutralization (6, 7), HSA is suggested to be a major

circulating antioxidant (8, 9). HSA can remarkably bind with a diverse array of drugs and toxins thus

controlling their bioavailability and pharmacologic effects (10). It has been previously shown that more than 70% of the free radical-trapping capacity of serum was due to HSA (reviewed in (11)). Importantly, several reports indicated that inflammation enhances vascular permeability of various tissues to HSA apparently to confer antioxidant beneficial effects against reactive species released by activated neutrophils (12-14). Although currently without direct experimental evidence, neutrophilia-mediated oxidative stress was implicated in the COVID-19 pathology and speculated to exacerbate the inflammatory immune response eventually causing multi organ failure and death (15). We hypothesized that COVID-19-mediated oxidative stress may be differentially reflected in HSA's structure and functions and employed electron paramagnetic resonance (EPR) spin labeling spectroscopy to explore HSA's structural changes in correlation with severity and mortality of critically ill COVID-19 patients.

Spin labeled long-chain fatty acids (SLFA) are established probes to explore structural and functional changes in albumin by EPR spectroscopy (16, 17). This approach relies on the well-studied ability of albumin to strongly and exclusively bind with fatty acids in blood. Albumin has at least seven different specific binding sites for long chain fatty acids located in different domains within the protein (18-20). Effectively, structural and functional changes in HSA may be assessed through the detection of parallel changes in mobility and binding affinity of SLFAs, in addition to the distribution of the spin labels on the albumin molecule (17). EPR spectra of spin labels bound to different domains of the protein provide information on the local fatty acids/protein interactions, which may probe changes in the overall structure of the protein under unfolding or damaging conditions; Fig. 1A, (18). Here we compare changes that occur to the mobility, binding affinity, and distribution of the HSA-bound SLFA in whole blood and plasma from COVID-19 patients in critical care unit relative to those observed in normal healthy individuals.

Methods

Study design and participants

The present study aims to analyze HSA protein configuration statuses in the most severe cases of COVID-19 in comparison with control subjects. This is a prospective observational cohort study of patients with confirmed RT-PCR positive COVID-19. Nasopharyngeal swab RT-PCR results and Lung CT scans were combined to classify severe symptomatic COVID-19 cases. All patients were recruited from Kasr Alainy Cairo University Hospital/ ICU-facility at the Internal Medicine Quarantine Hospital. Supportive therapy including supplemental oxygen and symptomatic treatment were administered as required. Patients with moderate to severe hypoxia (defined as requiring fraction of inspired oxygen [FiO2] ≥40%) were transferred to the intensive care for further management including invasive mechanical ventilation when necessary. Patients recruited in the current study were divided into two arms based on future mortality outcome: Those survived past 10 days following blood samples collection (Sev-R) and those who died within 10 days of samples collection (Sev-D).

No power analysis was done. Sample size was based on sample availability and collection continued until a total of N=25 has been reached for COVID-19 patients diagnosed as severely infected with SARS-CoV-2 and admitted to the ICU during the period from 13 October 2020 to 25 January 2021. Within the follow-up time 14 patients had died. However, in a few cases and due to leukopenia or blood samples insufficiency to run every experiment on each subject, a number of 6-11 were analyzed and reported per condition. Demographic and clinical data of the studied 25 COVID-19 confirmed positive cases were categorized according to mortality and used for correlative analyses (Table 1). No randomization was done and clinical data were made available after EPR data collection and analysis have been performed. Patients who survived 10-days post sample collections were considered survivors (Sev-R) while those died within this period were placed in the Sev-D group.

Written informed consents were obtained from participants in accordance with the principles of the Declaration of Helsinki. For COVID-19 and control blood/plasma collection, Children's Cancer Hospital's Institutional Review Board (IRB) has evaluated the study design and protocol, IRB number 31-2020 issued on July 6, 2020.

Electron paramagnetic resonance (EPR) measurements

We measured EPR spectra at 37 °C using a Benchtop Magnettech MiniScope MS5000 spectrometer (now Bruker Biospin, Berlin) equipped with Biotemperature control and computerized data acquisition and analysis capabilities. Typical instrumental parameters during these measurements were: microwave frequency 9.47 GHz, microwave power 10 mW, modulation frequency 100 kHz, modulation amplitude 0.2 mT, magnetic field range 332-342 mT. Each spectrum was the average of 5 scans with scan time of 60 s. Spin-labelled stearic acids (Fig. 1): 2-(3-carboxypropyl)-4,4-dimethyl-2-tridecyl-3-oxazolidinyloxy (5-doxyl-stearic acid, 5-DSA) and 2-(14-carboxytetradecyl)-2-ethyl-4,4-dimethyl-3-oxazolidinyloxy (16-doxyl-stearic acid, 16-DSA) were purchased from Aldrich Chemical Co. (Milwaukee, U.S.A.). Spin labeled fatty acid (SLFA)-serum albumin complexes were formed as recommended (17) in 0.01 M phosphate buffer (pH 7.4). Conditions were optimized to avoid aggregation of free SLFA and to minimize contributions from albumin-unbound SLFA. Signals were stable over the measurement time and at least up to 1 h afterward as observed by following time evolution of the recorded EPR spectra.

Analysis of biophysical EPR parameters: Fluidity measurement was carried out as we reported (21) and explained (22) previously. Nitroxyl radicals SLFA probes were used to determine local fluidity near the protein/aqueous interface (5-DSA) or the hydrophobic protein cores (16-DSA) of HSA. Whole blood or collected plasma from all subjects were labelled with 5-DSA or 16-DSA. Ethanolic solutions (0.02 M) of the spin labels were added to the blood/plasma in the ratio 1:100 and the cells were measured at 37 °C.

From the spectrum of 5-DSA an order parameter (S) was derived by measuring the outer and inner

hyperfine splitting $2T_\parallel$ and $2T_\perp$ as defined in Fig. 1B (23, 24) using the formula:

$$S = \frac{T_\parallel - T_\perp}{T_{zz} - T_{xx}} \cdot \frac{a_N}{a'_N}$$

where

$$a_N = \frac{1}{3(T_{zz} + 2T_{xx})} \; ; \; a'_N = \frac{1}{3(T_\parallel + 2T_\perp)}$$

T_{xx} and T_{zz} are principal values of hyperfine tensor T taken as 0.61 mT and 3.24 mT; respectively (24, 25).

To estimate fluidity of the inner lipophilic protein compartment, we used 16-DSA, i.e. a spin probe

containing the nitroxide group attached on C16 that is located on the opposite terminal relative to the

charged carboxyl fatty acid terminus. Rotational correlation time (τ_c), which is taken as a measure of

microenvironment fluidity in the fatty acid carrier sites in the HSA protein core was calculated as (26):

$$\tau_c = 6.6 \times 10^{-10} \times \Delta h_{+1}[\left(\frac{h_{+1}}{h_{-1}}\right)^{\frac{1}{2}} - 1]$$

Where the parameters Δh_{+1}, h_{+1}, and h_{-1} are determined from EPR spectra as shown in Fig. 1C.

16-DSA spectra (Fig. 1C) were also analyzed to obtain the ratio between weakly (W) and strongly (S)

bound subpopulations of the spin label in sterically hollow versus relatively jammed protein

microenvironments (22, 24).

Phenotyping of peripheral blood by flow cytometry

Citrate-anticoagulated whole peripheral blood was incubated with RBCs lysis buffer for 15 min. Lysed blood was then centrifuged at 500 X-g for 5 min. Cells were washed twice with phosphate buffered saline (PBS) and then resuspended in PBS. Distribution status of platelets, neutrophils, monocytes and lymphocytes was measured in whole blood samples by 13-color flow cytometry as described using CytoFLEX system (Beckman Coulter Life Sciences CytoFLEX benchtop flow cytometer). Suspended cells were incubated with combinations of anti-human monoclonal antibodies for subset identification as follows: CD-42b-PE (Beckman Coulter life sciences, IM1417U) for platelets, CD14-PC7 (Beckman Coulter life sciences, A22331) for monocytes, CD66b-APC-Alexa Fluor 750 for neutrophils (Beckman Coulter life sciences, B08756) and CD3-ECD (Beckman Coulter life sciences, IM2705U) for lymphocytes cells were incubated for 30 minutes in the dark at room temperature. At the end of the incubation period, cells were washed with PBS and resuspended in 300 μl PBS. Samples were then analyzed by flow cytometry for gating platelet specific CD42b-PE positive population, neutrophil specific CD66b-APC-Alexa Fluor 750 positive population, monocytes specific CD14-PE positive population and lymphocytes specific CD3-ECD positive population. A number of 20,000 events were acquired and analyzed using CytExpert software to determine the percent and mean fluorescence intensities (MFI) of analyzed cell subsets.

Measurement of ROS by flow cytometry

The intracellular ROS generation by different cell populations was measured in whole blood samples using 2',7'-dichlorofluorescein-diacetate (DCF, Sigma Aldrich, D6883). Suspended cells were incubated with DCF (20 μM) and combinations of monoclonal antibodies; CD-42b-PE (Platelets), CD14-PC7 (monocytes), CD66b-APC-Alexa Fluor 750 (neutrophils), and CD3-ECD (lymphocytes) for 30 minutes at room temperature in the dark. Cells were then washed with PBS and resuspended in 300 μl of PBS. 20,000 events were recorded and analyzed using CytExpert program.

Determination of hydrogen peroxide concentrations in plasma

Catalase was used to specifically and quantitatively determine levels of hydrogen peroxide in identical plasma volumes collected from the study subjects. Oxygen levels are monitored and recorded while 50 μL batches of plasma from control, Sev-R, and Sev-D subjects sequentially infused into tightly air-controlled O2k chamber containing catalase (315 Units/mL) in deoxygenated buffer. A continuous stream of pure nitrogen gas was blown over the chamber's sealing cap to prevent oxygen diffusion into the working solution. In addition to the initial rise due to residual oxygen in the added plasma samples, the decomposition of hydrogen peroxide in these samples produced oxygen in a quantitative manner; i.e. decomposition of one mole of H_2O_2 produced ½-mole O_2. To verify the assay, we measured the released oxygen upon adding increasing concentrations of standard hydrogen peroxide solution in PBS buffer. Linear fitting of the plotted $[O_2]$ versus $[H_2O_2]$ relation yielded a slope = 0.47 ± 0.03 (Pearson's r = 0.994, p = 5.6 x 10^{-4}), which is very close to the theoretically expected value of 0.5.

Measurement of ROS in isolated neutrophils by fluorescence imaging

The intracellular ROS generation of isolated neutrophils was detected and quantified using the ROS-sensitive DCF dye. Isolated cells were seeded and incubated for 30 minutes at 37°C followed by centrifugation to form a monolayer. Cells were then stained with DCF at a final concentration of 0.3 mM in PBS for 30 minutes. Cells were washed and stained with DAPI (Hoechst 33342 Solution, ThermoFischer Scientific, 62249) at a final concentration of 10 μM in PBS for 30 minutes at 37°C. Cytation 5 Cell Imaging Multi-Mode Reader (Agilent) was used to acquire images using 20X lens and the proper fluorescence filter cubes (λ_{ex} = 500 ± 12 nm and λ_{em} = 542 ± 14 nm). Images were processed to quantify fluorescence intensity using Gen5 software package 3.08.

Determination of plasma albumin concentrations

Albumin concentration in plasma was measured using bromocresol green (BCG) dye as the color intensity of the complex formed is proportional to albumin concentration. Standard curve was

generated using standard albumin and samples were measured in 96 well plate at 630 nm using

Cytation 5 (Agilent). Concentrations were calculated using the equation generated from standard curve

and presented in mg/mL.

Statistical analysis

Statistical analysis and data graphing were performed using OriginPro 2017 (OriginLab Corporation,

Northampton, USA). For comparisons of means between three or more independent groups, Tukey test

ANOVA for multiple comparison was performed. For correlation analysis, spearman rank correlation was

performed. Pearson's correlation coefficient was calculated for measuring the association between

variables of interest based on the method of covariance which also gives information about the

magnitude of the association, or correlation, as well as the direction of the relationship. p-values < 0.05

for correlations or means' comparisons were considered significant. Categorical variables are reported

as counts and percentages while continuous variables are expressed as mean ± standard deviation.

Differences between percentages were assessed by Pearson's χ^2 tests or Fisher exact tests when the

number of observations per group were less than 5. The χ^2 tests provided results that tested the

hypothesis that the mortality and a given variable (e.g. sex or a comorbidity) are independent. When p is

less than the significant level of 0.05, there is significant evidence of association between mortality and

the variable. Log-rank Kaplan–Meier survival analyses were carried out to estimate probability of

survival of COVID-19 patients in relation with cut-off thresholds arbitrarily stratified at mean values of

various parameters. The log-rank test for trends reports a χ^2-value and computes a p value testing the

null hypothesis that there is no linear trend between column order and median survival.

Data sharing statement

Original datasets and detailed protocols are available upon request from the corresponding author

sameh.ali@57357.org.

Results

Demographic, clinical, and laboratory hematologic characteristics of COVID-19 patients

Table 1 lists demographic data, comorbidities, ongoing medications, and administered anti-COVID-19 medications applied to treat current study participants that were divided into survivors (Sev-R) and deceased (Sev-D). Compared with recovered, deceased patients' mean age was slightly older (Tukey ANOVA, p<0.05). No other clinical or demographic characteristic showed statistically significant difference between Sev-R and Sev-D groups when analyzed by Pearson's Chi-Square test except that the frequency of remdesivir administration was significantly higher in Sev-D groups (p<0.05). In **Table 2** we show and statistically compare laboratory results of survivors vs. non-survivor COVID-19 groups. Although when comparing all parameters in the two COVID-19 groups we observed changes following the same reported trends in the literature, means' comparisons by Tukey test reported non-significant changes with strong trends observed only for C reactive protein (greater levels in Sev-D group, p = 0.050) and albumin (lower levels in Sev-D group, p = 0.067). Nevertheless, non-survivors' blood carried the frequently observed hallmarks of increased CRP, D-dimer, IL-6, ferritin, and the liver enzymes ALT and AST (Reviewed in: (27, 28)). However, it is concievable that the clinical severe category and the same ICU status of patients in the two groups in addition to relatively small sample size underlie the observed lack of robust statistical differences between these parameters.

Neutrophils are a major source of reactive oxygen species

It has been recently proposed that the high neutrophil-to-lymphocyte ratio observed in critically ill COVID-19 patients may tip the redox homeostasis due to increased reactive oxygen species production (15). Our hypothesis implicates elevated oxidative stress as a major cause of HSA damage in

severe COVID-19 patients. As a result, we started by following the dependence of clinical outcomes and mortality on ROS levels in blood cells. First, we used flowcytometry to assess percentages of neutrophils, lymphocytes, and platelets in all patients as described in Materials and Methods (Fig. 2A). Furthermore, we used the ROS-sensitive DCF dye to probe intracellular ROS levels in various cell populations in whole blood from all groups. Fig. 2 shows that while lympocyte counts decrease, a parallel dramatic increase in neutrophil counts (% total) was observable when going from Control (40.78 ± 14.0, n=9) to Sev-R (64.0 ± 20.0, n=10) to Sev-D (76.4 ± 6.8, n=11) groups (overall ANOVA p = 3.9 x 10^{-5}). Similar trend was clearly seen in the heat map depicting parameters for all patients analyzed by flow cytometry (Fig. 2B). It is also clear from Fig. 2B&C that changes in DCF-positive neutrophils follows similar trend observed for neutrophil counts. To confirm this relation we compared neutrophil counts with DCF-positive neutrophil counts and found that the two parameters were strongly correlated (Pearson's r = 0.8, p = 3 x 10^{-7}), Fig. 2C. Moreover, both parameters individually showed statistically significant increases in both of the studied COVID-19 groups when compared with control group (Fig. 2D&E). These results suggest that neutrophils are major sources of elevated oxidative stress in critically ill patients. Note that the observed trends in platelets, lymphocyte, neutrophils, and neutrophil-to-lymphocyte ratio (NLR) are similar to reported values (29, 30).

Hydrogen peroxide levels in plasma correlate with mortality

Next, we reasoned that elevated oxidative stress in both groups with critical COVID-19 infection would be echoed in plasma levels of hydrogen peroxide. We used a highly specific catalase-based assay that we developed and verified in our laboratory to quantify [H_2O_2] in plasma samples of all groups. The assay relies on high resolution detection and quantification of released oxygen due to hydrogen peroxide decompostion by catalase, Fig. 3A&B. We constructed a calibration curve to confirm the catalase-mediated H_2O_2 to O_2 stoicheometric conversion (Fig. 3B). A linear relation was obtained with zero intercept and slope of 0.47±0.03 which closely matches the theoretically expected value of 0.5

(95% CI: 0.37–0.56, p = 5.6 x 10^{-4}, Pearson's r = 0.994). Indeed, we detected striking differences between

groups even with relatively small sample sizes (Mean±SD, Control, n=7: 2.57±0.57, Sev-R, n=6: 5.37±1.0,

Sev-D, n=7: 8.8±1.7 ; overall ANOVA p-value = 1.15 x 10^{-7} ; Fig. 3C). The differences between groups have

reached statistical significance (Sev-R vs. Control, 95% CI: 1.08–4.51, p = 0.0017; Sev-D vs. Control, 95%

CI: 4.58–7.87, p = 0.0; Sev-D vs. Sev-R, 95% CI: 1.72–5.15, p = 2.3 x 10^{-4}). It appears from these results

that a measure of oxidative stress; *i.e.* [H_2O_2] in plasma is doubled in survivors and quadropled in

deceased COVID-19 patients relative to controls' plasma average levals.

To confirm this finding we performed DCF fluorescence imaging on freshly isolated neutrophils

of representative group of individuals from each group, Fig. 3D. We simultaneously stained neutrophils'

nuclei with DAPI (blue stain) to follow nuclear morphologic changes and DNA diffusion in all groups.

Although requiring more detailed studies, a closer look on the acquired images of DAPI-stained

neutrophils from a survivor patient showed significantly reduced average neutrophil size (control DNA

area, 144.8 ± 93.7 μm^2 (133 cells analyzed) vs. Sev-R DNA area, 36.0 ± 7.5 μm^2, (241 cells analyzed),

Welsh corrected two samples t test p = 0) with tendencies towards more condensed, more segmented

nuclei and frequent C shaped chromatin. However, neutrophils from a non-survivor exhibited diffused

chromatin and possessed in average a 26% larger DNA area than normal cells and roughly 5 times that

of Sev-R neutrophils (DNA area (97 cells analyzed), 182.7 ± 171 μm^2, p = 0 *vs*. both control and Sev-R

groups). Inspection of the DCF flourescence images indicated that the non-survivor's neutrophils

contained larger populations of what appears to be toxic granules and cytoplasmic vacuoles that are

highly ROS-positive. Analysis of mean DCF fluorescence intensities (MFI) per cell confirmed results

obtained by flow cytometry and catalase assay reporting increased levels of ROS in the order control <<

Sev-R < Sev-D (DCF MFI ± SD (x 10^3): Control (282 cells analyzed), 6.7 ± 0.7; Sev-R (351 cells analyzed),

10.9 ± 0.9; Sev-D (368 cells analyzed), 13.0 ± 1.1, Welsh corrected two samples t test, p = 0 for all

comparisons).

EPR-determined biophysical parameters reflecting albumin conformational changes are consistent

predictors of COVID-19 mortality

It has been previously shown that non-survivor COVID-19 patients exhibit mild but consistent

hypoalbuminemia relative to survivors (31). We started by assessing albumin levels in the studied

cohort of subjects to confirm if they follow similar trends. We found that [albumin] in plasma decreased

in the order Control > Sev-R >> Sev-D (Mean±SD, Control, n=8: 40.45±10.93 mg/mL, Sev-R, n=8:

30.66±10.99 mg/mL, Sev-D, n=10: 24.71±5.88 mg/mL; overall ANOVA *p*-value 0.006; Fig. 4A). We only

detected statistically significant decrease in [albumin] in plasma of Sev-D group (Sev-R vs. Control, 95%

CI: -21.44–1.87, *p* = 0.11; Sev-D vs. Control, 95% CI: -26.79– -4.67, *p* = 0.004; Sev-D vs. Sev-D, 95% CI: -

17.0–5.11, *p* = 0.38). The reference range of HSA concentration in serum is approximately 35-50 mg/mL,

but we and other groups found that COVID-19 associated mortality correlates with lowered [albumin] <

30 mg/mL (31). However, high prevelance of hypoalbuminemia in numerous disease states and the

age/sex-dependent wide dynamic range of this protein concentration limits its diagnostic utility (32). As

a result, we investigated biophysical parameters pertaining to HSA protein configuration in whole blood

and plasma of all groups as reflectors of this critical protein functions.

Previous studies reported that allosteric changes in HSA may be utilized to reflect critical

functional changes in albumin and explored diagnostic and prognostic values of these changes in cancer

(17, 33). It has also been found that long chain fatty acids binding alters the interactive binding of

ligands in the two major drug binding sites of HSA (34). We employed EPR spectroscopy to probe SLFAs'

binding statuses and protein configuration in all groups as detailed in the Methods' section above.

Analysis of the 5-DSA and 16-DSA EPR spectra revealed remarkable changes in spectral features

between control and COVID-19 groups (Fig. 4B, 5-DSA; Fig. 4C, 16-DSA in whole blood). For example, the

hyperfine coupling tensor element $2T_{\parallel}$ (defined in Fig. 1B and used to calculate the protein packing order

parameter *S*) is sensitive to microenvironmental effects (such as polarity, H-bonding, electrostatic

interactions, etc.) on the spin probe that is localized in one of the HSA native fatty acids binding pockets. Also, the S/W ratio (see Fig. 1C) corresponding to strongly-bound/weakly-bound populations of 16-DSA spin probe may reflect changes in protein folding that can alter protein-fatty acid interactions. Furthermore, the rotational correlation time τ_c which is a measure of the spin probe rotational mobility is also analyzed.

Calculated EPR spectral parameters for all groups are listed in Supplementary Table 1 including exact ANOVA and Tukey test *p* values along with the number of subjects analyzed. Both 5-DSA and 16-DSA were used to probe the degree of local interactions in sites where the spin labeled fatty acid is buried in the protein interior (high mobility, 16-DSA) and closer to the protein-aqueous interface (low mobility due to interactions with water molecules and polar amino acids, 5-DSA)(35). Indeed, 5-DSA reflected significantly greater *S* parameter values relative to 16-DSA both in plasma and in whole blood (Fig. 4D). However, independent of the spin probe and both in plasma and whole blood samples, the order parameter has been consistently lower in Sev-R which was further decreased in Sev-D patients relative to control group (Fig. 4D). In whole blood, similar results that showed more statistically robust differences have been observed.

Calculations of τ_c as described in Methods showed rotational mobility of 16-DSA is significantly faster when bound with HSA from COVID-19 patients relative to that from control subjects in plasma or whole blood. Finally, similar trends have been observed for the S/W parameter which signifies contributions of the strongly and weakly bound components of 16-DSA in different fatty acids pockets. Taken together, these results indicate that COVID-19 pathology is associated with extensive structural changes in the HSA protein that imply the prevelance of malfunctional derivatives of this critical protein.

Hampered water accessibility into HSA/fatty acid pockets in whole blood of COVID-19 patients

We followed water accessibility towards deep pockets carrying the spin labels through kinetic analysis of the nitroxide radical EPR silencing by the water-soluble ascorbate anion (36); Fig. 4(G-J). Under matching experimental conditions, 16-DSA and 5-DSA/HSA signals in whole blood of control subjects decayed remarkably faster when compared with both Sev-R and Sev-D (n=3 per group, p<0.05). Weaker and slower disappearance of the EPR signal of COVID-19 patients by ascorbate suggests less accessible space towards the nitroxide moiety of the spin label within the protein. It is clear from these data that HSA of COVID-19 patients is generally less water-accessible relative to control ones. However, the core of the HSA of both COVID-19 groups were not significantly different in terms of water-accessibility.

Association of EPR-determined HSA structural changes with neutrophil counts and measures of oxidative stress

In order to verify our proposed links between oxidative stress and albumin damage we applied linear correlations between plasma levels of hydrogen peroxide and EPR-calculated parameters pertaining to protein packing order parameter (S), fatty acid mobility (τ_c), and S/W ratio in all groups (n=19-26, Fig. 5A-D). Furthermore, to substantiate our suggestion that neutrophils are major sources of oxidative stress implicated in COVID-19 severity and mortality, we explored linear correlations between the populations of DCF positive neutrophils and τ_c (Fig. 5E) or S/W ratio (Fig. 5F). Despite relatively small numbers of subjects analyzed, correlations of biophysical parameters showed moderate-to-strong negative correlations with [H_2O_2] in plasma or with DCF positive populations (% of total); Pearson's r = -0.6– -0.76, p < 0.01 for all correlations. These observations may suggest that neutrophil-mediated oxidative stress in critically ill COVID-19 patients is a potential contributor to the observed structural changes in HSA of those patients.

Analysis of Kaplan–Meier estimates of time-to-mortality from blood sample collections during ICU hospitalization

Finally, we stratified patients into two groups, patients showing values below the means and patients with values above the means of each of the studied parameters: (*i.e.* S/W mean=0.1639 (0.090-0.261); plasma [albumin] mean = 0.26.69 mg/mL (16.42-51.48); and plasma [H_2O_2] mean = 7.1 μM (4.4-11.6)). Patients with lower values of the S/W ratio showed significantly higher in-hospital mortality (81.8% vs. 18.2%, logrank χ^2 = 6.95, p = 0.008, Fig. 5G). Similarly, patients with accumulated H_2O_2 in their plasma showed higher mortality (83.3% vs. 16.7%, logrank χ^2 = 3.85, p = 0.049, Fig. 5H). However, when we combined these two parameters to derive a risk score as the ratio ((S/W)/[H_2O_2]), the resultant risk score lower than the mean (< 0.0253) predicted mortality with 100% accuracy (100% vs. 0%, logrank χ^2 = 12.01, p = 5x10^{-4}, Fig. 5I). Under our current conditions, although showing weak statistical trends both of the order parameter (not shown) and the plasma albumin level (Fig. 5J) were not statistically significant determinants of in-hospital mortality.

Discussion

Human serum albumin is a pivotal protein with diverse multifaceted functions that are increasingly reported to reflect physiological and contribute to pathological states (32, 37). Closer to the context of COVID-19 pathophysiology, critical roles of HSA have been suggested as it acts as an anti-inflammatory and antioxidant protein (38), anticoagulant agent (39, 40), inhibitor of oxidative stress-mediated clotting and platelet activation (41, 42), drug-dependent allosteric carrier and regulator (43), and a potent heme scavenger that may also exhibit globin-like reactivity (44). In the light of these established functions it is natural to suggest albumin as a frontline protection against COVID-19-

associated lethality resulting from cytokine storm, oxidative stress, blood clotting, and the ensuing organ failure. Although hypoalbuminemia has been repeatidly reported as a predictor of mortality in COVID-19 (31, 45), low albumin levels may result from surgery, dialysis, abdominal infections, liver failure, pancreatitis, respiratory distress, bypass surgery, ovarian problems caused by fertility drugs, and many other conditions. High prevelance of hypoalbuminemia in numerous disease states and the age/sex-dependent wide dynamic range of this protein concentration limits its diagnostic utility (32). As a result, we investigated biophysical parameters pertaining to HSA protein structure in whole blood and plasma of all groups as reflectors of this critical protein functions.

We designed the present study to explore how COVID-19-caused mortality is related to oxidative stress and whether ROS-induced albumin damage can be used to predict such outcome. To avoid disparities related to clinical statuses and interventional protocols we restricted our subject recruitment to a homogeneous cohort of patients in terms of diseases severity, hospitalization, and interventions. Both sexes were represented in the overal COVID-19 subjects (44% females) and sex was not found to affect mortality (p = 0.86). The mean age of the studied subjects was 66.7 ± 8.9 y (42-81), and non-survivors were slightly orlder (p = 0.04). No other clinical or demographic characteristic showed statistically significant difference between survivors and non-survivors which further indicates the homogeniety of the studied pool of subjects.

Our current data provide converging and novel evidence that neutrophils are major players in oxidative stress associating inflammation in critically ill COVID-19 patients, especially those that don't survive the infection. We detected increased neutrophil count, increased ROS-positive neutrophil population, increased intracellular ROS level in these neutrophils, and a remarkably elevated hydrogen peroxide residue in plasma of non-survivor COVID-19 patients relative to control and survivor groups. Moreover, these oxidative stress measures were found to strongly correlate with EPR-detected strucrtural damages of HSA. Although we found a weak trend that albumin levels were inversely

associated with overall mortality, a biophysical measure of structural changes inflicted on albumin (S/W) was found to consistently predict overall mortality (p = 0.008) even with a relatively small n=22 sample size. Moreover, plasma levels of hydrogen peroxide was also found to predict in-ICU mortality with even smaller number of analyzed samples (n=12, p=0.049). This substantiates the utilization of these parameters as accurate mortality predictors of critically ill COVID-19 patients. Determined threshold values of these parameters can possibly help identifying patients in need for urgent medical attention, and may provide novel markers to assess new candidates for COVID-19 treatments targeting HSA replacements. We thus suggest the importance of studying the potential effects of albumin replacement therapy on clinical outcomes of Covid-19 patients.

Acknowledgments

The present work was funded by the Association of Friends of the National Cancer Institute and the Children's Cancer Hospital Foundation.

Authorship Contributions

MAB, BAY, EAA-R, AAE, and AK performed experiments, analyzed results, and assisted in the manuscript preparation. Subjects' recruitment, ethical approvals, clinical follow-up and assessments were coordinated by RE-M. AMS, RH, MZ, and NE assisted in experimental work. MH, MEl-A, AEl-H, AE, AH, and SA provided logistical support and critical manuscript revisions. SSA conceived and directed the project, designed experiments, analyzed data, and wrote the manuscript.

Disclosure of Conflicts of Interest

All authors declare no competing financial interests.

References

1. Singh-Zocchi M, Andreasen A, Zocchi G. Osmotic pressure contribution of albumin to colloidal interactions. Proceedings of the National Academy of Sciences of the United States of America. 1999;96(12):6711-5.

2. Bal W, Sokolowska M, Kurowska E, Faller P. Binding of transition metal ions to albumin: sites, affinities and rates. Biochimica et biophysica acta. 2013;1830(12):5444-55.

3. Stewart AJ, Blindauer CA, Berezenko S, Sleep D, Sadler PJ. Interdomain zinc site on human albumin. Proceedings of the National Academy of Sciences of the United States of America. 2003;100(7):3701-6.

4. Fujiwara S, Amisaki T. Fatty acid binding to serum albumin: molecular simulation approaches. Biochimica et biophysica acta. 2013;1830(12):5427-34.

5. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major update to the DrugBank database for 2018. Nucleic acids research. 2018;46(D1):D1074-D82.

6. Ascenzi P, Bocedi A, Notari S, Fanali G, Fesce R, Fasano M. Allosteric modulation of drug binding to human serum albumin. Mini reviews in medicinal chemistry. 2006;6(4):483-9.

7. Vorum H, Honore B. Influence of fatty acids on the binding of warfarin and phenprocoumon to human serum albumin with relation to anticoagulant therapy. The Journal of pharmacy and pharmacology. 1996;48(8):870-5.

8. Cha MK, Kim IH. Glutathione-linked thiol peroxidase activity of human serum albumin: a possible antioxidant role of serum albumin in blood plasma. Biochemical and biophysical research communications. 1996;222(2):619-25.

9. Loban A, Kime R, Powers H. Iron-binding antioxidant potential of plasma albumin. Clinical science. 1997;93(5):445-51.

10. Fasano M, Curry S, Terreno E, Galliano M, Fanali G, Narciso P, et al. The extraordinary ligand binding properties of human serum albumin. IUBMB life. 2005;57(12):787-96.

11. Roche M, Rondeau P, Singh NR, Tarnus E, Bourdon E. The antioxidant properties of serum albumin. FEBS letters. 2008;582(13):1783-7.

12. Cross CE, van der Vliet A, O'Neill CA, Louie S, Halliwell B. Oxidants, antioxidants, and respiratory tract lining fluids. Environmental health perspectives. 1994;102 Suppl 10:185-91.

13. Halliwell B. Albumin--an important extracellular antioxidant? Biochemical pharmacology. 1988;37(4):569-71.

14. Sitar ME, Aydin S, Cakatay U. Human serum albumin and its relation with oxidative stress. Clinical laboratory. 2013;59(9-10):945-52.

15. Laforge M, Elbim C, Frere C, Hemadi M, Massaad C, Nuss P, et al. Tissue damage from neutrophil-induced oxidative stress in COVID-19. Nature reviews Immunology. 2020;20(9):515-6.

16. Ge MT, Rananavare SB, Freed JH. ESR studies of stearic acid binding to bovine serum albumin. Biochimica et biophysica acta. 1990;1036(3):228-36.

17.   Haeri HH, Schunk B, Tomaszewski J, Schimm H, Gelos MJ, Hinderberger D. Fatty Acid Binding to Human Serum Albumin in Blood Serum Characterized by EPR Spectroscopy. ChemistryOpen. 2019;8(5):650-6.

18.   Bhattacharya AA, Grune T, Curry S. Crystallographic analysis reveals common modes of binding of medium and long-chain fatty acids to human serum albumin. Journal of molecular biology. 2000;303(5):721-32.

19.   Curry S, Brick P, Franks NP. Fatty acid binding to human serum albumin: new insights from crystallographic studies. Biochimica et biophysica acta. 1999;1441(2-3):131-40.

20.   Simard JR, Zunszain PA, Hamilton JA, Curry S. Location of high and low affinity fatty acid binding sites on human serum albumin revealed by NMR drug-competition analysis. Journal of molecular biology. 2006;361(2):336-51.

21.   Head BP, Peart JN, Panneerselvam M, Yokoyama T, Pearn ML, Niesman IR, et al. Loss of caveolin-1 accelerates neurodegeneration and aging. PloS one. 2010;5(12):e15697.

22.   E AA-R, Mahmoud AM, Khalifa AM, Ali SS. Physiological and pathophysiological reactive oxygen species as probed by EPR spectroscopy: the underutilized research window on muscle ageing. The Journal of physiology. 2016;594(16):4591-613.

23.   Gordon LM, Looney FD, Curtain CC. Fatty-acid spin probe interactions with erythrocyte ghosts and liposomes prepared from erythrocyte ghosts. The Journal of membrane biology. 1989;111(2):155-68.

24.   Marczak A, Kowalczyk A, Wrzesien-Kus A, Robak T, Jozwiak Z. Interaction of doxorubicin and idarubicin with red blood cells from acute myeloid leukaemia patients. Cell biology international. 2006;30(2):127-32.

25.   Sauerheber RD, Zimmermann TS, Esgate JA, VanderLaan WP, Gordon LM. Effects of calcium, lanthanum, and temperature on the fluidity of spin-labeled human platelets. The Journal of membrane biology. 1980;52(3):201-19.

26.   Schreier S, Polnaszek CF, Smith IC. Spin labels in membranes. Problems in practice. Biochimica et biophysica acta. 1978;515(4):395-436.

27.   Singh K, Mittal S, Gollapudi S, Butzmann A, Kumar J, Ohgami RS. A meta-analysis of SARS-CoV-2 patients identifies the combinatorial significance of D-dimer, C-reactive protein, lymphocyte, and neutrophil values as a predictor of disease severity. International journal of laboratory hematology. 2020.

28.   Velavan TP, Meyer CG. Mild versus severe COVID-19: Laboratory markers. International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases. 2020;95:304-7.

29.   Sun S, Cai X, Wang H, He G, Lin Y, Lu B, et al. Abnormalities of peripheral blood system in patients with COVID-19 in Wenzhou, China. Clinica chimica acta; international journal of clinical chemistry. 2020;507:174-80.

30.   Yang AP, Liu JP, Tao WQ, Li HM. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. International immunopharmacology. 2020;84:106504.

31.   Violi F, Cangemi R, Romiti GF, Ceccarelli G, Oliva A, Alessandri F, et al. Is Albumin Predictor of Mortality in COVID-19? Antioxidants & redox signaling. 2020.

32.   Levitt DG, Levitt MD. Human serum albumin homeostasis: a new look at the roles of synthesis, catabolism, renal and gastrointestinal excretion, and the clinical value of serum albumin measurements. International journal of general medicine. 2016;9:229-55.

33.   Kazmierczak SC, Gurachevsky A, Matthes G, Muravsky V. Electron spin resonance spectroscopy of serum albumin: a novel new test for cancer diagnosis and monitoring. Clinical chemistry. 2006;52(11):2129-34.

34.   Yamasaki K, Hyodo S, Taguchi K, Nishi K, Yamaotsu N, Hirono S, et al. Long chain fatty acids alter the interactive binding of ligands to the two principal drug binding sites of human serum albumin. PloS one. 2017;12(6):e0180404.

35.   Gantchev TG, Shopova MB. Characterization of spin-labelled fatty acids and hematoporphyrin binding sites interactions in serum albumin. Biochimica et biophysica acta. 1990;1037(3):422-34.

36.   Pavicevic AA, Popovic-Bijelic AD, Mojovic MD, Susnjar SV, Bacic GG. Binding of doxyl stearic spin labels to human serum albumin: an EPR study. The journal of physical chemistry B. 2014;118(37):10898-905.

37.   Coverdale JPC, Katundu KGH, Sobczak AIS, Arya S, Blindauer CA, Stewart AJ. Ischemia-modified albumin: Crosstalk between fatty acid and cobalt binding. Prostaglandins, leukotrienes, and essential fatty acids. 2018;135:147-57.

38.   Inoue M, Nakashima R, Enomoto M, Koike Y, Zhao X, Yip K, et al. Plasma redox imbalance caused by albumin oxidation promotes lung-predominant NETosis and pulmonary cancer metastasis. Nature communications. 2018;9(1):5116.

39.   Doweiko JP, Nompleggi DJ. The role of albumin in human physiology and pathophysiology, Part III: Albumin and disease states. JPEN Journal of parenteral and enteral nutrition. 1991;15(4):476-83.

40.   Ronit A, Kirkegaard-Klitbo DM, Dohlmann TL, Lundgren J, Sabin CA, Phillips AN, et al. Plasma Albumin and Incident Cardiovascular Disease: Results From the CGPS and an Updated Meta-Analysis. Arteriosclerosis, thrombosis, and vascular biology. 2020;40(2):473-82.

41.   Basili S, Carnevale R, Nocella C, Bartimoccia S, Raparelli V, Talerico G, et al. Serum Albumin Is Inversely Associated With Portal Vein Thrombosis in Cirrhosis. Hepatology communications. 2019;3(4):504-12.

42.   Tian W, Jiang W, Yao J, Nicholson CJ, Li RH, Sigurslid HH, et al. Predictors of mortality in hospitalized COVID-19 patients: A systematic review and meta-analysis. Journal of medical virology. 2020;92(10):1875-83.

43.   Fanali G, di Masi A, Trezza V, Marino M, Fasano M, Ascenzi P. Human serum albumin: from bench to bedside. Molecular aspects of medicine. 2012;33(3):209-90.

44.   Ascenzi P, di Masi A, Fanali G, Fasano M. Heme-based catalytic properties of human serum albumin. Cell death discovery. 2015;1:15025.

45.   Huang J, Cheng A, Kumar R, Fang Y, Chen G, Zhu Y, et al. Hypoalbuminemia predicts the outcome of COVID-19 independent of age and co-morbidity. Journal of medical virology. 2020;92(10):2152-8.

Table 1. Demographic and clinical characteristics of the studied subjects.

	Sev-R	Sev-D	Tukey 95%-CI	p
n	11	14		
Age (mean ± SD)	62.6 ± 9.7	69.8 ± 7.0	0.33-14.11	0.04
Male	54.54%	57.14%		0.896†
sO$_2$ (mean ± SD)	78.57 ± 13.6	73.71 ± 17.07	-20.42–10.7	0.52
Hypertension	27.27%	42.85%		0.65†
Diabetes	18.18%	50%		0.09†
Cardiovascular disease	0%	14.28%		0.2†
Cancer	0%	7.14%		0.36†
Bronchial asthma	0%	7.14%		0.36†
ACE inhibitors	0%	7.14%		0.36†
ARBs	9.09%	7.14%		0.85†
calcium channel blocker	9.09%	7.14%		0.85†
Beta Blockers	0%	7.14%		0.36†
Diuretics	0%	7.14%		0.36†
Sulphonylurea	9.09%	21.42		0.4†
Biguanides	0%	7.14%		0.36†
Insulin	18.18%	7.14%		0.39†
Anticoagulant	27.27%	35.71%		0.65†
Steroids	45.45%	57.14%		0.56†
Hydroxychloro-quine	18.18%	7.14%		0.4†
IL-6 receptor antibody	9.09%	35.71%		0.12†
Proton-pump inhibitor	9.09%	14.28%		0.69†
Azithromycin	9.09%	28.57%		0.22†
Cephalosporin	18.18%	14.28%		0.79†
Carbapenem	9.09%	42.85%		0.06†
Oxazolidinone	27.27%	35.71%		0.65†
Fluoro-quinolone	36.36%	35.71%		0.97†
Nitrofuran	9.09%	0%		0.25†
Remdesivir	0%	35.71%		0.026†
Ivermectin	0%	21.42%		0.10†

sO$_2$, blood oxygen saturation level, ACE, angiotensin converting enzyme; ARBs, angiotensin receptor blockers; IL-6, interleukin-6. † p values obtained through Pearson's χ^2 test.

Table 2. Laboratory parameters of the current study patients.

	Sev-R (mean ± SD)	Sev-D (mean ± SD)	Tukey 95%-CI	*p*
WBCs (x 10^3/mL)	10.85 ± 4.3	12.5 ± 5.1	-2.62–5.94	0.42
Platelets (x 10^6/mL)	274.3 ± 88.3	233.1 ± 92.2	-116.7–34.1	0.27
INR	1.34 ± 0.63	1.26 ± 0.23	-0.49–0.33	0.68
CRP (mg/L)	39.95 ± 42.6	90.6 ± 66.9	-0.1–101.4	0.05
D-dimer (µg/mL)	1.38 ± 1.7	3.79 ± 4.6	-0.81–5.64	0.13
IL-6 (pg/mL)	214.8 ± 442.6	393.0 ± 749.6	-427–783	0.54
Ferritin	715.6 ± 426	1081 ± 622	-94–826	0.11
Albumin (g/mL)	31.8 ± 4.8	26.9 ± 5.1	-1.0–0.04	0.067
Hemoglobin (g/dL)	12.18 ± 1.3	12.16 ± 2.0	-1.5–1.46	0.97
ALT (U/L)	29.6 ± 20.2	40.1 ± 37.3	-15.4–36.4	0.40
AST (U/L)	34.7 ± 20.5	44.3 ± 29.8	-12.2–31.4	0.37
Serum creatinine (mg/dL)	1.24 ± 0.8	1.78 ± 1.7	-0.64–1.73	0.35

WBCs, white blood cells; INR, international normalized ratio; CRP, high-sensitivity C reactive protein; ICU, intensive care unit; PLTs, platelets; ALT, alanine transaminase; AST, aspartate transaminase. The Tukey's calculated *p*-values as well as upper and lower 95%-confidence levels for the Sev-R vs. Sev-D means' comparisons are given.

Figure Legends

Fig. 1. A) HSA crystal structure containing 7 copies of stearic acid. B) Representative EPR spectra of free

and HSA-bound 5-DSA (B) and 16-DSA (C) in whole blood from the same COVID-9 recovered patient.

Chemical structures of the two spin-labeled fatty acids are given on the right side of the figure.

Fig. 2. Hematologic cellular counts and neutrophil-ROS levels reflect severity and mortality in COVID-19 patients. A) Representative flow cytometric diagrams comparing morphologic, hematologic, and ROS

levels in control (representative of n=9; upper row), Sev-R (representative of n=10; middle row), and

Sev-D (representative of n=11; lower row) groups. B) Heat diagram comparing lymphocyte, neutrophils,

platelets, and DCF-positive neutrophil counts as percentage of total cell counts in all of the studied

subjects. Yellow areas are either group separators or missing data due to insufficient sample size or

processing errors. C) A diagram showing statistically positive correlation between neutrophil count and

count of neutrophils stained positive for DCF dye in all groups (black dots denote controls; blue are Sev-

R; and red represent Sev-D patients). D) When neutrophil counts were compared for all groups, both

Sev-R and Sev-D groups showed statistically significant neutrophilia relative to control groups. However,

only a weak trend has been observed when comparing the two groups with COVID-19. E) DCF staining

revealed increased levels of ROS in Sev-R and Sev-D groups relative to control neutrophils. Sev-D

showed a trend of increased ROS level relative to Sev-R group. Multiple comparisons were carried out

using ANOVA followed by Tukey test and *p* values are given.

Fig. 3. Hydrogen peroxide levels in plasma and neutrophils reflect mortality in COVID-19 patients.

Catalase was used to specifically and quantitatively determine levels of hydrogen peroxide in identical

plasma volumes collected from control (n=7), Sev-R (n=6), and Sev-D (n=7) groups. A) Oxygen levels are

monitored and recorded while 50 mL batches of plasma from control, Sev-R, and Sev-D subjects

sequentially infused into tightly air-controlled O2k chamber containing catalase (315 Units/mL) in

deoxygenated buffer. In addition to the initial rise due to residual oxygen in the added plasma samples,

the decomposition of hydrogen peroxide in these samples produces oxygen quantitatively. B) To verify

the assay we measured the released oxygen upon adding an increasing volume of standard hydrogen

peroxide solution in PBS buffer with 0.2, 0.8, 1.2, and 1.6 μM final concentrations; inset. Linear fitting of

the plotted $[O_2]$ versus $[H_2O_2]$ relation yielded a slope = 0.47 ± 0.03 (Pearson's r = 0.994, p = 5.6 x 10^{-4}),

which is very close to the theoretically expected value of 0.5 as the catalase-mediated decomposition of

one mole of H_2O_2 produces ½-mole O_2. C) Plasma contents of H_2O_2 in plasma significantly increased in

the order Sev-D>Sev-R>Cont using ANOVA followed by Tukey test. D) Fluorescence imaging was used to

assess levels of ROS in freshly isolated neutrophils using DCF (2,7-Dichlorodihydrofluorescein diacetate,

green) staining in all groups. DAPI (4',6-diamidino-2-phenylindole, blue) binds strongly to adenine–

thymine-rich regions in DNA thus mapping nuclei through emitting blue fluorescence. Merged DCF and

DAPI images are shown on the third column. Images were acquired using Cytation 5 Cell Imaging Multi-

Mode Reader (Agilent) and analyzed using Gen5 Software package 3.08. Scale bar: 100 µm.

Fig. 4. EPR spectroscopic analyses of HSA-fatty acid binding reveal strong dependence of binding on

mortality in COVID-19 patients. A) Albumin level in plasma of control (n=8), Sev-R (n=8), and Sev-D

(n=10) groups showed that nonsurvivor COVID-19 patients exhibit statistically significant

hypoalbuminemia. Comparisons between representative spectra showing changes in line shape that are

related to mobility and microenvironmental statuses of HSA-bound 5-DSA (B) and 16-DSA (C) in whole

blood of a control (black trace), a Sev-R (blue trace), and a Sev-D (red trace) patients. Calculated

biophysical parameters including order parameter (D), rotational correlation time (E), and the ratio

between strongly bound to weakly bound spin labels (S/W, F) as defined in Fig. 1 and described in

Methods section. Statistical comparisons by ANOVA followed by Tukey tests were used for means'

comparisons and revealed remarkable decrease in the binding strengths and packing parameter of the

local microenvironment surrounding the spin labels. All calculated parameters along with exact p values

are given in the Supplementary Table 3. Water accessibility into albumin/fatty acids binding pockets are

followed by reacting with ascorbate, which reduces nitroxide radicals into the EPR silent

hydroxylamines. Representative EPR signal decays of 16-DSA in whole blood of control (G) and Sev-D (H)

are shown. Kinetic traces (n=3 per group) showing the reduction of 16-DSA (I) and 5-DSA (J) bound to

HSA by sodium ascorbate in whole blood. Kinetic traces are shown as the percentage loss of the signal

intensity of the middle peak (A/A_0). All samples contained 0.26 mM spin label and 3 mM sodium

ascorbate and measured at 37 °C. Weaker and slower disappearance of the EPR signal suggests

inaccessible space towards the nitroxide moiety of the spin label.

Fig. 5. Associations of oxidative stress measures and biophysical parameters with mortality outcome

in COVID-19 subjects. Linear correlations between plasma levels of hydrogen peroxide and EPR-

calculated parameters pertaining to protein packing order parameter (A,B), fatty acid mobility τ_c (C), and

S/W ratio (D) in Control (black circles), Sev-R (blue circles), and Sev-D (red circles) groups (n=19-26).

Neutrophils are major sources of oxidative stress as evident by linear correlations between % DCF

positive neutrophils and τ_c (E) or S/W ratio (F). On each correlation, Pearson's r and p values are

provided. **Kaplan–Meier estimates of time to mortality from blood sample collection during ICU**

hospitalization (G-J). Logrank Kaplan–Meier survival analyses were carried out to estimate probability of

survival of COVID-19 patients in relation with cut-off thresholds arbitrarily selected as the mean values

of the analyzed parameters. For S/W, plasma [H_2O_2], Risk Score defined as the {(S/W)/[H_2O_2]} ratio, and

plasma [Albumin], the number of analyzed COVID-19 patients were 22, 12, 12, and 21; respectively.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.