NETWORK DIFFUSION EMBEDDING REVEALS TRANSDIAGNOSTIC SUBNETWORK DISRUPTION AND POTENTIAL TREATMENT TARGETS IN INTERNALIZING PSYCHOPATHOLOGIES

Paul J. Thomas[1,2], Alex Leow[1,2], Heide Klumpp[2], K. Luan Phan[3], and Olusola Ajilore[1,*]

[1]Department of Psychiatry, University of Illinois at Chicago, Chicago, IL.
[2]Department of Bioengineering, University of Illinois at Chicago, Chicago, IL.
[3]Department of Psychiatry and Behavioral Health, The Ohio State University, Columbus, OH.

[*]**Corresponding Author**

Olusola Ajilore, MD, PhD
Department of Psychiatry
University of Illinois at Chicago
1603 W Taylor St.
Chicago, Illinois 60612
email: oajilore@uic.edu

Keywords: Neuroimaging; Network Diffusion; Internalizing Psychopathologies

Thu 1st Apr, 2021, 14:25

ABSTRACT

Network diffusion models are a common and powerful way to study the propagation of information through a complex system, they and offer straightforward approaches for studying multimodal brain network data. We developed an analytic framework to identify brain subnetworks with impaired information diffusion capacity using the structural basis that best maps to resting state functional connectivity and applied it towards a heterogenous internalizing psychopathology (IP) cohort. This research provides preliminary evidence of a transdiagnostic deficit characterized by information diffusion impairment of the right area 8BM, a key brain region involved in organizing a broad spectrum of cognitive tasks, that may underlie previously reported dysfunction of multiple brain circuits in the IPs. We also demonstrate that models of neuromodulation involving targeting this brain region normalize IP diffusion dynamics towards those of healthy controls. These analyses provide a framework for multimodal methods that identity diffusion disrupted subnetworks and potential targets for neuromodulatory intervention based on previously well-characterized methodology.

INTRODUCTION

The dynamics arising from the interaction of individual elements of a complex system are commonly investigated by representing such a system as a network or graph [13]. This has been applied extensively towards studying neural connectivity, where brain regions (represented by nodes) and the links between them (represented by edges) define a brain network, or connectome [71,72]. As in any network, the underlying structure of connectivity constrains the functions and processes that take place within it [78]. Indeed, the mechanisms of propagation of and neural response to brain pathologies have been closely tied to network topology [31].

A number of studies have found that the basis formed by the *anatomic* structure of the brain and the patterns of observed spatiotemporal neural activity are intimately related [1,2,5,23]. Many of these investigations use a network diffusion model to study the propagation of neural impulses, or information (given by functional connectivity), throughout the structure formed by the white matter tracts of the brain. These so-called network diffusion-based approaches all utilize the properties of the structural graph Laplacian, which encodes how a diffusive process spreads throughout a network over time [20]. The basis of diffusion given by the graph Laplacian, or its eigenmodes, can thus be used to model the flow of neural information [82], and has been shown to play important roles in healthy brain network organization [78]. More specifically, such an approach was used successfully to demonstrate that rs-fMRI functional connectivity is predictable from DTI structural eigenmodes [1,2]. Furthermore, resting state networks were found to closely match spatial patterns of structural eigenmodes [5]. Of note, many other approaches for investigating this interplay have been studied, including models based on epidemic-spreading [73], threshold models [55] and neural mass models [40] (for a review, see [6]).

Network diffusion-based methods have also been applied towards investigating the patho-physiology of neuropsychiatric diseases. For example, Raj and colleagues [65] show that white matter structural eigenmodes closely resemble known patterns of dementia and correlate strongly with regional atrophy. A closely related methodology to that of studying structural

eigenmodes is using the graph Laplacian-derived heat kernel of brain networks, a matrix which encodes how much information is transferred between every pair of network nodes after a given time, or diffusion depth. Heat kernel methods have been used to characterize brain network perturbations in brain networks information transfer in autism [68] and to predict future adverse motor function resulting from premature birth using white matter structural connectomes [19].

In the context of psychiatric conditions, the application of methods that integrate multimodal perspectives in their analytical approaches has great potential to elucidate the distributed perturbation of neurocircuitry that underlie the complex cognitive and behavioral disruption found in patients suffering from these disorders. This may hold particularly true for the internalizing psychopathologies (IPs), including mood (e.g., MDD, dysthymia) and anxiety (e.g., panic disorder (PD), social anxiety disorder (SAD), generalized anxiety disorder (GAD)) disorders. Many previous neuroimaging studies have been conducted on IPs, however, most of which use unimodal data analysis. Importantly, findings about potential neuropathological substrates are more often overlapping between the IPs than distinct to a specific IP. In addition, IPs, as traditionally categorized, are often comorbid with one another and present heterogeneously with a spectrum of related symptoms and disruptions to emotion regulation and negative valence system (NVS) processes [17, 44, 59]. Furthermore, the available first line treatments for the IPs, either cognitive behavioral therapy (CBT) or selective serotonin reuptake inhibitors (SSRI), are equally effective across the swath of IPs [26]. Many studies have concluded that similar structural and functional brain networks involving regions commonly implicated in the expression of fear, anxiety, negative affect and other NVS features are dysfunctional in these disorders [29, 38, 50]. In addition, previous findings also implicate the disruption of similar canonical resting state networks (RSNs), such as the default mode network (DMN), in both depression and anxiety using both structural and functional neuroimaging [45, 52, 60, 70, 76]. To address this pattern of findings, the National Institute of Mental Health's Research Domain Criteria (RDoC) initiative was developed [22, 41] in order to reorient the study of psychiatric disorders away from traditional diagnostic categories and

towards that of data-driven approaches such as identifying transdiagnostic cognitive domain disruptions, biomarkers of treatment response and targets for neuromodulatory intervention.

In line with the aims set forth by this initiative, we study a dataset from an RDoC clinical trial consisting of DTI and rs-fMRI neuroimaging scans of a treatment naive, heterogenous IP patient (PT) cohort and age and sex matched healthy controls (HC). PTs were then randomized to either 12 weeks of SSRI or CBT and completed Inventory of Depression and Anxiety Symptoms (IDAS-II) [79] self-reports pre- (Pre) and post-treatment (Post) to assess transdiagnostic dimensions of symptom burden. Several previous analyses have been conducted on this dataset but they are unimodal and have a hypothesis-driven focus on *a priori* brain region, often studied in association with specific task-based measures of NVS subsystems [14, 15, 32, 36, 48, 64, 77].

Because IPs share similarly dysfunctional brain networks and respond to similar treatments, we hypothesize that there exist pathophysiological features common to all IPs that result in impaired emotion regulation and the subsequent heterogenous expression of IP symptoms. In this paper we ask (1) can the presence of such unifying perturbations can be identified by incorporating both structural and functional connectivity data, (2) do these perturbations predict response to CBT and/or SSRI treatment and (3) can we identify candidate brain region as targets of neuromodulatory intervention to normalize connectivity dynamics to those found in healthy controls? To answer these questions, we use a multimodal data-driven approach based on network diffusion models to define structural diffusion distance (SDD) connectomes for each subject. We chose such a model for multimodal incorporation as it has both been well characterized in the study of structural functional mappings in brain networks [1] and allows for the computation of the diffusion-based network embedding given by the structural eigenbasis that most accurately underlies empirical functional connectivity. The parameters identified by this structural to functional mapping give the diffusion time, or depth (βt), at which this relationship is optimal. Thus, we achieve correspondence to functional connectivity by computing the embedding at the optimal diffusion depth for each subject (see Methods section for details). Furthermore, a network diffusion model easily allows for the simulation

of information flowing through the network. Simple mathematical properties of the graph Laplacian-derived heat kernel corresponding to the network embedding can be exploited to identify network nodes to supplement with additional 'heat' supplement in order to normalize deficient heat transfer throughout a PT subnetwork towards the desired diffusion dynamics of a target (mean HC) subnetwork. Our study applies this analytic framework to provide evidence for transdiagnostic IP subnetwork disruption and neuromodulation treatment targets, opening up application of multimodal network diffusion-based methods towards other brain disorders.

RESULTS

Data analyzed in this study are taken from a previously conducted Research Domain Criteria (RDoC) clinical trial on predictors of IP treatment outcomes (ClinicalTrials.gov identifier: NCT01903447) in which PT were randomized to either 12 weeks cognitive behavioral therapy (CBT) or selective serotonin reuptake inhibitor (SSRI) treatment. In addition, the patient population has heterogenous clinical presentations and IP comorbidity in order to study common pathological features without regard to traditional diagnostic categories. Baseline rs-fMRI and DTI scans as well as IDAS-Panic and -Depression Scores were assessed from both HC ($n = 22$) and PT ($n = 65$) subjects. The subset of PT ($n = 50$ total; $n = 28$ CBT cohort; $n = 22$ SSRI cohort) completed the IDAS-II self-reports following treatment are used for identifying correlates of treatment response. Of note, there were no group differences in age or sex between HC and PT. As expected, baseline IDAS-panic and -depression scores were significantly different between groups at baseline (table 1).

	HC	PT	Statistic
Age (years)	25.0 ± 10.7	28.1 ± 8.5	$t = -1.36$, $p = 0.18$
Sex (% female)	63.6	68.7	$\chi^2 = 0.01$, $p = 0.93$
IDAS-II Depression	26.3 ± 3.8	58.0 ± 9.8	$t = -14.6$, $p = 7.5e - 25$
IDAS-II Panic	8.14 ± 0.34	13.3 ± 4.6	$t = -5.18$, $p = 1.5e - 6$
S2F mapping r	0.26 ± 0.03	0.25 ± 0.03	$t = 1.96$, $p = 0.06$

Table 1: Baseline mean demographics, IDAS-II scales and structure to function mapping (S2F) Pearson r values by split healthy controls (HC) and patients (PT).

Figure 1 provides a visual overview of the analytic framework for computing the diffusion-based embeddings and subsequent structural diffusion distance (SDD) connectomes used in this study, discussed in detail in the Methods section. We first compute the structure to function mapping as described in [1] for each subject. The fit of the empirical rs-fMRI connectome to the rs-fMRI connectome estimated from the DTI structural connectome Laplacian eigenmodes is then determined by computing the Pearson r statistic between these adjacency matrices. The mean fit values for HC were slightly higher than PT ($r = 0.26 \pm 0.03$ and $r = 0.25 \pm 0.03$, respectively, table 1) but the difference between their mean values was not statistically significance via t-test ($p = 0.06$), indicating that the relation between resting state functional connectivity and the structural diffusion basis is likely preserved at the global scale. Finally, the distance matrix connectomes used for subnetwork discovery are computed from the structural diffusion basis at the diffusion depth, βt, that optimally maps structural to functional connectivity.

Identification of Diffusion Impaired Subnetwork

After SDD connectomes were computed, the Network-Based Statistic (NBS) algorithm [86] was used to discover subnetworks with altered diffusion properties in PT relative to HC. NBS was applied with one-sided t-tests used for edge comparison, over a range thresholds $t = 3.0, 3.5, 4.0, 4.5$. For a t-test contrast (left-sided t-test) of HC $<$ PT, the range of subnetwork sizes varied greatly as a function of t-statistic threshold, resulting in subnetworks

Figure 1: Visual overview of methodological pipeline for creating structural diffusion distance (SDD) connectomes using subject-specific optimal diffusion depths (βt) from the structure-to-function mapping as in [1] using subject DTI and rs-fMRI connectomes, respectively. NBS is then applied to SDD connectomes for identification of subnetworks with aberrant diffusion properties for subsequent analyses. See Methods section for details.

from 314 to 2 nodes with thresholds $t = 3$ and $t = 4.5$, respectively. With the threshold of

$t = 4$ a single significant subnetwork (SN1) consisting of 48 nodes and 73 edges was identified

(Figure 2A), which is used for subsequent analyses because of the biological interpretability

of its size, and for adequate estimated statistical power, $1 - \beta = 0.81$. Note that because

NBS was conducted on SDD connectomes, edges are pairwise distance between nodes in the

diffusion-based network embedding. This indicates that SN1 represents a subnetwork whose

nodes are significantly further apart in this diffusion space, i.e., diffusion of information occurs

more slowly between the nodes of SN1 along the edges of SN1 in PT relative to HC. For NBS

with a t-test contrast defined as HC > PT, no significant subnetworks were found.

To determine the brain regions that are most central to the diffusion impairment in SN1,

we calculated the mean strength, str_{SDD}, of each node using the SDD values of edges in

SN1. The nodes with the greatest str_{SDD} values are from brain regions found in the bilateral

anterior cingulate (ACC) and medial prefrontal (mPFC), right inferior parietal (IPC) and

left insular (INS) and frontal opercular (FOP) cortices (Figure 2A). Specifically, the brain

region with the greatest average str_{SDD} value in both HC and PT is the right area 8BM,

which has recently been reported to be a core region of a 'multiple demand' (MD) subnetwork

that is active during a broad spectrum of tasks and may play an important role in general

cognitive control [4]. Additionally, we found that the hubness of area 8BM was not present

in structural connectomes, determined by computing graph theoretical metrics, indicating

that the centrality of this brain region is unique to the SDD representations of brain networks

(Figure S1).

Figure 2: Visualizations of the significant subnetwork (SN1) found via NBS with a t-statistic threshold of $t = 4$ and contrast of HC $<$ PT, where information diffusion is impaired in PT relative to HC. **A** Wire and ball plots of SN1 subgraphs of mean SDD connectomes in HC (top row) and PT (bottom row). Nodes indicate brain regions of SN1, and edges are colored by mean distance in from SDD connectomes (lighter/more yellow indicates greater distance). **B & C** Bar plots of mean strength (str_{SDD}) of SN1 nodes **B** and nodes averaged by cortical region **C**, as determined by cortex assignment in the MMP1.0 parcellation [34], calculated using edges within SN1 and the corresponding edge values of mean SDD connectomes HC (orange bars) and PT (blue bars). A greater strength indicates a greater diffusion distance from other nodes in SN1.

Baseline Heat Kernel Correlates of Symptom Severity and Treatment Response

To investigate the association of information diffusion between brain regions of SN1 and clinical scales, we compute heat kernel matrices which encodes the amount of heat transferred between each node after a given amount of time has passed. We use each subject's diffusion depth, βt, as the time at which their heat kernel is calculated for so that we may study the diffusion dynamics at the time scale best associated with reconstructing functional connectivity. We took a hypothesis-driven approach by focusing our correlation analyses on the heat transfer between the bilateral ACC and all other SN1 cortical regions, as the nodes in these cortical regions are central to the diffusion impairment of SN1. The heat kernel values (HKVs) are averaged by cortical region and are then used for computing Spearman correlations with IDAS-II Depression and Panic subscales, which map to distress and fear, respectively, a previously used approach to broadly divide and assess symptom domains of IPs (Figure 3) [63, 64]. In addition, we use only PT ($n = 65$) for these analyses, as mean IDAS-II subscale values expectedly differ highly significantly between HC and PT, and correlations would likely be due to group differences. Finally, we report at most the top three most significant correlations that survive FDR correction ($n = 44$ comparisons) in this section of the manuscript, but tables of all significant correlations are available in Table S1.

Figure 3: Correlation of baseline heat transfer at subject diffusion depths βt with pre-treatment and percentage improvement post-treatment IDAS-II subscales. The brain network figure at the end of each row shows the corresponding pair of brain for each respective correlation in the row, indicated by the color of the edge connecting the regions and the correlation plot of the same color. Cortical regions shown in the visualization are plotted according to the average position of the individual ROIs that they consist of, as per the MMP1.0 parcellation [34]. Correlations of heat transfer with: **A** pre-treatment IDAS-II Depression in PT; **B** post-treatment IDAS-II Depression percent improvement in all PT; **C** post-treatment IDAS-II Panic percent improvement in SSRI cohort only; **D** post-treatment IDAS-II Depression percent improvement in CBT cohort only; **E** post-treatment IDAS-II Depression percent improvement in CBT cohort only.

First, we determine the HKV correlates of baseline IDAS-II subscales (Figure 3A). Heat transfer between the left ACC/mPFC and INS/FOP ($\rho = -0.37, p = 0.003, q = 0.04$) and the right ACC/mPFC and left inferior frontal cortex (IFC) ($\rho = -0.36, p = 0.004, q = 0.04$) was negatively correlated with IDAS-II Depression scales, indicating that less information diffusion between these brain regions is associated with higher IDAS-II Depression values. A positive correlation was found with heat transfer between the left ACC/mPFC and posterior cingulate cortex (PCC) ($\rho = 0.39, p = 0.001, q = 0.04$). No significant correlations were found using IDAS-II Panic.

To determine whether baseline HKVs could predict response to treatment, we determined correlations between heat transfer and percent improvement in IDAS-II subscales in the subset of PT ($n = 50$) that completes post-treatment IDAS-II self-reports (calculated as $\frac{\text{pre}-\text{post}}{\text{pre}}$, where pre and post are the IDAS-II subscales before and after 12 weeks of treatment, respectively). We first grouped both SSRI and CBT therapy cohorts of PT together to study common correlates of treatment response. Significant negative correlations with IDAS-II Depression and heat transfer between the left ACC/mPFC and IFC ($\rho = -0.45, p = 0.001, q = 0.04$), left auditory association cortex ($\rho = -0.42, p = 0.002, q = 0.04$) and INS/FOP ($\rho = -0.41, p = 0.003, q = 0.04$) were found (Figure 3B). Similar to baseline symptom correlations, no significant associations were found using IDAS-II Panic.

Next, we segregated PT by cohort to investigate treatment specific HKV predictors of therapeutic response. Interestingly, in the SSRI cohort ($n = 22$), significant negative correlations were found only with IDAS-II Panic improvement and heat transfer between the left ACC/mPFC and IFC ($\rho = -0.60, p = 0.003, q = 0.15$), INS/FOP ($\rho = -0.54, p = 0.009, q = 0.17$) and right NS/FOP ($\rho = -0.53, p = 0.012, q = 0.17$) (Figure 3C). In the CBT cohort ($n = 28$), we observed significant negative associations with IDAS-II Depression percent improvement and heat transfer between the right ACC/mPFC and superior parietal (SPC) ($\rho = -0.51, p = 0.006, q = 0.09$), inferior parietal (IPC) ($\rho = -0.51, p = 0.006, q = 0.09$) and left auditory association ($\rho = -0.50, p = 0.008, q = 0.09$) cortices (Figure 3D). We also found significant negative correlations with IDAS-II Panic percent improvement and heat

transfer between the right ACC/mPFC and SPC ($\rho = -0.55, p = 0.003, q = 0.12$) and the left ACC/mPFC and right lateral temporal (LTC) cortex ($\rho = -0.50, p = 0.007, q = 0.14$) (Figure 3E). Interestingly, the brain regions involved significant correlations of treatment response in all PT and the SSRI cohort are all Frontal cortical regions that have been tied to emotion regulation in MDD [39, 66, 67], while the regions found with the CBT cohort are members of canonical RSNs. For example, the IPC and LTC are members of the default mode network (DMN), while the SPC is part of the dorsal affective network (DAN) [12, 27]. Please note that we increased the threshold for significance after FDR correction from $q < 0.05$ as for baseline and percentage improvement of IDAS-II subscales in all PT to $q < 0.2$ given the decreased sample size and decreased power of these correlations. As such, these results should be understood as exploratory and preliminary.

Identification of Subnetwork Targets for Modulation

Once we had characterized the disrupted information diffusion of SN1 in IP PTs, we next sought to determine brain subnetwork regions that may be targets for modulation that would normalize these diffusion properties towards those of HCs. To this end, we computed the mean HC heat kernel matrix, $\mathcal{H}_{HC}$, and found the difference between $\mathcal{H}_{HC}$ and each k^{th} PT's heat kernel matrix, $\mathcal{H}_{PT}^k$. We then computed a heat kernel modifier matrix, $\mathcal{M}_i^k$, for each k^{th} subject and i^{th} brain region such that when added to $\mathcal{H}_{PT}^k$ it minimizes the norm of the difference between this sum and $\mathcal{H}_{HC}$. As $\mathcal{M}_i^k$ gives the effect of supplementing the diffusion dynamics of a subnetwork with a constant amount of additional heat, we choose this method to model the potential effects of neuromodulatory stimulation on the diffusion dynamics of a subnetwork (see Methods section for details). For each brain region, we obtain a heat and error value (given by the squared frobenius norm). As the error values were empirically observed to be uniform in distribution with small regional variance (Figure 4B), we defined optimality as the brain region requiring the smallest amount of supplemental heat. Furthermore, this allows us to identify brain region neuromodulatory targets that potentially more efficiently disseminate stimulation with fewer off-target effects from a greater modulatory energy.

We first conducted this analysis on the mean cortical regions of SN1 as target nodes and

found the right ACC/mPFC, which was identified as a hub of SN1 by mean str_{SDD}, to be the optimal region to target for heat supplementation, with the lowest mean heat across all PT (Figure 4A and B). In addition, we determined subject-specific optimal brain regions and the right ACC/mPFC was also found to be the modal optimal target (34 out of 65 PT) (Figure 4C). Next, we applied the same target node identification procedure to the individual brain regions of the right ACC/mPFC in SN1. We found that the right 8BM was the optimal nodal modulatory target. Note that the right 8BM was an SN1 hub with the greatest str_{SDD} of all individual subnetwork nodes (Figure S2A and B).

Figure 4: Overview of the identification of SN1 brain region targets for heat-based modulation and subsequent correction of information diffusion deficits. **A** Connectome plot visualizations of baseline (first row) and post-heat supplementation (second row) significant differences in heat kernel values with right ACC/mPFC as modulation target. Nodes are brain regions of SN1 and edges connecting them are present if a significant difference (defined as t-test p-values that survive FDR correction with $q < 0.05$) exists between HC and PT corresponding heat kernel values. Larger/darker node is the target for modulation. Edge colors indicate the type of significance (dark blue: present at baseline (first row) or remains after modulation (second row) (HC > PT); light blue: present at baseline but corrected following modulation; orange: newly significant following modulation (PT > HC)). **B** barplots for mean PT heat (left) and norm of error (right) values for all mean cortical regions in SN1. **C** Scatter plot indicating subject-specific optimal modulation target heat values grouped by brain mean cortical regions in SN1.

Correction of Subnetwork Diffusion Impairment

As a final step in our analysis, we sought to confirm whether the heat modulation approach identified as above successfully normalizes subnetwork information diffusion in PT to that of HC statistically. To carry this out, we performed left sided t-tests (contrast: HC > PT) at each heat kernel edge connecting the brain regions of SN1 at baseline in HC versus PT. Only this direction of t-test was assessed at baseline as this modulation model can only add heat, and is not presently able to correct hyperconnected HKVs in PT. We then modified PT heat kernels at the optimal target nodes and heat values as described above and repeated the edgewise t-test analysis with left and right sided t-tests in order identify significantly increased heat kernel values in PT relative to HC as a result of heat supplementation. A heat kernel value was defined as significant if it survived FDR correction ($q < 0.05$) with $n = \binom{N}{2}$ comparisons, where N is the number of brain regions.

We conducted this analysis first on mean cortical regions using the right ACC/mPFC ($N = 23$) as a target for modulation and identified 24 significant baseline HKV differences (Table S2). Following heat supplementation, 19 of the original 24 HKV differences were no longer significant (Figure 4A, Table S2). Three HKVs remained significant in the same direction as baseline: connections between the right dorsal visual stream and the right auditory association area, INS/FOP and LTC. On the other hand, 2 HKVs gained new significance (PT > HC) following modulation; connections between the right ACC/mPFC (target) and the left ventral visual stream and the right IPC.

We next conducted the same analysis on individual brain regions in SN1 ($N = 48$), using the right area 8BM as the modulatory target (Figure S2, Table S3). Of the 31 HKVs differences at baseline, 26 were corrected following heat modulation. Five HKVs remained significant following modulation, which predominantly follow the regional pattern as described above. All 6 of the newly significant (PT > HC) HKVs involved connections with the right area 8BM (target node).

DISCUSSION

This research provides preliminary evidence of a transdiagnostic deficit characterized by information transfer impairment of the right area 8BM, a key brain region involved in organizing a broad spectrum of cognitive tasks, that may underlie previously reported dysfunction of multiple brain circuits in the IPs. In addition, this is also the first report of using network diffusion models to study psychiatric disorders. We also demonstrate that models of neuromodulation involving targeting this brain region normalize PT diffusion dynamics towards those of healthy controls. These analyses provide a framework for multimodal methods that identity diffusion disrupted subnetworks and potential targets for neuromodulatory intervention based on previously well-characterized methodology.

From our analyses we discovered a subnetwork (SN1) with increased diffusion embedding distance, i.e., decreased information diffusion capacity at the time scale required for optimally capturing functional connectivity patterns, in HC relative to IP PT at baseline. Hub regions (as defined by group mean str_{SDD}, indicating regions with the greatest total diffusion impairment) of SN1 include the bilateral ACC/mPFC and INS/FOP, which bears resemblance to the cortical aspect of the cingulo-opercular network (CON) (Figure 2). The CON, which includes the dorsal ACC, anterior INS, PFC, hypothalamus, thalamus and amygdala has been shown to contribute to many brain functions, including the processing of pain and negative affect, as well as the maintaining general cognitive control during goal-oriented behaviors [24]. Dysfunction of the CON has also been demonstrated in IPs, including MDD [37] and anxiety disorders [75]. The dorsal ACC, a core hub of the CON, has been shown to be critical in both MDD [81] and anxiety disorders [75]. In particular, investigations of emotion regulation in anxiety disorders have demonstrated that increased ACC activity [16] and coupling to anterior INS activity [46,49] is present in HC relative to PT, indicating a potential regulatory role for the ACC during cognitive control of emotion processing within the CON.

SN1 also includes brain regions that are part of other functional brain networks, such as the LTC and IPC (a hub of SN1), part of the DMN, and the SPC, part of the DAN,

in addition to the IFC, which is important for the coordination of executive function and emotion processing [66, 69]. Of note, the IPC is also a member of the frontal parietal network (FPN), an executive function network that antagonistically deactivates the DMN [7]. Increased activity of the DMN, a network defined by functional connectivity at rest associated with both rumination and worry, relative to other RSNs has been implicated in MDD [27] and anxiety disorders [45]. Studies of MDD have also indicated the widespread disruption in the salience and central executive networks [35, 43]. Indeed, accumulating evidence suggests that dysfunctional coordination between multiple brain networks may better explain the pathophysiology of IPs.

Our findings from interrogating interregional information transfer within SN1 indicate that connections between the bilateral ACC/mPFC and brain regions involved with canonical brain networks are associated with baseline symptom severity. IDAS-II Depression scores correlated positively with heat transfer between the left ACC/mPFC and the PCC. Reduced connectivity between the PCC, a DMN hub, and DLPFC [51] and other frontal regions [84] has been previously reported in MDD. In the macaque, retrograde tracing studies have revealed afferent connections to the 8BM subregions of the ACC/mPFC from the PCC [28]. Taken together, our results may indicate that depressive symptoms in the IPs at least partly result from the hyperconnectivity of the DMN to the MDN and CON, which may skew cognitive resources away from executive control of emotion processing. On the other hand, negative correlations were found with IDAS-II Depression and heat transfer between ACC/mPFC and the IFC and INS/FOP. These results are similar to those from previous studies, where the ACC activity was found to negatively correlate with depression symptom scales [81] and with anxiety symptom scales in the context of its potentially antagonistic role with the INS in the CON [46, 49]. Overall, these baseline associations provide evidence for ACC/mPFC activity and connectivity to regions important for emotion regulation as critical to transdiagnostic depression symptoms in the IPs.

Our results also indicate that internodal information transfer of the bilateral ACC/mPFC in SN1 is associated with therapeutic response in both treatment modality agnostic and

specific manners. Interestingly, predictors of general and SSRI specific treatment response involved similar brain regions to those associated with baseline symptom severity. In all PT, negative correlations were found with IDAS-Depression symptom improvement and the heat transfer between the left ACC/mPFC and the IFC, INS/FOP and auditory association cortex. As with all PT at baseline, no associations were found with IDAS-II Panic. In the SSRI cohort, however, significant correlations were only found using IDAS-II Panic improvement, yielding negative associations with connections between similar regions as with all PT: heat transfer between the left ACC/mPFC and the IFC and bilateral INS/FOP. These findings indicate that lower baseline information transfer between the ACC/mPFC and CON or IFC are predictive of general and SSRI specific responsiveness to treatment, and these connections may be the substrate of treatment action. Interestingly, associations of baseline internodal information transfer and treatment response in the CBT cohort were found with unique brain regions compared to all PT and the SSRI cohort. Significant associations were also found using both IDAS-II Depression and Panic scales. Depression improvement was negatively associated with information transfer between the right ACC/mPFC and SPC and IPC, regions of the dorsal affective network and DMN. Using IDAS-II Panic improvement, we found significant negative correlations with information transfer between the right ACC/mPFC and SPC the left ACC/mPFC and right LTC, another DMN subnetwork region. These results indicate the presence of CBT specific neural substrates of treatment prediction, and again, that lower baseline information transfer between these regions is indicative of greater treatment efficacy. In addition, these findings are in line with a previous study of CBT for social anxiety disorder, where increased baseline activation of the ACC and LTC [47] was predictive of treatment response.

Central to the diffusion impairment of SN1 is a subregion of the right ACC/mPFC cortical region, area 8BM, the caudal aspect of the dorsomedial PFC (dmPFC), which borders the dorsal aspect of the ACC. Retrograde tracing studied in the macaque monkey have revealed both afferent and efferent neuronal connections between 8BM and the ACC, indicating likely functional synergy of these areas [28,57]. Importantly, 8BM is brain region that has recently

been found to be a key region of the multiple-demand subnetwork (MDN), a fronto-parietal system that is co-activated during a broad range of cognitively demanding tasks [4]. As such, the MDN is likely critical for the organization and recruitment of multiple task-specific subsystems in the brain towards current cognitive needs. Additionally, our network-diffusion model identified the most effective brain regions as putative targets for neuromodulatory stimulation on a subject-specific level. Examining the results of this investigation provide evidence that the right ACC/mPFC is an optimal target for neuromodulation to correct SN1 information diffusion dynamics of PT towards those observed in HC, and, further, that area 8BM is the most efficient subregion of the ACC/mPFC for such an intervention.

These preliminary findings have important clinical implications for the therapeutic application of neuromodulatory interventions for IPs, such as repetitive transcranial magnetic stimulation (rTMS), a focal, non-invasive brain stimulation method. rTMS, which is typically delivered to the dorsolateral PFC (dlPFC), is a first line treatment for SSRI-refractory MDD [33] but has also been shown to significantly reduce anxiety [18,25] and nicotine dependence symptoms [3]. Although rTMS is generally effective, MDD remission following treatment is 30-40% [33], indicating a need for developing better treatment paradigm, such as developing personalized treatment protocols. In line with these goals, a study that used MRI guidance to enhance targeting of specific regions of the dlPFC, all rTMS-resistant MDD PTs responded to this new approach [58]. Another study used fMRI guidance to individually targeted cortical emotion regulation systems to improve rTMS efficacy [53]. Our results offer another strategy for identifying brain network targets of rTMS treatment for correcting subject-specific structural deficits that may underlie functional expression of IP symptoms.

Limitations and Future Directions

A significant question about the implications of our findings is whether the addition and propagation of heat on a subnetwork sufficiently models the effects of rTMS stimulation. It has been shown that the propagation of direct cortical electrode [74] and rTMS stimulation and subsequent functional activity are best predicted by structural connectome topology [11,56], providing encouraging evidence that network diffusion-based analyses such as those proposed

in this study are an appropriate model for the effects of rTMS neuromodulation. Nevertheless, further studies are necessary to verify the utility of the proposed model. In addition, while this is the first report of using a multimodal information diffusion model towards a transdiagnostic sample of IPs, the findings presented in this paper pertaining to the pattern of diffusion impairment of SN1 should be replicated in a larger sample of IP PT and HC to generalizability of our results. This is especially true for the analyses obtained by segregating PT into SSRI and CBT cohorts, as the sample sizes of these groups are very small, and as such these findings must be interpreted as preliminary and exploratory. Further, availability of post-treatment brain imaging would be required to make reliable conclusions about network diffusion-based substrates of treatment response, and, as such, should be included in future studies.

Although our structural to functional mapping quality, as measured by Pearson correlation between the predicted and empirical functional connectomes, is in the range of values obtained in a replication study of related mappings, other relatively simple and network-diffusion related models have been developed that result in better predictive mappings [23]. Future studies of these proposed methods should investigate how the mapping parameters from these related methods can be incorporated into the current model for better structural-functional fusion. Another area for further development in our proposed network diffusion methods is that neuromodulation target identification and simulation can only model the effects of the *addition* of heat to subnetwork's diffusion dynamics. Therefore, if a brain circuit is found to be pathologically hyperconnected, other strategies will have to be developed in order model either direct or indirect inhibition of brain activity.

Future implications of this research are broad. We present here an analytical framework assembled from well-characterized neuroimaging and graph theoretical methods that can be used as is to study multimodal brain networks for other brain disorders. Aside from the application to other datasets and to larger similar datasets for replication, this model offers the ability to determine the diffusion embedding from functional connectivity other than that observed during resting-state. For example, functional connectomes can be generated from tasks that are known to be implicated in a certain disorder, and as a result the

diffusion embedding will be formed from the structural basis that best maps to the functional
connectivity observed during this task. This allows for the researcher to investigate the
structural connectivity potentially most pertinent to forming the functional brain activity
observed during specific tasks, and therefore permits a more granular interrogation of complex
features of brain disorders and states. Perhaps the most promising and immediate application
of our proposed methodology is that of identifying brain regions best suited for therapeutic
for neuromodulatory intervention, such as with rTMS. As the proposed methods provide both
subject-specific regional targets and magnitudes of stimulation that best modify subnetwork
dynamics towards those of a desired (HC) network, future work is in line with improving the
outcomes of rTMS intervention by personalizing treatment features.

Conclusion

There are two major outcomes of this study. First, we report impaired information diffusion
between area 8BM and other SN1 regions, many of which previously implicated in the
pathology of multiple IPs and the function of the CON, DMN and DAN. Second, we found
that hubs of SN1, found to be critical for organizing brain function for a variety of cognitive
and emotional processes, are optimal targets for modelled neuromodulatory intervention.
Taken together, our results may indicate a concerted disruption of multiple brain networks
pertinent to cognitive control of emotion regulation. This dysregulation of connectivity could
result in a loss of "top-down" executive control of emotion processing via connections between
the MDN and other task-specific (and task-negative) brain networks. Such a perturbation in
brain network dynamics involving the integration of multiple complex subsystems via a hub
of the MDN may represent the underlying pathophysiological brain network features that are
common to all IPs and give rise to the heterogenous expression of transdiagnostic symptoms.

ACKNOWLEDGEMENTS

This study was supported by funding from the National Institute of Mental Health of the

National Institutes of Health (NIMH-NIH) grants R01MH101497 (to KLP) and the NIMH

grant 5T32MH067631-14 (support for PJT).

METHODS

Resource table

RESOURCE	SOURCE	IDENTIFIER
Data		
UIC RDoC Study rs-fMRI and DTI connectomes and subject IDAS-II scales and demographics	This paper	https://github.com/pauljasonthomas/brainnetdiff
Software and Algorithms		
Brain network diffusion-based analysis software	This paper	https://github.com/pauljasonthomas/brainnetdiff
Python 3	Python Software Foundation	RRID SCR_008394 https://www.python.org
FSL	[42]	RRID SCR_002823 https://fsl.fmrib.ox.ac.uk/fsl/fslwiki
DSI Studio	[85]	RRID SCR_009557 http://dsi-studio.labsolver.org/
CONN	[80]	RRID SCR_009550 www.nitrc.org/projects/conn
MMP1.0 parcellation	[34]	https://balsa.wustl.edu/study/show/RVVG

Clinical trial and research participants

Subjects were recruited from the greater Chicago area through advertisements and through University of Illinois at Chicago (UIC) outpatient clinics and counselling centers as part of a larger Research Domain Criteria (RDoC) [22] investigation on predictors of IP treatment outcomes (ClinicalTrials.gov identifier: NCT01903447). A heterogeneous study population was recruited in order to obtain a sample with a broad range of symptom severity and functioning. Details regarding inclusion/exclusion criteria, participant recruitment, clinical characteristics and treatment have been previously described [36]. In brief, this study was approved by the UIC Institutional Review Board, and written informed consent was obtained for each participant. The inclusion criteria for subjects were age between 18 and 65 years, and the need for randomization to 12 weeks of treatment with SSRI or CBT, as determined by a consensus panel consisting of at least three trained clinicians or study staff. Subjects were excluded from the study if they have a history of current or past manic/hypomanic episodes or psychotic symptoms, active suicidal ideation, presence of contraindications or history of SSRI resistance (no response to >2 SSRIs despite adequate duration and dose), psychopathology not appropriate for the treatment algorithm, or current cognitive dysfunction or impairment. The SCID-5 [30] was used to determine current and lifetime Axis I diagnoses. The study was a parallel group randomized control trial with 1:1 allocation ratio to either 12 weeks of CBT or SSRI. For the SSRI cohort, PTs were administered one of 5 drugs (sertraline, fluoxetine, paroxetine, escitalopram or citalopram) with a flexible dosing schedule with a goal of obtaining target dose by 8 weeks. SSRI PTs met at 0, 2, 4, 8 and 12 weeks with their study psychiatrist for medication management. For the CBT cohort, PTs received 12 once-weekly 60 min sessions led by a PhD-level clinical psychologist. CBT procedures were based on the PT's principal diagnosis and predominant symptoms [15]. Each participant was scanned at enrollment and IP subject scans were acquired before treatment was administered.

At the time of enrollment (Pre) and after 12 weeks of treatment (Post), severity of IP symptoms was assessed in all subjects using the Inventory of Depression and Anxiety Symptoms (IDAS-II) [79]. Subjects responded to each of the 99 items in this inventory using a 5-point

Likert-type scale ranging from 1 (not at all) to 5 (extremely), yielding 17 empirically derived and symptom-specific scales (Suicidality, Lassitude, Insomnia, Appetite Loss, Appetite Gain, Ill-Temper, Well-Being, Panic, Social Anxiety, Traumatic Intrusions, Traumatic Avoidance, Mania, Euphoria, Claustrophobia, Checking, Ordering, Cleaning, General Depression and Dysphoria). As we are interested in transdiagnostic NVS construct disruptions in IPs, we use the IDAS-II Panic and Depression subscales, as these scales have been shown to map well to 'fear' and 'distress' dimensions, respectively, which is a previously used approach for broadly dividing and assessing these symptom domains [63,64,79].

Image acquisition and processing

Structural MRI

A diffusion tensor imaging (DTI) scheme was used, and a total of 32 diffusion sampling directions with 4 b0 images were acquired on a 3 Tesla GE Discovery MR750 System (Milwaukee, WI) at the UIC Center for Magnetic Resonance Research. The b-value was 1000 s/mm2. The in-plane resolution was 0.9375 mm. The slice thickness was 2.5 mm. Two sets of DTI images were acquired, with opposite phase encoding directions. For preprocessing, the two sets of DTI images were realigned and corrected for motion artifacts and eddy current induced distortions using the FMRIB Software Library (FSL) [42]. White matter tract reconstruction was done using DSI Studio software [85]. A deterministic fiber tracking algorithm was used with whole brain seeding with a total of 10000000 seeds, an angular of 70 degrees, step size of 1 mm and anisotropy threshold of 0.1. The fiber trajectories were smoothed by averaging the propagation direction with 10% of the previous direction. Tracks with length shorter than 10 or longer than 300 mm were discarded. Cortical areas were defined by the MMP1.0 parcellation comprising of 360 regions of interest (ROIs), 180 per hemisphere, from Glasser et al. [34]. A 360 x 360 structural connectivity matrix or connectome was created for each subject where each edge is given by the fiber count of reconstructed white matter tracts, normalized by median tract length, between ROIs.

 Functional MRI

Whole-brain blood-oxygen-level dependent (BOLD) functional images were acquired using a T2* weighted gradient-echo echo-planar imaging (EPI; 2s TR, 25ms TE, 82° flip, 64 × 64 matrix, 200 mm FOV, 3 mm slice thickness, 0 mm gap, with 44 axial slices) sequence optimized to reduced susceptibility artifact, along with a high-resolution T1 structural scan. During this scan, subjects were asked to view a fixation cross on a blank background for 5 minutes. Subjects were instructed to keep their eyes open and focused on the cross, and to try not to think of anything in particular for the duration of the scan. Functional connectivity was measured using the resting-state fMRI toolbox, CONN [80], using standard preprocessing and denoising pipelines as described in [62]. The principal components of the white matter, CSF signal, realignment parameters, and scrubbing were regressed out of the signal using the CompCor method in CONN [8]. BOLD signal data was passed through a band-pass filter of.008 to.09 Hz. Using 360 regions of interest (ROIs) defined by the MMP1.0 multimodal parcellation [34], functional connectivity measures were derived using pairwise Pearson correlations on the BOLD time series data between each ROI. A 360 x 360 functional connectivity matrix or connectome was created using these Pearson r statistics for each subject and used for the structure-to-function mapping.

Network notation

A brain network may be represented as a graph with nodes being grey matter regions and edges being the connection between these regions. For structural connectomes, edge weights are assigned based on the number of fiber counts between nodes, normalized by the median fiber length. For functional connectomes, edge weights are the Pearson correlation r of the time series of BOLD signals between nodes. Formally, a graph is defined as $G = (V, E)$ where V is the set of nodes of size N and E is the set of edges linking nodes in V. In addition, $w : E \to \mathbb{R}$, is a weight function that assigns weights to the E according to the modality of imaging from which the brain graph is constructed. G can then be encoded as an adjacency

514   matrix, $A \in \mathbb{R}^{NxN}$, where

$$A_{i,j} = \begin{cases} w_{i,j} & \text{if } (i,j) \in E \\ 0 & \text{otherwise} \end{cases}$$

each entry $A_{i,j}$ corresponds to the connection weight between nodes v_i and v_j. The diagonal

strength matrix is then defined as $D_{i,i} = \sum_{j=1}^{N} A_{i,j}$, which can be used to define the graph

Laplacian matrix, $L = D - A$, and, the normalized graph Laplacian is defined as $\mathcal{L} =$

$D^{\frac{-1}{2}} L D^{\frac{-1}{2}}$. The eigendecomposition of the graph Laplacian is given by $\mathcal{L} = U \Lambda U^T$, where U

is a matrix with columns as eigenvectors and Λ is the diagonal matrix of eigenvalues. The

spectral properties of the normalized graph Laplacian have been extensively studied [21].

Diffusion in networks

Diffusion in networks over time can be determined analytically using the heat equation given

by

$$\frac{\delta \mathcal{H}(t)}{\delta t} = -\mathcal{L} \mathcal{H}(t)$$

524   where $\mathcal{H}(t)$ is the heat kernel and fundamental solution to the heat equation, and t is time.

525   The heat kernal can be understood as describing the flow of information between all nodes of

526   a network over time, taking into account all possible pathways of information flow into and

527   out of all nodes in the network. Throughout this manuscript, heat or information with be

528   used interchangeably in this context. The heat kernel is then given by

$$\mathcal{H}(t) = e^{-\mathcal{L}t}$$

529   which can then be used to solve for the heat distribution on network nodes, $h(t)$ after time

530    $= t$, given an initial condition $h(0)$:

$$h(t) = \mathcal{H}(t)h(0)$$

531   The product of an element of the heat kernel and initial condition vector, $\mathcal{H}(t)_{i,j} * h(0)_i$, then

532   represents the amount of heat at the j^{th} node at time $= t$ that has diffused from the i^{th} node.

The heat kernel can be computed as a sum of the outer product of eigenvectors of the graph
Laplacian, scaled by exponentiating the corresponding eigenvalues with time:

$$\mathcal{H}(t) = \sum_{i=1}^{N} u_i u_i^T e^{-\lambda_i t} = U e^{-\Lambda t} U^T$$

Structure to function mapping

In this study, we use the methods proposed by Abdelnour and colleagues [2] [1] to identify the
graph diffusion-based mapping of functional connectivity from the structural basis. Briefly,
the observed functional activity in rs-fMRI, i.e., the transfer of information from brain region
(node) i to region j as measured by pairwise correlation of temporal BOLD signals, can
modeled by first order diffusion-like dynamics given by:

$$\frac{\delta h_i(t)}{\delta t} = \beta \left(D_{i,i}^{\frac{-1}{2}} \sum_j^N A_{i,j} D_{j,j}^{\frac{-1}{2}} h_j(t) - h_i(t) \right)$$

where β is a diffusion constant. This extends to the entire network then as the heat equation,
$\frac{\delta h(t)}{\delta t} = -\beta \mathcal{L} h(t)$. The mapping from the structural diffusion dynamics, yielding the estimated
functional connectome, $\mathcal{C}_{est}$, is then given by [2]

$$\mathcal{F}_{est} = e^{-\beta \mathcal{L} t}$$

and has been updated as [1]

$$\mathcal{F}_{est} = a e^{-\beta \mathcal{L} t} + bI = a \left(\sum_i^N u_i u_i^T e^{-\beta \lambda_i t} \right) + bI$$

where a and b are additional model parameters and I is the identity matrix. Note that the
summation allows for the exclusion of eigenvector-eigenvalue pairs. As in [1], we leave out the
first eigenvector with corresponding zero valued eigenvalue, as it largely represents uniform
background connectivity (captured by b parameter above) and is typically regressed out of
rs-fMRI data. The model is then fit by minimizing the normalized predictive error with

550   respect to the model parameters given by:

$$\underset{a,b,\beta t}{\text{minimize}} \frac{||\mathcal{F}_{est}(a,b,\beta t) - \mathcal{F}||_2^2}{||\mathcal{F}||_2^2}$$

where $\mathcal{F}$ is the empirical rs-fMRI connectivity matrix. This model is fit for each individual to

obtain subject specific βt parameters, used in the diffusion-based embedding discussed below.

Diffusion-based network embedding

The embedding of nodes based on the diffusion properties of a graph has been studied

extensively for dimensionality reduction and clustering of multi-dimensional data [9,54,61].

These approaches center around the embedding of network vertices via the eigenvectors of

the graph Laplacian. Each element of an eigenvector corresponds to the coordinate of the

corresponding node such that nodes that are closer together by geodesic distance on the

underlying graph or manifold have more similar coordinate values, i.e., are closer together in

Euclidean distance, in the embedding. A subset of these eigenvectors of size $k = 1, 2, ..., N$

can then be used to embed nodes in $\mathbb{R}^k$. In the context of this manuscript, then, brain nodes

that are able to more efficiently pass information between one another are embedded closer

together via the Laplacian eigenmodes of a brain network [78].

To compute the temporally dependent diffusion-based embedding for a network, we follow

the methods discussed by Xiao and colleagues, where node coordinates are obtained from

the Young-Householder decomposition of the heat kernel [82,83]. The embedding matrix,

$Y = (y_1|y_2|...|Y_N)$, with columns as embedding coordinate vectors for network nodes can

determined with the heat kernel by $\mathcal{H}(t) = Y^T Y$. The Euclidean distance between nodes i

and j is then

$$d_E[i,j] = \sqrt{\sum_k^N e^{-\lambda_i t}(u_k[i] - u_k[j])} = \sqrt{(y_i - y_j)^T(y_i - y_j)}$$

where the distance in each dimension is scaled by exponentiating the product of the cor-

responding eigenvalue and diffusion time, t. A pairwise Euclidean distance matrix, $\mathcal{D}$, of

embedded nodes is computed for each subject with the time parameter βt obtained via the

structure to function mapping as described above. This provides a newly defined structural

connectome where edges are distances in the embedding space spanned by the eigenvectors of

the graph Laplacian, and the scale of distance in each embedding dimension is dependent on

the time parameter, βt, for each subject:

$$\text{SDD}[i,j] = \sqrt{\sum_k^N e^{-\lambda_i \beta t}(u_k[i] - u_k[j])}$$

Subnetwork identification

The structural diffusion distance (SDD) connectome edges then represent the distance between

brain regions in diffusion embedding at the diffusion depth that best represents empirical

resting-state functional connectivity. Many previous findings from studies of IPs implicate

perturbations in resting state functional connectivity, especially within the DMN, which is

defined by activity at rest. As such, SDD connectomes then used for further analysis to

identify structural subnetworks with aberrant diffusion characteristics pertinent to rs-fMRI

dynamics using the Network-based Statistic (NBS) algorithm [86]. Briefly, NBS is carried

out by computing a t-statistic at each network edge, applying a pre-determined threshold

to the resulting t-statistics and determining the connected components formed by supra-

threshold edges. To determine the significance of each identified subnetwork, the size of

each component is compared to a null distribution of maximum sized connected components

obtained by shuffling group labels for t-statistic calculation. NBS was carried out using SDD

connectomes in HC versus PT at baseline, using a left sided t-test with t-statistic thresholds

of $t = 3.0, 3.5, 4.0, 4.5$. In doing this, a subnetwork, $S \subset G$, where G is the full brain network

graph, with significantly greater embedding distances, or significantly decreased diffusion

capacity, in PT relative to HC was.

 NBS subnetwork hubs

595 Hub node identification for brain regions within the significant NBS subnetwork was performed

596 using edges from both the SDD and standard fiber count structural connectomes, and hub or

597 centrality metric values were then averaged by group. For SDD connectomes, strength, or

598 weighted degree, $\text{str}_{SDD}(n)$, for the i^{th} node, n_i, in S is defined as the sum of all edge weights

599 (distances) that are within the identified subnetwork, S, i.e.,

$$\text{str}_{SDD}(n_i) = \sum_{j \in S} \text{SDD}[i,j]$$

A high strength value would then indicate that a node has relatively less diffusion between

other nodes in the subnetwork, thus identifying nodes that may have the greatest diffusion

impairment. For structural connectomes, standard measures of nodal strength, betweenness

centrality, local efficiency and clustering coefficient [13, 72] were computed on the whole brain

network graph and subgraph, $S \subset G$, for each node in S. For each of the above scenarios,

metrics were also determined by cortical region by averaging region of interest (ROI) level

metrics by their cortex assignment as defined by the MMP1.0 360-node parcellation [34].

Heat kernel edge-based analyses

In order to study more granular characteristics of subnetwork diffusion, heat kernel values

between nodes within the significant NBS subnetwork were investigated using t-tests in HC

vs PT at baseline. Correlates of baseline symptom severity (using PT only) of and treatment

response (defined as $\frac{\text{scale}_{pre} - \text{scale}_{post}}{\text{scale}_{pre}}$ were found by computing non-parametric Spearmen rho

statistics between symptom scales and heat kernel values. As above, where cortex-averaged

metrics were computed, heat kernel edge-based analyses were conducted cortex-averaged

heat kernels in order to simplify interpretation and reduce the number of mass univariate

tests performed. To further focus this analysis, statistics other than baseline t-statistics were

only computed on heat kernel values corresponding to pairwise links between SSD hubs (as

determined above) and all other subnetwork nodes. P-values for all statistics were corrected

for multiple comparisons using False discovery rate correction (FDR) [10].

Subnetwork targets for supplemental heat

To identify potential brain regions for neuromodulatory treatment, we model a brain subnet-

work receiving supplemental heat by simply adding a heat kernel modifying matrix, whose

rows are made by repeating rows of the original heat kernel, to the original heat kernel. As

discussed previously, the product of an entry of the heat kernel matrix, $\mathcal{H}(t)[i, j]$, and the

heat at node i at time $t = 0$, $h(0)_i$, indicates the amount of heat transferred from node i to

node j at time t. The product of the i^{th} row of the heat kernel matrix, $\mathcal{H}(t)[i, :]$, and $h(0)_i$

then yields the distribution of heat values for each j^{th} node in the network. If the i^{th} node of

a network is supplied with supplemental heat independent of the initial distribution of heat

on the network, $h(0)$, the network heat distribution at time t is given by

$$h(t) = \mathcal{H}(t)h(0) + \mathcal{H}(t)\bar{h}$$

where $\bar{h}$ is a vector of all zeros except for the i^{th} whose value is the amount of supplemental

heat at node i. Using this model, we can then identify both a node and supplemental heat

value that may best 'normalize' a heat kernel representing impaired diffusion processes to a

reference heat kernel. In this study, we use the mean HC heat kernel matrix as the reference

and identify patient-specific nodes and supplemental heat values for optimal correction of the

diffusion dynamics encoded in the heat kernel, described as follows.

Given the mean HC heat kernel, $\mathcal{H}_{HC}$, and the k^{th} patient's heat kernel, $\mathcal{H}_{PT}^k$, we compute

the residual heat kernel matrix as $\bar{\mathcal{H}}_i^k = \mathcal{H}_{HC} - \mathcal{H}_{PT}^k$. For the i^{th} node in the subnetwork, we

construct the heat kernel modifier matrix as $\mathcal{M}_i^k = \mathbf{1}(\mathcal{H}_{PT}^k[i, :])^T$ as the outer product of the

all ones vector and the i^{th} row of the k^{th} patient's heat kernel matrix. In order to find the

optimal supplementary heat value, c_i^k for a given node and subject, we minimize the following

objective function.

$$\underset{c_i^k}{\text{minimize}} ||\bar{\mathcal{H}}_i^k - c_i^k \mathcal{M}_i^k||_2$$

641 Which can be easily solved analytically by

$$c_i^k = \frac{\mathrm{tr}((\mathcal{M}_i^k)^T \bar{\mathcal{H}}_i^k)}{\mathrm{tr}((\mathcal{M}_i^k)^T \mathcal{M}_i^k)}$$

where $\mathrm{tr}(\cdot)$ denotes the matrix trace.

It is important to note, that the above model is also constrained such that only positive
values in the residual matrix are used for optimization, as this simple network diffusion model
only allows for the addition of heat.

This process is repeated for each unique node-patient pair, and the mean or patient-
specific optimal node and corresponding supplemental heat value can be determined. We have
experimentally observed that the minimum error between residual and heat kernel modifier
matrices has very little variation between brain regions, and we define optimal region as the
node with the lowest value supplemental heat value, such that neuromodulatory efficacy could
be achieved with lower amounts of stimulation and thus potentially yield fewer undesired
effects.

Visualizing effects of heat supplementation

To assess whether the significant group differences at baseline between HC and PT heat
kernel values are corrected with heat supplementation, we add the product of the optimal
supplemental heat value, c_i^k, at the i^{th} node, and the heat kernel modifier matrix, $\mathcal{M}_i^k$ to
each k^{th} patient's heat kernel. We then compute a t-statistic for each heat kernel entry, as
discussed above.

References

[1] F. Abdelnour, M. Dayan, O. Devinsky, T. Thesen, and A. Raj. Functional brain connectivity is predictable from anatomic network's laplacian eigen-structure. *NeuroImage*, 172:728–739, 2018.

[2] F. Abdelnour, H. U. Voss, and A. Raj. Network diffusion accurately models the relationship between structural and functional brain connectivity networks. *Neuroimage*, 90:335–347, 2014.

[3] A. A. Abdelrahman, M. Noaman, M. Fawzy, A. Moheb, A. A. Karim, and E. M. Khedr. A double-blind randomized clinical trial of high frequency rtms over the dlpfc on nicotine dependence, anxiety and depression. *Scientific Reports*, 11(1):1–10, 2021.

[4] M. Assem, M. F. Glasser, D. C. Van Essen, and J. Duncan. A domain-general cognitive core defined in multimodally parcellated human cortex. *Cerebral Cortex*, 30(8):4361–4380, 2020.

[5] S. Atasoy, I. Donnelly, and J. Pearson. Human brain networks function in connectome-specific harmonic waves. *Nature communications*, 7(1):1–10, 2016.

[6] A. Avena-Koenigsberger, B. Misic, and O. Sporns. Communication dynamics in complex brain networks. *Nature Reviews Neuroscience*, 19(1):17, 2018.

[7] F. Barkhof, S. Haller, and S. A. Rombouts. Resting-state functional mr imaging: a new window to the brain. *Radiology*, 272(1):29–49, 2014.

[8] Y. Behzadi, K. Restom, J. Liau, and T. T. Liu. A component based noise correction method (compcor) for bold and perfusion based fmri. *Neuroimage*, 37(1):90–101, 2007.

[9] M. Belkin and P. Niyogi. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural computation*, 15(6):1373–1396, 2003.

[10] Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.

[11] L. Beynel, L. Deng, C. A. Crowell, M. Dannhauer, H. Palmer, S. Hilbig, A. V. Peterchev, B. Luber, S. H. Lisanby, R. Cabeza, et al. Structural controllability predicts functional patterns and brain stimulation benefits associated with working memory. *Journal of Neuroscience*, 40(35):6770–6778, 2020.

[12] S. Brooks, J. Ipser, and D. Stein. Chronic and acute nicotine exposure versus placebo in smokers and nonsmokers: A systematic review of resting-state fmri studies. *Addictive Substances and Neurological Disease*, pages 319–338, 2017.

[13] E. Bullmore and O. Sporns. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature reviews neuroscience*, 10(3):186–198, 2009.

[14] K. L. Burkhouse, S. M. Gorka, H. Klumpp, A. E. Kennedy, S. Karich, J. Francis, O. Ajilore, M. G. Craske, S. A. Langenecker, and S. A. Shankman. Neural responsiveness to reward as an index of depressive symptom change following cognitive-behavioral therapy and ssri treatment. *The Journal of clinical psychiatry*, 79(4):0–0, 2018.

[15] K. L. Burkhouse, J. Jimmy, N. Defelice, H. Klumpp, O. Ajilore, B. Hosseini, K. D. Fitzgerald, C. S. Monk, and K. L. Phan. Nucleus accumbens volume as a predictor of anxiety symptom improvement following cbt and ssri treatment in two independent samples. *Neuropsychopharmacology*, 45(3):561–569, 2020.

[16] K. L. Burkhouse, A. Kujawa, B. Hosseini, H. Klumpp, K. D. Fitzgerald, S. A. Langenecker, C. S. Monk, and K. L. Phan. Anterior cingulate activation to implicit threat before and after treatment for pediatric anxiety disorders. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 84:250–256, 2018.

[17] J. L. Cecilione, L. M. Rappaport, S. E. Hahn, A. E. Anderson, L. E. Hazlett, J. R. Burchett, A. A. Moore, J. E. Savage, J. M. Hettema, and R. Roberson-Nay. Genetic and

environmental contributions of negative valence systems to internalizing pathways. *Twin research and human genetics: the official journal of the International Society for Twin Studies*, 21(1):12, 2018.

[18] L. Chen, A.-R. Hudaib, K. E. Hoy, and P. B. Fitzgerald. Is rtms effective for anxiety symptoms in major depressive disorder? an efficacy analysis comparing left-sided high-frequency, right-sided low-frequency, and sequential bilateral rtms protocols. *Depression and anxiety*, 36(8):723–731, 2019.

[19] A. W. Chung, M. Schirmer, M. L. Krishnan, G. Ball, P. Aljabar, A. D. Edwards, and G. Montana. Characterising brain network topologies: a dynamic analysis approach using heat kernels. *Neuroimage*, 141:490–501, 2016.

[20] F. R. Chung and F. C. Graham. *Spectral graph theory*. Number 92. American Mathematical Soc., 1997.

[21] F. R. Chung and F. C. Graham. *Spectral graph theory*. Number 92. American Mathematical Soc., 1997.

[22] B. N. Cuthbert. The rdoc framework: facilitating transition from icd/dsm to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry*, 13(1):28–35, 2014.

[23] S. Deslauriers-Gauthier, M. Zucchelli, M. Frigo, and R. Deriche. A unified framework for multimodal structure–function mapping based on eigenmodes. *Medical Image Analysis*, 66:101799, 2020.

[24] N. U. Dosenbach, D. A. Fair, F. M. Miezin, A. L. Cohen, K. K. Wenger, R. A. Dosenbach, M. D. Fox, A. Z. Snyder, J. L. Vincent, M. E. Raichle, et al. Distinct brain networks for adaptive and stable task control in humans. *Proceedings of the National Academy of Sciences*, 104(26):11073–11078, 2007.

[25] L. Du, H. Liu, W. Du, F. Chao, L. Zhang, K. Wang, C. Huang, Y. Gao, and Y. Tang. Stimulated left dlpfc-nucleus accumbens functional connectivity predicts the anti-depression and anti-anxiety effects of rtms for depression. *Translational psychiatry*, 7(11):1–10, 2018.

[26] B. W. Dunlop, E. B. Binder, J. F. Cubells, M. M. Goodman, M. E. Kelley, B. Kinkead, M. Kutner, C. B. Nemeroff, D. J. Newport, M. J. Owens, et al. Predictors of remission in depression to individual and combined treatments (predict): study protocol for a randomized controlled trial. *Trials*, 13(1):1–18, 2012.

[27] A. Dutta, S. McKie, and J. W. Deakin. Resting state networks in major depressive disorder. *Psychiatry Research: Neuroimaging*, 224(3):139–151, 2014.

[28] M. K. Eradath, H. Abe, M. Matsumoto, K. Matsumoto, K. Tanaka, and N. Ichinohe. Anatomical inputs to sulcal portions of areas 9m and 8bm in the macaque monkey. *Frontiers in neuroanatomy*, 9:30, 2015.

[29] A. Etkin and T. D. Wager. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in ptsd, social anxiety disorder, and specific phobia. *American journal of Psychiatry*, 164(10):1476–1488, 2007.

[30] M. First, J. Williams, R. Karg, R. Spitzer, et al. Structured clinical interview for dsm-5—research version (scid-5 for dsm-5, research version; scid-5-rv). *Arlington, VA: American Psychiatric Association*, pages 1–94, 2015.

[31] A. Fornito, A. Zalesky, and M. Breakspear. The connectomics of brain disorders. *Nature Reviews Neuroscience*, 16(3):159–172, 2015.

[32] J. Francis, O. Ajilore, S. A. Langenecker, S. A. Shankman, and M. G. Craske. Emotion-based brain mechanisms and predictors for ssri and cbt treatment of anxiety and depression: A randomized trial. 2019.

[33] M. S. George, J. J. Taylor, and E. B. Short. The expanding evidence base for rtms treatment of depression. *Current opinion in psychiatry*, 26(1):13, 2013.

[34] M. F. Glasser, T. S. Coalson, E. C. Robinson, C. D. Hacker, J. Harwell, E. Yacoub, K. Ugurbil, J. Andersson, C. F. Beckmann, M. Jenkinson, et al. A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615):171–178, 2016.

[35] Q. Gong and Y. He. Depression, neuroimaging and connectomics: a selective overview. *Biological psychiatry*, 77(3):223–235, 2015.

[36] S. M. Gorka, C. B. Young, H. Klumpp, A. E. Kennedy, J. Francis, O. Ajilore, S. A. Langenecker, S. A. Shankman, M. G. Craske, M. B. Stein, et al. Emotion-based brain mechanisms and predictors for ssri and cbt treatment of anxiety and depression: a randomized trial. *Neuropsychopharmacology*, 44(9):1639–1648, 2019.

[37] J. P. Hamilton, M. C. Chen, and I. H. Gotlib. Neural systems approaches to understanding major depressive disorder: an intrinsic functional organization perspective. *Neurobiology of disease*, 52:4–11, 2013.

[38] J. P. Hamilton, A. Etkin, D. J. Furman, M. G. Lemus, R. F. Johnson, and I. H. Gotlib. Functional neuroimaging of major depressive disorder: a meta-analysis and new integration of baseline activation and neural response data. *American Journal of Psychiatry*, 169(7):693–703, 2012.

[39] K. Helm, K. Viol, T. M. Weiger, P. A. Tass, C. Grefkes, D. Del Monte, and G. Schiepek. Neuronal connectivity in major depressive disorder: a systematic review. *Neuropsychiatric disease and treatment*, 2018.

[40] C. J. Honey, O. Sporns, L. Cammoun, X. Gigandet, J.-P. Thiran, R. Meuli, and P. Hagmann. Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences*, 106(6):2035–2040, 2009.

[41] T. Insel, B. Cuthbert, M. Garvey, R. Heinssen, D. S. Pine, K. Quinn, C. Sanislow, and P. Wang. Research domain criteria (rdoc): toward a new classification framework for research on mental disorders, 2010.

[42] M. Jenkinson, C. F. Beckmann, T. E. Behrens, M. W. Woolrich, and S. M. Smith. Fsl. *Neuroimage*, 62(2):782–790, 2012.

[43] R. H. Kaiser, J. R. Andrews-Hanna, T. D. Wager, and D. A. Pizzagalli. Large-scale network dysfunction in major depressive disorder: a meta-analysis of resting-state functional connectivity. *JAMA psychiatry*, 72(6):603–611, 2015.

[44] R. C. Kessler, W. T. Chiu, O. Demler, and E. E. Walters. Prevalence, severity, and comorbidity of 12-month dsm-iv disorders in the national comorbidity survey replication. *Archives of general psychiatry*, 62(6):617–627, 2005.

[45] Y.-K. Kim and H.-K. Yoon. Common and distinct brain networks underlying panic and social anxiety disorders. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 80:115–122, 2018.

[46] H. Klumpp, M. Angstadt, and K. L. Phan. Insula reactivity and connectivity to anterior cingulate cortex when processing threat in generalized social anxiety disorder. *Biological Psychology*, 89(1):273–276, 2012.

[47] H. Klumpp, D. A. Fitzgerald, and K. L. Phan. Neural predictors and mechanisms of cognitive behavioral therapy on threat processing in social anxiety disorder. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 45:83–91, 2013.

[48] H. Klumpp, J. Jimmy, K. L. Burkhouse, R. Bhaumik, J. Francis, M. G. Craske, K. L. Phan, and O. Ajilore. Brain response to emotional faces in anxiety and depression: neural predictors of cognitive behavioral therapy outcome and predictor-based subgroups following therapy. *Psychological Medicine*, pages 1–11, 2020.

[49] H. Klumpp, D. Post, M. Angstadt, D. A. Fitzgerald, and K. L. Phan. Anterior cingulate cortex and insula response during indirect and direct processing of emotional faces in generalized social anxiety disorder. *Biology of mood & anxiety disorders*, 3(1):1–9, 2013.

[50] M. S. Korgaonkar, A. Fornito, L. M. Williams, and S. M. Grieve. Abnormal structural networks characterize major depressive disorder: a connectome analysis. *Biological psychiatry*, 76(7):567–574, 2014.

[51] R. Leech and D. J. Sharp. The role of the posterior cingulate cortex in cognition and disease. *Brain*, 137(1):12–32, 2014.

[52] J. Liu, X. Xu, C. Zhu, L. Luo, Q. Wang, B. Xiao, B. Feng, L. Hu, and L. Liu. Disrupted structural brain network organization behind depressive symptoms in major depressive disorder. *Frontiers in psychiatry*, 11:1008, 2020.

[53] B. M. Luber, S. Davis, E. Bernhardt, A. Neacsiu, L. Kwapil, S. H. Lisanby, and T. J. Strauman. Reprint of "using neuroimaging to individualize tms treatment for depression: Toward a new paradigm for imaging-guided intervention". *Neuroimage*, 151:65–71, 2017.

[54] B. Luo, R. C. Wilson, and E. R. Hancock. Spectral embedding of graphs. *Pattern recognition*, 36(10):2213–2230, 2003.

[55] B. Mišić, R. F. Betzel, A. Nematzadeh, J. Goni, A. Griffa, P. Hagmann, A. Flammini, Y.-Y. Ahn, and O. Sporns. Cooperative and competitive spreading dynamics on the human connectome. *Neuron*, 86(6):1518–1529, 2015.

[56] D. Momi, R. A. Ozdemir, E. Tadayon, P. Boucher, M. M. Shafi, A. Pascual-Leone, and E. Santarnecchi. Network-level macroscale structural connectivity predicts propagation of transcranial magnetic stimulation. *NeuroImage*, 229:117698, 2021.

[57] R. Morecraft, K. Stilwell-Morecraft, P. Cipolloni, J. Ge, D. McNeal, and D. Pandya. Cytoarchitecture and cortical connections of the anterior cingulate and adjacent somatomotor fields in the rhesus monkey. *Brain research bulletin*, 87(4-5):457–497, 2012.

[58] M. Moreno-Ortega, A. Kangarlu, S. Lee, T. Perera, J. Kangarlu, T. Palomo, M. Glasser, and D. Javitt. Parcel-guided rtms for depression. *Translational psychiatry*, 10(1):1–6, 2020.

[59] J. S. Moser, C. E. Durbin, C. J. Patrick, and N. B. Schmidt. Combining neural and behavioral indicators in the assessment of internalizing psychopathology in children and adolescents. *Journal of Clinical Child & Adolescent Psychology*, 44(2):329–340, 2015.

[60] V. I. Müller, E. C. Cieslik, A. R. Laird, P. T. Fox, and S. B. Eickhoff. Dysregulated left inferior parietal activity in schizophrenia and depression: functional connectivity and characterization. *Frontiers in human neuroscience*, 7:268, 2013.

[61] A. Y. Ng, M. I. Jordan, Y. Weiss, et al. On spectral clustering: Analysis and an algorithm. *Advances in neural information processing systems*, 2:849–856, 2002.

[62] A. Nieto-Castanon. *Handbook of fcMRI methods in CONN*. Hilbert Press, 2020.

[63] S. Ofrat and R. F. Krueger. How research on the meta-structure of psychopathology aids in understanding biological correlates of mood and anxiety disorders. *Biology of mood & anxiety disorders*, 2(1):1–4, 2012.

[64] M. Radoman, K. L. Phan, and S. M. Gorka. Neural correlates of predictable and unpredictable threat in internalizing psychopathology. *Neuroscience letters*, 701:193–201, 2019.

[65] A. Raj, A. Kuceyeski, and M. Weiner. A network diffusion model of disease progression in dementia. *Neuron*, 73(6):1204–1215, 2012.

[66] G. Roberts, A. Lord, A. Frankland, A. Wright, P. Lau, F. Levy, R. K. Lenroot, P. B. Mitchell, and M. Breakspear. Functional dysconnection of the inferior frontal gyrus in young people with bipolar disorder or at genetic high risk. *Biological psychiatry*, 81(8):718–727, 2017.

[67] E. T. Rolls, W. Cheng, J. Du, D. Wei, J. Qiu, D. Dai, Q. Zhou, P. Xie, and J. Feng. Functional connectivity of the right inferior frontal gyrus and orbitofrontal cortex in depression. *Social cognitive and affective neuroscience*, 15(1):75–86, 2020.

[68] M. D. Schirmer and A. W. Chung. Heat kernels with functional connectomes reveal atypical energy transport in peripheral subnetworks in autism. In *International Workshop on Connectomics in Neuroimaging*, pages 54–63. Springer, 2019.

[69] W. W. Seeley, V. Menon, A. F. Schatzberg, J. Keller, G. H. Glover, H. Kenna, A. L. Reiss, and M. D. Greicius. Dissociable intrinsic connectivity networks for salience processing and executive control. *Journal of Neuroscience*, 27(9):2349–2356, 2007.

[70] Y. I. Sheline, D. M. Barch, J. L. Price, M. M. Rundle, S. N. Vaishnavi, A. Z. Snyder, M. A. Mintun, S. Wang, R. S. Coalson, and M. E. Raichle. The default mode network and self-referential processes in depression. *Proceedings of the National Academy of Sciences*, 106(6):1942–1947, 2009.

[71] O. Sporns. *Networks of the Brain.* MIT press, 2010.

[72] O. Sporns. Graph theory methods: applications in brain networks. *Dialogues in clinical neuroscience*, 20(2):111, 2018.

[73] C. Stam, E. Van Straaten, E. Van Dellen, P. Tewarie, G. Gong, A. Hillebrand, J. Meier, and P. Van Mieghem. The relation between structural and functional connectivity patterns in complex brain networks. *International Journal of Psychophysiology*, 103:149–160, 2016.

[74] J. Stiso, A. N. Khambhati, T. Menara, A. E. Kahn, J. M. Stein, S. R. Das, R. Gorniak, J. Tracy, B. Litt, K. A. Davis, et al. White matter network architecture guides direct electrical stimulation through optimal state transitions. *Cell reports*, 28(10):2554–2566, 2019.

[75] C. M. Sylvester, M. Corbetta, M. Raichle, T. Rodebaugh, B. Schlaggar, Y. Sheline, C. Zorumski, and E. Lenze. Functional network dysfunction in anxiety and anxiety disorders. *Trends in neurosciences*, 35(9):527–535, 2012.

[76] Y. Tao, B. Liu, X. Zhang, J. Li, W. Qin, C. Yu, and T. Jiang. The structural connectivity pattern of the default mode network and its association with memory and anxiety. *Frontiers in neuroanatomy*, 9:152, 2015.

[77] P. J. Thomas, S. Panchamukhi, J. Nathan, J. Francis, S. Langenecker, S. Gorka, A. Leow, H. Klumpp, K. L. Phan, and O. A. Ajilore. Graph theoretical measures of the uncinate fasciculus subnetwork as predictors and correlates of treatment response in a transdiagnostic psychiatric cohort. *Psychiatry Research: Neuroimaging*, 299:111064, 2020.

[78] M. B. Wang, J. P. Owen, P. Mukherjee, and A. Raj. Brain network eigenmodes provide a robust and compact representation of the structural connectome in health and disease. *PLoS computational biology*, 13(6):e1005550, 2017.

[79] D. Watson, M. W. O'Hara, K. Naragon-Gainey, E. Koffel, M. Chmielewski, R. Kotov, S. M. Stasik, and C. J. Ruggero. Development and validation of new anxiety and bipolar symptom scales for an expanded version of the idas (the idas-ii). *Assessment*, 19(4):399–420, 2012.

[80] S. Whitfield-Gabrieli and A. Nieto-Castanon. Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain connectivity*, 2(3):125–141, 2012.

[81] X. Wu, P. Lin, J. Yang, H. Song, R. Yang, and J. Yang. Dysfunction of the cingulo-opercular network in first-episode medication-naive patients with major depressive disorder. *Journal of affective disorders*, 200:275–283, 2016.

[82] B. Xiao, E. R. Hancock, and R. C. Wilson. Geometric characterization and clustering of graphs using heat kernel embeddings. *Image and Vision Computing*, 28(6):1003–1021, 2010.

[83] B. Xiao, R. C. Wilson, and E. R. Hancock. Characterising graphs using the heat kernel. In *Proc. BMVC*, 2005.

[84] R. Yang, C. Gao, X. Wu, J. Yang, S. Li, and H. Cheng. Decreased functional connectivity to posterior cingulate cortex in major depressive disorder. *Psychiatry Research: Neuroimaging*, 255:15–23, 2016.

[85] F.-C. Yeh, T. D. Verstynen, Y. Wang, J. C. Fernández-Miranda, and W.-Y. I. Tseng. Deterministic diffusion fiber tracking improved by quantitative anisotropy. *PloS one*, 8(11):e80713, 2013.

[86] A. Zalesky, A. Fornito, and E. T. Bullmore. Network-based statistic: identifying differences in brain networks. *Neuroimage*, 53(4):1197–1207, 2010.