Antibodies against type-I Interferon: detection and association with severe clinical outcome in COVID-19 patients

Goncalves David[1], Mezidi Mehdi[2], Bastard Paul[3,4,5], Perret Magali[1,6], Saker Kahina[7], Fabien Nicole[1], Pescarmona Rémi[1,6], Lombard Christine[1], Walzer Thierry[6], Casanova Jean-Laurent[3,4,5], Belot Alexandre[6,8,9], Richard Jean-Christophe[2], Trouillet-Assant Sophie[6,7]

[1]Immunology Department, Lyon Sud Hospital, Hospices Civils de Lyon, Pierre-Bénite, France.
[2]Lyon University, CREATIS, CNRS UMR5220, Inserm U1044, INSA, Lyon, France, Intensive Care Unit, Hospices Civils de Lyon, Lyon, France
[3]Laboratory of Human Genetics of Infectious Diseases, Necker Branch, INSERM U1163, Necker Hospital for Sick Children, Paris, France.
[4]University of Paris, Imagine Institute, Paris, France.
[5]St. Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The Rockefeller University, New York, NY, USA.
[6]International Center of Research in Infectiology, Lyon University, INSERM U1111, CNRS UMR 5308, ENS, UCBL, Lyon, France
[7]Infective Agents Institute, Hospices Civils de Lyon, Lyon, France
[8] National Referee Centre for Rheumatic and AutoImmune and Systemic diseases in childrEn (RAISE), France
[9] Pediatric Nephrology, Rheumatology, Dermatology Unit, Hospices Civils de Lyon, France

Corresponding author:
Sophie Trouillet-Assant, PhD
Centre Hospitalier Lyon Sud, Pierre-Bénite
69495 Lyon, France.
Phone: + 33 (0) 472 00 37 70. sophie.assant@chu-lyon.fr

Running head: Anti-interferon antibodies in COVID-19 patients

Abstract

Objectives Impairment of type I interferon (IFN-I) immunity has been reported in critically-ill COVID-19 patients. This defect can be explained in a subset of patients by the presence of circulating autoantibodies (auto-Abs) against IFN-I. We set out to improve the detection and the quantification of IFN-I auto-Abs in a cohort of critically-ill COVID-19 patients, in order to better evaluate the prevalence of these Abs as the pandemic progresses, and how they correlate with the clinical course of the disease.

Methods The concentration of anti-IFN-α_2 Abs was determined in the serum of 84 critically-ill COVID-19 patients who were admitted to ICU in *Hospices Civils de Lyon*, France using a commercially available kit (Thermo-Fisher, Catalog #BMS217).

Results A total of 21/84 (25%) critically-ill COVID-19 patients had circulating anti-IFN-α_2 Abs above cut-off (>34 ng.mL^{-1}). Among them, 15/21 had Abs with neutralizing activity against IFN-α_2, *i.e.* 15/84 (18%) of critically-ill patients. In addition, we noticed an impairment of the IFN-I response in the majority of patients with neutralizing anti-IFN-α_2 Abs. There was no significant difference in the clinical characteristics or outcome of with or without neutralizing anti-IFN-α_2 auto-Abs. We detected anti-IFN-α_2 auto-Abs in COVID-19 patients' sera throughout their ICU stay. Finally, we also found auto-Abs against multiple subtypes of IFN-I including IFN-ω.

Conclusions We reported that 18% of critically-ill COVID-19 patients were positive for IFN-I auto-Abs, confirming that the presence of these antibodies is associated with higher risk of developing a criticall COVID-19 form.

Keywords — Type I interferon; COVID-19; SARS-CoV-2 virus; Intensive care unit; viral infection; auto-antibodies.

Introduction

Severe acute respiratory syndrome coronavirus-2 (SARS-CoV2) infection leads to coronavirus disease 19 (COVID-19), whose spectrum of clinical presentations is wide and includes severe pneumonia. The anti-SARS-CoV-2 immune response has been extensively studied and defects in antiviral mechanisms have been linked to disease severity. In particular, impairment of type I interferon (IFN-I) immunity has been reported in critically-ill COVID-19 patients. Such defect can be due to either inherited genetic deficiencies in the IFN-I pathway or the occurrence of circulating auto-antibodies (auto-Abs) directed against 14 or the 17 individual IFN-I [1-4]. These auto-Abs have also been detected in a third of patients from a small international cohort who had suffered from severe adverse events following Yellow Fever vaccination (YFV-17D)[5]. These findings advocate for the development of diagnostic tools for the detection of IFN-I auto-Abs in routine laboratories, in order to identify early patients at risk of developing severe forms of COVID-19 and to analyze the prevalence of IFN-I Abs as the pandemic progresses and the virus evolves. To this aim, we tested a commercially available kit measuring IFN-I auto-Abs levels in the serum of COVID-19 patients.

Results

A total of 84 critically-ill COVID-19 patients, 11 patients with autoimmune polyendocrinopathy type 1

syndrome (APS-1), 10 mildly-symptomatic COVID-19 healthcare workers, and 76 healthy controls

were included in the study. The critically-ill COVID-19 patients were admitted to ICU in the Lyon

University Hospital, France, between September and December 2020. The presence of anti IFN-α_2

Abs was investigated: we first sought to determine a positive cut-off value for Abs detection by

performing measurements in 76 putative control sera, *i.e.* from healthy donors retrieved before the

COVID-19 outbreak. The mean value +3 standard deviation of these measurements provided a cut-

off value at 34 ng.mL^{-1}. We then assessed the presence of IFN-α2 Abs in putative positive sera, *i.e.*

sera from patients with autoimmune polyendocrinopathy type 1 syndrome (APS-1), a condition

known to be associated with anti-cytokine auto-Abs. All APS-1 patients tested (n=11) had high titers

of circulating anti-IFN-α2 auto-Abs (>100ng.mL^{-1}). We also evaluated the presence of anti-IFN-α_2

auto-Abs in 10 mildly-symptomatic COVID-19 healthcare workers and none of them was found

positive.

We then measured anti-IFN-α_2 Abs levels in the sera from the critically-ill COVID-19 patients: 21/84

(25%) were positive and had values above the cut-off (>34 ng.mL^{-1}). The neutralizing capacity of their

sera against IFN-α was then evaluated as previously described[5]. A neutralizing activity was observed

in 15/21 positive sera; in other words, 15/84 (18%) critically-ill COVID-19 patients had neutralizing

anti-IFN-α auto-Abs (Figure 1**a**). Importantly, all sera with a titer of anti-IFN-α_2 auto-Abs above 1

µg.mL^{-1} potently neutralized IFN-α *in vitro*. In addition, in most patients with neutralizing IFN-I Abs,

we noticed an impairment of the IFN-I response, which was determined by the measurement of i)

plasma IFN-α_2 levels using the new digital ELISA technology single-molecule arrays (Simoa) and ii)

blood Interferon Stimulating Genes (ISG) expression using the Nanostring nCounter technology in

blood samples collected in the first 15 days after symptom onset (Figure 1**b-c**). Moreover, there was

no significant difference in the clinical characteristics (age, sex ratio, co-morbidity) or outcome

(death, O_2 support) of critically-il COVID-19 patients with or without neutralizing anti-IFN-I auto-Abs

(Table 1). Then, serial measurement of IFN-I auto-Abs level during ICU stay was performed for 7

positive patients. The level of IFN-α_2 auto-Abs remained relatively stable across measurements

performed up to 40 days apart (Figure 1**d**). Finally, we assessed the presence of auto-Abs against

other IFNs-I in all sera positive for IFN-α2 auto-Abs (n=21). Consistently with previous results[1], we

observed that sera with anti-IFN-α_2 auto-Ab titers above 1 µg.mL^{-1} also contained auto-Abs against

other subtypes of IFN-α and 10/12 contained anti-IFN-ω auto-Abs (Figure 2). Of note, these sera

were able to neutralize IFN-ω *in vitro*. Only one serum contained auto-Abs targeting IFN-β at a low

titer, and none contained auto-Abs against IFN-ε or IFN-κ.

Discussion and conclusions

A previous study has reported that IFN-I auto-Abs were present in 10.2% of life-threatening COVID-

19 patients, undetectable in 663 individuals with asymptomatic or mild COVID-19, and detected in

only 0.33% of healthy individuals [1]. Here, 18% of critically-ill COVID-19 patients were positive for IFN-I

auto- Abs whereas all mild-COVID19 patients were negative. We noticed that only a part of auto-Abs

detected were able to neutralize IFN-I in the conditions we used, which confirms previous studies in

COVID-19 patients and systemic lupus erythematosus subjects [1, 6]. This finding further confirms the

deleterious role of IFN-I autoAbs in the antiviral immune response and the importance of the IFN-I

pathway in the defense against SARS-CoV2 infection. Based on its antiviral properties, recombinant

IFN-I has been tested as therapy for severe COVID-19, but the treatment showed little or no benefit

7-

8. Yet, the potential of such treatment may have been hindered by the presence of IFN-I auto-Abs in

patient sera, and this question could therefore be revisited by determining the level of these Abs, for

example using the ELISA method we used here. Moreover, patients could be treated with

recombinant

IFN-I that are not targeted by auto-Abs (*e.g.* IFN-β).

Finally, the detection of anti IFN-I auto-Abs in COVID-19 patients could be useful in routine to identify

patients at risk of developing a severe form of the disease. The technique we described here is

adequate for this purpose as it can rapidly provide quantitative measurements, and has a cut-off

correlated with neutralization assays (1 $\mu g.mL^{-1}$). However, the presence of auto-Abs was not

associated with poorer outcome in critically-ill patients and does not explain all the severe forms of

COVID-19, other causes should therefore be sought (*e.g.* cytokine release syndrome, presence of

other risk factors such as obesity, hypertension).

Methods

Participants *Critically-ill COVID-19 patients*

Plasma samples and Paxgene® tubes were collected from COVID-19 patients hospitalized in the

university hospital of Lyon (Hospices Civils de Lyon), France. Diagnosis of COVID-19 was established in

all patients by RT-PCR.

All critically-ill patients positive for SARS-CoV-2 virus, admitted to ICU (Croix-Rousse Hospital,

Hospices Civils de Lyon), were included in the MIR-COVID study. This study was registered to the

Commission nationale de l'informatique et des libertés (CNIL, French data protection agency) under

the number 20-097 and was approved by an ethics committee for biomedical research (*Comité de*

Protection des Personnes HCL) under the number N°20-41. In agreement with the General Data

Protection Regulation (Regulation (EU) 2016/679 and Directive 95/46/EC) and the French data

protection law (Law n°78-17 on 06/01/1978 and Décret n°2019-536 on 29/05/2019), we obtained

consent from each patient or his/her next of kin.

Mildly-symptomatic COVID-19 patients

Plasma samples and Paxgene® tubes were collected from symptomatic healthcare workers upon

COVID-19 diagnosis. Written informed consent was obtained from all participants. The study was

approved by the national review board for biomedical research in April 2020 (Comité de Protection

des Personnes Sud Méditerranée I, Marseille, France; ID RCB 2020-A00932-37). The study was

registered on ClinicalTrials.gov (NCT04341142) where the eligibility, inclusion, and exclusion criteria

are previously described [9].

Healthy controls

Prepandemic serum were selected from healthy controls who were recruited among donors to the

Lyon blood transfusion centre (Etablissement Français du Sang, EFS). According to French

procedures, a written non-opposition to the use of donated blood for research purposes was

obtained from HCs. The donors' personal data were anonymized before transfer to our research

laboratory. We obtained approval from the local ethical committee and the French ministry of

research (DC-2008-64) for handling and conservation of these samples.

Auto-Abs anti IFN-I

The presence of anti IFN-α2 auto-Abs was assessed in the plasma using a commercially available kit

(Thermo-Fisher, Catalog # BMS217). The positive cut-off value for Ab detection was 34 ng.mL^{-1}. The

presence of auto-Abs against other IFNs-I was assessed using an ELISA technique, as previously

described [1]. Briefly, 96-well ELISA plates were coated with different cytokines (rhIFN-α2 (Milteny

Biotec, ref. number 130-108-984), rhIFN-ω (Merck, ref. number SRP3061) or cytokines from PBL

Assay Science (catalog #11002-1), or IFN-β (Milteny Biotech, ref. number: 130-107-888)) and

incubated overnight at 4°C. Plates were then washed (PBS 0.005% Tween), then incubated with 5%

nonfat milk powder in the same buffer, washed again, and again incubated with 1:50 dilutions of

plasma from patients or controls for 2 h at room temperature (or with specific mAbs as positive

controls). Horseradish peroxidase (HRP)–conjugated Fc-specific IgG fractions from polyclonal goat

antiserum against human IgG, IgM, or IgA (Nordic Immunological Laboratories) were added to a final

concentration of 2 µg/mL after thorough washing. Plates were then incubated for 1 h at room

temperature and washed. HRP substrate was added and the optical density (OD) was measured.

The neutralization capacity of antibodies against IFN-α2 and −ω was determined as previously

described [5]. Briefly, HEK-293T cells were transfected with a plasmid encoding the firefly Luciferase

under the control of human *ISRE* promoters. Cells were then cultured in Dulbecco's modified Eagle

medium (DMEM, Thermo Fisher Scientific) supplemented with 10% healthy control or patient

serum/plasma, and were either left unstimulated or were stimulated with IFN-α2, IFN-ω, or IFN-β (10

ng.mL^{-1}) for 16 hours at 37°C. Finally, Luciferase level was measured using the Dual-Glo reagent,

according to the manufacturer's instructions (Promega).

Plasma protein quantification

The concentration of plasmatic IFN-α (fg.mL^{-1}) was measured using single molecule array (Simoa)

using a commercial kit for IFN-α$_2$ quantification (Quanterix™, Lexington, MA, USA). The assay was

based on a 3- step protocol and an HD- 1 Analyzer (Quanterix).

IFN score assessment

Total RNA was extracted from whole blood stored in Paxgene® tubes (Kit PreAnalytix, Qiagen©, SW)

and was quantified using a spectrophotometer (Nanodrop 2000, Thermo Scientific™, MA, USA). RNA

integrity was then assessed using the Agilent RNA microarray (Agilent Technologies©, Santa Clara,

CA, USA). The expression of 6 ISGs (*interferon alpha inducible protein 27 (IFI27), interferon induced*

protein 44 like (IFI44L), Interferon Induced Protein With Tetratricopeptide Repeats 1 (IFIT1) , ISG15

Ubiquitin Like Modifier (ISG15), Radical S-Adenosyl Methionine Domain Containing 2 (RSAD2), Sialic

Acid Binding Ig Like Lectin 1 (SIGLEC1)) and 3 housekeeping genes (*Actin Beta (ACTB), Hypoxanthine*

Phosphoribosyltransferase 1 (HPRT1), RNA Polymerase II Subunit A (POLR2A)) was quantified at the

transcript level using the Nanostring technology (Nanostring Technologies©, WA, USA). Data

standardization was performed using the geometric mean of internal control and housekeeping

genes counts. The ISG score was calculated as previously described [10].

Conflict of interest: All authors declare no conflict of interest

Funding: This research was supported by the *Hospices Civils de Lyon* and by the *Fondation des*
Hospices Civils de Lyon.

References

1.   Bastard P, Rosen LB, Zhang Q, *et al.* Autoantibodies against type I IFNs in patients with life-
threatening COVID-19. *Science* 2020; **370**: eabd4585.
2.   Hadjadj J, Yatim N, Barnabei L, *et al.* Impaired type I interferon activity and inflammatory
responses in severe COVID-19 patients. *Science* 2020; **369**: 718-724.
3.   Trouillet-Assant S, Viel S, Gaymard A, *et al.* Type I IFN immunoprofiling in COVID-19 patients.
J Allergy Clin Immunol 2020; **146**: 206-208.
4.   Zhang Q, Bastard P, Liu Z, *et al.* Inborn errors of type I IFN immunity in patients with life-
threatening COVID-19. *Science* 2020; **370**: eabd4570.
5.   Bastard P, Michailidis E, Hoffmann HH, *et al.* Auto-antibodies to type I IFNs can underlie
adverse reactions to yellow fever live attenuated vaccine. *J Exp Med* 2021; **218**: e20202486.
6.   Gupta S, Nakabo S, Chu J, Hasni S, Kaplan M. Association between anti-interferon-*α*
autoantibodies and COVID-19 in systemic lupus erythematosus. *medrxiv* 2020; doi:
10.1101/2020.10.29.20222000
7.   Pan H, Peto R, Henao-Restrepo AM, *et al.* Repurposed Antiviral Drugs for Covid-19 - Interim
WHO Solidarity Trial Results. *N Engl J Med* 2021; **384**: 497-511.
8.   Alavi Darazam I, Shokouhi S, Pourhoseingholi MA, *et al.* Role of interferon therapy in severe
COVID-19: the COVIFERON randomized controlled trial. *Sci Rep* 2021; **11**: 8059.
9.   Trouillet-Assant S, Albert Vega C, Bal A, *et al.* Assessment of serological techniques for
screening patients for COVID-19 (COVID-SER): a prospective, multicentric study. *BMJ Open* 2020; **10**:
e041268.
10.   Pescarmona R, Belot A, Villard M, *et al.* Comparison of RT-qPCR and Nanostring in the
measurement of blood interferon response for the diagnosis of type I interferonopathies. *Cytokine*
2018; **113**: 446-452.
224

Figure 1- Anti-Type I IFN antibodies (Abs) in patients with life-threatening COVID-19

(a) Concentration of auto-Abs against IFN-α_2 (ng.mL^{-1}) determined by a Thermofisher kit (Catalog

BMS217) in serum samples collected from COVID-19 patients admitted in ICU (n=84) and COVID-19

patients with mild respiratory symptoms (n=10). Auto-Abs concentrations with neutralizing capacity

against IFN-α. (b-c) IFN-α_2 concentration (fg.mL^{-1}) (b) and ISG score (c) in plasma and whole blood

collected from COVID-19 patients in the first 15 days after symptom onset (critically-ill COVID-19

patients (n=54) and mildly-symptomatic COVID-19 patients (n=10)). (d) Longitudinal detection of

auto-Abs against IFN-α_2 in COVID-19 patients' serum during their ICU stay according to the delay

post-symptom. Dotted lines represent positive cut-off value (threshold), lower limit of quantification

(LLOQ), and upper limit of quantification (ULOQ). Solid black lines represent median. Comparisons

were performed using the Kruskal Wallis test followed by Dunn's test. ***P-value ≤ 0.001; **** P-

value ≤ 0.0001.

Figure 2 - Auto-Abs against other subtypes of IFN-I.

The presence of auto-Abs against other subtypes of IFN-I was assessed by ELISA in all sera with anti-

IFN-α_2 auto-Abs (n=21, from left to right, increasing order of concentration of anti-IFN-α_2). APS-1

patient's serum was used as positive control and sera from 2 healthy controls were used as negative

controls.

Clinical features	Auto-nAb Negative (n=69)	Auto-nAb Positive (n=15)	P-value
Age (years)	67 [58-72]	65 [55-74]	0.75
Male sex	54 (78%)	13 (87%)	0.72
BMI (kg/m^2)	29 [26-34]	29 [25-32]	0.49
Auto-immune disease	7 (10%)	2 (13%)	0.66
Time between 1st symptoms and ICU admission (days)	9 [7-12]	10 [7-11]	0.80
Maximal ventilatory support			0.33
• Standard oxygen only	4 (6%)	0 (0%)	
• High flow oxygen only	19 (28%)	7 (47%)	
• Invasive mechanical ventilation	46 (67%)	8 (53%)	
ARDS criteria	44 (64%)	8 (53%)	0.65
Worst PaO2/FiO2 day 1 in ICU (mmHg)	75 [62-103]	89 [62-116]	0.40
ECMO	15 (22%)	2 (13%)	0.72
SOFA Day 1 in ICU	4 [3-8]	3 [2-5]	0.11
SAPS2 Day 1 in ICU	40 [31-47]	43 [38-48]	0.41
Vasopressor requirement in ICU	45 (65%)	9 (60%)	0.77
Renal replacement therapy in ICU	27 (39%)	4 (27%)	0.56
ICU length of stay (days)	13 [7-35]	13 [6-24]	0.74
ICU mortality	29 (42%)	5 (33%)	0.58

247 **Table 1 - Clinical characteristics of critically-ill COVID-19 patients admitted in intensive care unit**

248 Data are expressed as median [IQR] or count (percentage). Mann-Whitney and Fisher tests were used for quantitative and qualitative variables, respectively.
249 ICU - Intensive Care Unit, BMI - Body Mass Index, nAb – neutralizing Auto-Abs against IFN-α, ECMO- Extracorporeal membrane oxygenation, SOFA -
250 sequential organ failure assessment score, SAPS2 - Simplified Acute Physiology Score II, ARDS - Acute respiratory distress syndrome