Cohort profile: The Australian Parkinson's Genetics Study (APGS)

Svetlana Bivol[1], George D. Mellick[2], Jacob Gratten[3], Richard Parker[1], Aoibhe Mulcahy[1,4], Philip Mosley[1,5], Peter C. Poortvliet[2], Adrian I. Campos[1,6], Brittany L. Mitchell[1,4], Luis M. Garcia-Marin[1,6], Simone Cross[1], Mary Ferguson[1], Penelope A. Lind[1,4,6], Danuta Z. Loesch[7], Sarah E. Medland[1,8], Peter M. Visscher[9], Clemens R. Scherzer[10,11,12,13], Nicholas G. Martin[1] and Miguel E. Rentería[1,4,6,10,14,□]

1. QIMR Berghofer Medical Research Institute, Brisbane QLD Australia
2. Griffith Institute for Drug Discovery (GRIDD), Griffith University, Brisbane, Australia
3. Mater Research, Translational Research Institute, Brisbane, Australia
4. School of Biomedical Sciences, Faculty of Health, Queensland University of Technology, Brisbane, QLD, Australia
5. Queensland Brain Institute, The University of Queensland, Brisbane, Australia
6. School of Biomedical Sciences, Faculty of Medicine, The University of Queensland, Brisbane QLD Australia
7. School of Psychology and Public Health, La Trobe University, Melbourne, VIC, Australia
8. School of Psychology, University of Queensland, Brisbane, Australia
9. Institute for Molecular Bioscience, The University of Queensland, Brisbane, Australia
10. Center for Advanced Parkinson Research, Harvard Medical School and Brigham & Women's Hospital, Boston, MA, USA
11. Precision Neurology Program, Harvard Medical School and Brigham & Women's Hospital, Boston, MA, USA
12. Department of Neurology, Massachusetts General Hospital, Boston, MA, USA
13. Program in Neuroscience, Harvard Medical School, Boston, MA, USA
14. Global Brain Health Institute and Memory and Aging Center, University of California, San Francisco, San Francisco, CA, USA
□Correspondence: Dr Miguel E. Rentería (miguel.renteria@qimrberghofer.edu.au)

Abstract

Purpose: Parkinson's disease (PD) is a neurodegenerative disorder associated with progressive disability. While the precise aetiology is unknown, there is evidence of significant genetic and environmental influences on individual risk. The Australian Parkinson's Genetics Study (APGS) seeks to study patient-reported data from a large cohort of individuals with PD in Australia to understand the sociodemographic, genetic, and environmental basis of PD susceptibility, symptoms and progression.

Participants: In the pilot phase reported here, 1,819 participants were recruited through assisted mailouts facilitated by *Services Australia* (formerly known as the Australian Government Department of Human Services), based on having three or more prescriptions for anti-PD medications in their Pharmaceutical Benefits Scheme (PBS) records. The average age at the time of the questionnaire was 64 ± 6 years. We collected patient-reported PD information and socio-demographic variables via an online (93% of the cohort) or paper-based (7%) questionnaire. 1,532 participants (84.2%) had a current PD diagnosis and consented to provide a DNA sample via a saliva sampling kit sent by traditional post. Of these, 1435 (94%) returned the saliva samples for genotyping.

Findings to date: 65% of participants were male, and 92% identified as being of European descent. A previous traumatic brain injury was reported by 16% of participants and was associated with a younger age of symptom onset. At the time of the questionnaire, constipation (36% of participants), depression (34%), anxiety (17%), melanoma (16%) and diabetes (10%) were the most commonly reported comorbid conditions.

Future plans: We are actively seeking funding to expand the project's scope and reach, including recruiting unaffected controls, and a follow-up questionnaire focused on non-motor symptoms and cognitive function assessment. Future work will explore the role of genetic and environmental factors in the aetiology of PD susceptibility, onset, symptoms and progression, including as part of international PD research consortia.

Keywords: Parkinson's disease, Cohort study, Australia, Genetics, Neurology.
Word count: 4,568

Article summary

Strengths and limitations of this study

- We showed that recruiting Australian participants for PD research through the PBS database is feasible and a highly efficient method and enables us to reach people all over Australia who have been prescribed medications used to treat Parkinson's symptoms.
- The Australian Parkinson's Genetics Study (APGS) is the most extensive nation-wide genetic study of Parkinson's disease (PD) in Australia (n = 1,532 in its pilot phase). It will serve as a platform for advancing PD research in the country.
- As part of this pilot, we collected patient-reported outcomes and variables relevant to disease onset, diagnosis, symptoms, medical comorbidities and family history. Information on environmental exposures, lifestyle factors, ethnicity and socio-demographic variables were also included.
- A general limitation is that traditional motor-related PD symptoms can most reliably be assessed during a clinical visit. Also, patient-reported data may be subject to a degree of attribution or recall bias for items concerning the age of symptom onset or the environmental and occupational exposures.

Background

Parkinson's disease (PD) is a progressive neurodegenerative disorder that affects 2-3% of the global population over 65 years [1]. At present, over 100,000 Australians have PD, and approximately 32 new cases are diagnosed every day [2]. The economic cost of Parkinson's disease in Australia was estimated at almost $10 billion per year in 2015, but the prevalence and total financial cost associated with the disease have increased dramatically since then [3]. The number of newly diagnosed PD cases is expected to double over the coming decades due to population ageing, posing substantial challenges to the healthcare system, the economy, and society [4].

The pathological hallmarks of PD include nigrostriatal degeneration and widespread intracellular accumulation of Lewy bodies in multiple brain regions. The aetiological pathways of PD are mostly unknown. A vast range of risk factors have been explored, including certain drugs, head trauma, lifestyle factors, comorbid diseases, and exposure to toxic environmental agents [5–7]. In particular, age, mood disorders, consumption of well water [8], head injury [9], and exposure to environmental toxins [10,11] have been consistently associated with an increased PD risk. On the contrary, smoking and caffeine have been inversely related to PD risk and positively correlated with a later age of onset [12,13], suggesting a putative protective effect that needs further exploration. Associations with alcohol remain controversial [14].

The clinical manifestations of PD are highly heterogeneous. Diagnosis is based on the presence of cardinal motor symptoms, including resting tremor, bradykinesia, muscle rigidity, gait difficulty, and postural instability [1]. Non-motor symptoms are also common, particularly in late-stage PD, and involve cognitive impairment (e.g., executive dysfunction and attentional deficits), psychotic symptoms (e.g., visual hallucinations), autonomic dysfunction (e.g., orthostatic hypotension, urogenital incontinence, and constipation), disorders of sleep and mood, hyposmia, fatigue and chronic pain [15]. Notably, some comorbid medical conditions such as constipation, hyposmia, REM sleep behaviour disorder, diabetes and mood disorders may precede the onset of motor disability by years or even decades and serve as pre-diagnostic PD symptoms [16]. Thus, it is worth noting that PD onset, clinical presentation, and progression differ substantially among patients [17].

PD has a complex genetic architecture, with both monogenic and sporadic forms. A twin study in the Swedish Twin Registry estimated heritability at 34%, with concordance rates for PD of 11% for monozygotic twins and 4% for same-sexed dizygotic twin pairs [18]. Rare coding variants detected in the *LRRK2*, *VPS12C*, *GBA* and *SNCA* genes are associated with monogenic (familial) PD forms [19], whereas common variants around the same *GBA*, *SNCA* and *LRRK2* genes have also been

associated with sporadic (non-familial) PD [20]. A recent GWAS meta-analysis of 17 cohorts identified 90 independent single-nucleotide polymorphisms (SNPs) across 78 genomic regions associated with sporadic Parkinson's risk [21]. Despite this remarkable progress, common genetic variants only explain around 22% (95% CI 18% - 26%) of PD risk on the liability scale, implying that there are still many more unidentified genetic variants contributing to PD susceptibility. It has been pointed out that the inclusion of diverse populations in genetic studies improves the utility of genetic discovery [22]. However, until now, genetic studies of PD have focused mainly on European ancestry individuals, limiting their capacity to extrapolate results to other ethnicities. Australia is an ethnically diverse country, but the disease epidemiology among various ethnic groups has not been investigated [23,24].

Treatment response in PD is complex and poorly understood [25]. Although there is currently no cure for PD, a combination of pharmacological (e.g., levodopa, dopamine receptor agonists, anticholinergics) and non-pharmacological (e.g., physiotherapy, deep brain stimulation) therapies are the only available treatment options. However, such interventions only ameliorate the symptoms of the illness, and their long term use is associated with adverse side effects in some patients [26,27], which in turn result in suboptimal medication compliance and low quality of life. Research seeking to identify the disease's genetic components that contribute to drug response variability could lead to the discovery of personalised disease-modifying therapies with great potential to improve PD patients' well-being and quality of life.

Few studies have assessed the prevalence, risk factors, sociodemographics and genetics of PD in Australia. Similarly, details about disease onset, symptoms and progression and other medical comorbidities have not been collected and studied at a large-scale in a nation-wide sample. Long-running studies, such as the Griffith University Queensland Parkinson's Project, have made significant contributions over the past 20 years in describing risk factors and mechanisms underlying PD's pathological development and its disease progression [28–30]. Despite continued global scientific efforts, many questions remain around early and accurate diagnosis, differentiation, prognostication, and better traditional management methods and new treatment efforts such as neuroprotection and disease modification. With an aging population, the number of people with PD in Australia is set to increase dramatically. Thus, it is crucial to generate reliable evidence about the epidemiology and genetic aetiology of PD to inform policy and clinical practice. Ultimately, a more extensive representation of diverse Australian participants in worldwide PD studies may also lead to discovering novel therapeutic targets and developing new therapies and interventions to prevent, stop, or modify PD's clinical course in Australia and the rest of the world.

Hypothesis: We hypothesise that individual differences in PD susceptibility, symptomatology, progression and treatment response result, in part, from underlying individual genetic variability. Thus, large longitudinal cohorts of individuals with PD will be required to unravel the complex aetiology of PD heterogeneity.

Aim of the study: The Australian Parkinson's Genetics Study aims to recruit and follow-up a large cohort of Australians with PD (N=10,000) and characterise their phenotypic, genetic and environmental diversity. Here we describe the results of a pilot study comprising the first ~1,500 participants.

Objectives

1) To recruit and follow up a cohort of 10,000 Australians with PD longitudinally and collect and genotype their DNA samples.

2) To collect patient-reported data on disease onset, symptoms, comorbidities and progression; treatment response; environmental, lifestyle and sociodemographic factors; family and medical history; and cognitive function.

3) Discover and validate genetic and environmental markers of PD susceptibility and heterogeneity and contribute to identifying and validating therapeutic molecular targets.

4) To contribute to ongoing international collaborative PD research efforts.

Methodology

Here we describe the design of the Australian Parkinson's Genetics study and baseline characteristics of the first wave of participants, including data collection methods, sociodemographic variables, environmental and lifestyle exposures, and self-reported history of medical conditions. Participants in the present study were Australian residents diagnosed with PD. The pilot project was initially conceived by researchers at the QIMR Berghofer Medical Research Institute (QIMRB) and approved by QIMRB's Human Research Ethics Committee, with all participants providing informed consent for participation in the study and DNA genotyping.

Recruitment method

Participants were invited to participate in the study via assisted mailouts through *Services Australia* (formerly known as the Australian Government Department of Human Services) based on their pharmaceutical prescription history. *Services Australia* is an official government agency responsible for delivering a range of welfare, health, child support payments and other services to eligible

Australian citizens and permanent residents, including managing the Pharmaceutical Benefits Scheme (PBS), Australia's national medication subsidy program. *Services Australia* retains records for the most recent five years' PBS-subsidized medicines and considers requests for assisted mailouts from external organisations for health-related projects. External organisations must have approval from a Human Research Ethics Committee, as per the National Health and Medical Research Council's guidelines. *Services Australia* then sends the approach letters to individuals meeting specific selection criteria on behalf of the research organisation. Personal data of letters recipients is not shared with the researchers requesting the mailout and thus remains secure and confidential at all times.

In this pilot phase, 20,000 invitation letters were sent in June 2020 to individuals who: (1) were Australian residents (Australian citizens or permanent residents), aged 40-75 years; (2) had at least three PBS claims in the last two years for medications commonly used in the management of symptoms of Parkinson's disease (see **Table 1** for a list of medications and their corresponding PBS codes) according to their records in the PBS database. The invitation letters directed potential participants to a secure study website (https://www.qimrberghofer.edu.au/parkinsonsgenetics) with more detailed information about the study. Participants who provided informed consent completed an online or paper-based questionnaire. Participants indicated having a current clinical diagnosis for PD and the type of clinician who initially diagnosed them (e.g., neurologist, geriatrician, general practitioner). Prospective participants were also asked to indicate their consent to donate a saliva sample for DNA extraction and genotyping. Those willing to provide a saliva sample were sent a saliva DNA collection kit via mail. A small percentage of prospective participants (~ 10%) requested to be mailed a paper-based version of the questionnaire and consent form. Of the 1,819 individuals who filled out the registration form and provided informed consent to participate (9.1% response rate for the mailouts), 1,532 (84%) confirmed having a current PD diagnosis and were willing to provide a DNA sample. Of those who consented to donate their saliva for genetic analysis, 1,435 (94%) have returned saliva kits by mail as of February 2021. Since information about the letter recipients remained confidential, we could not follow up with the 18,181 nonrespondents, some of whom may have died or moved to a new address. Interestingly, the 9% response rate in this assisted mailout is considerably higher than for similar studies targeted at patients with depression, bipolar disorder, alcohol dependence and Alzheimer's disease, which have attained an average of 5% response rate [31]. Refer to **Table 2** for the percentage of APGS invitations sent and participant registration/response rates per state or territory of Australia.

Data collection

The online instruments consisted of several sections, covering sociodemographic, clinical, occupational, lifestyle and environmental factors, with a schematic of the APGS and questionnaire content shown in **Figure 1**. As part of the informed consent process, all participants agreed to be recontacted for future related studies, understanding that their participation in future studies is voluntary. The second wave of data collection is planned for the first half of 2021. It will include questions assessing heterogeneity in PD onset and motor and non-motor disability, including the following instruments: ~80% of items in the Movement Disorder Society United Parkinson's Disease Rating Scale (MDS-UPDRS) [32]; a short scale for the assessment of motor impairments and disability in Parkinson's disease (The SPES/SCOPA); the Apathy Screening Tool; Geriatric Anxiety Inventory [32,33]; Geriatric Depression Scale (GDS); Assessment of Psychiatric Complications in Parkinson's disease (The SCOPA-PC) [34]; and the SCOPA-sleep and PDSS scales [35] to identify sleep disorders in PD.

Results

We recruited 1,532 participants (average age = 64 years; SD = 6 years; range 40 – 88 years) for the Australian Genetics of Parkinson's Disease Study carried out at QIMR Berghofer. Sociodemographic data were collected via web-based or paper questionnaires, with most participants (93%; N = 1,424) responding via the online questionnaire. Of our sample, 65% (N = 995) were males, with the sample mean self-reported BMI of 28 kg/m2 (SD = 5 kg/m2; range 17 – 55 kg/m2).

Most participants report being diagnosed with PD by a neurologist (83%; N = 1,260), having children (87%; N = 1,338), being married or in a de facto relationship (81%; N = 1,245) and are of European ancestry (92%; N = 1,397). Nearly 60% (N = 903) have a post-high school qualification and 15% (N = 228) speak more than one language. Tremor was the most common first symptom of PD reported in 788 respondents (54% of the sample group). The next most commonly reported symptoms were bradykinesia and muscle rigidity (20%, N = 291), followed by gait disorder (10%; N = 151), with an average age at onset of the first PD symptom of 57 years (SD = 8 years; range 27 – 86 years). The frequency of self-reported first symptom of PD in participants is presented in **Figure 2**.

For 1,296 participants (85%), PD symptoms started on one side of the body, and more than half (44%) experienced the symptoms on the right side of the body. A large number of respondents (94%; N = 1,424) were currently on an anti-PD medication, 42% (N = 631) have experienced PD-related falls, and 12% (N = 177) reported having had a brain surgery for PD. The frequencies and percentage of all self-reported social, demographic and PD-related characteristics are outlined in **Table 3**.

Occupational and environmental risk factors of PD

Previously reported occupational (e.g., welding, metal melting, agriculture, petrochemistry) and environmental risk factors associated with PD were examined in this cohort and included exposure to pesticides or herbicides, head injury, well water drinking, working in high-risk occupations, living on a farm for over a year, and having metal poisoning or brain infection at any point across the life course. Approximately one-third of the sample (34%) reported being exposed to pesticides or herbicides at some point in their lives. Of those exposed to pesticides/herbicides, 390 were male, and 127 were female, with most of these cases (82%) reporting pesticide/herbicide exposure before 37 years of age.

Two hundred and forty-four participants (16% of the sample) had experienced head trauma, and the average age of head injury was 26 years (SD = 18 years; range 2 – 82 years). Of those with a previous head injury, 165 (67%) reported traumatic head injury with loss of consciousness. There was a significant positive association between the age of traumatic head injury and the age at onset of the first symptom of PD (Pearson correlation coefficient r = 0.3, $p < 0.001$) (**Figure 3**).

Eighty-two participants (5%) reported having frequently consumed water from a well for 19 years on average (SD = 12 years; range 2 – 50 years). A third of the sample (33%; N = 504) disclosed having worked in a high-risk occupation throughout their life, including welding, metal melting, galvanisation, milling, petrochemistry, agriculture, and wood processing, textile or industrial painting. Of these participants, 42% of respondents (N = 211) reported exposure to more than one occupational risk factor. Approximately 24% of the cohort (N = 360) worked or lived on a farm for more than a year, 1% (N = 22) acquired metal poisoning and 2% (N = 31) encephalitis (brain infection). The occupational and environmental risk factors of PD, summarised by sex, are shown in **Table 4**.

Variables correlated with lower PD risk

As shown in **Table 5**, we investigated a range of variables that have been correlated in previous PD studies with lower PD risk. These included alcohol drinking, smoking, exposure to second-hand smoke, and drinking caffeinated coffee, tea, and other beverages. More than two-thirds of participants (71%) currently drink an average of 8 standard alcoholic drinks per week (SD = 8), and 20% reported consuming alcohol heavily in the past. There was a significant positive association between the number of alcoholic drinks consumed per week and the age at onset of the first PD symptom (Pearson correlation coefficient r = 0.1, $p < 0.05$).

Fifty participants (3% of the cohort) are current cigarette, cigar, or pipe smokers and smoke on average 13 cigarettes/cigars/pipes per day (SD = 8; range 2 – 40). Five hundred and fifty-four participants (36% of the sample) reported being former smokers and disclosed smoking 15 cigarettes/cigars/pipes a day in the past (SD = 11; range 1 – 60). For current smokers, there was a significant negative relationship between the number of cigarettes/cigars/pipes smoked daily and the age at onset of the first symptom of PD (Pearson correlation coefficient r = - 0.2, $p < 0.05$).

Many participants reported being exposed to second-hand smoke at home (56%; N = 858), at work (46%; N = 701) and in other areas (38%; N = 586). For the total sample, the average number of hours per day exposed to second-hand smoke at home was 4.5 hours (SD = 4 hours; range 0.5 – 20 hours). The mean for those who reported exposure to smoke at work was 5 hours (SD = 3 hours; range 0.5 – 12 hours), whereas the mean for those who were exposed to smoke in other areas was 2 hours (SD = 1 hour; range 0.5 – 14 hours).

Sixty eight percent of participants (N = 1,043) drink tea, 73% (N = 1,120) drink coffee, and 24% (N = 371) consume other caffeinated beverages. On average, participants drink 2 cups of caffeinated tea or coffee daily, with a range of half a cup to 12 cups a day. There was a significant positive correlation between the number of teacups consumed daily and the age at onset of PD symptoms (Pearson correlation coefficient r = 0.1, $p < 0.05$). Those who reported drinking other caffeinated beverages indicated that they consumed on average 330 ml of other caffeinated drinks per day (SD = 335 ml; range 10 – 2000 ml). For sex differences in the reported outcomes related to PD's potential protective factors, refer to **Table 3**.

Parkinson's disease comorbidities

Back problems were identified as the most common comorbid medical condition (722 participants) at a frequency of 47%. PD co-occurring disorders among participants included constipation (36%; N = 533), depression (34%; N = 514), high blood pressure (33%; N = 506), high cholesterol (28%; N = 430), frequent acid reflux (25%; N = 377) and cold sores (22%; N = 336). Other medical conditions prevalent in the cohort included migraine (17%; N = 264), anxiety (17%; N = 262), melanoma (16%; N = 243), sleep apnoea (16%; N = 241), insomnia (13%; N = 202), cancer (13%; N = 201), asthma or COPD (13%; N = 194), pneumonia (12%; N = 192) and REM Sleep behaviour disorder (12%; N = 177). Diabetes, osteoporosis, and eczema were common in 10% (N = 156), 8% (N = 126) and 7% (N = 107) of the respondents, respectively (**Table 6**).

Discussion

The present study describes sample characteristics of 1,532 Australians with PD recruited for the Australian Parkinson's Genetics Study to unravel environmental and genetic associations with PD, along with PD heterogeneity, anti-PD medication response and adverse drug reactions.

Parkinson's disease is a multifactorial disorder, and the estimated incidence of the disease varies with gender, age, and race or ethnicity [36,37]. We observed a higher prevalence of male participants than female (65% of males versus 35% of females), in line with most previous PD studies in various populations worldwide that found a significantly higher incidence rate of PD in males [36–38]. The causes of higher male-to-female ratios in PD are unknown, and it is still unclear whether male sex is an independent risk factor for PD. Estrogens are believed to be neuroprotective in females [39]. Nonetheless, whether males could also be at greater risk for developing PD due to higher exposure to environmental or occupational risk factors, future studies leveraging the APGS will be aimed at filling out these gaps in knowledge and pinpointing the role of sex hormones in PD.

The present study is the first to employ an assisted mailout approach to recruit PD participants from all Australian states. We found that individuals of European descent (92% of the sample group) were the most prevalent in the APGS, with a lower proportion of participants from Asian background (4%), Indigenous Australians and Torres Strait Islanders (1%), Africans (0.1%), Pacific Islanders (0.6%), Hispanics (0.3%), and in individuals of mixed ancestry (5%). We are unable to estimate differences in PD prevalence across ethnicities, given the lack of unaffected controls. Previous Australian epidemiological studies have reported PD prevalence rates between 66 and 415 per 100,000 Australians [40–42]. Similarly, worldwide PD prevalence has been consistently reported between 60 and 335 per 100,000 individuals, with a broad spectrum of ethnic backgrounds identified. For instance, the Israel population, Native Americans and Alaska Natives, have the highest prevalence (335/100 000 for Jewish Israeli and 355/100 000 for Native Americans and Alaska Natives) [43,44]. At the same time, Asian and African communities report the lowest prevalence rates [45–47]. The significantly increased prevalence of PD among distinct ethnic groups is partly attributable to a higher occurrence of PD-linked mutations in these populations (e.g., *LRRK2* G2019S and *GBA* in *Ashkenazi* Jews) [48,49]. However, other factors such as variation in age groups surveyed, sampling methods, and diagnostic and screening criteria cannot be ruled out.

Parkinson's disease is rare before the age of 50, and most patients present with symptoms in the late fifties [1]. Therefore, aging remains the primary risk factor for developing idiopathic PD [50–52], with the incidence of new cases increasing up to 10-fold between 60 and 90 years of age [36,53,54].

In our study, the mean age of onset for PD symptoms was 56 years, suggesting that self-selected PD participants may be among the younger age group, which tends to be more technology-savvy and motivated to participate in research. A possible explanation is that However, aging alone is insufficient to cause PD, and a complex relationship between aging and genetic and environmental factors prompts the disease onset [55]. Aging-associated impairments [56,57] and degeneration of dopaminergic neurons [58] may explain the steep rise in PD prevalence with age in cellular pathways.

Previous research suggests that exposure to toxic environmental agents (e.g., pesticides [10,59]), work-related iron or manganese exposure [60], farming or agricultural work [61] and traumatic head injury [62–64] are associated with increased risk of PD. However, the strength of these associations varies across studies and remains unclear. In light of these uncertainties, we investigated environmental and occupational exposures to assess their potential association with PD. Our results suggest that pesticide or herbicide exposure, working in known high-risk occupations, and working or living on a farm for over a year and suffering a head trauma are prevalent risk factors in the APGS. Interestingly, of those with traumatic brain injury, two-thirds reported having lost consciousness following head trauma.

Previous studies have shown that a traumatic brain injury with loss of consciousness early in life is associated with late-life neurodegenerative conditions, accumulation of Lewy bodies, PD progression, and increased PD risk [65]. However, the pathogenesis of brain injury and its link with neurodegenerative disorders has not been fully elucidated. When exploring the relationship between head trauma and the onset of PD, we identified a significant positive association between the age of head injury and the age at onset of PD symptoms. Participants who had a head injury early in life also reported an earlier age at which the first PD symptoms were manifested. However, only a small percentage of the cohort reported a lifetime traumatic head injury. Thus, our results are aligned with the hypothesis in which traumatic brain injury may initiate or exacerbate PD, perhaps due to persistent inflammation, metabolic dysfunction and abnormal protein aggregation [66].

Lifestyle may affect PD susceptibility. For instance, alcohol [14,67,68], smoking [69,70] and caffeine [71,72] are negatively correlated with PD risk. Previous studies have found that alcohol users have a lower PD risk than non-alcohol users, and those who drink more are less susceptible to PD than those who drink less [68]. Excessive alcohol consumption has been found to reverse this protective effect, increasing PD susceptibility independent of gender [73]. We observed a positive association between alcohol use and PD age of onset in the AGPD sample whereby participants with

PD who had more alcoholic drinks consumed per week reported a later age of onset of PD compared to those who consumed less alcoholic drinks weekly. This interim observation agrees with previous findings and may imply that alcohol has a protective effect and delays the onset of PD. However, this needs to be explored further due to the possibility of ascertainment bias (e.g., alcohol consumption has negative effects on health and increases risk of mortality risk).

An inverse association has also been reported between smoking and PD in several longitudinal studies. A lower PD risk was observed among current and past smokers compared with never smokers [69,74]. Notably, smoking duration and intensity have been inversely related to PD risk, with PD susceptibility decreasing up to 70% when the duration of smoking in years increased [69]. In contrast, we observed an earlier age of symptoms onset in those who smoked more cigarettes/cigars/pipes per day than those who smoked fewer. This should be interpreted with caution, given the small sample size. Also, a recent Mendelian randomisation study by us [75] indicated that the continuation of the smoking habit rather than smoking heaviness (i.e., numbers of cigarettes smoked per day) is what may exert the protective effect of smoking over PD risk.

Similarly, prospective studies assessing lifetime coffee and tea consumption and PD highlight a negative correlation between caffeine in coffee or tea and PD risk [76], and the effect is more pronounced among men than women who regularly drink coffee or tea [71,77,78]. Importantly, there is substantial between-study heterogeneity across reported effect estimates. Our results on the correlation between caffeinated tea consumption and age of onset of PD are similar to those previously reported by others [79], with regular tea consumers reporting a later age of symptom onset than non-habitual tea consumers. The precise mechanisms underlying this relationship between lifestyle variables and PD susceptibility are poorly understood.

While there is strong evidence for the contribution of environmental factors in PD risk, it remains unclear why some individuals are more susceptible to PD. To advance our understanding of individual variability in vulnerability to environmental and lifestyle exposures, APGS will explore gene-environment interactions to detect potential genetic modifiers of PD susceptibility.

Comorbid medical conditions in PD patients could result from the underlying neurodegenerative processes occurring in the mesolimbic pathways, locus coeruleus and raphe nuclei, or could be due to changes in the dopaminergic, noradrenergic and serotonergic systems [80–82]. These prominent non-motor symptoms significantly reduce the quality of life of PD patients and lead to poor medication adherence and deterioration in their physical and mental status. Thus, it is urgently

needed to identify disabling medical conditions co-occurring with PD to improve health outcomes and clinical management of PD patients in Australia. Consistent with previous findings we report high rates of comorbidities including constipation [83,84], depression [85,86], anxiety disorder [87], melanoma [88,89] and diabetes [90]. Although the prevalence estimates of common comorbidities in our cohort are higher than in the general population [91–94], our results support the body of evidence that these particular conditions are substantially more common among people with PD and thus may share a common aetiology.

In summary, we report the baseline characteristics of the APGS pilot cohort. The APGS aims to characterise and improve our understanding of the sociodemographic, genetic, and environmental basis of PD susceptibility, symptoms and progression in Australia. We used an innovative and highly-efficient recruitment approach through assisted mailouts, establishing feasibility and laying the groundwork for expanding and extending the study's scale and reach in the future. With a 9% response rate to our mailout approach, the results of this study will provide an unprecedented opportunity for generating robust evidence on the epidemiology and genetic aetiology of PD to inform policy and clinical practice in the country. Ultimately, a more extensive representation of diverse Australian participants in worldwide PD studies may lead to a better understanding of the causes of PD and help in the discovery and validation of novel therapeutic targets and the development of new therapies and interventions to prevent, stop or modify the clinical course of PD in Australia and the rest of the world.

Twitter: @QIMRBerghofer, @mrenteria_

Acknowledgments: We thank all of the participants for kindly giving their time to contribute to this study. We thank Steve Gilbert, Rhonda Charlesworth, and the Health Data Analysis and Strategy Branch team at Services Australia for their assistance in conducting the mailout. MER thanks the support of NHMRC and the Australian Research Council (ARC) through an NHMRC-ARC Dementia Research Development Fellowship (GNT1102821). AIC and LMGM are supported by UQ Research Training Scholarships from The University of Queensland (UQ). JG thanks the NHMRC (1127440) and Mater Foundation for support. SEM is supported by an NHMRC Investigator grant (APP1172917). The views expressed are those of the authors and not necessarily those of the affiliated or funding institutions.

Author's contributions: MER designed the APGS study with input from GDM, JG, PM, PCP, PL, DZL, SEM, NGM and CRS. AM, AIC, BLM, LMGM revised and tested the questionnaire and provided intellectual input into the content. RP coordinated the project and sample collection with

help from SC and MF. SV cleaned and analysed the data for this cohort profile manuscript. SV and MER drafted the manuscript. All coauthors revised the article for intellectual content. All authors have read and approved the final version of the manuscript.

Funding: The APGS mailout was primarily funded by National Health and Medical Research Council (NHMRC) and the Australian Research Council (ARC) through an NHMRC-ARC Dementia Research Development Fellowship award to MER (GNT1102821).

Competing interests: The authors declare no competing interests.

Patient consent for publication: Obtained.

Disclosure statement: CRS has collaborated with Pfizer, Opko, Proteome Sciences, Genzyme Inc; has consulted for Genzyme; has served as Advisor to the Michael J. Fox Foundation, NIH, Department of Defense; is on the Scientific Advisory Board of the American Parkinson Disease Association; has received funding from the NIH, the U.S. Department of Defense, the Harvard NeuroDiscovery Center, the Michael J. Fox Foundation, and American Parkinson Disease Association. XD has received funding from the NIH, the Michael J. Fox Foundation, and American Parkinson Disease Association.

Ethics approval: This study has been approved by the Human Research Ethics Committee of the QIMR Berghofer Medical Research Institute.

Data availability statement: Data is available upon reasonable request for research purposes, in anonymised form, to preserve participants' identity. Inquiries from potential scientific collaborators can be directed to the corresponding author (Miguel.Renteria@qimrberghofer.edu.au).

ORCID IDs: Svetlana Bivol (0000-0002-7085-8366) George D. Mellick (0000-0002-7211-4651), Jacob Gratten (0000-0003-1293-409X), Richard Parker (0000-0003-1451-5622), Philip Mosley (0000-0003-1721-3419), Adrian I. Campos (0000-0003-3468-8619), Brittany L. Mitchell (0000-0002-9050-1516), Luis M. García-Marín (0000-0003-4731-6558), Penelope A. Lind (0000-0002-3887-2598), Sarah E. Medland (0000-0003-1382-380X), Peter M. Visscher (0000-0002-2143-8760), Clemens R. Scherzer (0000-0002-0567-9193), Nicholas G. Martin (0000-0003-4069-8020), Miguel E. Rentería (0000-0003-4626-7248).

Tables

Tables 1 to 6 can be found here:

https://docs.google.com/document/d/1zI_EMPzNZuMnw_auXLR6sMZ9w_jRs-EwsqqNSJZKQPc/edit?usp=sharing

Figures

Figures 1 to 3 can be found here:

https://docs.google.com/document/d/1pyzSTS5omsITXqvhqvf0zl11_lFAZ2AncELUNuaIKog/edit?usp=sharing

References

Poewe W, Seppi K, Tanner CM, *et al.* Parkinson disease. *Nat Rev Dis Primers* 2017;**3**:17013.

Deloitte Access Economics. Living with Parkinson's Disease. 2016. https://www2.deloitte.com/au/en/pages/economics/articles/living-with-parkinsons-disease.html (accessed 30 Jan 2021).

Bohingamu Mudiyanselage S, Watts JJ, Abimanyi-Ochom J, *et al.* Cost of Living with Parkinson's Disease over 12 Months in Australia: A Prospective Cohort Study. *Parkinsons Dis* 2017;**2017**:5932675.

GBD 2016 Parkinson's Disease Collaborators. Global, regional, and national burden of Parkinson's disease, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2018;**17**:939–53.

Bellou V, Belbasis L, Tzoulaki I, *et al.* Environmental risk factors and Parkinson's disease: An umbrella review of meta-analyses. *Parkinsonism Relat Disord* 2016;**23**:1–9.

Menegon A, Board PG, Blackburn AC, *et al.* Parkinson's disease, pesticides, and glutathione transferase polymorphisms. *Lancet* 1998;**352**:1344–6.

Dissanayaka NNW, Sellbach A, Matheson S, *et al.* Anxiety disorders in Parkinson's disease: prevalence and risk factors. *Mov Disord* 2010;**25**:838–45.

Noyce AJ, Bestwick JP, Silveira-Moriyama L, *et al.* Meta-analysis of early nonmotor features and risk factors for Parkinson disease. *Ann Neurol* 2012;**72**:893–901.

Jafari S, Etminan M, Aminzadeh F, *et al.* Head injury and risk of Parkinson disease: a systematic review and meta-analysis. *Mov Disord* 2013;**28**:1222–9.

Pezzoli G, Cereda E. Exposure to pesticides or solvents and risk of Parkinson disease. *Neurology* 2013;**80**:2035–41.

van der Mark M, Brouwer M, Kromhout H, *et al.* Is pesticide use related to Parkinson disease? Some clues to heterogeneity in study results. *Environ Health Perspect* 2012;**120**:340–7.

Domínguez-Baleón C, Ong JS, Scherzer CR. Genetic evidence for protective effects of smoking and drinking behavior on Parkinson's disease: A Mendelian Randomization study. *medRxiv* Published Online First: 2020.https://www.medrxiv.org/content/10.1101/2020.04.20.20073247v1.abstract

Checkoway H, Nielsen SS, Racette BA. The search for environmental risk factors for Parkinson disease. Current Topics in Occupational Epidemiology. 2013;:31–41. doi:10.1093/med/9780199683901.003.0003

Bettiol SS, Rose TC, Hughes CJ, *et al.* Alcohol Consumption and Parkinson's Disease Risk: A Review of Recent Findings. *J Parkinsons Dis* 2015;**5**:425–42.

Przedborski S. The two-century journey of Parkinson disease research. *Nat Rev Neurosci* 2017;**18**:251–9.

Santiago JA, Bottero V, Potashkin JA. Biological and Clinical Implications of Comorbidities in Parkinson's Disease. Frontiers in Aging Neuroscience. 2017;**9**. doi:10.3389/fnagi.2017.00394

Poortvliet PC, O'Maley K, Silburn PA, *et al.* Perspective: Current Pitfalls in the Search for Future Treatments and Prevention of Parkinson's Disease. *Front Neurol* 2020;**11**:686.

Wirdefeldt K, Gatz M, Reynolds CA, *et al.* Heritability of Parkinson disease in Swedish twins: a longitudinal study. *Neurobiol Aging* 2011;**32**:1923.e1–8.

Hernandez DG, Reed X, Singleton AB. Genetics in Parkinson disease: Mendelian versus non-Mendelian inheritance. *J Neurochem* 2016;**139 Suppl 1**:59–74.

Billingsley KJ, Bandres-Ciga S, Saez-Atienzar S, *et al.* Genetic risk factors in Parkinson's disease. *Cell Tissue Res* 2018;**373**:9–20.

Nalls MA, Blauwendraat C, Vallerga CL, *et al.* Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 2019;**18**:1091–102.

Hindorff LA, Bonham VL, Brody LC, *et al.* Prioritizing diversity in human genomics research. *Nat Rev Genet* 2018;**19**:175–85.

Mehta P, Kifley A, Wang JJ, *et al.* Population prevalence and incidence of Parkinson's disease in an Australian community. *Intern Med J* 2007;**37**:812–4.

Peters CM, Gartner CE, Silburn PA, *et al.* Prevalence of Parkinson's disease in metropolitan and rural Queensland: a general practice survey. *J Clin Neurosci* 2006;**13**:343–8.

Politi C, Ciccacci C, Novelli G, *et al.* Genetics and Treatment Response in Parkinson's Disease: An Update on Pharmacogenetic Studies. *Neuromolecular Med* 2018;**20**:1–17.

Li B-D, Bi Z-Y, Liu J-F, *et al.* Adverse effects produced by different drugs used in the treatment of Parkinson's disease: A mixed treatment comparison. *CNS Neurosci Ther* 2017;**23**:827–42.

Deng H, Wang P, Jankovic J. The genetics of Parkinson disease. *Ageing Res Rev* 2018;**42**:72–85.

Gao Y, Wilson GR, Salce N, *et al.* Genetic Analysis of in an Early-Onset Parkinson's Disease Cohort. *Front Neurol* 2020;**11**:523.

Bentley SR, Bortnick S, Guella I, *et al.* Pipeline to gene discovery - Analysing familial Parkinsonism in the Queensland Parkinson's Project. *Parkinsonism Relat Disord* 2018;**49**:34–41.

Poortvliet PC, Gluch A, Silburn PA, *et al.* The Queensland Parkinson's Project: An Overview of 20 Years of Mortality from Parkinson's Disease. *J Mov Disord* 2021;**14**:34–41.

Byrne EM, Kirk KM, Medland SE, *et al.* Cohort profile: the Australian genetics of depression study. *BMJ Open* 2020;**10**:e032580.

Goetz CG, Fahn S, Martinez-Martin P, *et al.* Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Process, format, and clinimetric testing plan. *Mov Disord* 2007;**22**:41–7.

Pachana NA, Byrne GJ, Siddle H, *et al.* Development and validation of the Geriatric Anxiety Inventory. *Int Psychogeriatr* 2007;**19**:103–14.

Visser M, Verbaan D, van Rooden SM, *et al.* Assessment of psychiatric complications in Parkinson's disease: The SCOPA-PC. *Mov Disord* 2007;**22**:2221–8.

Martinez-Martin P, Visser M, Rodriguez-Blazquez C, *et al.* SCOPA-sleep and PDSS: two scales for assessment of sleep disorder in Parkinson's disease. *Mov Disord* 2008;**23**:1681–8.

Van Den Eeden SK, Tanner CM, Bernstein AL, *et al.* Incidence of Parkinson's disease: variation by age, gender, and race/ethnicity. *Am J Epidemiol* 2003;**157**:1015–22.

Baldereschi M, Di Carlo A, Rocca WA, *et al.* Parkinson's disease and parkinsonism in a longitudinal study: two-fold higher incidence in men. ILSA Working Group. Italian Longitudinal Study on Aging. *Neurology* 2000;**55**:1358–63.

Solla P, Cannas A, Ibba FC, *et al.* Gender differences in motor and non-motor symptoms among Sardinian patients with Parkinson's disease. *J Neurol Sci* 2012;**323**:33–9.

Cereda E, Barichella M, Cassani E, *et al.* Reproductive factors and clinical features of Parkinson's disease. *Parkinsonism Relat Disord* 2013;**19**:1094–9.

Jenkins AC. Epidemiology of parkinsonism in Victoria. *Med J Aust* 1966;**2**:496–502.

McCann SJ, LeCouteur DG, Green AC, *et al.* The epidemiology of Parkinson's disease in an Australian population. *Neuroepidemiology* 1998;**17**:310–7.

Chan DK, Dunne M, Wong A, *et al.* Pilot study of prevalence of Parkinson's disease in Australia. *Neuroepidemiology* 2001;**20**:112–7.

Chillag-Talmor O, Giladi N, Linn S, *et al.* Use of a refined drug tracer algorithm to estimate prevalence and incidence of Parkinson's disease in a large israeli population. *J Parkinsons Dis* 2011;**1**:35–47.

Gordon PH, Mehal JM, Holman RC, *et al.* Parkinson's disease among American Indians and Alaska natives: a nationwide prevalence study. *Mov Disord* 2012;**27**:1456–9.

Bower JH. Understanding Parkinson disease in sub Saharan Africa: A call to action for the international neurologic community. Parkinsonism Relat. Disord. 2017;**41**:1–2.

McInerney-Leo A, Gwinn-Hardy K, Nussbaum RL. Prevalence of Parkinson's disease in

populations of African ancestry: a review. *J Natl Med Assoc* 2004;**96**:974–9.

Wang SJ, Fuh JL, Teng EL, *et al.* A door-to-door survey of Parkinson's disease in a Chinese population in Kinmen. *Arch Neurol* 1996;**53**:66–71.

Orr-Urtreger A, Shifrin C, Rozovski U, *et al.* The LRRK2 G2019S mutation in Ashkenazi Jews with Parkinson disease: is there a gender effect? *Neurology* 2007;**69**:1595–602.

Sidransky E, Nalls MA, Aasly JO, *et al.* Multicenter analysis of glucocerebrosidase mutations in Parkinson's disease. *N Engl J Med* 2009;**361**:1651–61.

Bennett DA, Beckett LA, Murray AM, *et al.* Prevalence of parkinsonian signs and associated mortality in a community population of older people. *N Engl J Med* 1996;**334**:71–6.

Morens DM, Davis JW, Grandinetti A, *et al.* Epidemiologic observations on Parkinson's disease: incidence and mortality in a prospective study of middle-aged men. *Neurology* 1996;**46**:1044–50.

Tanner CM, Goldman SM. Epidemiology of Parkinson's disease. *Neurol Clin* 1996;**14**:317–35.

Twelves D, Perkins KSM, Counsell C. Systematic review of incidence studies of Parkinson's disease. *Mov Disord* 2003;**18**:19–31.

Savica R, Grossardt BR, Bower JH, *et al.* Incidence and pathology of synucleinopathies and tauopathies related to parkinsonism. *JAMA Neurol* 2013;**70**:859–66.

Pang SY-Y, Ho PW-L, Liu H-F, *et al.* The interplay of aging, genetics and environmental factors in the pathogenesis of Parkinson's disease. *Transl Neurodegener* 2019;**8**:23.

Kaushik S, Cuervo AM. Proteostasis and aging. *Nat Med* 2015;**21**:1406–15.

Collier TJ, Lipton J, Daley BF, *et al.* Aging-related changes in the nigrostriatal dopamine system and the response to MPTP in nonhuman primates: diminished compensatory mechanisms as a prelude to parkinsonism. *Neurobiol Dis* 2007;**26**:56–65.

Collier TJ, Kanaan NM, Kordower JH. Aging and Parkinson's disease: Different sides of the same coin? *Mov Disord* 2017;**32**:983–90.

Baldi I, Lebailly P, Mohammed-Brahim B, *et al.* Neurodegenerative diseases and exposure to pesticides in the elderly. *Am J Epidemiol* 2003;**157**:409–14.

Nandipati S, Litvan I. Environmental Exposures and Parkinson's Disease. *Int J Environ Res Public Health* 2016;**13**. doi:10.3390/ijerph13090881

Tüchsen F, Jensen AA. Agricultural work and the risk of Parkinson's disease in Denmark, 1981-1993. *Scand J Work Environ Health* 2000;**26**:359–62.

Goldman SM, Kamel F, Ross GW, *et al.* Head injury, α-synuclein Rep1, and Parkinson's disease. *Ann Neurol* 2012;**71**:40–8.

Goldman SM, Tanner CM, Oakes D, *et al.* Head injury and Parkinson's disease risk in twins. *Ann Neurol* 2006;**60**:65–72.

Lee P-C, Bordelon Y, Bronstein J, *et al.* Traumatic brain injury, paraquat exposure, and their relationship to Parkinson disease. *Neurology* 2012;**79**:2061–6.

Crane PK, Gibbons LE, Dams-O'Connor K, *et al.* Association of Traumatic Brain Injury With Late-Life Neurodegenerative Conditions and Neuropathologic Findings. *JAMA Neurol* 2016;**73**:1062–9.

Delic V, Beck KD, Pang KCH, *et al.* Biological links between traumatic brain injury and Parkinson's disease. *Acta Neuropathol Commun* 2020;**8**:45.

Eriksson A-K, Löfving S, Callaghan RC, *et al.* Alcohol use disorders and risk of Parkinson's disease: findings from a Swedish national cohort study 1972-2008. *BMC Neurol* 2013;**13**:190.

Zhang D, Jiang H, Xie J. Alcohol intake and risk of Parkinson's disease: a meta-analysis of observational studies. *Mov Disord* 2014;**29**:819–22.

Thacker EL, O'Reilly EJ, Weisskopf MG, *et al.* Temporal relationship between cigarette smoking and risk of Parkinson disease. *Neurology* 2007;**68**:764–8.

Gallo V, Vineis P, Cancellieri M, *et al.* Exploring causality of the association between smoking and Parkinson's disease. *Int J Epidemiol* 2019;**48**:912–25.

Ross GW, Abbott RD, Petrovitch H, *et al.* Association of coffee and caffeine intake with the risk of Parkinson disease. *JAMA* 2000;**283**:2674–9.

Costa J, Lunet N, Santos C, *et al.* Caffeine exposure and the risk of Parkinson's disease: a systematic review and meta-analysis of observational studies. *J Alzheimers Dis* 2010;**20 Suppl 1**:S221–38.

Eriksson A, Löfving S, Callaghan RC, *et al.* Alcohol use disorders and risk of Parkinson's disease: findings from a Swedish national cohort study 1972-2008. *Eur J Public Health* 2013;**23**. doi:10.1093/eurpub/ckt126.181

Hernán MA, Zhang SM, Rueda-deCastro AM, *et al.* Cigarette smoking and the incidence of Parkinson's disease in two prospective studies. *Ann Neurol* 2001;**50**:780–6.

Domínguez-Baleón C, Ong J-S, Scherzer CR, *et al.* Genetic evidence for protective effects of smoking and drinking behavior on Parkinson's disease: A Mendelian Randomization study. *medRxiv* 2020;:2020.04.20.20073247.

Hu G, Bidel S, Jousilahti P, *et al.* Coffee and tea consumption and the risk of Parkinson's disease. *Mov Disord* 2007;**22**:2242–8.

Ascherio A, Zhang SM, Hernán MA, *et al.* Prospective study of caffeine consumption and risk of Parkinson's disease in men and women. *Ann Neurol* 2001;**50**:56–63.

Ascherio A, Weisskopf MG, O'Reilly EJ, *et al.* Coffee consumption, gender, and Parkinson's disease mortality in the cancer prevention study II cohort: the modifying effects of estrogen. *Am J Epidemiol* 2004;**160**:977–84.

Kandinov B, Giladi N, Korczyn AD. Smoking and tea consumption delay onset of Parkinson's disease. *Parkinsonism Relat Disord* 2009;**15**:41–6.

Braak H, Del Tredici K, Rüb U, *et al.* Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 2003;**24**:197–211.

Prakash KG, Bannur BM, Chavan MD, *et al.* Neuroanatomical changes in Parkinson's disease in relation to cognition: An update. *J Adv Pharm Technol Res* 2016;**7**:123–6.

Fontoura JL, Baptista C, Pedroso F de B, *et al.* Depression in Parkinson's Disease: The Contribution from Animal Studies. *Parkinsons Dis* 2017;**2017**:9124160.

Cheon S-M, Ha M-S, Park MJ, *et al.* Nonmotor symptoms of Parkinson's disease: prevalence and awareness of patients and families. *Parkinsonism Relat Disord* 2008;**14**:286–90.

Saleem TZ, Higginson IJ, Chaudhuri KR, *et al.* Symptom prevalence, severity and palliative care needs assessment using the Palliative Outcome Scale: a cross-sectional study of patients with Parkinson's disease and related neurological conditions. *Palliat Med* 2013;**27**:722–31.

Aarsland D, Påhlhagen S, Ballard CG, *et al.* Depression in Parkinson disease--epidemiology, mechanisms and management. *Nat Rev Neurol* 2011;**8**:35–47.

Schrag A, Barone P, Brown RG, *et al.* Depression rating scales in Parkinson's disease: critique and recommendations. *Mov Disord* 2007;**22**:1077–92.

Richard IH. Anxiety disorders in Parkinson's disease. *Adv Neurol* 2005;**96**:42–55.

Dalvin LA, Damento GM, Yawn BP, *et al.* Parkinson Disease and Melanoma: Confirming and Reexamining an Association. *Mayo Clin Proc* 2017;**92**:1070–9.

Ferreira JJ, Neutel D, Mestre T, *et al.* Skin cancer and Parkinson's disease. *Mov Disord* 2010;**25**:139–48.

Santiago JA, Potashkin JA. Shared dysregulated pathways lead to Parkinson's disease and diabetes. *Trends Mol Med* 2013;**19**:176–86.

World Health Organization. *Global Report on Diabetes*. World Health Organization 2016.

Forootan M, Bagheri N, Darvishi M. Chronic constipation: A review of literature. *Medicine* 2018;**97**:e10631.

Organization WH, Others. Depression and other common mental disorders: global health estimates. World Health Organization 2017. https://apps.who.int/iris/bitstream/handle/10665/254610/WHO-MSD-MER-2017.2-eng.pdf

Ward WH, Farma JM, editors. *Cutaneous Melanoma: Etiology and Therapy*. Brisbane (AU): : Codon Publications 2018.

Table 1 Anti-Parkinson's disease medication name and corresponding PBS code

Medication	PBS code
Levodopa + carbidopa	1242J, 1245M, 1255C, 8970D, 9743T, 9744W
Levodopa + benserazide	2225D, 2226E, 2227F, 2228G, 2229H, 2231K, 8218M, 8219N
Levodopa + carbidopa + entacapone	8797B, 8798C, 8799D, 9292C, 9344T, 9345W
Amantadine hydrochloride	3016R
Apomorphine hydrochloride hemihydrate	10950H, 10971K, 11083H, 11093W, 5609F, 5610G, 9607P, 9640J
Cabergoline	8393R, 8394T
Pramipexole dihydrochloride monohydrate	3418X, 3419Y, 3420B, 3421C, 3422D, 5143Q, 5145T, 9151P, 9152Q, 9153R, 9393J, 9394K
Rotigotine	1140H, 2384L, 2385M, 2410W

Table 2 APGS Invitations and respondents per state or territory of Australia

State or Territory	Letters sent by state (%)	Respondents by state (%)
New South Wales	31.1	30.2
Victoria	25.1	22.2
Queensland	21.1	24.3
South Australia	9.4	9.4
Western Australia	8.7	9.3
Tasmania	3.2	2.3
Australian Capital Territory	1.1	1.7
Northern Territory	0.3	0.6

Table 3 Sociodemographic characteristics of the study sample

	Total N (%)	Male N (%)	Female N (%)	P-values Male vs Female
Response method				0.87
On-line questionnaire	1424 (92.9)	924 (60.3)	499 (32.6)	
Paper questionnaire	108 (7.1)	71 (4.6)	37 (2.4)	
Age in years				
Mean (SD)	63.9 (5.9)	64.2 (5.9)	63.5 (6.0)	0.04
Range	40 – 88	40 - 86	42 - 88	
Sex				
Female	536 (35.0)			-
Male	995 (64.9)			-
MTF	1 (0.06)			-
Current height in cm				
Mean (SD)	170.7 (9.9)	175.4 (7.8)	162.0 (7.2)	<0.001
Range	124 - 200	150 - 200	124 - 178	
Current weight in kg				
Mean (SD)	80.8 (16.9)	85.5 (15.3)	72.3 (16.5)	<0.001
Range	40 - 164	45 - 164	40 - 140	

Current BMI (kg/m^2)				
Mean (SD)	27.6 (5.3)	27.7 (4.8)	27.5 (6.0)	0.29
Range	16.6 – 55.1	17.5 – 50.7	16.6 – 55.1	
Have children	1338 (87.3)	870 (56.8)	467 (30.5)	0.86
Marital status				<0.001
Single / never married	66 (4.3)	45 (2.9)	21 (1.4)	
Married / de factor relationship	1245 (81.3)	848 (55.4)	396 (25.8)	
Separated / divorced	145 (9.5)	70 (4.5)	75 (4.9)	
Widowed	48 (3.1)	17 (1.1)	31 (2.0)	
In a relationship but not leaving together	26 (1.7)	13 (0.8)	13 (0.8)	
Other	2 (0.1)	2 (0.1)	0 (0.0)	
Ethnicity				0.73
European	1397 (91.6)	910 (59.4)	486 (31.7)	
East Asian	40 (2.6)	23 (1.5)	17 (1.1)	
Indigenous Australian / Torres Strait Islander	12 (0.8)	8 (0.5)	4 (0.3)	
African	2 (0.1)	2 (0.1)	0 (0.0)	
South Asian	26 (1.7)	14 (0.9)	12 (0.8)	
Pacific Islander	9 (0.6)	6 (0.4)	3 (0.2)	
Hispanics	5 (0.3)	4 (0.3)	1 (0.06)	
Mixed	21 (1.4)	16 (1.0)	5 (0.5)	
Other	13 (0.8)	8 (0.5)	5 (0.5)	
Information not provided	7 (0.4)	4 (0.3)	3 (0.2)	
Education				0.02
Junior high school or less	348 (23.2)	218 (14.2)	130 (8.5)	
Senior high school	276 (18.1)	184 (12.1)	92 (6.0)	
Certificate III/IV	196 (12.8)	124 (8.0)	72 (4.7)	
Diploma	190 (12.4)	128 (8.3)	62 (4.0)	
Bachelor's degree	235 (15.4)	163 (10.6)	71 (4.6)	
Graduate diploma or certificate	134 (8.8)	77 (5.0)	57 (3.7)	
Postgraduate	148 (9.7)	97 (6.3)	51 (3.3)	
Information not provided	5 (0.3)	5 (0.3)	0 (0.0)	
Hand preference				0.78
Left hand	156 (10.2)	103 (6.8)	53 (3.4)	
Right hand	1323 (86.5)	856 (55.8)	467 (30.5)	
Both hands	50 (3.3)	34 (2.2)	15 (0.9)	
Not sure	1 (0.06)	1 (0.06)	0 (0.0)	
Chronotype				<0.001
Morning-type person	800 (52.3)	518 (33.8)	282 (18.4)	
Evening-type person	376 (24.6)	212 (13.8)	163 (10.6)	
Neither	355 (23.2)	264 (17.2)	91 (5.9)	
Consider themselves spiritual or religious	599 (39.2)	347 (22.6)	251 (16.4)	<0.001
Diet type				0.004
Vegan	8 (0.5)	7 (0.4)	1 (0.06)	
Vegetarian	42 (2.7)	18 (1.1)	22 (1.4)	
Speak more than one language	228 (14.9)	147 (9.5)	81 (5.2)	0.88
Age at onset of the first PD symptom				
Mean (SD)	56.1 (8.3)	56.4 (8.2)	55.6 (8.3)	0.10

Range	27 - 86	30 – 77	27 - 86	
Healthcare provider who made PD diagnosis				0.001
Neurologist	1260 (82.5)	790 (51.5)	469 (30.6)	
Geriatrician	16 (1.0)	12 (0.7)	4 (0.3)	
General practitioner	237 (15.5)	178 (11.6)	59 (3.8)	
Other	15 (0.9)	12 (0.7)	3 (0.2)	
Information not provided	4 (0.3)	3 (0.2)	1 (0.06)	
Had brain surgery for PD	177 (11.7)	119 (7.7)	58 (3.7)	0.53
Take medication for PD	1424 (93.9)	930 (60.7)	493 (32.1)	0.43
Experienced falls related to PD	631 (41.5)	383 (25.0)	247 (16.1)	0.004
PD signs started on one side of the body	1296 (84.5)	833 (54.3)	463 (30.2)	0.12
Side of the body where PD symptoms started				0.69
Left	625 (40.8)	405 (26.4)	220 (14.3)	
Right	671 (43.8)	427 (27.8)	243 (15.8)	

Note: MTF – male-to-female transgender person; SD – standard deviation; BMI – body mass index; *P*-values were calculated using a t-test for comparing mean values or a Chi-square test for comparing frequencies; Sample N=1532.

Table 4 Occupational and environmental risk factors

Risk factor	Total N (%)	Male N (%)	Female N %
Exposure to pesticides or herbicides	517 (33.6)	390 (25.4)	127 (8.2)
Age of exposure to pesticides or herbicides			
< 18 years of age	185 (12.1)	128 (8.3)	57 (3.7)
18 – 36	239 (15.6)	190 (12.4)	49 (3.1)
37 – 55	84 (5.5)	64 (4.2)	20 (1.3)
≥ 56	9 (0.6)	8 (0.5)	1 (0.01)
Head injury	244 (16.1)	186 (12.1)	58 (3.7)
Age of head injury			
Mean (SD)	26.3 (17.6)	26.2 (16.5)	26.9 (20.8)
Range	2 - 82	3 - 67	2 - 82
Head injury with loss of consciousness	165 (10.7)	128 (8.4)	37 (2.4)
Well water drinking	82 (5.3)	55 (3.5%)	27 (1.7)
Number of years drinking well water			
Mean (SD)	18.7 (12.4)	18.3 (12.9)	19.4 (11.6)
Range	2 - 50	2 - 50	4 - 40
Worked in high-risk occupations	504 (32.9)	425 (27.7)	79 (5.1)
Number of high-risk occupations worked throughout of life			
One	293 (19.1)	226 (14.7)	67 (4.3)
Two	115 (7.5)	107 (6.9)	8 (0.5)
Three	52 (3.4)	48 (3.1)	4 (0.2)

	Total N (%)	Male N (%)	Female N (%)
Four	29 (1.8)	29 (1.8)	0 (0.0)
Five or more	15 (1.0)	15 (1.0)	0 (0.0)
Worked or lived on a farm for over 1 year	360 (23.5)	236 (15.4)	124 (8.1)
Had metal poisoning	22 (1.4)	17 (1.1)	5 (0.3)
Had brain infection (encephalitis)	31 (2.0)	20 (1.3)	11 (0.7)

Note: High-risk occupations included welding, metal melting, galvanization, milling, petrochemistry, agriculture, wood processing, textile, or industrial painting; SD – standard deviation; N = 1532

Table 5 Lifestyle variables

	Total N (%)	Male N (%)	Female N (%)
Currently drink alcohol	1090 (71.1)	751 (49.0)	339 (22.1)
Mean number of drinks per week (SD)	7.6 (7.7)	8.3 (8.4)	5.7 (5.3)
Range	1 - 50	1 - 50	1 - 30
Consumed alcohol heavily in the past	313 (20.4)	252 (16.4)	61 (3.9)
Current cigarette/cigar/pipe smokers	50 (3.3)	28 (1.8)	22 (1.4)
Mean number of cigarettes/cigar/pipes smoked per day (SD)	13.3 (8.3)	14.3 (8.9)	12.0 (7.6)
Range	2 - 40	2 - 40	2 - 30
Former smokers	554 (36.2)	368 (24.0)	186 (12.1)
Mean number of cigarettes smoked per day (SD)	15.3 (11.2)	18.1 (4.6)	17.9 (3.5)
Range	1 - 60	10 – 49	10 - 31
Exposure to second-hand smoke at home	858 (56.0)	510 (33.3)	348 (22.7)
Mean number of hours per day (SD)	4.5 (3.7)	4.3 (3.6)	4.9 (3.9)
Range	0.5 - 20	0.5 - 18	0.5 – 20
Exposure to second-hand smoke at work	701 (45.7)	521 (34.0)	179 (11.7)
Mean number of hours per day (SD)	4.9 (3.1)	4.9 (3.2)	4.9 (2.9)
Range	0.5 - 12	0.5 - 12	0.5 – 12
Exposure to second-hand smoke in other areas	586 (38.2)	405 (26.4)	181 (11.8)
Mean number of hours per day (SD)	2.0 (1.7)	2.1 (1.8)	1.9 (1.6)
Range	0.5 - 14	0.5 - 14	0.5 – 12
Drink caffeinated tea	1043 (68.1)	668 (43.6)	375 (24.4)
Mean number of cups per day (SD)	2 (1.4)	2 (1.3)	2.3 (1.6)
Range	0.5 - 12	0.5 - 8	0.5 - 12
Drink caffeinated coffee	1120 (73.1)	761 (49.7)	359 (2.2)
Mean number of cups per day (SD)	2 (1.2)	2.1 (1.3)	1.8 (1.2)
Range	0.5 - 10	0.5 - 10	0.5 - 10
Drink other caffeinated beverages	371 (24.2)	271 (17.7)	100 (6.5)
Average volume in ml per day (SD)	330.8 (336.6)	335.8 (340.2)	317.3 (330.7)
Range	10 - 2000	10 - 2000	10 - 2000

Note: SD – standard deviation; Sample N=1532

Table 6 Self-reported history of comorbidities in the study sample

Condition / Disorder	N	%
Back problems	722	47.1
Constipation	533	36.1
Depression	514	33.6
High blood pressure	506	33.0
High cholesterol	430	28.1
Frequent acid reflux	377	24.6
Cold sores	336	21.9
Migraine	264	17.2
Anxiety	262	17.1
Melanoma	243	15.9
Sleep apnoea	241	15.7
Insomnia	202	13.2
Cancer	201	13.1
Asthma / COPD	194	12.7
Pneumonia	192	12.5
REM Sleep behaviour disorder	177	11.6
Diabetes	156	10.2
Osteoporosis	126	8.2
Eczema	107	7.0

Note: COPD – chronic obstructive pulmonary disease; Sample N=1532

Figure 1. **Schematic of the APGS and questionnaire content**

Note: APGS, Australian Parkinson's Disease Study; PBS, Pharmaceutical Benefits Scheme medication prescription database; PD, Parkinson's disease; COPD, chronic obstructive pulmonary disease.

Figure 2 Frequency of self-reported first symptom of PD in participants

Note: 'Other' symptoms included numbness, falls, olfactory dysfunction, small handwriting, loss of taste, slurred speech.

Figure 3 Relationship between the age of traumatic head injury and age at onset of the first symptom of Parkinson's disease in participants